

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nchsed@cdc.gov](mailto:nchsed@cdc.gov), or call 301-458-4688







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental discharge of firearms (W32-W34)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental explosion (W39-W40)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Accidental explosion (W39-W40)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Accidental explosion (W39-W40)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Accidental explosion (W39-W40)

New Hampshire

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Accidental explosion (W39-W40)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental explosion (W39-W40)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Accidental explosion (W39-W40)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Accidental explosion (W39-W40)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental explosion (W39-W40)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental explosion (W39-W40)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental explosion (W39-W40)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental explosion (W39-W40)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental explosion (W39-W40)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Arkansas												
All races, both sexes.....	42	-	9	5	1	9	2	5	-	3	2	2
Male.....	33	-	6	4	-	6	2	5	-	2	2	2
Female.....	9	-	3	1	1	3	-	-	-	1	-	-
White, both sexes.....	32	-	7	4	-	7	1	4	-	3	1	2
Male.....	26	-	5	3	-	5	1	4	-	2	1	2
Female.....	6	-	2	1	-	2	-	-	-	1	-	-
Black, both sexes.....	8	-	2	-	1	1	1	1	-	-	1	-
Male.....	6	-	1	-	-	1	1	1	-	-	1	-
Female.....	2	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	1	-	1	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
California												
All races, both sexes.....	372	10	54	8	9	20	29	22	15	16	26	32
Male.....	283	7	34	6	8	14	27	20	11	14	23	21
Female.....	89	3	20	2	1	6	2	2	4	2	3	11
White, both sexes.....	294	9	48	5	6	13	22	13	10	11	20	29
Male.....	224	6	31	4	5	9	21	11	8	11	19	19
Female.....	70	3	17	1	1	4	1	2	2	-	1	10
Black, both sexes.....	28	1	1	3	2	2	2	1	2	2	3	2
Male.....	21	1	1	2	2	1	2	1	1	1	2	2
Female.....	7	-	-	1	-	1	-	-	1	1	1	-
American Indian 1/, both sexes.....	4	-	1	-	-	1	-	-	1	1	-	-
Male.....	4	-	1	-	-	1	-	-	1	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	46	-	4	-	1	4	5	8	2	2	3	1
Male.....	34	-	1	-	1	3	4	8	1	1	2	-
Female.....	12	-	3	-	-	1	1	-	1	1	1	1





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Florida												
All races, both sexes.....	390	6	72	10	8	27	39	10	9	18	20	29
Male.....	298	4	51	6	5	22	33	10	8	16	18	23
Female.....	92	2	21	4	3	5	6	-	1	2	2	6
White, both sexes.....	305	4	53	4	3	14	32	5	8	15	17	23
Male.....	228	3	34	1	2	10	27	5	7	13	16	18
Female.....	77	1	19	3	1	4	5	-	1	2	1	5
Black, both sexes.....	70	2	16	6	5	12	7	5	1	3	1	4
Male.....	58	1	14	5	3	11	6	5	1	3	-	4
Female.....	12	1	2	1	2	1	1	-	-	-	1	-
American Indian 1/, both sexes.....	3	-	1	-	-	-	-	-	-	-	-	1
Male.....	3	-	1	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	12	-	2	-	-	1	-	-	-	-	2	1
Male.....	9	-	2	-	-	1	-	-	-	-	2	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes.....	125	2	25	4	1	12	15	9	8	6	11	8
Male.....	90	-	15	2	1	11	14	8	6	4	4	6
Female.....	35	2	10	2	-	1	1	1	2	2	7	2
White, both sexes.....	95	-	21	3	-	6	13	8	2	5	10	6
Male.....	69	-	12	2	-	6	12	7	1	4	4	4
Female.....	26	-	9	1	-	-	1	1	1	1	6	2
Black, both sexes.....	26	2	4	1	-	5	2	1	6	-	1	2
Male.....	19	-	3	-	-	4	2	1	5	-	-	2
Female.....	7	2	1	1	-	1	-	-	1	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	1	1	-	-	-	1	-	-
Male.....	2	-	-	-	1	1	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	-	-



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental drowning and submersion (W65-W74)

Nebraska

All races, both sexes.....	12	-	-	-	1	2	3	1	-	-	2	-
Male.....	12	-	-	-	1	2	3	1	-	-	2	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	12	-	-	-	1	2	3	1	-	-	2	-
Male.....	12	-	-	-	1	2	3	1	-	-	2	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	38	1	8	1	1	4	5	1	1	2	3	1
Male.....	29	-	7	-	1	4	3	1	1	2	2	1
Female.....	9	1	1	1	-	-	2	-	-	-	1	-
White, both sexes.....	30	1	8	-	-	3	3	1	-	-	3	1
Male.....	23	-	7	-	-	3	2	1	-	-	2	1
Female.....	7	1	1	-	-	-	1	-	-	-	1	-
Black, both sexes.....	2	-	-	1	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	1	-	2	-	-	2	-	-
Male.....	4	-	-	-	1	-	1	-	-	2	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
New Mexico												
All races, both sexes.....	22	-	2	-	-	3	2	-	-	1	1	8
Male.....	17	-	-	-	-	3	2	-	-	1	1	7
Female.....	5	-	2	-	-	-	-	-	-	-	-	1
White, both sexes.....	18	-	1	-	-	2	2	-	-	1	-	7
Male.....	14	-	-	-	-	2	2	-	-	1	-	6
Female.....	4	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	1	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	1	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	116	3	12	4	5	14	6	5	5	12	9	8
Male.....	85	1	9	3	3	10	4	4	5	9	7	6
Female.....	31	2	3	1	2	4	2	1	-	3	2	2
White, both sexes.....	91	2	10	3	1	10	5	3	2	10	8	7
Male.....	67	-	7	3	1	7	4	3	2	7	6	5
Female.....	24	2	3	-	-	3	1	-	-	3	2	2
Black, both sexes.....	20	1	-	1	3	4	1	1	3	1	1	1
Male.....	15	1	-	-	2	3	-	1	3	1	1	1
Female.....	5	-	-	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	2	-	1	-	-	1	-	1	-	-
Male.....	3	-	2	-	-	-	-	-	-	1	-	-
Female.....	2	-	-	-	1	-	-	1	-	-	-	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Arkansas

All races, both sexes.....	-	1	1	-	1	-	1	-	-	-	-	-
Male.....	-	1	1	-	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	1	-	-	1	-	1	-	-	-	-	-
Male.....	-	1	-	-	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	24	21	22	11	8	18	9	11	3	-	-	4
Male.....	16	15	18	7	8	15	7	7	1	-	-	4
Female.....	8	6	4	4	-	3	2	4	2	-	-	-
White, both sexes.....	19	16	21	9	5	14	8	9	3	-	-	4
Male.....	12	10	17	6	5	11	7	7	1	-	-	4
Female.....	7	6	4	3	-	3	1	2	2	-	-	-
Black, both sexes.....	1	3	-	1	-	-	-	2	-	-	-	-
Male.....	1	3	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	2	1	1	3	4	1	-	-	-	-	-
Male.....	3	2	1	-	3	4	-	-	-	-	-	-
Female.....	1	-	-	1	-	-	1	-	-	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

New Hampshire

All races, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	4	1	3	1	5	4	3	2	-	-	-	-
Male.....	3	1	1	1	3	1	1	2	-	-	-	-
Female.....	1	-	2	-	2	3	2	-	-	-	-	-
White, both sexes.....	3	1	3	-	3	2	3	2	-	-	-	-
Male.....	3	1	1	-	1	1	1	2	-	-	-	-
Female.....	-	-	2	-	2	1	2	-	-	-	-	-
Black, both sexes.....	-	-	-	1	2	1	-	-	-	-	-	-
Male.....	-	-	-	1	2	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Alaska

All races, both sexes.....	6	1	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	5	1	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	82	1	1	-	-	-	-	-	2	5	3	2
Male.....	41	-	-	-	-	-	-	-	1	4	-	2
Female.....	41	1	1	-	-	-	-	-	1	1	3	-
White, both sexes.....	70	1	1	-	-	-	-	-	2	-	2	1
Male.....	32	-	-	-	-	-	-	-	1	-	-	1
Female.....	38	1	1	-	-	-	-	-	1	-	2	-
Black, both sexes.....	3	-	-	-	-	-	-	-	-	1	-	-
Male.....	2	-	-	-	-	-	-	-	-	1	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	8	-	-	-	-	-	-	-	-	3	1	1
Male.....	7	-	-	-	-	-	-	-	-	3	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-





























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Arkansas

All races, both sexes.....	7	8	1	4	7	13	9	10	4	4	-	-
Male.....	5	5	-	2	5	10	7	5	2	1	-	-
Female.....	2	3	1	2	2	3	2	5	2	3	-	-
White, both sexes.....	4	7	1	4	7	12	9	9	4	4	-	-
Male.....	3	4	-	2	5	9	7	4	2	1	-	-
Female.....	1	3	1	2	2	3	2	5	2	3	-	-
Black, both sexes.....	3	1	-	-	-	1	-	1	-	-	-	-
Male.....	2	1	-	-	-	1	-	1	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	17	16	12	17	15	20	29	42	12	7	1	-
Male.....	7	9	8	13	11	9	11	24	6	2	-	-
Female.....	10	7	4	4	4	11	18	18	6	5	1	-
White, both sexes.....	17	14	10	11	13	16	25	39	11	5	-	-
Male.....	7	9	7	8	10	7	8	23	5	1	-	-
Female.....	10	5	3	3	3	9	17	16	6	4	-	-
Black, both sexes.....	-	1	1	5	1	-	1	-	-	-	-	-
Male.....	-	-	-	4	-	-	1	-	-	-	-	-
Female.....	-	1	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	1	4	3	3	1	2	1	-
Male.....	-	-	1	1	1	2	2	1	1	1	-	-
Female.....	-	-	-	-	-	2	1	2	-	1	1	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Maine

All races, both sexes.....	1	-	2	1	1	3	7	3	1	3	1	-
Male.....	1	-	2	1	1	2	4	1	-	-	-	-
Female.....	-	-	-	-	-	1	3	2	1	3	1	-
White, both sexes.....	1	-	2	1	1	3	7	3	1	3	1	-
Male.....	1	-	2	1	1	2	4	1	-	-	-	-
Female.....	-	-	-	-	-	1	3	2	1	3	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	4	2	-	3	1	5	4	7	3	-	-	-
Male.....	4	1	-	2	-	-	2	4	1	-	-	-
Female.....	-	1	-	1	1	5	2	3	2	-	-	-
White, both sexes.....	2	1	-	1	1	4	3	5	2	-	-	-
Male.....	2	1	-	1	-	-	2	3	1	-	-	-
Female.....	-	-	-	-	1	4	1	2	1	-	-	-
Black, both sexes.....	2	1	-	2	-	1	1	2	-	-	-	-
Male.....	2	-	-	1	-	-	-	1	-	-	-	-
Female.....	-	1	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

New Mexico

All races, both sexes.....	1	2	-	2	1	3	3	5	3	-	1	-
Male.....	-	1	-	2	1	2	1	4	3	-	1	-
Female.....	1	1	-	-	-	1	2	1	-	-	-	-
White, both sexes.....	1	2	-	2	-	3	2	4	3	-	-	-
Male.....	-	1	-	2	-	2	-	4	3	-	-	-
Female.....	1	1	-	-	-	1	2	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	1	1	-	-	1	-
Male.....	-	-	-	-	1	-	1	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	12	9	9	11	17	22	31	36	24	7	3	-
Male.....	10	5	7	6	9	12	17	17	15	2	1	-
Female.....	2	4	2	5	8	10	14	19	9	5	2	-
White, both sexes.....	10	8	9	10	16	20	31	33	24	7	3	-
Male.....	9	5	7	6	8	10	17	16	15	2	1	-
Female.....	1	3	2	4	8	10	14	17	9	5	2	-
Black, both sexes.....	2	1	-	1	1	2	-	2	-	-	-	-
Male.....	1	-	-	-	1	2	-	1	-	-	-	-
Female.....	1	1	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Oregon

All races, both sexes.....	1	4	3	3	3	3	6	8	2	1	-	-
Male.....	-	4	1	2	1	2	2	4	2	-	-	-
Female.....	1	-	2	1	2	1	4	4	-	1	-	-
White, both sexes.....	1	4	3	3	3	3	6	8	2	1	-	-
Male.....	-	4	1	2	1	2	2	4	2	-	-	-
Female.....	1	-	2	1	2	1	4	4	-	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	5	7	12	11	21	39	63	59	40	7	2	-
Male.....	4	2	9	6	16	20	38	34	14	3	-	-
Female.....	1	5	3	5	5	19	25	25	26	4	2	-
White, both sexes.....	3	5	10	9	17	32	60	47	38	5	2	-
Male.....	3	1	8	5	13	16	38	29	14	3	-	-
Female.....	-	4	2	4	4	16	22	18	24	2	2	-
Black, both sexes.....	2	2	2	2	3	6	2	11	2	2	-	-
Male.....	1	1	1	1	2	4	-	5	-	-	-	-
Female.....	1	1	1	1	1	2	2	6	2	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	1	1	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-





























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to electric current (W85-W87)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to electric current (W85-W87)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to electric current (W85-W87)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental exposure to electric current (W85-W87)

Vermont

Table with 13 columns (Age groups) and 14 rows (Race categories) for Vermont. All values are dashes (-).

Virginia

Table with 13 columns (Age groups) and 14 rows (Race categories) for Virginia. Values are dashes (-) or 1.











## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Arkansas												
All races, both sexes.....	66	-	6	-	-	4	1	4	2	2	4	6
Male.....	37	-	6	-	-	3	1	2	1	2	3	5
Female.....	29	-	-	-	-	1	-	2	1	-	1	1
White, both sexes.....	49	-	3	-	-	4	1	4	2	2	3	5
Male.....	28	-	3	-	-	3	1	2	1	2	2	4
Female.....	21	-	-	-	-	1	-	2	1	-	1	1
Black, both sexes.....	17	-	3	-	-	-	-	-	-	-	1	1
Male.....	9	-	3	-	-	-	-	-	-	-	1	1
Female.....	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	234	3	5	9	2	9	11	11	11	11	8	13
Male.....	144	2	3	3	1	6	9	6	7	10	7	11
Female.....	90	1	2	6	1	3	2	5	4	1	1	2
White, both sexes.....	189	2	2	7	2	7	9	9	11	9	8	10
Male.....	118	1	2	2	1	5	7	4	7	8	7	8
Female.....	71	1	-	5	1	2	2	5	4	1	1	2
Black, both sexes.....	28	1	1	-	-	1	1	1	-	1	-	1
Male.....	15	1	-	-	-	1	1	1	-	1	-	1
Female.....	13	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	14	-	2	2	-	1	1	1	-	1	-	2
Male.....	8	-	1	1	-	-	1	1	-	1	-	2
Female.....	6	-	1	1	-	1	-	-	-	-	-	-































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
New Mexico												
All races, both sexes.....	19	1	1	2	-	-	-	-	-	1	-	-
Male.....	12	1	1	1	-	-	-	-	-	-	-	-
Female.....	7	-	-	1	-	-	-	-	-	1	-	-
White, both sexes.....	18	1	1	2	-	-	-	-	-	-	-	-
Male.....	12	1	1	1	-	-	-	-	-	-	-	-
Female.....	6	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	179	6	12	8	6	2	9	7	6	5	11	12
Male.....	101	4	5	7	5	2	5	1	3	2	7	7
Female.....	78	2	7	1	1	-	4	6	3	3	4	5
White, both sexes.....	123	4	4	1	4	1	8	4	4	3	6	8
Male.....	68	3	-	1	3	1	4	1	2	1	4	5
Female.....	55	1	4	-	1	-	4	3	2	2	2	3
Black, both sexes.....	48	2	6	7	2	1	1	2	2	2	4	4
Male.....	26	1	3	6	2	1	1	-	1	1	2	2
Female.....	22	1	3	1	-	-	-	2	1	1	2	2
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	-	2	-	-	-	-	1	-	-	1	-
Male.....	6	-	2	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Oregon												
All races, both sexes.....	33	1	2	2	2	-	-	-	-	-	-	4
Male.....	16	-	2	1	-	-	-	-	-	-	-	2
Female.....	17	1	-	1	2	-	-	-	-	-	-	2
White, both sexes.....	31	1	2	2	2	-	-	-	-	-	-	4
Male.....	16	-	2	1	-	-	-	-	-	-	-	2
Female.....	15	1	-	1	2	-	-	-	-	-	-	2
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	181	3	19	15	4	7	5	9	5	8	6	9
Male.....	110	2	16	8	-	3	3	6	3	3	5	6
Female.....	71	1	3	7	4	4	2	3	2	5	1	3
White, both sexes.....	142	2	13	8	4	7	4	7	4	6	5	8
Male.....	85	1	12	3	-	3	3	5	2	3	4	6
Female.....	57	1	1	5	4	4	1	2	2	3	1	2
Black, both sexes.....	38	1	6	7	-	-	1	2	1	1	1	1
Male.....	25	1	4	5	-	-	-	1	1	-	1	-
Female.....	13	-	2	2	-	-	1	1	-	1	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Arkansas

All races, both sexes.....	5	6	1	5	4	1	5	5	3	1	1	-
Male.....	1	2	1	3	1	-	2	2	1	-	1	-
Female.....	4	4	-	2	3	1	3	3	2	1	-	-
White, both sexes.....	3	4	1	4	4	-	3	4	2	-	-	-
Male.....	1	1	1	2	1	-	1	2	1	-	-	-
Female.....	2	3	-	2	3	-	2	2	1	-	-	-
Black, both sexes.....	2	2	-	1	-	1	2	1	1	1	1	-
Male.....	-	1	-	1	-	-	1	-	-	-	1	-
Female.....	2	1	-	-	-	1	1	1	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	19	18	19	18	9	16	20	14	6	-	1	1
Male.....	12	8	13	12	5	10	9	6	3	-	-	1
Female.....	7	10	6	6	4	6	11	8	3	-	1	-
White, both sexes.....	18	14	14	14	7	12	16	12	5	-	-	1
Male.....	11	8	10	8	4	8	8	5	3	-	-	1
Female.....	7	6	4	6	3	4	8	7	2	-	-	-
Black, both sexes.....	1	4	3	3	2	3	3	1	1	-	-	-
Male.....	1	-	1	3	1	1	1	1	-	-	-	-
Female.....	-	4	2	-	1	2	2	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	1	1	-	1	-	-	-	-	-	-
Male.....	-	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	1	1	-	-	1	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	1	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Maine

All races, both sexes.....	-	-	-	2	-	1	-	-	-	-	-	-
Male.....	-	-	-	2	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	2	-	1	-	-	-	-	-	-
Male.....	-	-	-	2	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	5	5	-	6	4	5	3	7	-	1	-	-
Male.....	3	4	-	5	2	1	-	4	-	1	-	-
Female.....	2	1	-	1	2	4	3	3	-	-	-	-
White, both sexes.....	1	4	-	3	3	2	2	5	-	-	-	-
Male.....	-	3	-	2	1	-	-	3	-	-	-	-
Female.....	1	1	-	1	2	2	2	2	-	-	-	-
Black, both sexes.....	4	1	-	3	1	2	-	2	-	1	-	-
Male.....	3	1	-	3	1	1	-	1	-	1	-	-
Female.....	1	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Nebraska

All races, both sexes.....	3	-	-	-	-	-	-	-	-	-	1	-
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes.....	3	-	-	-	-	-	-	-	-	-	1	-
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	1	2	4	4	3	3	2	1	-	-	-	-
Male.....	1	1	3	4	2	-	-	-	-	-	-	-
Female.....	-	1	1	-	1	3	2	1	-	-	-	-
White, both sexes.....	1	2	4	3	2	2	2	-	-	-	-	-
Male.....	1	1	3	3	1	-	-	-	-	-	-	-
Female.....	-	1	1	-	1	2	2	-	-	-	-	-
Black, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental contact with heat and hot substances (X10-X19)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental contact with heat and hot substances (X10-X19)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental contact with heat and hot substances (X10-X19)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental contact with heat and hot substances (X10-X19)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental contact with heat and hot substances (X10-X19)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Iowa

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Maine

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maine. All values are dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maryland. Values are dashes (-) or 1.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Nebraska

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Ohio

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Oregon

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental contact with heat and hot substances (X10-X19)

Vermont

Table with 13 columns (Age groups) and 13 rows (Race categories) for Vermont. All cells contain dashes (-).

Virginia

Table with 13 columns (Age groups) and 13 rows (Race categories) for Virginia. All cells contain dashes (-).



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Victim of lightning (X33)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Victim of lightning (X33)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Victim of lightning (X33)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Victim of lightning (X33)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Victim of lightning (X33)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Victim of lightning (X33)

Washington

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Victim of lightning (X33)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for Maine (All races, White, Black, American Indian, Asian or Pacific Islander).

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for Maryland (All races, White, Black, American Indian, Asian or Pacific Islander).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Delaware

Table with 13 columns (Age groups) and 13 rows (Race categories) for Delaware. All cells contain dashes (-).

Dist. of Columbia

Table with 13 columns (Age groups) and 13 rows (Race categories) for District of Columbia. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Florida

Table with 13 columns (Age groups) and 13 rows (Race categories) for Florida. All cells contain dashes (-).

Georgia

Table with 13 columns (Age groups) and 13 rows (Race categories) for Georgia. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race categories) for Idaho. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Illinois

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Massachusetts

Table with 13 columns (Age groups) and 13 rows (Race categories) for Massachusetts. All cells contain dashes (-).

Michigan

Table with 13 columns (Age groups) and 13 rows (Race categories) for Michigan. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Minnesota

Table with 13 columns (Age groups) and 13 rows (Race categories) for Minnesota. All cells contain dashes (-).

Mississippi

Table with 13 columns (Age groups) and 13 rows (Race categories) for Mississippi. All cells contain dashes (-).







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

North Carolina

Table with 13 columns for age groups and one for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All cells contain dashes.

North Dakota

Table with 13 columns for age groups and one for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All cells contain dashes.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cataclysmic storm and flood (X37-X38)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Delaware

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Hawaii

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Maine

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Vermont

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cataclysmic storm and flood (X37-X38)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Idaho. All cells contain dashes (-).







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cataclysmic storm and flood (X37-X38)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

South Dakota

Table with 13 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for South Dakota.

Tennessee

Table with 13 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for Tennessee.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cataclysmic storm and flood (X37-X38)

Washington

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

West Virginia

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Alaska												
All races, both sexes.....	87	-	-	-	1	-	10	11	13	13	4	19
Male.....	54	-	-	-	-	-	8	8	6	10	3	11
Female.....	33	-	-	-	1	-	2	3	7	3	1	8
White, both sexes.....	59	-	-	-	-	-	5	10	7	10	2	13
Male.....	42	-	-	-	-	-	4	7	5	8	2	9
Female.....	17	-	-	-	-	-	1	3	2	2	-	4
Black, both sexes.....	2	-	-	-	1	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	1	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	26	-	-	-	-	-	5	1	6	2	2	6
Male.....	11	-	-	-	-	-	4	1	1	1	1	2
Female.....	15	-	-	-	-	-	1	-	5	1	1	4
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	757	-	2	1	2	36	50	58	75	85	107	127
Male.....	515	-	1	-	-	24	40	47	47	56	70	91
Female.....	242	-	1	1	2	12	10	11	28	29	37	36
White, both sexes.....	679	-	1	1	1	35	44	56	68	76	88	116
Male.....	458	-	1	-	-	24	36	45	43	50	55	83
Female.....	221	-	-	1	1	11	8	11	25	26	33	33
Black, both sexes.....	36	-	1	-	-	-	2	-	4	3	9	5
Male.....	26	-	-	-	-	-	1	-	2	2	6	4
Female.....	10	-	1	-	-	-	1	-	2	1	3	1
American Indian 1/, both sexes.....	38	-	-	-	-	1	4	2	3	4	9	6
Male.....	29	-	-	-	-	-	3	2	2	3	8	4
Female.....	9	-	-	-	-	1	1	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	1	-	-	-	-	2	1	-
Male.....	2	-	-	-	-	-	-	-	-	1	1	-
Female.....	2	-	-	-	1	-	-	-	-	1	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Arkansas												
All races, both sexes.....	162	-	-	-	1	6	15	14	9	26	19	29
Male.....	104	-	-	-	1	5	13	12	6	15	13	16
Female.....	58	-	-	-	-	1	2	2	3	11	6	13
White, both sexes.....	147	-	-	-	1	5	12	12	9	25	18	27
Male.....	92	-	-	-	1	4	10	10	6	14	13	15
Female.....	55	-	-	-	-	1	2	2	3	11	5	12
Black, both sexes.....	15	-	-	-	-	1	3	2	-	1	1	2
Male.....	12	-	-	-	-	1	3	2	-	1	-	1
Female.....	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	3,372	2	2	-	6	72	176	207	207	338	458	615
Male.....	2,213	1	-	-	4	51	131	154	143	231	296	396
Female.....	1,159	1	2	-	2	21	45	53	64	107	162	219
White, both sexes.....	2,898	1	2	-	4	66	157	186	182	291	402	536
Male.....	1,918	-	-	-	2	48	122	144	128	201	262	340
Female.....	980	1	2	-	2	18	35	42	54	90	140	196
Black, both sexes.....	351	-	-	-	1	3	10	11	17	35	33	62
Male.....	220	-	-	-	1	1	4	6	9	21	21	46
Female.....	131	-	-	-	-	2	6	5	8	14	12	16
American Indian 1/, both sexes.....	48	-	-	-	-	2	1	3	3	2	11	11
Male.....	27	-	-	-	-	1	1	1	3	1	6	6
Female.....	21	-	-	-	-	1	-	2	-	1	5	5
Asian or Pacific Islander 2/, both sexes.....	75	1	-	-	1	1	8	7	5	10	12	6
Male.....	48	1	-	-	1	1	4	3	3	8	7	4
Female.....	27	-	-	-	-	-	4	4	2	2	5	2



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Nebraska

All races, both sexes.....	68	1	-	-	-	2	3	8	5	14	8	8
Male.....	41	1	-	-	-	2	3	7	3	7	3	5
Female.....	27	-	-	-	-	-	-	1	2	7	5	3
White, both sexes.....	61	1	-	-	-	2	2	7	4	14	7	7
Male.....	37	1	-	-	-	2	2	7	2	7	3	4
Female.....	24	-	-	-	-	-	-	-	2	7	4	3
Black, both sexes.....	2	-	-	-	-	-	1	-	-	-	1	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	4	-	-	-	-	-	-	1	-	-	-	1
Male.....	2	-	-	-	-	-	-	-	-	-	-	1
Female.....	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	417	-	-	-	3	6	41	40	42	40	65	58
Male.....	255	-	-	-	2	3	28	29	30	27	37	34
Female.....	162	-	-	-	1	3	13	11	12	13	28	24
White, both sexes.....	379	-	-	-	3	5	38	36	37	36	57	55
Male.....	232	-	-	-	2	2	26	28	27	24	33	33
Female.....	147	-	-	-	1	3	12	8	10	12	24	22
Black, both sexes.....	33	-	-	-	-	1	3	3	4	4	7	3
Male.....	20	-	-	-	-	1	2	1	2	3	3	1
Female.....	13	-	-	-	-	-	1	2	2	1	4	2
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	1	-
Male.....	2	-	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	1	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
New Hampshire												
All races, both sexes.....	157	-	-	-	-	6	20	18	12	19	25	20
Male.....	104	-	-	-	-	5	17	13	11	12	15	11
Female.....	53	-	-	-	-	1	3	5	1	7	10	9
White, both sexes.....	155	-	-	-	-	6	20	17	12	19	25	20
Male.....	102	-	-	-	-	5	17	12	11	12	15	11
Female.....	53	-	-	-	-	1	3	5	1	7	10	9
Black, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	-
Male.....	2	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	662	1	-	1	2	13	58	84	70	90	122	88
Male.....	481	1	-	1	-	8	46	70	53	68	95	57
Female.....	181	-	-	-	2	5	12	14	17	22	27	31
White, both sexes.....	541	-	-	-	1	12	55	78	59	71	98	75
Male.....	403	-	-	-	-	8	44	66	45	58	76	50
Female.....	138	-	-	-	1	4	11	12	14	13	22	25
Black, both sexes.....	115	1	-	1	-	-	2	6	10	18	24	13
Male.....	76	1	-	1	-	-	1	4	8	9	19	7
Female.....	39	-	-	-	-	-	1	2	2	9	5	6
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	-	-	1	1	1	-	1	1	-	-
Male.....	2	-	-	-	-	-	1	-	-	1	-	-
Female.....	4	-	-	-	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
New Mexico												
All races, both sexes.....	401	-	1	-	-	13	21	32	45	35	60	70
Male.....	288	-	1	-	-	10	16	28	28	31	40	49
Female.....	113	-	-	-	-	3	5	4	17	4	20	21
White, both sexes.....	365	-	1	-	-	8	20	27	41	30	55	64
Male.....	263	-	1	-	-	7	15	23	26	26	37	46
Female.....	102	-	-	-	-	1	5	4	15	4	18	18
Black, both sexes.....	8	-	-	-	-	1	-	1	-	-	2	2
Male.....	5	-	-	-	-	-	-	1	-	-	1	1
Female.....	3	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes.....	26	-	-	-	-	4	1	4	4	5	3	3
Male.....	19	-	-	-	-	3	1	4	2	5	2	1
Female.....	7	-	-	-	-	1	-	-	2	-	1	2
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	1,379	-	3	-	1	24	85	122	131	137	202	232
Male.....	968	-	1	-	-	20	68	102	97	92	145	147
Female.....	411	-	2	-	1	4	17	20	34	45	57	85
White, both sexes.....	1,103	-	-	-	1	20	79	113	115	111	171	173
Male.....	783	-	-	-	-	16	63	94	85	76	120	119
Female.....	320	-	-	-	1	4	16	19	30	35	51	54
Black, both sexes.....	258	-	3	-	-	4	3	6	12	24	30	56
Male.....	172	-	1	-	-	4	2	5	9	15	25	26
Female.....	86	-	2	-	-	-	1	1	3	9	5	30
American Indian 1/, both sexes.....	2	-	-	-	-	-	1	-	-	1	-	-
Male.....	2	-	-	-	-	-	1	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	16	-	-	-	-	-	2	3	4	1	1	3
Male.....	11	-	-	-	-	-	2	3	3	-	-	2
Female.....	5	-	-	-	-	-	-	-	1	1	1	1







## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Oregon												
All races, both sexes.....	373	1	-	1	2	8	21	36	33	38	43	55
Male.....	239	1	-	-	2	6	19	22	25	24	28	34
Female.....	134	-	-	1	-	2	2	14	8	14	15	21
White, both sexes.....	356	1	-	1	1	8	21	34	32	37	41	51
Male.....	226	1	-	-	1	6	19	22	24	23	28	30
Female.....	130	-	-	1	-	2	2	12	8	14	13	21
Black, both sexes.....	9	-	-	-	-	-	-	1	-	1	1	1
Male.....	7	-	-	-	-	-	-	-	-	1	-	1
Female.....	2	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes.....	6	-	-	-	1	-	-	1	1	-	1	1
Male.....	4	-	-	-	1	-	-	-	1	-	-	1
Female.....	2	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	2
Male.....	2	-	-	-	-	-	-	-	-	-	-	2
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	1,566	-	3	-	1	46	154	179	150	157	209	261
Male.....	1,090	-	2	-	1	33	120	133	112	107	155	159
Female.....	476	-	1	-	-	13	34	46	38	50	54	102
White, both sexes.....	1,358	-	3	-	1	43	144	175	137	141	175	224
Male.....	946	-	2	-	1	30	113	132	105	95	133	137
Female.....	412	-	1	-	-	13	31	43	32	46	42	87
Black, both sexes.....	202	-	-	-	-	3	9	4	12	16	32	37
Male.....	142	-	-	-	-	3	6	1	7	12	22	22
Female.....	60	-	-	-	-	-	3	3	5	4	10	15
American Indian 1/, both sexes.....	3	-	-	-	-	-	1	-	-	-	1	-
Male.....	2	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	1	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	1	-	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

South Dakota

All races, both sexes.....	24	-	-	-	1	-	2	4	4	2	3	5
Male.....	13	-	-	-	1	-	1	3	1	1	1	4
Female.....	11	-	-	-	-	-	1	1	3	1	2	1
White, both sexes.....	15	-	-	-	1	-	2	2	1	1	2	4
Male.....	7	-	-	-	1	-	1	1	-	-	1	3
Female.....	8	-	-	-	-	-	1	1	1	1	1	1
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	8	-	-	-	-	-	-	2	3	1	1	-
Male.....	5	-	-	-	-	-	-	2	1	1	-	-
Female.....	3	-	-	-	-	-	-	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	798	-	-	-	2	21	50	73	83	93	127	127
Male.....	482	-	-	-	2	14	35	45	58	59	78	76
Female.....	316	-	-	-	-	7	15	28	25	34	49	51
White, both sexes.....	683	-	-	-	2	20	48	68	74	83	112	103
Male.....	400	-	-	-	2	14	35	42	50	53	67	58
Female.....	283	-	-	-	-	6	13	26	24	30	45	45
Black, both sexes.....	111	-	-	-	-	1	1	4	8	10	15	24
Male.....	80	-	-	-	-	-	-	3	7	6	11	18
Female.....	31	-	-	-	-	1	1	1	1	4	4	6
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	1	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Vermont												
All races, both sexes.....	46	-	-	-	-	-	3	6	3	8	8	5
Male.....	28	-	-	-	-	-	1	5	2	3	4	5
Female.....	18	-	-	-	-	-	2	1	1	5	4	-
White, both sexes.....	45	-	-	-	-	-	3	6	3	8	8	5
Male.....	27	-	-	-	-	-	1	5	2	3	4	5
Female.....	18	-	-	-	-	-	2	1	1	5	4	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	562	-	-	1	2	12	60	70	60	75	85	90
Male.....	367	-	-	-	1	8	45	53	43	45	48	58
Female.....	195	-	-	1	1	4	15	17	17	30	37	32
White, both sexes.....	470	-	-	1	2	12	54	62	52	59	71	75
Male.....	308	-	-	-	1	8	40	46	39	34	42	47
Female.....	162	-	-	1	1	4	14	16	13	25	29	28
Black, both sexes.....	87	-	-	-	-	-	5	7	7	16	14	14
Male.....	55	-	-	-	-	-	4	6	4	11	6	10
Female.....	32	-	-	-	-	-	1	1	3	5	8	4
American Indian 1/, both sexes.....	2	-	-	-	-	-	1	1	-	-	-	-
Male.....	2	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	1	-	-	1
Male.....	2	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	1	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Arkansas

All races, both sexes.....	13	7	8	5	4	2	3	-	1	-	-	-
Male.....	7	4	4	4	2	-	1	-	1	-	-	-
Female.....	6	3	4	1	2	2	2	-	-	-	-	-
White, both sexes.....	11	6	6	5	4	2	3	-	1	-	-	-
Male.....	5	3	3	4	2	-	1	-	1	-	-	-
Female.....	6	3	3	1	2	2	2	-	-	-	-	-
Black, both sexes.....	2	1	2	-	-	-	-	-	-	-	-	-
Male.....	2	1	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	618	373	164	51	29	15	17	16	2	3	-	1
Male.....	408	236	99	27	14	5	6	8	-	2	-	1
Female.....	210	137	65	24	15	10	11	8	2	1	-	-
White, both sexes.....	514	302	140	45	22	13	14	15	2	3	-	1
Male.....	342	188	86	24	11	4	5	8	-	2	-	1
Female.....	172	114	54	21	11	9	9	7	2	1	-	-
Black, both sexes.....	88	56	21	6	5	2	1	-	-	-	-	-
Male.....	55	37	12	3	3	1	-	-	-	-	-	-
Female.....	33	19	9	3	2	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	7	6	2	-	-	-	-	-	-	-	-	-
Male.....	4	4	-	-	-	-	-	-	-	-	-	-
Female.....	3	2	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	9	1	-	2	-	2	1	-	-	-	-
Male.....	7	7	1	-	-	-	1	-	-	-	-	-
Female.....	2	2	-	-	2	-	1	1	-	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

New Hampshire

All races, both sexes.....	14	9	6	2	1	4	1	-	-	-	-	-
Male.....	7	5	4	2	-	1	1	-	-	-	-	-
Female.....	7	4	2	-	1	3	-	-	-	-	-	-
White, both sexes.....	14	8	6	2	1	4	1	-	-	-	-	-
Male.....	7	4	4	2	-	1	1	-	-	-	-	-
Female.....	7	4	2	-	1	3	-	-	-	-	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	66	32	12	4	5	5	4	2	2	1	-	-
Male.....	41	21	9	3	5	1	1	1	-	-	-	-
Female.....	25	11	3	1	-	4	3	1	2	1	-	-
White, both sexes.....	48	20	9	3	2	2	4	1	2	1	-	-
Male.....	29	15	7	2	2	-	1	-	-	-	-	-
Female.....	19	5	2	1	-	2	3	1	2	1	-	-
Black, both sexes.....	18	12	3	1	3	2	-	1	-	-	-	-
Male.....	12	6	2	1	3	1	-	1	-	-	-	-
Female.....	6	6	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Alaska

All races, both sexes.....	63	-	-	-	-	-	6	10	9	11	1	14
Male.....	39	-	-	-	-	-	4	7	5	9	1	8
Female.....	24	-	-	-	-	-	2	3	4	2	-	6
White, both sexes.....	52	-	-	-	-	-	5	10	7	9	1	10
Male.....	36	-	-	-	-	-	4	7	5	7	1	7
Female.....	16	-	-	-	-	-	1	3	2	2	-	3
Black, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	10	-	-	-	-	-	1	-	2	1	-	4
Male.....	2	-	-	-	-	-	-	-	-	1	-	1
Female.....	8	-	-	-	-	-	1	-	2	-	-	3
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	697	-	1	1	1	32	44	55	71	78	100	122
Male.....	474	-	-	-	-	24	35	44	46	51	65	86
Female.....	223	-	1	1	1	8	9	11	25	27	35	36
White, both sexes.....	637	-	-	1	1	32	42	53	64	71	83	114
Male.....	431	-	-	-	-	24	34	42	42	47	52	81
Female.....	206	-	-	1	1	8	8	11	22	24	31	33
Black, both sexes.....	36	-	1	-	-	-	2	-	4	3	9	5
Male.....	26	-	-	-	-	-	1	-	2	2	6	4
Female.....	10	-	1	-	-	-	1	-	2	1	3	1
American Indian 1/, both sexes.....	21	-	-	-	-	-	-	2	3	2	7	3
Male.....	15	-	-	-	-	-	-	2	2	1	6	1
Female.....	6	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	2	1	-
Male.....	2	-	-	-	-	-	-	-	-	1	1	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Arkansas

All races, both sexes.....	129	-	-	-	1	3	15	13	7	23	14	21
Male.....	80	-	-	-	1	2	13	11	5	13	9	10
Female.....	49	-	-	-	-	1	2	2	2	10	5	11
White, both sexes.....	116	-	-	-	1	3	12	11	7	22	13	19
Male.....	70	-	-	-	1	2	10	9	5	12	9	9
Female.....	46	-	-	-	-	1	2	2	2	10	4	10
Black, both sexes.....	13	-	-	-	-	-	3	2	-	1	1	2
Male.....	10	-	-	-	-	-	3	2	-	1	-	1
Female.....	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	3,059	1	1	-	5	62	154	186	183	307	411	569
Male.....	1,963	1	-	-	3	44	112	137	122	207	258	353
Female.....	1,096	-	1	-	2	18	42	49	61	100	153	216
White, both sexes.....	2,625	-	1	-	4	56	137	170	161	266	361	492
Male.....	1,691	-	-	-	2	41	104	131	110	178	226	298
Female.....	934	-	1	-	2	15	33	39	51	88	135	194
Black, both sexes.....	327	-	-	-	1	3	8	10	15	31	31	61
Male.....	209	-	-	-	1	1	3	5	7	21	21	46
Female.....	118	-	-	-	-	2	5	5	8	10	10	15
American Indian 1/, both sexes.....	46	-	-	-	-	2	1	2	3	1	11	11
Male.....	27	-	-	-	-	1	1	1	3	1	6	6
Female.....	19	-	-	-	-	1	-	1	-	-	5	5
Asian or Pacific Islander 2/, both sexes.....	61	1	-	-	-	1	8	4	4	9	8	5
Male.....	36	1	-	-	-	1	4	-	2	7	5	3
Female.....	25	-	-	-	-	-	4	4	2	2	3	2

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Nebraska

All races, both sexes.....	50	1	-	-	-	1	3	5	3	12	7	6
Male.....	28	1	-	-	-	1	3	5	2	5	2	3
Female.....	22	-	-	-	-	-	-	-	1	7	5	3
White, both sexes.....	46	1	-	-	-	1	2	5	3	12	6	5
Male.....	25	1	-	-	-	1	2	5	2	5	2	2
Female.....	21	-	-	-	-	-	-	-	1	7	4	3
Black, both sexes.....	2	-	-	-	-	-	1	-	-	-	1	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	1
Male.....	2	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	399	-	-	-	1	6	40	40	42	36	62	56
Male.....	241	-	-	-	-	3	27	29	30	23	35	33
Female.....	158	-	-	-	1	3	13	11	12	13	27	23
White, both sexes.....	362	-	-	-	1	5	37	36	37	32	54	54
Male.....	219	-	-	-	-	2	25	28	27	20	31	33
Female.....	143	-	-	-	1	3	12	8	10	12	23	21
Black, both sexes.....	32	-	-	-	-	1	3	3	4	4	7	2
Male.....	19	-	-	-	-	1	2	1	2	3	3	-
Female.....	13	-	-	-	-	-	1	2	2	1	4	2
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	1	-
Male.....	2	-	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	1	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

New Hampshire

All races, both sexes.....	149	-	-	-	-	5	19	18	12	18	24	20
Male.....	100	-	-	-	-	4	17	13	11	12	14	11
Female.....	49	-	-	-	-	1	2	5	1	6	10	9
White, both sexes.....	147	-	-	-	-	5	19	17	12	18	24	20
Male.....	98	-	-	-	-	4	17	12	11	12	14	11
Female.....	49	-	-	-	-	1	2	5	1	6	10	9
Black, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	-
Male.....	2	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	627	1	-	-	1	13	53	80	70	90	116	83
Male.....	456	1	-	-	-	8	41	69	53	68	89	52
Female.....	171	-	-	-	1	5	12	11	17	22	27	31
White, both sexes.....	512	-	-	-	1	12	50	75	59	71	93	70
Male.....	382	-	-	-	-	8	39	65	45	58	71	45
Female.....	130	-	-	-	1	4	11	10	14	13	22	25
Black, both sexes.....	110	1	-	-	-	-	2	5	10	18	23	13
Male.....	72	1	-	-	-	-	1	4	8	9	18	7
Female.....	38	-	-	-	-	-	1	1	2	9	5	6
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	1	1	-	1	1	-	-
Male.....	2	-	-	-	-	-	1	-	-	1	-	-
Female.....	3	-	-	-	-	1	-	-	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
New Mexico												
All races, both sexes.....	369	-	1	-	-	11	20	29	40	32	57	64
Male.....	263	-	1	-	-	10	15	25	24	28	37	44
Female.....	106	-	-	-	-	1	5	4	16	4	20	20
White, both sexes.....	347	-	1	-	-	7	19	27	39	29	53	60
Male.....	248	-	1	-	-	7	14	23	24	25	35	42
Female.....	99	-	-	-	-	-	5	4	15	4	18	18
Black, both sexes.....	7	-	-	-	-	1	-	1	-	-	2	1
Male.....	4	-	-	-	-	-	-	1	-	-	1	-
Female.....	3	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes.....	13	-	-	-	-	3	1	1	1	3	2	2
Male.....	10	-	-	-	-	3	1	1	-	3	1	1
Female.....	3	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	1,296	-	-	-	1	20	79	117	124	135	197	220
Male.....	911	-	-	-	-	16	63	98	92	91	142	138
Female.....	385	-	-	-	1	4	16	19	32	44	55	82
White, both sexes.....	1,034	-	-	-	1	17	75	109	109	109	166	162
Male.....	737	-	-	-	-	13	59	91	81	75	117	111
Female.....	297	-	-	-	1	4	16	18	28	34	49	51
Black, both sexes.....	245	-	-	-	-	3	1	5	11	24	30	56
Male.....	162	-	-	-	-	3	1	4	8	15	25	26
Female.....	83	-	-	-	-	-	-	1	3	9	5	30
American Indian 1/, both sexes.....	2	-	-	-	-	-	1	-	-	1	-	-
Male.....	2	-	-	-	-	-	1	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	15	-	-	-	-	-	2	3	4	1	1	2
Male.....	10	-	-	-	-	-	2	3	3	-	-	1
Female.....	5	-	-	-	-	-	-	-	1	1	1	1









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

South Dakota

All races, both sexes.....	19	-	-	-	1	-	1	4	4	1	2	4
Male.....	9	-	-	-	1	-	-	3	1	-	1	3
Female.....	10	-	-	-	-	-	1	1	3	1	1	1
White, both sexes.....	12	-	-	-	1	-	1	2	1	1	1	3
Male.....	5	-	-	-	1	-	-	1	-	-	1	2
Female.....	7	-	-	-	-	-	1	1	1	1	-	1
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	6	-	-	-	-	-	-	2	3	-	1	-
Male.....	3	-	-	-	-	-	-	2	1	-	-	-
Female.....	3	-	-	-	-	-	-	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	761	-	-	-	2	19	48	71	81	90	119	121
Male.....	454	-	-	-	2	12	33	44	56	57	74	71
Female.....	307	-	-	-	-	7	15	27	25	33	45	50
White, both sexes.....	653	-	-	-	2	18	46	67	73	81	106	97
Male.....	377	-	-	-	2	12	33	41	49	52	64	53
Female.....	276	-	-	-	-	6	13	26	24	29	42	44
Black, both sexes.....	105	-	-	-	-	1	1	4	7	9	13	24
Male.....	75	-	-	-	-	-	-	3	6	5	10	18
Female.....	30	-	-	-	-	1	1	1	1	4	3	6
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	1	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Vermont

All races, both sexes.....	44	-	-	-	-	-	3	6	3	8	8	5
Male.....	27	-	-	-	-	-	1	5	2	3	4	5
Female.....	17	-	-	-	-	-	2	1	1	5	4	-
White, both sexes.....	43	-	-	-	-	-	3	6	3	8	8	5
Male.....	26	-	-	-	-	-	1	5	2	3	4	5
Female.....	17	-	-	-	-	-	2	1	1	5	4	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	522	-	-	1	1	11	56	68	58	71	80	83
Male.....	337	-	-	-	1	7	41	51	41	42	46	53
Female.....	185	-	-	1	-	4	15	17	17	29	34	30
White, both sexes.....	436	-	-	1	1	11	50	60	50	56	67	71
Male.....	283	-	-	-	1	7	36	44	37	32	41	44
Female.....	153	-	-	1	-	4	14	16	13	24	26	27
Black, both sexes.....	82	-	-	-	-	-	5	7	7	15	13	12
Male.....	51	-	-	-	-	-	4	6	4	10	5	9
Female.....	31	-	-	-	-	-	1	1	3	5	8	3
American Indian 1/, both sexes.....	2	-	-	-	-	-	1	1	-	-	-	-
Male.....	2	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Arkansas

All races, both sexes.....	12	5	5	5	3	-	1	-	1	-	-	-
Male.....	6	2	1	4	2	-	-	-	1	-	-	-
Female.....	6	3	4	1	1	-	1	-	-	-	-	-
White, both sexes.....	11	4	3	5	3	-	1	-	1	-	-	-
Male.....	5	1	-	4	2	-	-	-	1	-	-	-
Female.....	6	3	3	1	1	-	1	-	-	-	-	-
Black, both sexes.....	1	1	2	-	-	-	-	-	-	-	-	-
Male.....	1	1	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	567	339	157	44	25	14	16	12	2	3	-	1
Male.....	367	209	94	24	13	5	6	5	-	2	-	1
Female.....	200	130	63	20	12	9	10	7	2	1	-	-
White, both sexes.....	471	274	133	38	19	12	13	11	2	3	-	1
Male.....	307	165	81	21	10	4	5	5	-	2	-	1
Female.....	164	109	52	17	9	8	8	6	2	1	-	-
Black, both sexes.....	82	51	21	6	4	2	1	-	-	-	-	-
Male.....	51	34	12	3	3	1	-	-	-	-	-	-
Female.....	31	17	9	3	1	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	7	6	2	-	-	-	-	-	-	-	-	-
Male.....	4	4	-	-	-	-	-	-	-	-	-	-
Female.....	3	2	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	8	1	-	2	-	2	1	-	-	-	-
Male.....	5	6	1	-	-	-	1	-	-	-	-	-
Female.....	2	2	-	-	2	-	1	1	-	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

New Hampshire

All races, both sexes.....	13	8	5	2	1	3	1	-	-	-	-	-
Male.....	6	5	3	2	-	1	1	-	-	-	-	-
Female.....	7	3	2	-	1	2	-	-	-	-	-	-
White, both sexes.....	13	7	5	2	1	3	1	-	-	-	-	-
Male.....	6	4	3	2	-	1	1	-	-	-	-	-
Female.....	7	3	2	-	1	2	-	-	-	-	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	61	31	10	2	4	5	3	1	2	1	-	-
Male.....	38	20	8	2	4	1	1	1	-	-	-	-
Female.....	23	11	2	-	-	4	2	-	2	1	-	-
White, both sexes.....	44	19	7	2	1	2	3	-	2	1	-	-
Male.....	27	14	6	2	1	-	1	-	-	-	-	-
Female.....	17	5	1	-	-	2	2	-	2	1	-	-
Black, both sexes.....	17	12	3	-	3	2	-	1	-	-	-	-
Male.....	11	6	2	-	3	1	-	1	-	-	-	-
Female.....	6	6	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other and unspecified solid or liquid substances (X45-X46,X48-X49)

Alaska

All races, both sexes.....	21	-	-	-	-	-	3	1	4	2	3	4
Male.....	13	-	-	-	-	-	3	1	1	1	2	2
Female.....	8	-	-	-	-	-	-	-	3	1	1	2
White, both sexes.....	6	-	-	-	-	-	-	-	-	1	1	2
Male.....	5	-	-	-	-	-	-	-	-	1	1	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	15	-	-	-	-	-	3	1	4	1	2	2
Male.....	8	-	-	-	-	-	3	1	1	-	1	1
Female.....	7	-	-	-	-	-	-	-	3	1	1	1
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	50	-	1	-	1	3	6	2	2	7	7	4
Male.....	36	-	1	-	-	-	5	2	-	5	5	4
Female.....	14	-	-	-	1	3	1	-	2	2	2	-
White, both sexes.....	33	-	1	-	-	3	2	2	2	5	5	1
Male.....	22	-	1	-	-	-	2	2	-	3	3	1
Female.....	11	-	-	-	-	3	-	-	2	2	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	16	-	-	-	-	-	4	-	-	2	2	3
Male.....	14	-	-	-	-	-	3	-	-	2	2	3
Female.....	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-









































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Accidental poisoning by and exposure to other and unspecified solid or liquid substances (X45-X46,X48-X49)

## Oregon

All races, both sexes.....	16	-	-	-	2	2	2	1	1	1	1	-
Male.....	14	-	-	-	2	2	2	1	1	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	1	-
White, both sexes.....	15	-	-	-	1	2	2	1	1	1	1	-
Male.....	13	-	-	-	1	2	2	1	1	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Pennsylvania

All races, both sexes.....	57	-	-	-	-	2	4	-	6	12	8	9
Male.....	35	-	-	-	-	-	4	-	4	8	3	7
Female.....	22	-	-	-	-	2	-	-	2	4	5	2
White, both sexes.....	49	-	-	-	-	2	3	-	5	12	6	7
Male.....	29	-	-	-	-	-	3	-	4	8	2	5
Female.....	20	-	-	-	-	2	-	-	1	4	4	2
Black, both sexes.....	6	-	-	-	-	-	1	-	-	-	2	2
Male.....	5	-	-	-	-	-	1	-	-	-	1	2
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other and unspecified solid or liquid substances (X45-X46,X48-X49)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other and unspecified solid or liquid substances (X45-X46,X48-X49)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other gases and vapors (X47)

New Hampshire

All races, both sexes.....	3	-	-	-	-	1	-	-	-	-	-	-
Male.....	2	-	-	-	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	3	-	-	-	-	1	-	-	-	-	-	-
Male.....	2	-	-	-	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	12	-	-	1	1	-	-	1	-	-	2	-
Male.....	7	-	-	1	-	-	-	-	-	-	2	-
Female.....	5	-	-	-	1	-	-	1	-	-	-	-
White, both sexes.....	8	-	-	-	-	-	-	-	-	-	2	-
Male.....	5	-	-	-	-	-	-	-	-	-	2	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	3	-	-	1	-	-	-	1	-	-	-	-
Male.....	2	-	-	1	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Alaska

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Accidental poisoning by and exposure to other gases and vapors (X47)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Arkansas

All races, both sexes.....	110	1	-	1	-	5	1	1	2	2	4	10
Male.....	68	1	-	1	-	4	1	1	1	1	3	8
Female.....	42	-	-	-	-	1	-	-	1	1	1	2
White, both sexes.....	92	1	-	1	-	5	1	-	1	2	3	8
Male.....	58	1	-	1	-	4	1	-	-	1	2	8
Female.....	34	-	-	-	-	1	-	-	1	1	1	-
Black, both sexes.....	17	-	-	-	-	-	-	1	1	-	1	2
Male.....	10	-	-	-	-	-	-	1	1	-	1	-
Female.....	7	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	613	4	12	5	6	14	14	24	19	28	40	41
Male.....	405	2	7	3	5	13	11	17	17	21	36	31
Female.....	208	2	5	2	1	1	3	7	2	7	4	10
White, both sexes.....	520	2	11	3	4	10	13	23	16	22	29	35
Male.....	339	1	6	2	3	10	10	16	14	16	26	27
Female.....	181	1	5	1	1	-	3	7	2	6	3	8
Black, both sexes.....	45	-	-	2	-	2	-	1	1	5	7	4
Male.....	36	-	-	1	-	2	-	1	1	4	6	2
Female.....	9	-	-	1	-	-	-	-	-	1	1	2
American Indian 1/, both sexes.....	8	-	-	-	1	-	-	-	1	1	-	1
Male.....	5	-	-	-	1	-	-	-	1	1	-	1
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	40	2	1	-	1	2	1	-	1	-	4	1
Male.....	25	1	1	-	1	1	1	-	1	-	4	1
Female.....	15	1	-	-	-	1	-	-	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Illinois

All races, both sexes.....	490	2	2	2	-	5	5	7	10	10	12	23
Male.....	265	2	1	2	-	3	4	5	8	7	11	18
Female.....	225	-	1	-	-	2	1	2	2	3	1	5
White, both sexes.....	440	2	1	2	-	4	4	5	9	9	7	16
Male.....	235	2	-	2	-	3	3	4	8	7	6	12
Female.....	205	-	1	-	-	1	1	1	1	2	1	4
Black, both sexes.....	42	-	1	-	-	1	1	2	1	1	4	7
Male.....	24	-	1	-	-	-	1	1	-	-	4	6
Female.....	18	-	-	-	-	1	-	1	1	1	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	-	-	-	-	-	1	-
Male.....	6	-	-	-	-	-	-	-	-	-	1	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-

Indiana

All races, both sexes.....	301	-	2	2	4	3	4	1	5	6	9	15
Male.....	145	-	2	2	2	2	2	1	3	5	7	13
Female.....	156	-	-	-	2	1	2	-	2	1	2	2
White, both sexes.....	277	-	2	-	3	2	4	1	5	6	7	11
Male.....	131	-	2	-	2	2	2	1	3	5	6	10
Female.....	146	-	-	-	1	-	2	-	2	1	1	1
Black, both sexes.....	20	-	-	1	-	-	-	-	-	-	2	3
Male.....	13	-	-	1	-	-	-	-	-	-	1	3
Female.....	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	1	1	1	-	-	-	-	-	1
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	1	1	-	-	-	-	-	1



















## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

## New Mexico

All races, both sexes.....	70	-	-	2	1	2	4	1	3	-	7	5
Male.....	55	-	-	-	-	2	4	1	3	-	6	5
Female.....	15	-	-	2	1	-	-	-	-	-	1	-
White, both sexes.....	54	-	-	1	1	2	2	1	2	-	4	4
Male.....	41	-	-	-	-	2	2	1	2	-	3	4
Female.....	13	-	-	1	1	-	-	-	-	-	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	16	-	-	1	-	-	2	-	1	-	3	1
Male.....	14	-	-	-	-	-	2	-	1	-	3	1
Female.....	2	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## New York

All races, both sexes.....	402	5	1	1	1	5	10	11	10	9	15	25
Male.....	239	4	1	1	1	3	7	11	8	8	11	22
Female.....	163	1	-	-	-	2	3	-	2	1	4	3
White, both sexes.....	340	4	1	1	1	4	7	10	4	8	14	19
Male.....	195	4	1	1	1	3	5	10	4	8	10	17
Female.....	145	-	-	-	-	1	2	-	-	-	4	2
Black, both sexes.....	53	1	-	-	-	-	3	1	5	1	1	6
Male.....	39	-	-	-	-	-	2	1	3	-	1	5
Female.....	14	1	-	-	-	-	1	-	2	1	-	1
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	-	-	-	-	1	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	1	-	-	-	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Arkansas

All races, both sexes.....	8	9	9	1	6	9	14	13	11	3	-	-
Male.....	8	7	8	1	4	6	4	6	2	1	-	-
Female.....	-	2	1	-	2	3	10	7	9	2	-	-
White, both sexes.....	7	7	6	-	6	9	12	12	8	3	-	-
Male.....	7	6	5	-	4	6	4	6	1	1	-	-
Female.....	-	1	1	-	2	3	8	6	7	2	-	-
Black, both sexes.....	1	1	3	1	-	-	2	1	3	-	-	-
Male.....	1	1	3	1	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	2	1	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	59	42	29	22	25	46	56	51	50	18	7	1
Male.....	51	33	21	18	13	25	32	20	23	3	2	1
Female.....	8	9	8	4	12	21	24	31	27	15	5	-
White, both sexes.....	50	35	26	18	22	31	51	48	47	18	6	-
Male.....	44	26	19	15	12	19	28	19	21	3	2	-
Female.....	6	9	7	3	10	12	23	29	26	15	4	-
Black, both sexes.....	4	5	1	4	2	4	3	-	-	-	-	-
Male.....	3	5	1	3	1	4	2	-	-	-	-	-
Female.....	1	-	-	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	-	-	1	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	1	1	-	1	10	2	3	3	-	1	1
Male.....	4	1	1	-	-	2	2	1	2	-	-	1
Female.....	-	-	-	-	1	8	-	2	1	-	1	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Nebraska

All races, both sexes.....	1	4	1	2	3	5	11	8	9	5	1	-
Male.....	1	3	1	2	2	4	7	4	6	-	-	-
Female.....	-	1	-	-	1	1	4	4	3	5	1	-
White, both sexes.....	1	3	1	2	2	5	11	8	9	5	1	-
Male.....	1	2	1	2	2	4	7	4	6	-	-	-
Female.....	-	1	-	-	-	1	4	4	3	5	1	-
Black, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	4	6	2	5	1	3	5	5	8	4	-	-
Male.....	3	5	1	5	-	2	2	2	4	2	-	-
Female.....	1	1	1	-	1	1	3	3	4	2	-	-
White, both sexes.....	2	6	2	5	1	2	4	5	8	4	-	-
Male.....	1	5	1	5	-	2	2	2	4	2	-	-
Female.....	1	1	1	-	1	-	2	3	4	2	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

New Hampshire

All races, both sexes.....	2	1	1	1	1	4	2	5	4	1	-	-
Male.....	2	1	1	1	1	3	1	2	1	-	-	-
Female.....	-	-	-	-	-	1	1	3	3	1	-	-
White, both sexes.....	2	1	1	1	1	4	2	5	4	1	-	-
Male.....	2	1	1	1	1	3	1	2	1	-	-	-
Female.....	-	-	-	-	-	1	1	3	3	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	15	8	8	19	8	29	41	70	46	25	1	-
Male.....	12	8	6	11	5	15	19	21	13	8	-	-
Female.....	3	-	2	8	3	14	22	49	33	17	1	-
White, both sexes.....	10	7	6	17	7	25	39	68	42	23	1	-
Male.....	8	7	5	10	4	13	18	21	12	8	-	-
Female.....	2	-	1	7	3	12	21	47	30	15	1	-
Black, both sexes.....	4	1	2	2	-	3	1	-	2	2	-	-
Male.....	4	1	1	1	-	2	1	-	1	-	-	-
Female.....	-	-	1	1	-	1	-	-	1	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	1	1	1	2	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

New Mexico

All races, both sexes.....	8	6	5	3	3	5	4	3	5	3	-	-
Male.....	7	6	4	3	1	3	3	3	3	1	-	-
Female.....	1	-	1	-	2	2	1	-	2	2	-	-
White, both sexes.....	5	5	4	3	3	3	4	3	5	2	-	-
Male.....	4	5	3	3	1	2	3	3	3	-	-	-
Female.....	1	-	1	-	2	1	1	-	2	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	1	1	-	-	2	-	-	-	1	-	-
Male.....	3	1	1	-	-	1	-	-	-	1	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	18	16	23	29	15	38	44	54	46	24	2	-
Male.....	15	11	19	18	10	22	22	26	12	7	-	-
Female.....	3	5	4	11	5	16	22	28	34	17	2	-
White, both sexes.....	11	9	16	25	11	34	40	51	45	23	2	-
Male.....	10	5	13	14	7	19	20	24	12	7	-	-
Female.....	1	4	3	11	4	15	20	27	33	16	2	-
Black, both sexes.....	6	6	7	3	3	3	3	3	-	1	-	-
Male.....	5	5	6	3	2	2	2	2	-	-	-	-
Female.....	1	1	1	-	1	1	1	1	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	1	1	1	1	-	-	-	-	-
Male.....	-	1	-	1	1	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Texas

All races, both sexes.....	62	56	46	48	41	57	101	115	84	48	13	1
Male.....	50	43	32	35	30	28	57	48	24	9	2	1
Female.....	12	13	14	13	11	29	44	67	60	39	11	-
White, both sexes.....	50	47	36	41	36	54	93	111	73	44	13	1
Male.....	42	36	26	29	27	27	54	47	21	8	2	1
Female.....	8	11	10	12	9	27	39	64	52	36	11	-
Black, both sexes.....	10	9	10	6	4	3	7	4	10	4	-	-
Male.....	7	7	6	5	2	1	3	1	3	1	-	-
Female.....	3	2	4	1	2	2	4	3	7	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	1	1	-	-	-	1	-	-	-
Male.....	1	-	-	1	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-

Utah

All races, both sexes.....	1	2	4	4	6	3	11	16	13	3	-	-
Male.....	1	1	2	2	4	-	5	4	5	-	-	-
Female.....	-	1	2	2	2	3	6	12	8	3	-	-
White, both sexes.....	1	2	4	4	6	3	9	16	13	3	-	-
Male.....	1	1	2	2	4	-	5	4	5	-	-	-
Female.....	-	1	2	2	2	3	4	12	8	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Alaska												
All races, both sexes.....	149	...	...	-	1	14	18	20	13	13	12	15
Male.....	123	...	...	-	1	12	14	17	12	9	9	10
Female.....	26	...	...	-	-	2	4	3	1	4	3	5
White, both sexes.....	94	...	...	-	-	3	8	10	7	12	8	10
Male.....	80	...	...	-	-	3	7	10	7	8	6	7
Female.....	14	...	...	-	-	-	1	-	-	4	2	3
Black, both sexes.....	4	...	...	-	-	1	2	1	-	-	-	-
Male.....	4	...	...	-	-	1	2	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	48	...	...	-	1	10	8	7	6	1	4	5
Male.....	36	...	...	-	1	8	5	4	5	1	3	3
Female.....	12	...	...	-	-	2	3	3	1	-	1	2
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	-	2	-	-	-	-
Male.....	3	...	...	-	-	-	-	2	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	1,016	...	...	-	5	39	96	102	84	76	93	113
Male.....	798	...	...	-	3	29	86	89	68	58	68	78
Female.....	218	...	...	-	2	10	10	13	16	18	25	35
White, both sexes.....	950	...	...	-	3	33	83	89	77	68	91	109
Male.....	744	...	...	-	2	25	75	77	62	50	66	78
Female.....	206	...	...	-	1	8	8	12	15	18	25	31
Black, both sexes.....	18	...	...	-	-	-	1	4	2	3	2	2
Male.....	15	...	...	-	-	-	1	4	1	3	2	-
Female.....	3	...	...	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes.....	36	...	...	-	1	5	9	6	4	5	-	1
Male.....	29	...	...	-	-	4	7	5	4	5	-	-
Female.....	7	...	...	-	1	1	2	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	12	...	...	-	1	1	3	3	1	-	-	1
Male.....	10	...	...	-	1	-	3	3	1	-	-	-
Female.....	2	...	...	-	-	1	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Arkansas												
All races, both sexes.....	402	...	...	-	2	16	24	38	31	34	59	50
Male.....	322	...	...	-	2	12	18	34	22	28	41	39
Female.....	80	...	...	-	-	4	6	4	9	6	18	11
White, both sexes.....	372	...	...	-	1	14	19	33	27	31	53	49
Male.....	296	...	...	-	1	10	14	29	18	25	37	38
Female.....	76	...	...	-	-	4	5	4	9	6	16	11
Black, both sexes.....	26	...	...	-	1	2	5	4	4	3	4	1
Male.....	23	...	...	-	1	2	4	4	4	3	3	1
Female.....	3	...	...	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes.....	2	...	...	-	-	-	-	1	-	-	1	-
Male.....	2	...	...	-	-	-	-	1	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	-	-	-	-	-	1	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
California												
All races, both sexes.....	3,602	...	...	1	14	123	252	265	249	306	350	431
Male.....	2,770	...	...	1	11	102	206	219	199	241	256	326
Female.....	832	...	...	-	3	21	46	46	50	65	94	105
White, both sexes.....	3,082	...	...	1	10	95	185	213	201	246	303	379
Male.....	2,384	...	...	1	9	79	154	180	163	200	221	284
Female.....	698	...	...	-	1	16	31	33	38	46	82	95
Black, both sexes.....	167	...	...	-	1	9	23	22	19	21	18	18
Male.....	131	...	...	-	1	7	19	17	17	15	13	15
Female.....	36	...	...	-	-	2	4	5	2	6	5	3
American Indian 1/, both sexes.....	24	...	...	-	-	-	4	4	2	3	2	1
Male.....	19	...	...	-	-	-	4	4	2	2	1	1
Female.....	5	...	...	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes.....	329	...	...	-	3	19	40	26	27	36	27	33
Male.....	236	...	...	-	1	16	29	18	17	24	21	26
Female.....	93	...	...	-	2	3	11	8	10	12	6	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Colorado												
All races, both sexes.....	811	...	...	-	6	34	54	74	61	71	96	88
Male.....	632	...	...	-	6	28	44	54	53	54	75	66
Female.....	179	...	...	-	-	6	10	20	8	17	21	22
White, both sexes.....	766	...	...	-	6	31	46	70	60	62	92	79
Male.....	597	...	...	-	6	26	38	50	52	49	71	60
Female.....	169	...	...	-	-	5	8	20	8	13	21	19
Black, both sexes.....	22	...	...	-	-	1	5	2	1	3	3	4
Male.....	17	...	...	-	-	-	4	2	1	1	3	3
Female.....	5	...	...	-	-	1	1	-	-	2	-	1
American Indian 1/, both sexes.....	10	...	...	-	-	-	1	1	-	4	-	3
Male.....	8	...	...	-	-	-	1	1	-	3	-	2
Female.....	2	...	...	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes.....	13	...	...	-	-	2	2	1	-	2	1	2
Male.....	10	...	...	-	-	2	1	1	-	1	1	1
Female.....	3	...	...	-	-	-	1	-	-	1	-	1
Connecticut												
All races, both sexes.....	271	...	...	-	-	17	16	18	14	17	33	40
Male.....	213	...	...	-	-	15	15	15	13	12	24	31
Female.....	58	...	...	-	-	2	1	3	1	5	9	9
White, both sexes.....	260	...	...	-	-	16	14	17	13	16	32	38
Male.....	203	...	...	-	-	14	14	14	12	11	23	29
Female.....	57	...	...	-	-	2	-	3	1	5	9	9
Black, both sexes.....	9	...	...	-	-	1	1	1	1	1	1	2
Male.....	9	...	...	-	-	1	1	1	1	1	1	2
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Delaware												
All races, both sexes.....	95	...	...	-	-	5	5	2	4	11	11	19
Male.....	71	...	...	-	-	2	5	1	2	9	7	16
Female.....	24	...	...	-	-	3	-	1	2	2	4	3
White, both sexes.....	86	...	...	-	-	4	5	1	3	10	9	18
Male.....	63	...	...	-	-	1	5	-	2	8	5	15
Female.....	23	...	...	-	-	3	-	1	1	2	4	3
Black, both sexes.....	7	...	...	-	-	-	-	-	1	1	2	1
Male.....	6	...	...	-	-	-	-	-	-	1	2	1
Female.....	1	...	...	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	1	-	1	-	-	-	-
Male.....	2	...	...	-	-	1	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes.....	36	...	...	-	-	1	1	3	4	3	2	4
Male.....	24	...	...	-	-	1	1	2	2	2	1	2
Female.....	12	...	...	-	-	-	-	1	2	1	1	2
White, both sexes.....	14	...	...	-	-	-	-	2	1	1	-	1
Male.....	12	...	...	-	-	-	-	2	-	1	-	1
Female.....	2	...	...	-	-	-	-	-	1	-	-	-
Black, both sexes.....	21	...	...	-	-	1	1	-	3	2	2	3
Male.....	12	...	...	-	-	1	1	-	2	1	1	1
Female.....	9	...	...	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Florida												
All races, both sexes.....	2,587	...	...	-	6	71	156	143	175	192	251	320
Male.....	2,001	...	...	-	3	64	127	115	133	145	179	237
Female.....	586	...	...	-	3	7	29	28	42	47	72	83
White, both sexes.....	2,406	...	...	-	3	62	127	123	152	174	231	304
Male.....	1,860	...	...	-	2	55	101	97	113	132	167	225
Female.....	546	...	...	-	1	7	26	26	39	42	64	79
Black, both sexes.....	134	...	...	-	3	6	24	17	17	13	12	10
Male.....	107	...	...	-	1	6	23	15	15	9	7	7
Female.....	27	...	...	-	2	-	1	2	2	4	5	3
American Indian 1/, both sexes.....	3	...	...	-	-	-	1	-	1	-	-	-
Male.....	2	...	...	-	-	-	-	-	1	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	44	...	...	-	-	3	4	3	5	5	8	6
Male.....	32	...	...	-	-	3	3	3	4	4	5	5
Female.....	12	...	...	-	-	-	1	-	1	1	3	1
Georgia												
All races, both sexes.....	997	...	...	-	3	29	76	67	78	98	120	126
Male.....	801	...	...	-	2	23	63	53	59	80	95	99
Female.....	196	...	...	-	1	6	13	14	19	18	25	27
White, both sexes.....	862	...	...	-	2	22	56	50	63	77	108	111
Male.....	686	...	...	-	2	16	45	40	47	64	84	86
Female.....	176	...	...	-	-	6	11	10	16	13	24	25
Black, both sexes.....	122	...	...	-	-	7	18	15	14	18	12	12
Male.....	105	...	...	-	-	7	16	11	12	13	11	11
Female.....	17	...	...	-	-	-	2	4	2	5	1	1
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	12	...	...	-	1	-	2	2	1	3	-	3
Male.....	9	...	...	-	-	-	2	2	-	3	-	2
Female.....	3	...	...	-	1	-	-	-	1	-	-	1









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Kentucky												
All races, both sexes.....	649	...	...	-	5	32	40	68	58	62	80	72
Male.....	526	...	...	-	4	23	35	56	48	50	60	59
Female.....	123	...	...	-	1	9	5	12	10	12	20	13
White, both sexes.....	615	...	...	-	4	29	37	60	53	59	76	69
Male.....	496	...	...	-	3	20	32	49	45	47	56	56
Female.....	119	...	...	-	1	9	5	11	8	12	20	13
Black, both sexes.....	29	...	...	-	1	3	3	5	5	1	4	3
Male.....	25	...	...	-	1	3	3	4	3	1	4	3
Female.....	4	...	...	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	...	...	-	-	-	-	2	-	2	-	-
Male.....	4	...	...	-	-	-	-	2	-	2	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes.....	522	...	...	-	9	25	45	66	47	51	51	55
Male.....	426	...	...	-	8	23	42	54	42	40	39	37
Female.....	96	...	...	-	1	2	3	12	5	11	12	18
White, both sexes.....	446	...	...	-	7	20	34	54	36	42	43	49
Male.....	366	...	...	-	7	19	32	45	32	34	34	33
Female.....	80	...	...	-	-	1	2	9	4	8	9	16
Black, both sexes.....	72	...	...	-	2	5	11	11	11	8	7	6
Male.....	59	...	...	-	1	4	10	9	10	6	5	4
Female.....	13	...	...	-	1	1	1	2	1	2	2	2
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	-	1	-	1	1	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	3	...	...	-	-	-	-	1	-	1	1	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Maine												
All races, both sexes.....	191	...	...	-	-	13	18	14	10	12	20	19
Male.....	156	...	...	-	-	9	16	11	10	8	15	18
Female.....	35	...	...	-	-	4	2	3	-	4	5	1
White, both sexes.....	186	...	...	-	-	12	16	14	10	12	19	19
Male.....	151	...	...	-	-	8	14	11	10	8	14	18
Female.....	35	...	...	-	-	4	2	3	-	4	5	1
Black, both sexes.....	2	...	...	-	-	1	1	-	-	-	-	-
Male.....	2	...	...	-	-	1	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	1	...	...	-	-	-	-	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	-	1	-	-	-	-	-
Male.....	2	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes.....	518	...	...	-	2	27	41	39	50	42	71	56
Male.....	405	...	...	-	-	22	33	33	40	35	54	43
Female.....	113	...	...	-	2	5	8	6	10	7	17	13
White, both sexes.....	417	...	...	-	-	21	32	29	37	27	53	51
Male.....	328	...	...	-	-	17	27	26	29	23	42	39
Female.....	89	...	...	-	-	4	5	3	8	4	11	12
Black, both sexes.....	84	...	...	-	2	4	7	8	11	14	16	5
Male.....	67	...	...	-	-	4	5	6	9	11	12	4
Female.....	17	...	...	-	2	-	2	2	2	3	4	1
American Indian 1/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	16	...	...	-	-	2	1	2	2	1	2	-
Male.....	9	...	...	-	-	1	-	1	2	1	-	-
Female.....	7	...	...	-	-	1	1	1	-	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Massachusetts												
All races, both sexes.....	516	...	...	-	2	19	33	40	40	60	60	50
Male.....	405	...	...	-	2	16	29	37	30	50	47	40
Female.....	111	...	...	-	-	3	4	3	10	10	13	10
White, both sexes.....	483	...	...	-	2	17	29	33	33	53	58	48
Male.....	374	...	...	-	2	14	25	30	24	43	46	38
Female.....	109	...	...	-	-	3	4	3	9	10	12	10
Black, both sexes.....	22	...	...	-	-	1	4	5	2	5	2	2
Male.....	21	...	...	-	-	1	4	5	2	5	1	2
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	11	...	...	-	-	1	-	2	5	2	-	-
Male.....	10	...	...	-	-	1	-	2	4	2	-	-
Female.....	1	...	...	-	-	-	-	-	1	-	-	-
Michigan												
All races, both sexes.....	1,131	...	...	-	6	52	80	86	78	103	115	162
Male.....	905	...	...	-	3	40	64	71	69	81	89	128
Female.....	226	...	...	-	3	12	16	15	9	22	26	34
White, both sexes.....	1,040	...	...	-	3	42	71	77	71	93	105	153
Male.....	835	...	...	-	1	33	56	64	63	74	80	120
Female.....	205	...	...	-	2	9	15	13	8	19	25	33
Black, both sexes.....	74	...	...	-	3	5	8	9	4	9	8	6
Male.....	60	...	...	-	2	4	7	7	4	7	8	6
Female.....	14	...	...	-	1	1	1	2	-	2	-	-
American Indian 1/, both sexes.....	8	...	...	-	-	2	-	-	2	-	1	3
Male.....	5	...	...	-	-	1	-	-	1	-	1	2
Female.....	3	...	...	-	-	1	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes.....	9	...	...	-	-	3	1	-	1	1	1	-
Male.....	5	...	...	-	-	2	1	-	1	-	-	-
Female.....	4	...	...	-	-	1	-	-	-	1	1	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Nebraska												
All races, both sexes.....	181	...	...	-	3	15	16	9	12	18	18	19
Male.....	139	...	...	-	-	12	15	9	10	13	15	12
Female.....	42	...	...	-	3	3	1	-	2	5	3	7
White, both sexes.....	175	...	...	-	3	15	13	8	12	18	17	19
Male.....	133	...	...	-	-	12	12	8	10	13	14	12
Female.....	42	...	...	-	3	3	1	-	2	5	3	7
Black, both sexes.....	4	...	...	-	-	-	2	-	-	-	1	-
Male.....	4	...	...	-	-	-	2	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	...	...	-	-	-	1	1	-	-	-	-
Male.....	2	...	...	-	-	-	1	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes.....	471	...	...	-	5	16	27	30	25	44	49	45
Male.....	369	...	...	-	2	12	25	24	18	35	37	36
Female.....	102	...	...	-	3	4	2	6	7	9	12	9
White, both sexes.....	432	...	...	-	2	15	23	25	24	40	46	43
Male.....	339	...	...	-	-	11	21	20	17	31	34	34
Female.....	93	...	...	-	2	4	2	5	7	9	12	9
Black, both sexes.....	18	...	...	-	2	1	2	3	1	2	1	1
Male.....	16	...	...	-	2	1	2	3	1	2	1	1
Female.....	2	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	4	...	...	-	-	-	-	2	-	1	-	-
Male.....	2	...	...	-	-	-	-	1	-	1	-	-
Female.....	2	...	...	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	17	...	...	-	1	-	2	-	-	1	2	1
Male.....	12	...	...	-	-	-	2	-	-	1	2	1
Female.....	5	...	...	-	1	-	-	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
New Hampshire												
All races, both sexes.....	158	...	...	-	1	5	10	4	12	9	19	23
Male.....	123	...	...	-	1	2	7	3	8	5	14	18
Female.....	35	...	...	-	-	3	3	1	4	4	5	5
White, both sexes.....	155	...	...	-	1	5	10	4	11	9	18	23
Male.....	122	...	...	-	1	2	7	3	8	5	14	18
Female.....	33	...	...	-	-	3	3	1	3	4	4	5
Black, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	1	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	1	-	-	-
New Jersey												
All races, both sexes.....	596	...	...	-	3	24	50	44	47	60	66	62
Male.....	472	...	...	-	1	22	37	38	38	48	48	46
Female.....	124	...	...	-	2	2	13	6	9	12	18	16
White, both sexes.....	530	...	...	-	3	21	39	40	39	54	57	56
Male.....	424	...	...	-	1	19	28	34	32	44	41	43
Female.....	106	...	...	-	2	2	11	6	7	10	16	13
Black, both sexes.....	35	...	...	-	-	2	7	3	3	5	6	3
Male.....	30	...	...	-	-	2	7	3	3	3	6	2
Female.....	5	...	...	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	31	...	...	-	-	1	4	1	5	1	3	3
Male.....	18	...	...	-	-	1	2	1	3	1	1	1
Female.....	13	...	...	-	-	-	2	-	2	-	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
New Mexico												
All races, both sexes.....	401	...	...	-	6	31	30	32	25	27	52	44
Male.....	326	...	...	-	6	26	25	25	20	24	39	37
Female.....	75	...	...	-	-	5	5	7	5	3	13	7
White, both sexes.....	342	...	...	-	2	25	20	26	18	21	42	42
Male.....	274	...	...	-	2	20	17	20	13	19	30	35
Female.....	68	...	...	-	-	5	3	6	5	2	12	7
Black, both sexes.....	4	...	...	-	1	-	2	-	-	-	-	-
Male.....	3	...	...	-	1	-	1	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	52	...	...	-	3	6	8	6	7	5	9	2
Male.....	48	...	...	-	3	6	7	5	7	4	9	2
Female.....	4	...	...	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	-	-	-	1	1	-
Male.....	1	...	...	-	-	-	-	-	-	1	-	-
Female.....	2	...	...	-	-	-	-	-	-	-	1	-
New York												
All races, both sexes.....	1,396	...	...	-	4	52	97	119	93	106	175	153
Male.....	1,065	...	...	-	2	42	80	92	71	81	140	122
Female.....	331	...	...	-	2	10	17	27	22	25	35	31
White, both sexes.....	1,195	...	...	-	3	37	72	88	71	88	153	140
Male.....	928	...	...	-	2	30	63	72	58	66	125	111
Female.....	267	...	...	-	1	7	9	16	13	22	28	29
Black, both sexes.....	126	...	...	-	1	11	15	23	12	11	13	11
Male.....	91	...	...	-	-	9	10	18	9	8	11	9
Female.....	35	...	...	-	1	2	5	5	3	3	2	2
American Indian 1/, both sexes.....	3	...	...	-	-	1	-	-	1	-	1	-
Male.....	1	...	...	-	-	-	-	-	1	-	-	-
Female.....	2	...	...	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	72	...	...	-	-	3	10	8	9	7	8	2
Male.....	45	...	...	-	-	3	7	2	3	7	4	2
Female.....	27	...	...	-	-	-	3	6	6	-	4	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Oregon												
All races, both sexes.....	594	...	...	-	1	21	37	38	51	47	52	60
Male.....	466	...	...	-	1	16	29	33	41	35	38	45
Female.....	128	...	...	-	-	5	8	5	10	12	14	15
White, both sexes.....	575	...	...	-	1	18	33	34	51	45	48	59
Male.....	455	...	...	-	1	14	26	30	41	35	36	44
Female.....	120	...	...	-	-	4	7	4	10	10	12	15
Black, both sexes.....	6	...	...	-	-	1	2	2	-	-	1	-
Male.....	6	...	...	-	-	1	2	2	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	...	...	-	-	2	-	-	-	-	1	-
Male.....	2	...	...	-	-	1	-	-	-	-	1	-
Female.....	1	...	...	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	10	...	...	-	-	-	2	2	-	2	2	1
Male.....	3	...	...	-	-	-	1	1	-	-	-	1
Female.....	7	...	...	-	-	-	1	1	-	2	2	-
Pennsylvania												
All races, both sexes.....	1,441	...	...	-	8	48	118	107	97	137	158	171
Male.....	1,188	...	...	-	7	38	105	94	82	116	133	126
Female.....	253	...	...	-	1	10	13	13	15	21	25	45
White, both sexes.....	1,346	...	...	-	8	43	103	96	86	127	149	161
Male.....	1,112	...	...	-	7	35	94	84	74	109	124	118
Female.....	234	...	...	-	1	8	9	12	12	18	25	43
Black, both sexes.....	78	...	...	-	-	5	11	6	10	10	7	9
Male.....	64	...	...	-	-	3	9	6	8	7	7	8
Female.....	14	...	...	-	-	2	2	-	2	3	-	1
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	16	...	...	-	-	-	4	4	1	-	2	1
Male.....	11	...	...	-	-	-	2	3	-	-	2	-
Female.....	5	...	...	-	-	-	2	1	1	-	-	1



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

South Dakota

All races, both sexes.....	102	...	...	-	3	12	12	12	8	5	2	11
Male.....	76	...	...	-	1	7	12	10	4	4	1	10
Female.....	26	...	...	-	2	5	-	2	4	1	1	1
White, both sexes.....	80	...	...	-	2	4	8	8	7	4	2	9
Male.....	60	...	...	-	1	2	8	6	4	3	1	9
Female.....	20	...	...	-	1	2	-	2	3	1	1	-
Black, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	20	...	...	-	1	8	4	2	1	1	-	2
Male.....	14	...	...	-	-	5	4	2	-	1	-	1
Female.....	6	...	...	-	1	3	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	844	...	...	1	3	29	54	61	67	85	88	99
Male.....	643	...	...	1	3	26	45	48	56	59	62	70
Female.....	201	...	...	-	-	3	9	13	11	26	26	29
White, both sexes.....	780	...	...	-	1	22	48	54	59	79	85	92
Male.....	591	...	...	-	1	19	40	41	49	56	61	64
Female.....	189	...	...	-	-	3	8	13	10	23	24	28
Black, both sexes.....	53	...	...	1	2	6	4	5	7	6	3	7
Male.....	44	...	...	1	2	6	4	5	7	3	1	6
Female.....	9	...	...	-	-	-	-	-	-	3	2	1
American Indian 1/, both sexes.....	3	...	...	-	-	-	1	2	-	-	-	-
Male.....	3	...	...	-	-	-	1	2	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	...	...	-	-	1	1	-	1	-	-	-
Male.....	5	...	...	-	-	1	-	-	-	-	-	-
Female.....	3	...	...	-	-	-	1	-	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Texas												
All races, both sexes.....	2,433	...	...	-	12	134	219	194	230	217	254	285
Male.....	1,932	...	...	-	10	108	190	153	189	167	190	214
Female.....	501	...	...	-	2	26	29	41	41	50	64	71
White, both sexes.....	2,261	...	...	-	9	121	199	178	207	199	232	268
Male.....	1,793	...	...	-	8	98	171	139	169	154	174	201
Female.....	468	...	...	-	1	23	28	39	38	45	58	67
Black, both sexes.....	120	...	...	-	3	10	13	12	19	11	13	14
Male.....	104	...	...	-	2	8	13	12	17	7	11	11
Female.....	16	...	...	-	1	2	-	-	2	4	2	3
American Indian 1/, both sexes.....	3	...	...	-	-	-	-	-	-	-	1	-
Male.....	2	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	49	...	...	-	-	3	7	4	4	7	8	3
Male.....	33	...	...	-	-	2	6	2	3	6	5	2
Female.....	16	...	...	-	-	1	1	2	1	1	3	1
Utah												
All races, both sexes.....	378	...	...	-	1	23	47	45	32	39	30	42
Male.....	302	...	...	-	-	21	37	40	27	28	18	33
Female.....	76	...	...	-	1	2	10	5	5	11	12	9
White, both sexes.....	362	...	...	-	1	19	42	45	30	37	29	42
Male.....	288	...	...	-	-	17	33	40	25	26	18	33
Female.....	74	...	...	-	1	2	9	5	5	11	11	9
Black, both sexes.....	2	...	...	-	-	-	-	-	1	1	-	-
Male.....	2	...	...	-	-	-	-	-	1	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	10	...	...	-	-	3	4	-	1	-	1	-
Male.....	9	...	...	-	-	3	4	-	1	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	4	...	...	-	-	1	1	-	-	1	-	-
Male.....	3	...	...	-	-	1	-	-	-	1	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-



## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Vermont												
All races, both sexes.....	89	...	...	-	2	3	7	8	7	4	11	5
Male.....	70	...	...	-	1	3	6	7	7	2	9	3
Female.....	19	...	...	-	1	-	1	1	-	2	2	2
White, both sexes.....	88	...	...	-	2	3	6	8	7	4	11	5
Male.....	69	...	...	-	1	3	5	7	7	2	9	3
Female.....	19	...	...	-	1	-	1	1	-	2	2	2
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	880	...	...	-	4	33	64	59	61	80	105	113
Male.....	651	...	...	-	4	27	56	42	44	54	75	73
Female.....	229	...	...	-	-	6	8	17	17	26	30	40
White, both sexes.....	764	...	...	-	3	28	46	45	56	65	96	94
Male.....	564	...	...	-	3	22	40	34	40	44	68	63
Female.....	200	...	...	-	-	6	6	11	16	21	28	31
Black, both sexes.....	88	...	...	-	1	5	12	12	4	10	5	17
Male.....	68	...	...	-	1	5	10	7	4	8	3	10
Female.....	20	...	...	-	-	-	2	5	-	2	2	7
American Indian 1/, both sexes.....	3	...	...	-	-	-	2	-	-	-	1	-
Male.....	3	...	...	-	-	-	2	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	25	...	...	-	-	-	4	2	1	5	3	2
Male.....	16	...	...	-	-	-	4	1	-	2	3	-
Female.....	9	...	...	-	-	-	-	1	1	3	-	2









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Arkansas

All races, both sexes.....	35	29	23	19	9	16	10	5	1	1	-	-
Male.....	30	21	19	17	7	16	10	4	1	1	-	-
Female.....	5	8	4	2	2	-	-	1	-	-	-	-
White, both sexes.....	35	28	22	18	9	16	10	5	1	1	-	-
Male.....	30	21	18	16	7	16	10	4	1	1	-	-
Female.....	5	7	4	2	2	-	-	1	-	-	-	-
Black, both sexes.....	-	1	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	433	343	242	141	90	149	105	68	33	4	1	2
Male.....	320	238	177	120	70	117	86	48	26	4	1	2
Female.....	113	105	65	21	20	32	19	20	7	-	-	-
White, both sexes.....	382	318	222	127	78	131	94	60	31	4	-	2
Male.....	281	217	164	108	63	108	78	42	26	4	-	2
Female.....	101	101	58	19	15	23	16	18	5	-	-	-
Black, both sexes.....	15	8	5	4	1	2	-	-	1	-	-	-
Male.....	10	7	3	4	1	2	-	-	-	-	-	-
Female.....	5	1	2	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	4	1	2	-	-	-	-	1	-	-	-	-
Male.....	3	1	1	-	-	-	-	-	-	-	-	-
Female.....	1	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	32	16	13	10	11	16	11	7	1	-	1	-
Male.....	26	13	9	8	6	7	8	6	-	-	1	-
Female.....	6	3	4	2	5	9	3	1	1	-	-	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Maine

All races, both sexes.....	18	19	11	9	7	10	4	5	2	-	-	-
Male.....	16	13	8	9	5	9	3	5	1	-	-	-
Female.....	2	6	3	-	2	1	1	-	1	-	-	-
White, both sexes.....	17	19	11	9	7	10	4	5	2	-	-	-
Male.....	15	13	8	9	5	9	3	5	1	-	-	-
Female.....	2	6	3	-	2	1	1	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	49	39	33	15	16	17	14	5	2	-	-	-
Male.....	39	32	21	10	11	14	11	5	2	-	-	-
Female.....	10	7	12	5	5	3	3	-	-	-	-	-
White, both sexes.....	40	34	28	14	16	16	12	5	2	-	-	-
Male.....	30	27	18	9	11	13	10	5	2	-	-	-
Female.....	10	7	10	5	5	3	2	-	-	-	-	-
Black, both sexes.....	7	5	3	1	-	-	1	-	-	-	-	-
Male.....	7	5	2	1	-	-	1	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	2	-	2	-	-	1	1	-	-	-	-	-
Male.....	2	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	1	-	-	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Nebraska

All races, both sexes.....	17	15	9	5	6	11	4	2	2	-	-	-
Male.....	13	11	7	3	5	9	3	1	1	-	-	-
Female.....	4	4	2	2	1	2	1	1	1	-	-	-
White, both sexes.....	16	15	9	5	6	11	4	2	2	-	-	-
Male.....	12	11	7	3	5	9	3	1	1	-	-	-
Female.....	4	4	2	2	1	2	1	1	1	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	51	57	34	31	15	26	6	7	3	-	-	-
Male.....	38	45	27	23	14	21	4	6	2	-	-	-
Female.....	13	12	7	8	1	5	2	1	1	-	-	-
White, both sexes.....	50	51	30	28	15	25	5	7	3	-	-	-
Male.....	37	42	24	22	14	20	4	6	2	-	-	-
Female.....	13	9	6	6	1	5	1	1	1	-	-	-
Black, both sexes.....	1	3	-	1	-	-	-	-	-	-	-	-
Male.....	1	2	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	4	2	-	1	1	-	-	-	-	-
Male.....	-	1	3	1	-	1	-	-	-	-	-	-
Female.....	-	1	1	1	-	-	1	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

New Mexico

All races, both sexes.....	45	28	25	19	7	14	4	7	4	1	-	-
Male.....	29	23	21	16	7	13	4	6	4	1	-	-
Female.....	16	5	4	3	-	1	-	1	-	-	-	-
White, both sexes.....	41	28	23	18	6	14	4	7	4	1	-	-
Male.....	27	23	19	15	6	13	4	6	4	1	-	-
Female.....	14	5	4	3	-	1	-	1	-	-	-	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	-	1	1	1	-	-	-	-	-	-	-
Male.....	2	-	1	1	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	149	124	85	66	46	55	39	25	6	2	-	-
Male.....	113	85	61	48	29	43	31	18	5	2	-	-
Female.....	36	39	24	18	17	12	8	7	1	-	-	-
White, both sexes.....	131	112	76	65	40	51	36	25	5	2	-	-
Male.....	101	79	55	47	25	40	30	18	4	2	-	-
Female.....	30	33	21	18	15	11	6	7	1	-	-	-
Black, both sexes.....	12	8	4	-	3	1	1	-	-	-	-	-
Male.....	7	4	2	-	2	1	1	-	-	-	-	-
Female.....	5	4	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	4	5	1	3	3	2	-	1	-	-	-
Male.....	5	2	4	1	2	2	-	-	1	-	-	-
Female.....	1	2	1	-	1	1	2	-	-	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Vermont

All races, both sexes.....	13	6	8	6	3	2	3	1	-	-	-	-
Male.....	9	5	5	5	3	1	3	1	-	-	-	-
Female.....	4	1	3	1	-	1	-	-	-	-	-	-
White, both sexes.....	13	6	8	6	3	2	3	1	-	-	-	-
Male.....	9	5	5	5	3	1	3	1	-	-	-	-
Female.....	4	1	3	1	-	1	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	92	58	63	34	31	36	27	12	6	2	-	-
Male.....	61	40	52	25	24	32	25	11	4	2	-	-
Female.....	31	18	11	9	7	4	2	1	2	-	-	-
White, both sexes.....	83	54	57	33	27	32	26	12	5	2	-	-
Male.....	52	37	48	24	20	29	24	11	3	2	-	-
Female.....	31	17	9	9	7	3	2	1	2	-	-	-
Black, both sexes.....	6	2	6	-	4	3	1	-	-	-	-	-
Male.....	6	2	4	-	4	3	1	-	-	-	-	-
Female.....	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	2	-	1	-	1	-	-	1	-	-	-
Male.....	3	1	-	1	-	-	-	-	1	-	-	-
Female.....	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Washington

All races, both sexes.....	89	71	50	35	30	26	30	24	15	1	-	-
Male.....	67	54	40	25	25	23	26	20	14	1	-	-
Female.....	22	17	10	10	5	3	4	4	1	-	-	-
White, both sexes.....	83	69	49	33	30	25	29	24	15	1	-	-
Male.....	63	53	39	23	25	22	25	20	14	1	-	-
Female.....	20	16	10	10	5	3	4	4	1	-	-	-
Black, both sexes.....	1	-	-	1	-	-	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	-	-	1	-	-	-	-	-	-
Male.....	-	1	1	-	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	1	-	1	-	-	1	-	-	-	-	-
Male.....	3	-	-	1	-	-	1	-	-	-	-	-
Female.....	1	1	-	-	-	-	-	-	-	-	-	-

West Virginia

All races, both sexes.....	25	27	17	14	10	7	13	7	1	-	-	-
Male.....	19	18	17	10	9	6	13	7	-	-	-	-
Female.....	6	9	-	4	1	1	-	-	1	-	-	-
White, both sexes.....	24	27	17	14	9	6	13	7	1	-	-	-
Male.....	19	18	17	10	8	6	13	7	-	-	-	-
Female.....	5	9	-	4	1	-	-	-	1	-	-	-
Black, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Arkansas

All races, both sexes.....	44	...	...	-	-	3	3	3	1	2	11	7
Male.....	21	...	...	-	-	1	2	1	-	-	6	4
Female.....	23	...	...	-	-	2	1	2	1	2	5	3
White, both sexes.....	40	...	...	-	-	3	3	3	1	2	8	7
Male.....	19	...	...	-	-	1	2	1	-	-	4	4
Female.....	21	...	...	-	-	2	1	2	1	2	4	3
Black, both sexes.....	3	...	...	-	-	-	-	-	-	-	2	-
Male.....	1	...	...	-	-	-	-	-	-	-	1	-
Female.....	2	...	...	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	1	...	...	-	-	-	-	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	629	...	...	-	-	9	26	32	37	67	72	101
Male.....	300	...	...	-	-	5	14	20	23	41	40	43
Female.....	329	...	...	-	-	4	12	12	14	26	32	58
White, both sexes.....	562	...	...	-	-	9	17	30	34	56	58	95
Male.....	271	...	...	-	-	5	11	19	20	36	33	42
Female.....	291	...	...	-	-	4	6	11	14	20	25	53
Black, both sexes.....	27	...	...	-	-	-	2	1	2	4	7	2
Male.....	11	...	...	-	-	-	1	-	2	2	2	1
Female.....	16	...	...	-	-	-	1	1	-	2	5	1
American Indian 1/, both sexes.....	6	...	...	-	-	-	1	-	-	1	-	-
Male.....	2	...	...	-	-	-	1	-	-	-	-	-
Female.....	4	...	...	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	34	...	...	-	-	-	6	1	1	6	7	4
Male.....	16	...	...	-	-	-	1	1	1	3	5	-
Female.....	18	...	...	-	-	-	5	-	-	3	2	4





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

New Hampshire

All races, both sexes.....	31	...	...	-	-	2	1	-	1	2	2	7
Male.....	16	...	...	-	-	1	1	-	-	1	1	5
Female.....	15	...	...	-	-	1	-	-	1	1	1	2
White, both sexes.....	31	...	...	-	-	2	1	-	1	2	2	7
Male.....	16	...	...	-	-	1	1	-	-	1	1	5
Female.....	15	...	...	-	-	1	-	-	1	1	1	2
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	75	...	...	-	1	-	6	3	8	5	12	10
Male.....	35	...	...	-	-	-	3	1	4	4	5	5
Female.....	40	...	...	-	1	-	3	2	4	1	7	5
White, both sexes.....	65	...	...	-	1	-	3	3	7	4	11	9
Male.....	28	...	...	-	-	-	1	1	3	3	5	4
Female.....	37	...	...	-	1	-	2	2	4	1	6	5
Black, both sexes.....	5	...	...	-	-	-	2	-	1	1	-	1
Male.....	5	...	...	-	-	-	2	-	1	1	-	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	-	-	-	1	-	-	-	1	-
Male.....	2	...	...	-	-	-	-	-	-	-	-	-
Female.....	3	...	...	-	-	-	1	-	-	-	1	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

## New Mexico

All races, both sexes.....	67	...	...	-	-	4	2	6	1	2	7	8
Male.....	33	...	...	-	-	2	2	3	-	1	2	5
Female.....	34	...	...	-	-	2	-	3	1	1	5	3
White, both sexes.....	62	...	...	-	-	3	1	6	1	1	7	8
Male.....	31	...	...	-	-	1	1	3	-	1	2	5
Female.....	31	...	...	-	-	2	-	3	1	-	5	3
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	4	...	...	-	-	1	1	-	-	1	-	-
Male.....	2	...	...	-	-	1	1	-	-	-	-	-
Female.....	2	...	...	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	-	-

## New York

All races, both sexes.....	233	...	...	-	-	1	8	16	10	26	31	32
Male.....	108	...	...	-	-	1	5	9	5	15	19	18
Female.....	125	...	...	-	-	-	3	7	5	11	12	14
White, both sexes.....	210	...	...	-	-	-	6	13	9	23	29	29
Male.....	99	...	...	-	-	-	4	8	5	14	18	16
Female.....	111	...	...	-	-	-	2	5	4	9	11	13
Black, both sexes.....	16	...	...	-	-	1	1	1	-	3	1	3
Male.....	8	...	...	-	-	1	1	1	-	1	1	2
Female.....	8	...	...	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	6	...	...	-	-	-	1	2	1	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	5	...	...	-	-	-	1	2	1	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

South Dakota

All races, both sexes.....	8	...	...	-	-	-	-	-	1	-	-	4
Male.....	4	...	...	-	-	-	-	-	-	-	-	3
Female.....	4	...	...	-	-	-	-	-	1	-	-	1
White, both sexes.....	7	...	...	-	-	-	-	-	1	-	-	3
Male.....	4	...	...	-	-	-	-	-	-	-	-	3
Female.....	3	...	...	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	1
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	102	...	...	-	-	3	4	8	8	14	10	16
Male.....	47	...	...	-	-	2	1	3	5	7	3	7
Female.....	55	...	...	-	-	1	3	5	3	7	7	9
White, both sexes.....	99	...	...	-	-	3	3	8	8	13	10	15
Male.....	47	...	...	-	-	2	1	3	5	7	3	7
Female.....	52	...	...	-	-	1	2	5	3	6	7	8
Black, both sexes.....	2	...	...	-	-	-	-	-	-	1	-	1
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	2	...	...	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Vermont

All races, both sexes.....	18	...	...	-	2	-	-	1	-	2	4	1
Male.....	6	...	...	-	1	-	-	-	-	-	3	-
Female.....	12	...	...	-	1	-	-	1	-	2	1	1
White, both sexes.....	18	...	...	-	2	-	-	1	-	2	4	1
Male.....	6	...	...	-	1	-	-	-	-	-	3	-
Female.....	12	...	...	-	1	-	-	1	-	2	1	1
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	148	...	...	-	-	2	6	8	14	17	20	33
Male.....	51	...	...	-	-	-	4	1	6	6	9	12
Female.....	97	...	...	-	-	2	2	7	8	11	11	21
White, both sexes.....	136	...	...	-	-	2	6	5	14	13	18	31
Male.....	48	...	...	-	-	-	4	-	6	4	9	12
Female.....	88	...	...	-	-	2	2	5	8	9	9	19
Black, both sexes.....	9	...	...	-	-	-	-	3	-	1	2	2
Male.....	2	...	...	-	-	-	-	1	-	1	-	-
Female.....	7	...	...	-	-	-	-	2	-	-	2	2
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	-	-	-	3	-	-
Male.....	1	...	...	-	-	-	-	-	-	1	-	-
Female.....	2	...	...	-	-	-	-	-	-	2	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Arkansas

All races, both sexes.....	5	3	4	2	-	-	-	-	-	-	-	-
Male.....	2	1	3	1	-	-	-	-	-	-	-	-
Female.....	3	2	1	1	-	-	-	-	-	-	-	-
White, both sexes.....	5	2	4	2	-	-	-	-	-	-	-	-
Male.....	2	1	3	1	-	-	-	-	-	-	-	-
Female.....	3	1	1	1	-	-	-	-	-	-	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	96	78	48	14	8	16	11	8	6	-	-	-
Male.....	36	29	27	8	3	7	2	-	2	-	-	-
Female.....	60	49	21	6	5	9	9	8	4	-	-	-
White, both sexes.....	86	74	45	14	6	16	11	7	4	-	-	-
Male.....	32	25	26	8	3	7	2	-	2	-	-	-
Female.....	54	49	19	6	3	9	9	7	2	-	-	-
Black, both sexes.....	6	1	1	-	-	-	-	-	1	-	-	-
Male.....	2	1	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	2	-	1	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	3	1	-	2	-	-	-	1	-	-	-
Male.....	1	3	1	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	2	-	-	-	1	-	-	-

























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Arkansas

All races, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
White, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	27	...	...	-	-	3	2	-	3	1	1	6
Male.....	16	...	...	-	-	2	1	-	1	1	1	1
Female.....	11	...	...	-	-	1	1	-	2	-	-	5
White, both sexes.....	17	...	...	-	-	1	-	-	3	-	1	4
Male.....	10	...	...	-	-	1	-	-	1	-	1	1
Female.....	7	...	...	-	-	-	-	-	2	-	-	3
Black, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	1
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	...	...	-	-	2	2	-	-	1	-	1
Male.....	6	...	...	-	-	1	1	-	-	1	-	-
Female.....	3	...	...	-	-	1	1	-	-	-	-	1















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Kentucky

All races, both sexes.....	4	...	...	-	-	1	-	-	-	-	-	2
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	3	...	...	-	-	1	-	-	-	-	-	2
White, both sexes.....	4	...	...	-	-	1	-	-	-	-	-	2
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	3	...	...	-	-	1	-	-	-	-	-	2
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	2	...	...	-	-	-	-	-	-	1	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	1	-	-
White, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	1	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	1	-	-







































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Delaware

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'Male', 'Female', 'White, both sexes', 'Male', 'Female', 'Black, both sexes', 'Male', 'Female', 'American Indian 1/, both sexes', 'Male', 'Female', and 'Asian or Pacific Islander 2/, both sexes', 'Male', 'Female'.

Dist. of Columbia

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'Male', 'Female', 'White, both sexes', 'Male', 'Female', 'Black, both sexes', 'Male', 'Female', 'American Indian 1/, both sexes', 'Male', 'Female', and 'Asian or Pacific Islander 2/, both sexes', 'Male', 'Female'.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Nebraska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Alaska

All races, both sexes.....	40	...	...	-	1	6	7	9	4	3	5	4
Male.....	28	...	...	-	1	4	3	8	3	1	4	3
Female.....	12	...	...	-	-	2	4	1	1	2	1	1
White, both sexes.....	16	...	...	-	-	1	3	3	1	3	3	1
Male.....	12	...	...	-	-	1	2	3	1	1	3	-
Female.....	4	...	...	-	-	-	1	-	-	2	-	1
Black, both sexes.....	1	...	...	-	-	1	-	-	-	-	-	-
Male.....	1	...	...	-	-	1	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	22	...	...	-	1	4	4	5	3	-	2	3
Male.....	14	...	...	-	1	2	1	4	2	-	1	3
Female.....	8	...	...	-	-	2	3	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	180	...	...	-	5	16	23	23	20	19	12	21
Male.....	149	...	...	-	3	10	19	21	19	17	8	18
Female.....	31	...	...	-	2	6	4	2	1	2	4	3
White, both sexes.....	156	...	...	-	3	13	20	18	16	14	12	20
Male.....	130	...	...	-	2	9	17	16	15	12	8	18
Female.....	26	...	...	-	1	4	3	2	1	2	4	2
Black, both sexes.....	1	...	...	-	-	-	-	-	-	1	-	-
Male.....	1	...	...	-	-	-	-	-	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	18	...	...	-	1	2	3	4	4	4	-	-
Male.....	15	...	...	-	-	1	2	4	4	4	-	-
Female.....	3	...	...	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	-	1	1	-	1	-	-	-	1
Male.....	3	...	...	-	1	-	-	1	-	-	-	-
Female.....	2	...	...	-	-	1	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Arkansas

All races, both sexes.....	78	...	...	-	2	5	6	11	11	10	16	10
Male.....	66	...	...	-	2	4	4	9	6	10	14	10
Female.....	12	...	...	-	-	1	2	2	5	-	2	-
White, both sexes.....	69	...	...	-	1	5	6	9	9	9	14	9
Male.....	58	...	...	-	1	4	4	7	4	9	13	9
Female.....	11	...	...	-	-	1	2	2	5	-	1	-
Black, both sexes.....	8	...	...	-	1	-	-	2	2	1	1	1
Male.....	8	...	...	-	1	-	-	2	2	1	1	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-

California

All races, both sexes.....	958	...	...	1	8	49	101	103	92	102	106	102
Male.....	750	...	...	1	6	36	81	78	74	82	80	92
Female.....	208	...	...	-	2	13	20	25	18	20	26	10
White, both sexes.....	756	...	...	1	4	36	78	82	69	79	87	84
Male.....	609	...	...	1	4	27	63	66	60	67	64	76
Female.....	147	...	...	-	-	9	15	16	9	12	23	8
Black, both sexes.....	47	...	...	-	1	4	7	7	6	7	6	4
Male.....	36	...	...	-	1	2	5	5	5	5	6	3
Female.....	11	...	...	-	-	2	2	2	1	2	-	1
American Indian 1/, both sexes.....	10	...	...	-	-	-	2	2	1	2	2	-
Male.....	9	...	...	-	-	-	2	2	1	2	1	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	145	...	...	-	3	9	14	12	16	14	11	14
Male.....	96	...	...	-	1	7	11	5	8	8	9	13
Female.....	49	...	...	-	2	2	3	7	8	6	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Colorado

All races, both sexes.....	167	...	...	-	2	19	21	22	11	13	25	19
Male.....	141	...	...	-	2	15	20	17	10	11	22	17
Female.....	26	...	...	-	-	4	1	5	1	2	3	2
White, both sexes.....	152	...	...	-	2	17	18	19	11	10	23	19
Male.....	128	...	...	-	2	14	17	14	10	9	20	17
Female.....	24	...	...	-	-	3	1	5	1	1	3	2
Black, both sexes.....	6	...	...	-	-	1	2	1	-	-	2	-
Male.....	5	...	...	-	-	-	2	1	-	-	2	-
Female.....	1	...	...	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	4	...	...	-	-	-	1	1	-	2	-	-
Male.....	3	...	...	-	-	-	1	1	-	1	-	-
Female.....	1	...	...	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	-	-	1	-	1	-	1	-	-
Male.....	5	...	...	-	-	1	-	1	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	106	...	...	-	-	10	11	10	5	7	16	16
Male.....	86	...	...	-	-	8	10	8	4	6	12	13
Female.....	20	...	...	-	-	2	1	2	1	1	4	3
White, both sexes.....	99	...	...	-	-	9	9	9	5	7	15	14
Male.....	80	...	...	-	-	7	9	7	4	6	11	11
Female.....	19	...	...	-	-	2	-	2	1	1	4	3
Black, both sexes.....	6	...	...	-	-	1	1	1	-	-	1	2
Male.....	6	...	...	-	-	1	1	1	-	-	1	2
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Delaware

All races, both sexes.....	23	...	...	-	-	2	1	1	1	7	1	4
Male.....	20	...	...	-	-	1	1	1	1	6	1	4
Female.....	3	...	...	-	-	1	-	-	-	1	-	-
White, both sexes.....	21	...	...	-	-	2	1	-	1	6	1	4
Male.....	18	...	...	-	-	1	1	-	1	5	1	4
Female.....	3	...	...	-	-	1	-	-	-	1	-	-
Black, both sexes.....	1	...	...	-	-	-	-	-	-	1	-	-
Male.....	1	...	...	-	-	-	-	-	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Dist. of Columbia

All races, both sexes.....	9	...	...	-	-	-	-	2	1	-	-	3
Male.....	5	...	...	-	-	-	-	1	-	-	-	2
Female.....	4	...	...	-	-	-	-	1	1	-	-	1
White, both sexes.....	5	...	...	-	-	-	-	1	-	-	-	1
Male.....	4	...	...	-	-	-	-	1	-	-	-	1
Female.....	1	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	3	...	...	-	-	-	-	-	1	-	-	2
Male.....	1	...	...	-	-	-	-	-	-	-	-	1
Female.....	2	...	...	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Florida

All races, both sexes.....	564	...	...	-	3	33	57	44	53	59	45	79
Male.....	463	...	...	-	-	29	47	38	42	53	37	64
Female.....	101	...	...	-	3	4	10	6	11	6	8	15
White, both sexes.....	513	...	...	-	1	31	49	39	43	51	43	75
Male.....	421	...	...	-	-	27	40	34	32	47	36	60
Female.....	92	...	...	-	1	4	9	5	11	4	7	15
Black, both sexes.....	36	...	...	-	2	-	5	4	8	4	1	3
Male.....	30	...	...	-	-	-	5	3	8	3	1	3
Female.....	6	...	...	-	2	-	-	1	-	1	-	-
American Indian 1/, both sexes.....	2	...	...	-	-	-	-	-	1	-	-	-
Male.....	2	...	...	-	-	-	-	-	1	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	13	...	...	-	-	2	3	1	1	4	1	1
Male.....	10	...	...	-	-	2	2	1	1	3	-	1
Female.....	3	...	...	-	-	-	1	-	-	1	1	-

Georgia

All races, both sexes.....	173	...	...	-	3	10	20	15	25	23	24	24
Male.....	145	...	...	-	2	7	15	11	20	20	22	24
Female.....	28	...	...	-	1	3	5	4	5	3	2	-
White, both sexes.....	148	...	...	-	2	8	16	12	23	19	19	23
Male.....	125	...	...	-	2	5	12	9	19	17	17	23
Female.....	23	...	...	-	-	3	4	3	4	2	2	-
Black, both sexes.....	20	...	...	-	-	2	4	2	1	3	5	-
Male.....	17	...	...	-	-	2	3	1	1	2	5	-
Female.....	3	...	...	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	-	1	-	-	1	1	1	-	1
Male.....	3	...	...	-	-	-	-	1	-	1	-	1
Female.....	2	...	...	-	1	-	-	-	1	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Maine

All races, both sexes.....	45	...	...	-	-	8	8	6	2	4	3	5
Male.....	38	...	...	-	-	5	7	6	2	3	3	5
Female.....	7	...	...	-	-	3	1	-	-	1	-	-
White, both sexes.....	42	...	...	-	-	7	8	6	2	4	2	5
Male.....	35	...	...	-	-	4	7	6	2	3	2	5
Female.....	7	...	...	-	-	3	1	-	-	1	-	-
Black, both sexes.....	1	...	...	-	-	1	-	-	-	-	-	-
Male.....	1	...	...	-	-	1	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	1	...	...	-	-	-	-	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	162	...	...	-	1	14	19	23	15	18	21	16
Male.....	127	...	...	-	-	11	14	19	10	16	16	14
Female.....	35	...	...	-	1	3	5	4	5	2	5	2
White, both sexes.....	120	...	...	-	-	11	14	17	12	13	15	14
Male.....	99	...	...	-	-	9	11	16	8	11	12	13
Female.....	21	...	...	-	-	2	3	1	4	2	3	1
Black, both sexes.....	29	...	...	-	1	1	4	4	3	4	4	2
Male.....	22	...	...	-	-	1	3	2	2	4	4	1
Female.....	7	...	...	-	1	-	1	2	1	-	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	13	...	...	-	-	2	1	2	-	1	2	-
Male.....	6	...	...	-	-	1	-	1	-	1	-	-
Female.....	7	...	...	-	-	1	1	1	-	-	2	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Nebraska

All races, both sexes.....	51	...	...	-	3	5	5	8	3	3	5	5
Male.....	36	...	...	-	-	3	4	8	2	2	5	1
Female.....	15	...	...	-	3	2	1	-	1	1	-	4
White, both sexes.....	45	...	...	-	3	5	2	7	3	3	4	5
Male.....	30	...	...	-	-	3	1	7	2	2	4	1
Female.....	15	...	...	-	3	2	1	-	1	1	-	4
Black, both sexes.....	4	...	...	-	-	-	2	-	-	-	1	-
Male.....	4	...	...	-	-	-	2	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	...	...	-	-	-	1	1	-	-	-	-
Male.....	2	...	...	-	-	-	1	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	68	...	...	-	4	3	6	6	4	11	6	9
Male.....	59	...	...	-	2	2	6	4	4	10	5	9
Female.....	9	...	...	-	2	1	-	2	-	1	1	-
White, both sexes.....	57	...	...	-	1	2	5	4	4	9	6	9
Male.....	51	...	...	-	-	1	5	3	4	8	5	9
Female.....	6	...	...	-	1	1	-	1	-	1	1	-
Black, both sexes.....	5	...	...	-	2	1	-	1	-	1	-	-
Male.....	5	...	...	-	2	1	-	1	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	-	1	-	1	-	-	1	-	-
Male.....	3	...	...	-	-	-	1	-	-	1	-	-
Female.....	2	...	...	-	1	-	-	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

New Hampshire

All races, both sexes.....	30	...	...	-	1	1	3	2	6	3	3	3
Male.....	22	...	...	-	1	-	1	1	3	3	2	3
Female.....	8	...	...	-	-	1	2	1	3	-	1	-
White, both sexes.....	28	...	...	-	1	1	3	2	5	3	3	3
Male.....	21	...	...	-	1	-	1	1	3	3	2	3
Female.....	7	...	...	-	-	1	2	1	2	-	1	-
Black, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	1	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	1	-	-	-

New Jersey

All races, both sexes.....	228	...	...	-	1	15	24	22	22	27	25	22
Male.....	182	...	...	-	-	14	17	21	19	22	19	16
Female.....	46	...	...	-	1	1	7	1	3	5	6	6
White, both sexes.....	201	...	...	-	1	14	21	20	19	25	22	21
Male.....	165	...	...	-	-	13	15	19	17	20	17	16
Female.....	36	...	...	-	1	1	6	1	2	5	5	5
Black, both sexes.....	11	...	...	-	-	1	2	2	-	1	2	-
Male.....	9	...	...	-	-	1	2	2	-	1	2	-
Female.....	2	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	16	...	...	-	-	-	1	-	3	1	1	1
Male.....	8	...	...	-	-	-	-	-	2	1	-	-
Female.....	8	...	...	-	-	-	1	-	1	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

New Mexico

All races, both sexes.....	112	...	...	-	5	10	7	7	14	12	23	11
Male.....	102	...	...	-	5	7	6	7	12	12	20	11
Female.....	10	...	...	-	-	3	1	-	2	-	3	-
White, both sexes.....	78	...	...	-	2	6	1	4	8	8	18	10
Male.....	69	...	...	-	2	3	1	4	6	8	15	10
Female.....	9	...	...	-	-	3	-	-	2	-	3	-
Black, both sexes.....	1	...	...	-	1	-	-	-	-	-	-	-
Male.....	1	...	...	-	1	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	32	...	...	-	2	4	6	3	6	3	5	1
Male.....	31	...	...	-	2	4	5	3	6	3	5	1
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	1	-	-
Male.....	1	...	...	-	-	-	-	-	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	411	...	...	-	4	26	43	49	37	30	46	36
Male.....	320	...	...	-	2	20	38	36	26	24	39	31
Female.....	91	...	...	-	2	6	5	13	11	6	7	5
White, both sexes.....	332	...	...	-	3	18	34	35	28	24	39	32
Male.....	268	...	...	-	2	13	31	28	23	19	35	27
Female.....	64	...	...	-	1	5	3	7	5	5	4	5
Black, both sexes.....	33	...	...	-	1	5	4	9	5	1	3	2
Male.....	23	...	...	-	-	4	3	7	3	-	2	2
Female.....	10	...	...	-	1	1	1	2	2	1	1	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	46	...	...	-	-	3	5	5	4	5	4	2
Male.....	29	...	...	-	-	3	4	1	-	5	2	2
Female.....	17	...	...	-	-	-	1	4	4	-	2	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Oregon

All races, both sexes.....	98	...	...	-	-	6	11	12	12	9	13	11
Male.....	75	...	...	-	-	3	5	10	9	7	12	9
Female.....	23	...	...	-	-	3	6	2	3	2	1	2
White, both sexes.....	92	...	...	-	-	5	9	10	12	9	12	11
Male.....	72	...	...	-	-	3	4	9	9	7	11	9
Female.....	20	...	...	-	-	2	5	1	3	2	1	2
Black, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	...	...	-	-	1	-	-	-	-	1	-
Male.....	1	...	...	-	-	-	-	-	-	-	1	-
Female.....	1	...	...	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	1	2	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	2	...	...	-	-	-	1	1	-	-	-	-

Pennsylvania

All races, both sexes.....	379	...	...	-	7	20	40	42	35	42	41	44
Male.....	321	...	...	-	7	15	33	36	32	38	38	33
Female.....	58	...	...	-	-	5	7	6	3	4	3	11
White, both sexes.....	341	...	...	-	7	17	33	38	29	38	38	40
Male.....	292	...	...	-	7	14	30	33	27	34	35	30
Female.....	49	...	...	-	-	3	3	5	2	4	3	10
Black, both sexes.....	28	...	...	-	-	3	5	1	5	4	3	3
Male.....	24	...	...	-	-	1	3	1	5	4	3	3
Female.....	4	...	...	-	-	2	2	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	...	...	-	-	-	2	2	1	-	-	1
Male.....	4	...	...	-	-	-	-	1	-	-	-	-
Female.....	5	...	...	-	-	-	2	1	1	-	-	1





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Texas

All races, both sexes.....	585	...	...	-	7	52	71	77	75	74	69	66
Male.....	471	...	...	-	5	40	62	58	64	57	57	58
Female.....	114	...	...	-	2	12	9	19	11	17	12	8
White, both sexes.....	537	...	...	-	5	50	65	73	68	64	62	63
Male.....	432	...	...	-	4	38	56	55	57	49	52	55
Female.....	105	...	...	-	1	12	9	18	11	15	10	8
Black, both sexes.....	27	...	...	-	2	1	4	2	5	5	3	3
Male.....	25	...	...	-	1	1	4	2	5	4	3	3
Female.....	2	...	...	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	1	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	20	...	...	-	-	1	2	2	2	5	3	-
Male.....	14	...	...	-	-	1	2	1	2	4	2	-
Female.....	6	...	...	-	-	-	-	1	-	1	1	-

Utah

All races, both sexes.....	85	...	...	-	1	11	23	10	8	8	5	7
Male.....	72	...	...	-	-	11	19	9	6	6	4	6
Female.....	13	...	...	-	1	-	4	1	2	2	1	1
White, both sexes.....	76	...	...	-	1	9	19	10	6	8	5	7
Male.....	64	...	...	-	-	9	16	9	4	6	4	6
Female.....	12	...	...	-	1	-	3	1	2	2	1	1
Black, both sexes.....	1	...	...	-	-	-	-	-	1	-	-	-
Male.....	1	...	...	-	-	-	-	-	1	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	6	...	...	-	-	1	3	-	1	-	-	-
Male.....	6	...	...	-	-	1	3	-	1	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	1	1	-	-	-	-	-
Male.....	1	...	...	-	-	1	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	1	-	-	-	-	-



## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

## Vermont

All races, both sexes.....	26	...	...	-	-	2	3	4	4	1	2	1
Male.....	23	...	...	-	-	2	2	4	4	1	2	1
Female.....	3	...	...	-	-	-	1	-	-	-	-	-
White, both sexes.....	25	...	...	-	-	2	2	4	4	1	2	1
Male.....	22	...	...	-	-	2	1	4	4	1	2	1
Female.....	3	...	...	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

## Virginia

All races, both sexes.....	164	...	...	-	3	9	17	19	21	19	20	18
Male.....	131	...	...	-	3	7	16	14	16	18	15	14
Female.....	33	...	...	-	-	2	1	5	5	1	5	4
White, both sexes.....	127	...	...	-	2	8	10	15	18	15	18	12
Male.....	102	...	...	-	2	6	10	12	14	15	13	9
Female.....	25	...	...	-	-	2	-	3	4	-	5	3
Black, both sexes.....	26	...	...	-	1	1	6	3	2	4	1	5
Male.....	23	...	...	-	1	1	5	2	2	3	1	5
Female.....	3	...	...	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	10	...	...	-	-	-	-	1	1	-	1	1
Male.....	5	...	...	-	-	-	-	-	-	-	1	-
Female.....	5	...	...	-	-	-	-	1	1	-	-	1









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Arkansas

All races, both sexes.....	3	1	1	-	-	1	1	-	-	-	-	-
Male.....	3	1	1	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	3	1	1	-	-	1	1	-	-	-	-	-
Male.....	3	1	1	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	101	60	39	24	14	20	19	12	4	1	-	-
Male.....	83	47	33	18	8	10	12	6	2	1	-	-
Female.....	18	13	6	6	6	10	7	6	2	-	-	-
White, both sexes.....	88	54	33	17	10	10	12	7	4	1	-	-
Male.....	72	42	30	12	6	6	8	2	2	1	-	-
Female.....	16	12	3	5	4	4	4	5	2	-	-	-
Black, both sexes.....	1	1	1	1	1	-	-	-	-	-	-	-
Male.....	1	1	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	12	4	5	6	3	10	7	5	-	-	-	-
Male.....	10	3	3	5	1	4	4	4	-	-	-	-
Female.....	2	1	2	1	2	6	3	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Maine

All races, both sexes.....	3	3	-	1	-	1	-	1	-	-	-	-
Male.....	3	1	-	1	-	1	-	1	-	-	-	-
Female.....	-	2	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	2	3	-	1	-	1	-	1	-	-	-	-
Male.....	2	1	-	1	-	1	-	1	-	-	-	-
Female.....	-	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	15	5	8	2	2	2	1	-	-	-	-	-
Male.....	14	4	6	2	-	1	-	-	-	-	-	-
Female.....	1	1	2	-	2	1	1	-	-	-	-	-
White, both sexes.....	10	4	5	2	2	1	-	-	-	-	-	-
Male.....	9	3	5	2	-	-	-	-	-	-	-	-
Female.....	1	1	-	-	2	1	-	-	-	-	-	-
Black, both sexes.....	3	1	2	-	-	-	-	-	-	-	-	-
Male.....	3	1	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	2	-	1	-	-	1	1	-	-	-	-	-
Male.....	2	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	1	-	-	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Nebraska

All races, both sexes.....	3	3	-	1	3	3	-	-	1	-	-	-
Male.....	3	3	-	-	3	2	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	-	1	-	-	-
White, both sexes.....	2	3	-	1	3	3	-	-	1	-	-	-
Male.....	2	3	-	-	3	2	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	-	1	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	6	6	2	-	-	2	2	1	-	-	-	-
Male.....	6	6	2	-	-	2	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes.....	6	6	1	-	-	2	1	1	-	-	-	-
Male.....	6	6	1	-	-	2	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

New Mexico

All races, both sexes.....	9	4	7	1	1	-	1	-	-	-	-	-
Male.....	8	4	7	1	1	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	8	4	7	1	-	-	1	-	-	-	-	-
Male.....	7	4	7	1	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	43	22	22	14	8	12	12	5	2	-	-	-
Male.....	33	16	17	11	6	8	9	2	2	-	-	-
Female.....	10	6	5	3	2	4	3	3	-	-	-	-
White, both sexes.....	36	18	19	14	7	9	10	5	1	-	-	-
Male.....	28	14	14	11	5	6	9	2	1	-	-	-
Female.....	8	4	5	3	2	3	1	3	-	-	-	-
Black, both sexes.....	1	2	-	-	-	-	-	-	-	-	-	-
Male.....	-	2	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	2	3	-	1	3	2	-	1	-	-	-
Male.....	5	-	3	-	1	2	-	-	1	-	-	-
Female.....	1	2	-	-	-	1	2	-	-	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Vermont

All races, both sexes.....	5	2	1	1	-	-	-	-	-	-	-	-
Male.....	3	2	1	1	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	5	2	1	1	-	-	-	-	-	-	-	-
Male.....	3	2	1	1	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	10	8	5	5	2	3	2	1	2	-	-	-
Male.....	8	5	5	4	1	2	1	1	1	-	-	-
Female.....	2	3	-	1	1	1	1	-	1	-	-	-
White, both sexes.....	6	5	5	5	2	2	2	1	1	-	-	-
Male.....	4	3	5	4	1	2	1	1	-	-	-	-
Female.....	2	2	-	1	1	-	1	-	1	-	-	-
Black, both sexes.....	2	1	-	-	-	-	-	-	-	-	-	-
Male.....	2	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	-	-	-	1	-	-	1	-	-	-
Male.....	2	1	-	-	-	-	-	-	1	-	-	-
Female.....	-	1	-	-	-	1	-	-	-	-	-	-









## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Intentional self-harm (suicide) by discharge of firearms (X72-X74)

## Arkansas

All races, both sexes.....	244	...	...	-	-	8	12	23	17	20	26	23
Male.....	206	...	...	-	-	7	10	23	14	16	17	15
Female.....	38	...	...	-	-	1	2	-	3	4	9	8
White, both sexes.....	229	...	...	-	-	6	9	20	15	18	25	23
Male.....	191	...	...	-	-	5	7	20	12	14	16	15
Female.....	38	...	...	-	-	1	2	-	3	4	9	8
Black, both sexes.....	13	...	...	-	-	2	3	2	2	2	1	-
Male.....	13	...	...	-	-	2	3	2	2	2	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

## California

All races, both sexes.....	1,422	...	...	-	6	37	89	83	78	81	111	144
Male.....	1,257	...	...	-	5	36	80	80	72	75	85	125
Female.....	165	...	...	-	1	1	9	3	6	6	26	19
White, both sexes.....	1,277	...	...	-	6	27	66	66	66	69	102	128
Male.....	1,121	...	...	-	5	26	59	64	61	64	77	109
Female.....	156	...	...	-	1	1	7	2	5	5	25	19
Black, both sexes.....	62	...	...	-	-	5	10	6	8	5	4	9
Male.....	60	...	...	-	-	5	9	6	7	5	4	9
Female.....	2	...	...	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	6	...	...	-	-	-	1	1	1	-	-	1
Male.....	6	...	...	-	-	-	1	1	1	-	-	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	77	...	...	-	-	5	12	10	3	7	5	6
Male.....	70	...	...	-	-	5	11	9	3	6	4	6
Female.....	7	...	...	-	-	-	1	1	-	1	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Colorado

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Kentucky

All races, both sexes.....	422	...	...	-	2	15	22	40	37	37	46	46
Male.....	359	...	...	-	2	11	21	35	32	28	36	38
Female.....	63	...	...	-	-	4	1	5	5	9	10	8
White, both sexes.....	406	...	...	-	2	14	21	38	33	36	44	45
Male.....	345	...	...	-	2	10	20	33	29	27	34	37
Female.....	61	...	...	-	-	4	1	5	4	9	10	8
Black, both sexes.....	15	...	...	-	-	1	1	1	4	1	2	1
Male.....	13	...	...	-	-	1	1	1	3	1	2	1
Female.....	2	...	...	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	335	...	...	-	5	16	30	35	26	31	28	35
Male.....	286	...	...	-	5	16	29	28	22	24	23	27
Female.....	49	...	...	-	-	-	1	7	4	7	5	8
White, both sexes.....	288	...	...	-	4	12	23	28	22	26	23	31
Male.....	244	...	...	-	4	12	22	23	18	21	18	24
Female.....	44	...	...	-	-	-	1	5	4	5	5	7
Black, both sexes.....	45	...	...	-	1	4	7	6	4	5	5	4
Male.....	41	...	...	-	1	4	7	5	4	3	5	3
Female.....	4	...	...	-	-	-	-	1	-	2	-	1
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	1	-	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Intentional self-harm (suicide) by discharge of firearms (X72-X74)

## Massachusetts

All races, both sexes.....	114	...	...	-	-	4	7	7	5	10	16	10
Male.....	110	...	...	-	-	4	7	7	5	8	16	10
Female.....	4	...	...	-	-	-	-	-	-	2	-	-
White, both sexes.....	107	...	...	-	-	4	4	5	4	9	16	10
Male.....	103	...	...	-	-	4	4	5	4	7	16	10
Female.....	4	...	...	-	-	-	-	-	-	2	-	-
Black, both sexes.....	5	...	...	-	-	-	3	1	-	1	-	-
Male.....	5	...	...	-	-	-	3	1	-	1	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	-	-	-	-	1	1	-	-	-
Male.....	2	...	...	-	-	-	-	1	1	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

## Michigan

All races, both sexes.....	558	...	...	-	-	23	30	39	35	41	47	75
Male.....	508	...	...	-	-	21	27	35	33	36	43	66
Female.....	50	...	...	-	-	2	3	4	2	5	4	9
White, both sexes.....	515	...	...	-	-	19	27	33	33	37	42	70
Male.....	471	...	...	-	-	18	24	30	31	33	38	61
Female.....	44	...	...	-	-	1	3	3	2	4	4	9
Black, both sexes.....	37	...	...	-	-	1	3	6	2	4	4	4
Male.....	32	...	...	-	-	1	3	5	2	3	4	4
Female.....	5	...	...	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes.....	3	...	...	-	-	1	-	-	-	-	1	1
Male.....	3	...	...	-	-	1	-	-	-	-	1	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	2	-	-	-	-	-	-
Male.....	2	...	...	-	-	1	-	-	-	-	-	-
Female.....	1	...	...	-	-	1	-	-	-	-	-	-

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Vermont

All races, both sexes.....	43	...	...	-	-	1	4	3	3	1	5	3
Male.....	39	...	...	-	-	1	4	3	3	1	4	2
Female.....	4	...	...	-	-	-	-	-	-	-	1	1
White, both sexes.....	43	...	...	-	-	1	4	3	3	1	5	3
Male.....	39	...	...	-	-	1	4	3	3	1	4	2
Female.....	4	...	...	-	-	-	-	-	-	-	1	1
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	487	...	...	-	1	18	33	27	24	42	47	51
Male.....	411	...	...	-	1	17	29	24	21	28	37	40
Female.....	76	...	...	-	-	1	4	3	3	14	10	11
White, both sexes.....	431	...	...	-	1	14	25	20	22	36	45	42
Male.....	365	...	...	-	1	13	22	19	19	24	35	35
Female.....	66	...	...	-	-	1	3	1	3	12	10	7
Black, both sexes.....	45	...	...	-	-	4	5	6	2	4	-	8
Male.....	37	...	...	-	-	4	4	4	2	3	-	5
Female.....	8	...	...	-	-	-	1	2	-	1	-	3
American Indian 1/, both sexes.....	2	...	...	-	-	-	1	-	-	-	1	-
Male.....	2	...	...	-	-	-	1	-	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	...	...	-	-	-	2	1	-	2	1	1
Male.....	7	...	...	-	-	-	2	1	-	1	1	-
Female.....	2	...	...	-	-	-	-	-	-	1	-	1











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Arkansas

All races, both sexes.....	24	23	15	16	7	15	9	5	1	-	-	-
Male.....	22	18	13	15	7	15	9	4	1	-	-	-
Female.....	2	5	2	1	-	-	-	1	-	-	-	-
White, both sexes.....	24	23	14	15	7	15	9	5	1	-	-	-
Male.....	22	18	12	14	7	15	9	4	1	-	-	-
Female.....	2	5	2	1	-	-	-	1	-	-	-	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	162	142	117	87	56	98	65	42	21	1	1	1
Male.....	140	118	94	81	50	91	63	38	21	1	1	1
Female.....	22	24	23	6	6	7	2	4	-	-	-	-
White, both sexes.....	149	131	109	82	53	94	64	42	21	1	-	1
Male.....	127	109	86	76	48	87	62	38	21	1	-	1
Female.....	22	22	23	6	5	7	2	4	-	-	-	-
Black, both sexes.....	4	3	3	3	-	2	-	-	-	-	-	-
Male.....	4	3	3	3	-	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	1	-	-	-	-	-	-	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	8	4	2	3	2	1	-	-	-	1	-
Male.....	8	6	4	2	2	2	1	-	-	-	1	-
Female.....	-	2	-	-	1	-	-	-	-	-	-	-















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Washington

All races, both sexes.....	35	37	32	24	19	21	22	18	11	1	-	-
Male.....	33	32	28	21	19	20	21	18	11	1	-	-
Female.....	2	5	4	3	-	1	1	-	-	-	-	-
White, both sexes.....	34	37	31	23	19	20	21	18	11	1	-	-
Male.....	32	32	27	20	19	19	20	18	11	1	-	-
Female.....	2	5	4	3	-	1	1	-	-	-	-	-
Black, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	1	-	-	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

West Virginia

All races, both sexes.....	15	19	14	12	8	7	12	7	1	-	-	-
Male.....	15	16	14	9	8	6	12	7	-	-	-	-
Female.....	-	3	-	3	-	1	-	-	1	-	-	-
White, both sexes.....	15	19	14	12	7	6	12	7	1	-	-	-
Male.....	15	16	14	9	7	6	12	7	-	-	-	-
Female.....	-	3	-	3	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Arkansas

All races, both sexes.....	2	...	...	-	-	-	-	-	-	-	-	1
Male.....	2	...	...	-	-	-	-	-	-	-	-	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
White, both sexes.....	2	...	...	-	-	-	-	-	-	-	-	1
Male.....	2	...	...	-	-	-	-	-	-	-	-	1
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	155	...	...	-	-	6	6	21	11	18	27	19
Male.....	125	...	...	-	-	6	4	16	10	15	24	15
Female.....	30	...	...	-	-	-	2	5	1	3	3	4
White, both sexes.....	121	...	...	-	-	5	4	15	8	14	24	17
Male.....	98	...	...	-	-	5	3	12	7	12	21	13
Female.....	23	...	...	-	-	-	1	3	1	2	3	4
Black, both sexes.....	13	...	...	-	-	-	-	4	2	1	1	-
Male.....	11	...	...	-	-	-	-	2	2	1	1	-
Female.....	2	...	...	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	21	...	...	-	-	1	2	2	1	3	2	2
Male.....	16	...	...	-	-	1	1	2	1	2	2	2
Female.....	5	...	...	-	-	-	1	-	-	1	-	-



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Massachusetts

All races, both sexes.....	12	...	...	-	-	1	1	2	1	1	2	1
Male.....	5	...	...	-	-	-	-	2	1	1	-	-
Female.....	7	...	...	-	-	1	1	-	-	-	2	1
White, both sexes.....	11	...	...	-	-	1	1	1	1	1	2	1
Male.....	4	...	...	-	-	-	-	1	1	1	-	-
Female.....	7	...	...	-	-	1	1	-	-	-	2	1
Black, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Michigan

All races, both sexes.....	15	...	...	-	-	1	3	1	1	5	1	2
Male.....	13	...	...	-	-	1	2	1	1	4	1	2
Female.....	2	...	...	-	-	-	1	-	-	1	-	-
White, both sexes.....	13	...	...	-	-	1	3	1	1	4	1	2
Male.....	12	...	...	-	-	1	2	1	1	4	1	2
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
Black, both sexes.....	1	...	...	-	-	-	-	-	-	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	-	-	-	-	-	-	1	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	1	...	...	-	-	-	-	-	-	1	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

New Mexico

All races, both sexes.....	2	...	...	-	-	-	-	1	-	-	-	-
Male.....	2	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
White, both sexes.....	2	...	...	-	-	-	-	1	-	-	-	-
Male.....	2	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	111	...	...	-	-	4	6	8	6	7	19	10
Male.....	77	...	...	-	-	4	4	6	4	4	14	7
Female.....	34	...	...	-	-	-	2	2	2	3	5	3
White, both sexes.....	86	...	...	-	-	4	4	6	2	3	15	9
Male.....	59	...	...	-	-	4	3	4	1	-	11	6
Female.....	27	...	...	-	-	-	1	2	1	3	4	3
Black, both sexes.....	17	...	...	-	-	-	1	2	2	2	3	1
Male.....	12	...	...	-	-	-	-	2	2	2	2	1
Female.....	5	...	...	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	-	1	-	-	-
Male.....	1	...	...	-	-	-	-	-	1	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	...	...	-	-	-	1	-	1	2	1	-
Male.....	5	...	...	-	-	-	1	-	-	2	1	-
Female.....	2	...	...	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by jumping from a high place (X80)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Arkansas

All races, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	16	13	2	6	2	3	3	1	-	1	-	-
Male.....	13	10	-	5	1	1	3	1	-	1	-	-
Female.....	3	3	2	1	1	2	-	-	-	-	-	-
White, both sexes.....	10	10	2	5	2	2	2	-	-	1	-	-
Male.....	8	7	-	5	1	1	2	-	-	1	-	-
Female.....	2	3	2	-	1	1	-	-	-	-	-	-
Black, both sexes.....	3	2	-	-	-	-	-	-	-	-	-	-
Male.....	3	2	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	1	-	1	-	1	1	1	-	-	-	-
Male.....	2	1	-	-	-	-	1	1	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Massachusetts

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by jumping from a high place (X80)

New Mexico

All races, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	11	7	11	8	7	1	1	2	2	1	-	-
Male.....	10	3	7	5	4	1	1	1	1	1	-	-
Female.....	1	4	4	3	3	-	-	1	1	-	-	-
White, both sexes.....	8	6	9	8	5	1	1	2	2	1	-	-
Male.....	8	3	6	5	3	1	1	1	1	1	-	-
Female.....	-	3	3	3	2	-	-	1	1	-	-	-
Black, both sexes.....	3	1	2	-	-	-	-	-	-	-	-	-
Male.....	2	-	1	-	-	-	-	-	-	-	-	-
Female.....	1	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intentional self-harm (suicide) by jumping from a high place (X80)

Vermont

Table with 13 columns (Age groups) and 13 rows (Race categories) for Vermont. All values are dashes (-).

Virginia

Table with 13 columns (Age groups) and 13 rows (Race categories) for Virginia. Values include 1 and 2 in the 80-84 and 85-89 age groups for 'All races, both sexes' and 'White, both sexes'.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (\*U03,X71,X75-X79,X81-X84,Y87.0)

Arkansas

All races, both sexes.....	18	...	...	-	-	-	2	1	1	2	2	5
Male.....	14	...	...	-	-	-	2	1	1	2	1	5
Female.....	4	...	...	-	-	-	-	-	-	-	1	-
White, both sexes.....	17	...	...	-	-	-	1	1	1	2	2	5
Male.....	13	...	...	-	-	-	1	1	1	2	1	5
Female.....	4	...	...	-	-	-	-	-	-	-	1	-
Black, both sexes.....	1	...	...	-	-	-	1	-	-	-	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	319	...	...	-	-	16	27	19	23	29	26	47
Male.....	245	...	...	-	-	15	25	18	16	20	20	38
Female.....	74	...	...	-	-	1	2	1	7	9	6	9
White, both sexes.....	265	...	...	-	-	14	19	14	17	21	25	41
Male.....	206	...	...	-	-	13	17	13	12	15	20	33
Female.....	59	...	...	-	-	1	2	1	5	6	5	8
Black, both sexes.....	15	...	...	-	-	-	4	3	1	3	-	2
Male.....	11	...	...	-	-	-	4	3	1	1	-	2
Female.....	4	...	...	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes.....	1	...	...	-	-	-	-	1	-	-	-	-
Male.....	1	...	...	-	-	-	-	1	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	38	...	...	-	-	2	4	1	5	5	1	4
Male.....	27	...	...	-	-	2	4	1	3	4	-	3
Female.....	11	...	...	-	-	-	-	-	2	1	1	1





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (\*U03,X71,X75-X79,X81-X84,Y87.0)

New Hampshire

All races, both sexes.....	10	...	...	-	-	1	1	-	1	1	-	2
Male.....	7	...	...	-	-	-	1	-	1	-	-	1
Female.....	3	...	...	-	-	1	-	-	-	1	-	1
White, both sexes.....	10	...	...	-	-	1	1	-	1	1	-	2
Male.....	7	...	...	-	-	-	1	-	1	-	-	1
Female.....	3	...	...	-	-	1	-	-	-	1	-	1
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	79	...	...	-	-	5	7	2	4	6	7	9
Male.....	60	...	...	-	-	4	5	2	3	3	4	7
Female.....	19	...	...	-	-	1	2	-	1	3	3	2
White, both sexes.....	69	...	...	-	-	4	4	2	3	4	6	8
Male.....	54	...	...	-	-	3	2	2	3	3	3	7
Female.....	15	...	...	-	-	1	2	-	-	1	3	1
Black, both sexes.....	3	...	...	-	-	-	1	-	-	2	-	-
Male.....	1	...	...	-	-	-	1	-	-	-	-	-
Female.....	2	...	...	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	...	...	-	-	1	2	-	1	-	1	1
Male.....	5	...	...	-	-	1	2	-	-	-	1	-
Female.....	2	...	...	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (\*U03,X71,X75-X79,X81-X84,Y87.0)

New Mexico

All races, both sexes.....	10	...	...	-	-	1	1	2	-	-	1	2
Male.....	9	...	...	-	-	1	-	2	-	-	1	2
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
White, both sexes.....	7	...	...	-	-	-	1	1	-	-	-	2
Male.....	6	...	...	-	-	-	-	1	-	-	-	2
Female.....	1	...	...	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	...	...	-	-	1	-	1	-	-	1	-
Male.....	3	...	...	-	-	1	-	1	-	-	1	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	153	...	...	-	-	6	11	13	13	14	18	18
Male.....	115	...	...	-	-	4	8	12	13	12	15	13
Female.....	38	...	...	-	-	2	3	1	-	2	3	5
White, both sexes.....	119	...	...	-	-	3	6	10	9	10	14	15
Male.....	90	...	...	-	-	2	6	9	9	8	12	11
Female.....	29	...	...	-	-	1	-	1	-	2	2	4
Black, both sexes.....	28	...	...	-	-	3	3	3	3	4	2	3
Male.....	21	...	...	-	-	2	1	3	3	4	2	2
Female.....	7	...	...	-	-	1	2	-	-	-	-	1
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	...	...	-	-	-	2	-	1	-	2	-
Male.....	4	...	...	-	-	-	1	-	1	-	1	-
Female.....	2	...	...	-	-	-	1	-	-	-	1	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (\*U03,X71,X75-X79,X81-X84,Y87.0)

South Dakota

All races, both sexes.....	5	...	...	-	-	-	1	1	-	1	-	-
Male.....	4	...	...	-	-	-	1	1	-	1	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	-	-
White, both sexes.....	5	...	...	-	-	-	1	1	-	1	-	-
Male.....	4	...	...	-	-	-	1	1	-	1	-	-
Female.....	1	...	...	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	24	...	...	-	-	-	2	2	3	1	5	4
Male.....	17	...	...	-	-	-	2	1	2	-	4	3
Female.....	7	...	...	-	-	-	-	1	1	1	1	1
White, both sexes.....	18	...	...	-	-	-	2	2	1	1	4	3
Male.....	14	...	...	-	-	-	2	1	1	-	4	2
Female.....	4	...	...	-	-	-	-	1	-	1	-	1
Black, both sexes.....	3	...	...	-	-	-	-	-	1	-	1	1
Male.....	2	...	...	-	-	-	-	-	1	-	-	1
Female.....	1	...	...	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	...	...	-	-	-	-	-	-	-	-	-
Male.....	-	...	...	-	-	-	-	-	-	-	-	-
Female.....	-	...	...	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	-	-	-	-	-	1	-	-	-
Male.....	1	...	...	-	-	-	-	-	-	-	-	-
Female.....	2	...	...	-	-	-	-	-	1	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (\*U03,X71,X75-X79,X81-X84,Y87.0)

Arkansas

All races, both sexes.....	-	1	2	-	2	-	-	-	-	-	-	-
Male.....	-	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	2	-	-	-	-	-	-	-
White, both sexes.....	-	1	2	-	2	-	-	-	-	-	-	-
Male.....	-	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	2	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	41	33	25	5	8	9	5	4	-	1	-	1
Male.....	33	18	17	5	6	5	4	3	-	1	-	1
Female.....	8	15	8	-	2	4	1	1	-	-	-	-
White, both sexes.....	34	32	23	4	6	6	4	3	-	1	-	1
Male.....	29	18	17	4	4	4	3	2	-	1	-	1
Female.....	5	14	6	-	2	2	1	1	-	-	-	-
Black, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	2	1	2	3	1	1	-	-	-	-
Male.....	4	-	-	1	2	1	1	1	-	-	-	-
Female.....	2	-	2	-	-	2	-	-	-	-	-	-























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Alaska												
All races, both sexes.....	50	3	1	-	1	5	6	2	5	9	4	8
Male.....	37	1	-	-	1	3	5	2	4	8	2	7
Female.....	13	2	1	-	-	2	1	-	1	1	2	1
White, both sexes.....	31	1	1	-	-	1	3	1	3	6	4	6
Male.....	23	-	-	-	-	-	3	1	2	6	2	5
Female.....	8	1	1	-	-	1	-	-	1	-	2	1
Black, both sexes.....	3	-	-	-	-	-	2	-	-	-	-	1
Male.....	3	-	-	-	-	-	2	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	15	1	-	-	1	4	1	1	2	3	-	1
Male.....	10	-	-	-	1	3	-	1	2	2	-	1
Female.....	5	1	-	-	-	1	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	528	14	9	5	4	60	109	78	67	41	39	21
Male.....	418	8	3	3	3	50	91	73	52	33	32	15
Female.....	110	6	6	2	1	10	18	5	15	8	7	6
White, both sexes.....	415	12	7	3	3	47	84	64	49	30	29	16
Male.....	331	8	2	2	2	39	70	62	39	26	23	10
Female.....	84	4	5	1	1	8	14	2	10	4	6	6
Black, both sexes.....	64	1	2	2	-	10	13	12	10	8	-	3
Male.....	48	-	1	1	-	9	10	10	7	5	-	3
Female.....	16	1	1	1	-	1	3	2	3	3	-	-
American Indian 1/, both sexes.....	45	1	-	-	1	3	12	2	6	3	9	2
Male.....	37	-	-	-	1	2	11	1	6	2	8	2
Female.....	8	1	-	-	-	1	1	1	-	1	1	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	-	-	2	-	1	-
Male.....	2	-	-	-	-	-	-	-	-	-	1	-
Female.....	2	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Arkansas												
All races, both sexes.....	243	4	12	2	5	19	34	26	27	23	22	22
Male.....	182	3	6	1	4	16	26	24	23	18	14	16
Female.....	61	1	6	1	1	3	8	2	4	5	8	6
White, both sexes.....	108	3	6	-	2	4	11	10	15	7	9	12
Male.....	72	2	2	-	1	2	6	8	13	4	9	10
Female.....	36	1	4	-	1	2	5	2	2	3	-	2
Black, both sexes.....	132	1	6	2	3	14	23	16	12	15	13	9
Male.....	107	1	4	1	3	13	20	16	10	13	5	5
Female.....	25	-	2	1	-	1	3	-	2	2	8	4
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	1	-	-	-	1	-	-
Male.....	2	-	-	-	-	1	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	2,376	35	30	11	28	391	487	337	271	195	164	158
Male.....	2,000	24	13	4	22	357	448	291	233	154	124	124
Female.....	376	11	17	7	6	34	39	46	38	41	40	34
White, both sexes.....	1,529	24	16	8	20	237	318	229	173	118	103	102
Male.....	1,284	17	9	3	16	219	294	197	149	91	75	76
Female.....	245	7	7	5	4	18	24	32	24	27	28	26
Black, both sexes.....	703	11	11	1	7	132	138	93	88	65	48	45
Male.....	610	7	4	-	6	119	127	84	78	55	40	39
Female.....	93	4	7	1	1	13	11	9	10	10	8	6
American Indian 1/, both sexes.....	14	-	1	-	-	3	3	2	1	-	2	1
Male.....	9	-	-	-	-	3	2	-	-	-	2	1
Female.....	5	-	1	-	-	-	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	130	-	2	2	1	19	28	13	9	12	11	10
Male.....	97	-	-	1	-	16	25	10	6	8	7	8
Female.....	33	-	2	1	1	3	3	3	3	4	4	2





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Florida												
All races, both sexes.....	1,324	23	22	7	9	145	233	195	125	135	116	99
Male.....	1,041	15	9	3	6	121	205	160	105	105	96	66
Female.....	283	8	13	4	3	24	28	35	20	30	20	33
White, both sexes.....	674	17	12	5	3	44	95	84	57	74	76	63
Male.....	489	9	4	1	2	34	78	66	44	53	62	41
Female.....	185	8	8	4	1	10	17	18	13	21	14	22
Black, both sexes.....	634	6	10	2	6	100	138	107	64	61	39	34
Male.....	541	6	5	2	4	86	127	92	58	52	33	24
Female.....	93	-	5	-	2	14	11	15	6	9	6	10
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	1	1
Male.....	2	-	-	-	-	-	-	-	-	-	1	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	14	-	-	-	-	1	-	4	4	-	-	1
Male.....	9	-	-	-	-	1	-	2	3	-	-	-
Female.....	5	-	-	-	-	-	-	2	1	-	-	1
Georgia												
All races, both sexes.....	771	17	17	9	9	89	124	105	79	84	60	66
Male.....	603	13	5	7	3	71	114	85	66	63	48	40
Female.....	168	4	12	2	6	18	10	20	13	21	12	26
White, both sexes.....	264	7	-	4	2	19	30	34	23	35	23	34
Male.....	201	4	-	4	2	14	27	28	17	27	18	20
Female.....	63	3	-	-	-	5	3	6	6	8	5	14
Black, both sexes.....	499	10	17	5	7	70	92	69	56	48	37	30
Male.....	397	9	5	3	1	57	85	55	49	36	30	19
Female.....	102	1	12	2	6	13	7	14	7	12	7	11
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	-	2	2	-	1	-	2
Male.....	5	-	-	-	-	-	2	2	-	-	-	1
Female.....	3	-	-	-	-	-	-	-	-	1	-	1









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Kentucky												
All races, both sexes.....	213	5	6	3	4	20	21	29	19	17	16	27
Male.....	161	1	3	3	2	17	15	26	15	14	10	22
Female.....	52	4	3	-	2	3	6	3	4	3	6	5
White, both sexes.....	141	4	6	3	4	7	9	12	11	12	11	21
Male.....	100	1	3	3	2	6	5	9	7	10	7	18
Female.....	41	3	3	-	2	1	4	3	4	2	4	3
Black, both sexes.....	70	1	-	-	-	13	12	17	6	5	5	6
Male.....	59	-	-	-	-	11	10	17	6	4	3	4
Female.....	11	1	-	-	-	2	2	-	-	1	2	2
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes.....	627	8	11	6	7	79	116	99	76	54	50	35
Male.....	512	5	5	5	6	72	98	86	64	40	35	26
Female.....	115	3	6	1	1	7	18	13	12	14	15	9
White, both sexes.....	161	3	1	3	2	9	23	15	16	12	16	11
Male.....	119	2	1	2	1	7	17	11	12	9	12	7
Female.....	42	1	-	1	1	2	6	4	4	3	4	4
Black, both sexes.....	452	5	9	3	5	69	93	81	57	39	32	23
Male.....	382	3	3	3	5	64	81	73	49	28	23	18
Female.....	70	2	6	-	-	5	12	8	8	11	9	5
American Indian 1/, both sexes.....	3	-	-	-	-	1	-	-	2	-	-	-
Male.....	3	-	-	-	-	1	-	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	11	-	1	-	-	-	-	3	1	3	2	1
Male.....	8	-	1	-	-	-	-	2	1	3	-	1
Female.....	3	-	-	-	-	-	-	1	-	-	2	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Massachusetts												
All races, both sexes.....	188	5	3	6	7	25	49	32	12	9	12	7
Male.....	139	3	1	1	5	23	45	29	10	2	8	3
Female.....	49	2	2	5	2	2	4	3	2	7	4	4
White, both sexes.....	102	3	3	5	3	8	25	14	8	5	8	6
Male.....	66	3	1	-	2	7	22	12	7	-	4	3
Female.....	36	-	2	5	1	1	3	2	1	5	4	3
Black, both sexes.....	83	2	-	1	4	16	23	18	4	4	4	1
Male.....	71	-	-	1	3	15	22	17	3	2	4	-
Female.....	12	2	-	-	1	1	1	1	1	2	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	1	1	-	-	-	-	-
Male.....	2	-	-	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes.....	704	6	19	6	9	86	109	93	90	74	56	48
Male.....	567	4	13	3	9	75	99	79	74	60	36	37
Female.....	137	2	6	3	-	11	10	14	16	14	20	11
White, both sexes.....	203	4	9	4	3	16	21	17	18	20	21	20
Male.....	137	3	6	2	3	12	16	13	10	14	10	15
Female.....	66	1	3	2	-	4	5	4	8	6	11	5
Black, both sexes.....	493	2	10	2	5	70	87	76	70	53	35	28
Male.....	427	1	7	1	5	63	83	66	63	46	26	22
Female.....	66	1	3	1	-	7	4	10	7	7	9	6
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	1	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	-	-	1	-	1	-	1	-	-	-
Male.....	3	-	-	-	1	-	-	-	1	-	-	-
Female.....	3	-	-	-	-	-	1	-	-	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
New Hampshire												
All races, both sexes.....	14	-	-	-	1	-	2	1	1	1	1	1
Male.....	7	-	-	-	-	-	2	1	-	-	-	1
Female.....	7	-	-	-	1	-	-	-	1	1	1	-
White, both sexes.....	14	-	-	-	1	-	2	1	1	1	1	1
Male.....	7	-	-	-	-	-	2	1	-	-	-	1
Female.....	7	-	-	-	1	-	-	-	1	1	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	403	9	3	6	4	52	94	68	46	30	25	24
Male.....	329	5	1	3	3	52	79	60	38	18	18	18
Female.....	74	4	2	3	1	-	15	8	8	12	7	6
White, both sexes.....	136	4	1	5	-	9	28	16	10	9	13	10
Male.....	94	1	-	3	-	9	20	12	6	6	7	7
Female.....	42	3	1	2	-	-	8	4	4	3	6	3
Black, both sexes.....	260	5	2	1	4	43	64	51	34	21	12	13
Male.....	231	4	1	-	3	43	58	48	31	12	11	10
Female.....	29	1	1	1	1	-	6	3	3	9	1	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	-	-	-	-	-	2	1	2	-	-	1
Male.....	4	-	-	-	-	-	1	-	1	-	-	1
Female.....	3	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
New Mexico												
All races, both sexes.....	160	6	4	1	3	19	23	23	15	16	11	11
Male.....	121	2	3	1	1	16	19	20	11	15	7	7
Female.....	39	4	1	-	2	3	4	3	4	1	4	4
White, both sexes.....	123	4	3	1	3	15	14	16	11	13	11	9
Male.....	90	-	2	1	1	13	12	13	9	12	7	5
Female.....	33	4	1	-	2	2	2	3	2	1	4	4
Black, both sexes.....	10	-	-	-	-	3	2	1	1	1	-	1
Male.....	9	-	-	-	-	3	2	1	-	1	-	1
Female.....	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	24	2	1	-	-	-	7	6	3	2	-	1
Male.....	21	2	1	-	-	-	5	6	2	2	-	1
Female.....	3	-	-	-	-	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes.....	840	18	17	4	8	106	162	121	95	95	64	44
Male.....	680	9	9	2	7	98	147	104	82	75	47	27
Female.....	160	9	8	2	1	8	15	17	13	20	17	17
White, both sexes.....	338	14	9	3	6	28	58	43	40	36	23	24
Male.....	248	8	5	2	5	25	48	32	35	28	15	14
Female.....	90	6	4	1	1	3	10	11	5	8	8	10
Black, both sexes.....	485	3	8	1	2	77	103	76	51	56	39	19
Male.....	422	1	4	-	2	72	99	70	45	46	31	12
Female.....	63	2	4	1	-	5	4	6	6	10	8	7
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	1	1	-	-
Male.....	2	-	-	-	-	-	-	-	1	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	15	1	-	-	-	1	1	2	3	2	2	1
Male.....	8	-	-	-	-	1	-	2	1	-	1	1
Female.....	7	1	-	-	-	-	1	-	2	2	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Ohio												
All races, both sexes.....	639	12	25	6	4	66	96	111	66	49	54	55
Male.....	494	9	13	3	3	57	79	94	55	38	44	30
Female.....	145	3	12	3	1	9	17	17	11	11	10	25
White, both sexes.....	277	11	13	3	1	17	31	33	18	24	32	28
Male.....	186	8	6	1	1	12	20	25	13	18	25	13
Female.....	91	3	7	2	-	5	11	8	5	6	7	15
Black, both sexes.....	357	1	12	3	3	49	65	78	47	23	22	26
Male.....	304	1	7	2	2	45	59	69	41	18	19	17
Female.....	53	-	5	1	1	4	6	9	6	5	3	9
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	-	-	-	1	2	-	1
Male.....	4	-	-	-	-	-	-	-	1	2	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
Oklahoma												
All races, both sexes.....	251	11	6	-	1	31	35	34	25	19	18	25
Male.....	184	6	4	-	-	26	23	30	20	13	14	19
Female.....	67	5	2	-	1	5	12	4	5	6	4	6
White, both sexes.....	145	5	4	-	-	16	16	18	11	11	12	16
Male.....	102	2	3	-	-	13	10	15	9	8	8	13
Female.....	43	3	1	-	-	3	6	3	2	3	4	3
Black, both sexes.....	81	4	2	-	1	13	19	11	8	5	5	7
Male.....	65	3	1	-	-	12	13	11	7	3	5	5
Female.....	16	1	1	-	1	1	6	-	1	2	-	2
American Indian 1/, both sexes.....	23	2	-	-	-	2	-	4	6	3	1	1
Male.....	16	1	-	-	-	1	-	4	4	2	1	-
Female.....	7	1	-	-	-	1	-	2	2	1	-	1
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Oregon												
All races, both sexes.....	81	2	3	1	1	4	10	14	8	6	4	10
Male.....	61	-	3	-	1	2	5	12	8	5	3	9
Female.....	20	2	-	1	-	2	5	2	-	1	1	1
White, both sexes.....	67	1	3	1	-	4	9	12	6	4	4	7
Male.....	51	-	3	-	-	2	4	10	6	4	3	7
Female.....	16	1	-	1	-	2	5	2	-	-	1	-
Black, both sexes.....	9	1	-	-	-	-	1	2	1	1	-	2
Male.....	6	-	-	-	-	-	1	2	1	-	-	1
Female.....	3	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes.....	5	-	-	-	1	-	-	-	1	1	-	1
Male.....	4	-	-	-	1	-	-	-	1	1	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	750	11	12	5	5	104	131	120	82	69	56	50
Male.....	605	3	9	3	2	97	120	106	65	52	35	38
Female.....	145	8	3	2	3	7	11	14	17	17	21	12
White, both sexes.....	257	5	6	4	1	13	34	24	32	26	29	20
Male.....	165	1	5	2	-	10	28	18	20	16	14	12
Female.....	92	4	1	2	1	3	6	6	12	10	15	8
Black, both sexes.....	477	5	5	1	4	91	96	94	49	42	25	27
Male.....	430	2	3	1	2	87	92	87	44	35	21	24
Female.....	47	3	2	-	2	4	4	7	5	7	4	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	16	1	1	-	-	-	1	2	1	1	2	3
Male.....	10	-	1	-	-	-	-	1	1	1	-	2
Female.....	6	1	-	-	-	-	1	1	-	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Rhode Island												
All races, both sexes.....	24	-	-	-	-	3	7	3	2	1	3	1
Male.....	19	-	-	-	-	2	6	3	1	1	2	1
Female.....	5	-	-	-	-	1	1	-	1	-	1	-
White, both sexes.....	19	-	-	-	-	2	5	2	2	1	3	-
Male.....	14	-	-	-	-	1	4	2	1	1	2	-
Female.....	5	-	-	-	-	1	1	-	1	-	1	-
Black, both sexes.....	4	-	-	-	-	1	1	1	-	-	-	1
Male.....	4	-	-	-	-	1	1	1	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes.....	382	7	9	5	3	41	72	52	37	33	25	25
Male.....	296	6	5	3	2	35	65	41	28	25	17	15
Female.....	86	1	4	2	1	6	7	11	9	8	8	10
White, both sexes.....	143	1	2	2	1	3	17	14	14	13	13	16
Male.....	101	1	1	2	1	1	16	10	10	8	8	10
Female.....	42	-	1	-	-	2	1	4	4	5	5	6
Black, both sexes.....	235	5	7	3	2	38	55	37	22	20	12	9
Male.....	194	5	4	1	1	34	49	31	18	17	9	5
Female.....	41	-	3	2	1	4	6	6	4	3	3	4
American Indian 1/, both sexes.....	2	1	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
South Dakota												
All races, both sexes.....	15	2	1	-	-	2	2	1	-	1	1	3
Male.....	7	1	1	-	-	1	1	1	-	-	-	1
Female.....	8	1	-	-	-	1	1	-	-	1	1	2
White, both sexes.....	11	2	1	-	-	-	-	1	-	1	1	3
Male.....	5	1	1	-	-	-	-	1	-	-	-	1
Female.....	6	1	-	-	-	-	-	-	-	1	1	2
Black, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	3	-	-	-	-	2	1	-	-	-	-	-
Male.....	2	-	-	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes.....	475	8	6	-	3	49	89	68	56	44	41	33
Male.....	387	2	5	-	3	47	76	62	47	37	22	24
Female.....	88	6	1	-	-	2	13	6	9	7	19	9
White, both sexes.....	213	5	-	-	-	21	30	21	23	19	24	20
Male.....	161	2	-	-	-	20	22	17	18	16	12	15
Female.....	52	3	-	-	-	1	8	4	5	3	12	5
Black, both sexes.....	258	3	6	-	3	28	58	47	32	24	17	13
Male.....	224	-	5	-	3	27	53	45	29	21	10	9
Female.....	34	3	1	-	-	1	5	2	3	3	7	4
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	1	-	1	-	-	-
Male.....	2	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Vermont												
All races, both sexes.....	13	2	-	-	-	1	2	2	2	2	1	1
Male.....	9	1	-	-	-	1	2	-	2	2	1	-
Female.....	4	1	-	-	-	-	-	2	-	-	-	1
White, both sexes.....	13	2	-	-	-	1	2	2	2	2	1	1
Male.....	9	1	-	-	-	1	2	-	2	2	1	-
Female.....	4	1	-	-	-	-	-	2	-	-	-	1
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	420	9	5	1	4	54	78	72	30	34	34	29
Male.....	328	5	2	-	1	46	60	60	28	30	23	20
Female.....	92	4	3	1	3	8	18	12	2	4	11	9
White, both sexes.....	156	5	3	1	3	14	26	18	11	14	11	11
Male.....	103	3	2	-	-	8	17	13	11	10	5	9
Female.....	53	2	1	1	3	6	9	5	-	4	6	2
Black, both sexes.....	252	4	2	-	1	39	51	54	17	19	22	17
Male.....	216	2	-	-	1	38	42	47	15	19	18	10
Female.....	36	2	2	-	-	1	9	7	2	-	4	7
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	11	-	-	-	-	1	1	-	2	1	1	1
Male.....	8	-	-	-	-	-	1	-	2	1	-	1
Female.....	3	-	-	-	-	1	-	-	-	-	1	-









## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Arkansas												
All races, both sexes.....	16	10	8	5	2	2	3	1	-	-	-	-
Male.....	8	7	6	5	2	1	2	-	-	-	-	-
Female.....	8	3	2	-	-	1	1	1	-	-	-	-
White, both sexes.....	11	4	5	4	1	1	2	1	-	-	-	-
Male.....	5	1	3	4	1	-	1	-	-	-	-	-
Female.....	6	3	2	-	-	1	1	1	-	-	-	-
Black, both sexes.....	5	6	3	1	1	1	1	-	-	-	-	-
Male.....	3	6	3	1	1	1	1	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	93	70	39	25	11	8	10	6	1	-	1	5
Male.....	72	54	29	23	9	5	8	2	-	-	1	3
Female.....	21	16	10	2	2	3	2	4	1	-	-	2
White, both sexes.....	64	48	22	14	9	6	9	3	1	-	1	4
Male.....	52	36	15	13	7	4	7	1	-	-	1	2
Female.....	12	12	7	1	2	2	2	2	1	-	-	2
Black, both sexes.....	25	14	12	7	1	1	1	3	-	-	-	-
Male.....	18	11	11	7	1	1	1	1	-	-	-	-
Female.....	7	3	1	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	7	5	4	1	1	-	-	-	-	-	1
Male.....	2	6	3	3	1	-	-	-	-	-	-	1
Female.....	2	1	2	1	-	1	-	-	-	-	-	-





















































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Arkansas

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Arkansas. All values are dashes (-).

California

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for California. Values are 1 or dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race categories) for Idaho. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

New Mexico

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Oregon

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All data points are dashes.

Pennsylvania

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All data points are dashes.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Rhode Island

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Washington

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'.

West Virginia

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (\*U01.6-\*U01.7,X85-X90)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

New Mexico

All races, both sexes.....	8	1	1	-	-	-	3	-	-	-	-	1
Male.....	7	1	1	-	-	-	3	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	6	-	1	-	-	-	2	-	-	-	-	1
Male.....	5	-	1	-	-	-	2	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	42	5	2	-	2	-	5	4	4	2	2	6
Male.....	13	2	1	-	1	-	1	2	1	1	-	1
Female.....	29	3	1	-	1	-	4	2	3	1	2	5
White, both sexes.....	25	5	2	-	2	-	3	2	1	-	1	4
Male.....	7	2	1	-	1	-	1	-	1	-	-	1
Female.....	18	3	1	-	1	-	2	2	-	-	1	3
Black, both sexes.....	15	-	-	-	-	-	2	1	2	2	1	2
Male.....	5	-	-	-	-	-	-	1	-	1	-	-
Female.....	10	-	-	-	-	-	2	-	2	1	1	2
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	1	1	-	-	-
Male.....	1	-	-	-	-	-	-	1	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

South Dakota

All races, both sexes.....	2	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	-	-
White, both sexes.....	2	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	14	1	-	-	-	1	2	-	1	1	3	-
Male.....	5	1	-	-	-	1	-	-	-	-	-	-
Female.....	9	-	-	-	-	-	2	-	1	1	3	-
White, both sexes.....	8	1	-	-	-	1	2	-	-	1	-	-
Male.....	4	1	-	-	-	1	-	-	-	-	-	-
Female.....	4	-	-	-	-	-	2	-	-	1	-	-
Black, both sexes.....	5	-	-	-	-	-	-	-	-	-	3	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Iowa

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All values are dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. Values are 1 or dashes (-).







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Nebraska

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Oregon

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by hanging, strangulation and suffocation (X91)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Arkansas												
All races, both sexes.....	153	-	-	2	5	16	29	21	19	18	14	10
Male.....	126	-	-	1	4	14	25	20	18	14	9	8
Female.....	27	-	-	1	1	2	4	1	1	4	5	2
White, both sexes.....	58	-	-	-	2	2	8	7	9	6	7	5
Male.....	44	-	-	-	1	1	5	6	9	4	7	5
Female.....	14	-	-	-	1	1	3	1	-	2	-	-
Black, both sexes.....	93	-	-	2	3	13	21	14	10	11	7	5
Male.....	80	-	-	1	3	12	20	14	9	9	2	3
Female.....	13	-	-	1	-	1	1	-	1	2	5	2
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	1	-	-	-	1	-	-
Male.....	2	-	-	-	-	1	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	1,701	3	5	4	23	341	412	261	219	126	103	89
Male.....	1,494	2	1	-	20	311	386	233	193	103	81	72
Female.....	207	1	4	4	3	30	26	28	26	23	22	17
White, both sexes.....	1,052	3	2	4	17	196	259	176	133	72	59	53
Male.....	912	2	-	-	15	181	243	154	116	56	44	39
Female.....	140	1	2	4	2	15	16	22	17	16	15	14
Black, both sexes.....	563	-	3	-	6	128	131	78	78	49	33	33
Male.....	509	-	1	-	5	116	122	73	71	43	28	30
Female.....	54	-	2	-	1	12	9	5	7	6	5	3
American Indian 1/, both sexes.....	7	-	-	-	-	2	2	-	1	-	2	-
Male.....	6	-	-	-	-	2	2	-	-	-	2	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	79	-	-	-	-	15	20	7	7	5	9	3
Male.....	67	-	-	-	-	12	19	6	6	4	7	3
Female.....	12	-	-	-	-	3	1	1	1	1	2	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Florida												
All races, both sexes.....	935	1	4	3	6	122	197	157	97	94	87	62
Male.....	788	1	1	1	5	108	178	138	85	80	73	45
Female.....	147	-	3	2	1	14	19	19	12	14	14	17
White, both sexes.....	420	-	4	2	1	31	74	54	39	48	57	39
Male.....	324	-	1	-	1	25	62	47	31	38	48	27
Female.....	96	-	3	2	-	6	12	7	8	10	9	12
Black, both sexes.....	506	1	-	1	5	90	123	100	56	46	30	21
Male.....	457	1	-	1	4	82	116	89	52	42	25	17
Female.....	49	-	-	-	1	8	7	11	4	4	5	4
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	1	-	3	2	-	-	1
Male.....	6	-	-	-	-	1	-	2	2	-	-	-
Female.....	2	-	-	-	-	-	-	1	-	-	-	1
Georgia												
All races, both sexes.....	557	1	3	3	4	76	114	85	67	63	46	39
Male.....	469	1	1	2	2	64	106	73	58	52	38	29
Female.....	88	-	2	1	2	12	8	12	9	11	8	10
White, both sexes.....	173	-	-	-	2	16	25	25	16	26	17	23
Male.....	137	-	-	-	2	12	22	22	12	22	14	15
Female.....	36	-	-	-	-	4	3	3	4	4	3	8
Black, both sexes.....	377	1	3	3	2	60	87	59	51	36	29	14
Male.....	328	1	1	2	-	52	82	50	46	30	24	13
Female.....	49	-	2	1	2	8	5	9	5	6	5	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	-	-	-	-	-	2	1	-	1	-	2
Male.....	4	-	-	-	-	-	2	1	-	-	-	1
Female.....	3	-	-	-	-	-	-	-	-	1	-	1







## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Kentucky												
All races, both sexes.....	152	-	-	1	2	18	17	27	14	12	10	20
Male.....	122	-	-	1	2	15	13	24	12	10	7	17
Female.....	30	-	-	-	-	3	4	3	2	2	3	3
White, both sexes.....	93	-	-	1	2	7	6	11	7	7	9	15
Male.....	69	-	-	1	2	6	3	8	5	6	7	13
Female.....	24	-	-	-	-	1	3	3	2	1	2	2
Black, both sexes.....	58	-	-	-	-	11	11	16	6	5	1	5
Male.....	52	-	-	-	-	9	10	16	6	4	-	4
Female.....	6	-	-	-	-	2	1	-	-	1	1	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes.....	488	3	-	2	6	71	102	87	60	40	44	19
Male.....	417	3	-	2	6	65	90	79	52	30	31	15
Female.....	71	-	-	-	-	6	12	8	8	10	13	4
White, both sexes.....	105	2	-	1	1	7	16	14	9	6	15	5
Male.....	80	2	-	1	1	5	13	10	7	5	11	4
Female.....	25	-	-	-	-	2	3	4	2	1	4	1
Black, both sexes.....	372	1	-	1	5	63	86	70	49	32	27	13
Male.....	329	1	-	1	5	59	77	67	43	23	20	10
Female.....	43	-	-	-	-	4	9	3	6	9	7	3
American Indian 1/, both sexes.....	3	-	-	-	-	1	-	-	2	-	-	-
Male.....	3	-	-	-	-	1	-	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	-	-	3	-	2	2	1
Male.....	5	-	-	-	-	-	-	2	-	2	-	1
Female.....	3	-	-	-	-	-	-	1	-	-	2	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Massachusetts												
All races, both sexes.....	115	-	-	1	5	20	38	21	6	7	7	3
Male.....	95	-	-	1	4	19	36	20	5	2	4	1
Female.....	20	-	-	-	1	1	2	1	1	5	3	2
White, both sexes.....	46	-	-	-	1	4	16	7	3	3	4	3
Male.....	34	-	-	-	1	4	15	7	3	-	1	1
Female.....	12	-	-	-	-	-	1	-	-	3	3	2
Black, both sexes.....	67	-	-	1	4	15	21	14	3	4	3	-
Male.....	59	-	-	1	3	14	20	13	2	2	3	-
Female.....	8	-	-	-	1	1	1	1	1	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	1	1	-	-	-	-	-
Male.....	2	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes.....	506	-	3	3	9	77	98	75	83	56	37	22
Male.....	434	-	3	2	9	69	91	64	71	47	29	18
Female.....	72	-	-	1	-	8	7	11	12	9	8	4
White, both sexes.....	105	-	-	1	3	10	15	13	15	10	12	9
Male.....	75	-	-	1	3	8	11	9	9	7	8	8
Female.....	30	-	-	-	-	2	4	4	6	3	4	1
Black, both sexes.....	396	-	3	2	5	67	82	62	67	45	25	13
Male.....	357	-	3	1	5	61	80	55	61	40	21	10
Female.....	39	-	-	1	-	6	2	7	6	5	4	3
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	1	-	1	-	1	-	-	-
Male.....	2	-	-	-	1	-	-	-	1	-	-	-
Female.....	2	-	-	-	-	-	1	-	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

New Hampshire

All races, both sexes.....	6	-	-	-	-	-	1	1	1	1	-	1
Male.....	3	-	-	-	-	-	1	1	-	-	-	1
Female.....	3	-	-	-	-	-	-	-	1	1	-	-
White, both sexes.....	6	-	-	-	-	-	1	1	1	1	-	1
Male.....	3	-	-	-	-	-	1	1	-	-	-	1
Female.....	3	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	276	-	-	-	3	47	78	56	34	18	12	11
Male.....	247	-	-	-	3	47	69	51	32	15	9	10
Female.....	29	-	-	-	-	-	9	5	2	3	3	1
White, both sexes.....	65	-	-	-	-	6	21	8	3	4	7	3
Male.....	50	-	-	-	-	6	17	6	3	4	4	3
Female.....	15	-	-	-	-	-	4	2	-	-	3	-
Black, both sexes.....	208	-	-	-	3	41	57	47	30	14	5	7
Male.....	195	-	-	-	3	41	52	45	28	11	5	6
Female.....	13	-	-	-	-	-	5	2	2	3	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	1	1	-	-	1
Male.....	2	-	-	-	-	-	-	-	1	-	-	1
Female.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

New Mexico

All races, both sexes.....	78	-	-	-	1	13	13	12	8	10	7	4
Male.....	60	-	-	-	1	11	11	11	6	9	4	1
Female.....	18	-	-	-	-	2	2	1	2	1	3	3
White, both sexes.....	70	-	-	-	1	11	11	10	7	9	7	4
Male.....	53	-	-	-	1	9	9	9	6	8	4	1
Female.....	17	-	-	-	-	2	2	1	1	1	3	3
Black, both sexes.....	5	-	-	-	-	2	-	1	1	1	-	-
Male.....	4	-	-	-	-	2	-	1	-	1	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	3	-	-	-	-	-	2	1	-	-	-	-
Male.....	3	-	-	-	-	-	2	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	522	-	-	1	2	90	121	93	70	58	38	13
Male.....	467	-	-	-	2	84	115	83	63	48	30	12
Female.....	55	-	-	1	-	6	6	10	7	10	8	1
White, both sexes.....	157	-	-	1	1	23	34	28	28	16	8	3
Male.....	128	-	-	-	1	20	29	23	24	12	5	3
Female.....	29	-	-	1	-	3	5	5	4	4	3	-
Black, both sexes.....	359	-	-	-	1	67	87	64	40	41	29	9
Male.....	335	-	-	-	1	64	86	59	37	36	25	8
Female.....	24	-	-	-	-	3	1	5	3	5	4	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	-	-	1	1	1	1	1
Male.....	3	-	-	-	-	-	-	1	1	-	-	1
Female.....	2	-	-	-	-	-	-	-	-	1	1	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Oregon												
All races, both sexes.....	43	-	-	-	1	2	7	8	5	3	3	5
Male.....	33	-	-	-	1	1	4	7	5	3	2	4
Female.....	10	-	-	-	-	1	3	1	-	-	1	1
White, both sexes.....	35	-	-	-	-	2	6	6	4	2	3	3
Male.....	26	-	-	-	-	1	3	5	4	2	2	3
Female.....	9	-	-	-	-	1	3	1	-	-	1	-
Black, both sexes.....	6	-	-	-	-	-	1	2	1	-	-	2
Male.....	5	-	-	-	-	-	1	2	1	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	2	-	-	-	1	-	-	-	-	1	-	-
Male.....	2	-	-	-	1	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	554	-	1	2	2	88	118	111	64	55	32	27
Male.....	485	-	-	1	1	83	113	98	55	45	25	24
Female.....	69	-	1	1	1	5	5	13	9	10	7	3
White, both sexes.....	154	-	1	2	1	8	28	21	21	18	13	10
Male.....	113	-	-	1	-	7	25	16	14	13	8	8
Female.....	41	-	1	1	1	1	3	5	7	5	5	2
Black, both sexes.....	395	-	-	-	1	80	90	88	43	37	19	15
Male.....	368	-	-	-	1	76	88	81	41	32	17	14
Female.....	27	-	-	-	-	4	2	7	2	5	2	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	-	-	2	-	-	-	2
Male.....	4	-	-	-	-	-	-	1	-	-	-	2
Female.....	1	-	-	-	-	-	-	1	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Vermont

All races, both sexes.....	9	-	-	-	-	1	1	2	1	2	1	1
Male.....	6	-	-	-	-	1	1	-	1	2	1	-
Female.....	3	-	-	-	-	-	-	2	-	-	-	1
White, both sexes.....	9	-	-	-	-	1	1	2	1	2	1	1
Male.....	6	-	-	-	-	1	1	-	1	2	1	-
Female.....	3	-	-	-	-	-	-	2	-	-	-	1
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	308	1	1	1	3	51	59	59	25	24	16	21
Male.....	254	-	-	-	-	44	48	52	23	22	12	16
Female.....	54	1	1	1	3	7	11	7	2	2	4	5
White, both sexes.....	104	-	-	1	3	13	19	13	6	9	6	8
Male.....	69	-	-	-	-	8	12	9	6	7	2	6
Female.....	35	-	-	1	3	5	7	4	-	2	4	2
Black, both sexes.....	194	1	1	-	-	37	39	46	17	15	10	12
Male.....	177	-	-	-	-	36	35	43	15	15	10	9
Female.....	17	1	1	-	-	1	4	3	2	-	-	3
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	-	-	-	-	1	1	-	2	-	-	1
Male.....	7	-	-	-	-	-	1	-	2	-	-	1
Female.....	2	-	-	-	-	1	-	-	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Illinois

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Iowa

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Wisconsin

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by sharp object (X99)												
Arkansas												
All races, both sexes.....	25	-	-	-	-	1	1	1	3	4	3	6
Male.....	17	-	-	-	-	1	-	1	2	3	2	5
Female.....	8	-	-	-	-	-	1	-	1	1	1	1
White, both sexes.....	8	-	-	-	-	-	-	-	2	1	1	2
Male.....	5	-	-	-	-	-	-	-	2	-	1	2
Female.....	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes.....	16	-	-	-	-	1	1	1	1	3	2	3
Male.....	11	-	-	-	-	1	-	1	-	3	1	2
Female.....	5	-	-	-	-	-	1	-	1	-	1	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	281	-	1	3	4	34	40	42	20	31	25	25
Male.....	226	-	-	1	2	34	38	36	14	23	19	18
Female.....	55	-	1	2	2	-	2	6	6	8	6	7
White, both sexes.....	203	-	-	2	2	26	31	33	14	20	17	18
Male.....	171	-	-	1	1	26	30	29	11	16	13	14
Female.....	32	-	-	1	1	-	1	4	3	4	4	4
Black, both sexes.....	56	-	-	-	1	3	4	7	6	10	7	3
Male.....	43	-	-	-	1	3	4	6	3	7	6	2
Female.....	13	-	-	-	-	-	-	1	3	3	1	1
American Indian 1/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	21	-	1	1	1	4	5	2	-	1	1	4
Male.....	11	-	-	-	-	4	4	1	-	-	-	2
Female.....	10	-	1	1	1	-	1	1	-	1	1	2































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by sharp object (X99)												
New Mexico												
All races, both sexes.....	23	-	-	-	-	4	4	5	4	2	2	-
Male.....	20	-	-	-	-	3	3	5	4	2	1	-
Female.....	3	-	-	-	-	1	1	-	-	-	1	-
White, both sexes.....	14	-	-	-	-	3	-	3	2	2	2	-
Male.....	13	-	-	-	-	3	-	3	2	2	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	7	-	-	-	-	-	3	2	2	-	-	-
Male.....	6	-	-	-	-	-	2	2	2	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes.....	122	-	-	1	3	11	25	14	12	21	10	7
Male.....	95	-	-	-	3	9	23	11	10	16	6	4
Female.....	27	-	-	1	-	2	2	3	2	5	4	3
White, both sexes.....	58	-	-	-	2	4	12	8	6	13	2	5
Male.....	45	-	-	-	2	4	12	5	5	10	1	3
Female.....	13	-	-	-	-	-	-	3	1	3	1	2
Black, both sexes.....	59	-	-	1	1	6	12	6	6	7	7	2
Male.....	46	-	-	-	1	4	11	6	5	5	4	1
Female.....	13	-	-	1	-	2	1	-	1	2	3	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	1	1	-	-	-	1	-
Male.....	3	-	-	-	-	1	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by sharp object (X99)												
Rhode Island												
All races, both sexes.....	2	-	-	-	-	1	-	-	-	-	1	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes.....	48	-	-	-	-	-	6	5	7	9	4	1
Male.....	32	-	-	-	-	-	5	3	6	8	2	-
Female.....	16	-	-	-	-	-	1	2	1	1	2	1
White, both sexes.....	20	-	-	-	-	-	2	1	3	3	2	-
Male.....	12	-	-	-	-	-	1	1	3	2	1	-
Female.....	8	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes.....	27	-	-	-	-	-	4	3	4	6	2	1
Male.....	20	-	-	-	-	-	4	2	3	6	1	-
Female.....	7	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-















































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by blunt object (Y00)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Assault (homicide) by blunt object (Y00)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by blunt object (Y00)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Assault (homicide) by blunt object (Y00)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Assault (homicide) by bodily force (Y04)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by bodily force (Y04)

Massachusetts

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Assault (homicide) by bodily force (Y04)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Assault (homicide) by bodily force (Y04)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Assault (homicide) by bodily force (Y04)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Delaware

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Missouri

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Nebraska

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Sexual assault (homicide) by bodily force (Y05)

New Hampshire

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

North Carolina

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Rhode Island

Table with 13 columns representing age groups and 13 rows representing race and sex categories for Rhode Island. All cells contain dashes (-).

South Carolina

Table with 13 columns representing age groups and 13 rows representing race and sex categories for South Carolina. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Vermont

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sexual assault (homicide) by bodily force (Y05)

Washington

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Rhode Island

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

South Dakota

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for South Dakota. All cells contain dashes (-).

Tennessee

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Tennessee. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Sexual assault (homicide) by bodily force (Y05)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Idaho. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. Most cells contain dashes (-), but the 85-89 age group for 'All races, both sexes' and 'White, both sexes' contains the number 1.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Washington

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Alaska

All races, both sexes.....	15	1	1	-	-	1	2	1	-	5	1	1
Male.....	9	1	-	-	-	1	1	1	-	4	-	1
Female.....	6	-	1	-	-	-	1	-	-	1	1	-
White, both sexes.....	5	-	1	-	-	-	-	-	-	2	1	-
Male.....	2	-	-	-	-	-	-	-	-	2	-	-
Female.....	3	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	8	-	-	-	-	1	1	1	-	3	-	1
Male.....	5	-	-	-	-	1	-	1	-	2	-	1
Female.....	3	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	95	9	6	1	-	4	10	7	13	7	6	5
Male.....	61	7	3	1	-	2	7	6	7	5	5	4
Female.....	34	2	3	-	-	2	3	1	6	2	1	1
White, both sexes.....	74	9	4	-	-	4	9	7	7	3	3	4
Male.....	48	7	2	-	-	2	7	6	4	3	3	3
Female.....	26	2	2	-	-	2	2	1	3	-	-	1
Black, both sexes.....	6	-	2	1	-	-	-	-	1	2	-	-
Male.....	3	-	1	1	-	-	-	-	-	1	-	-
Female.....	3	-	1	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes.....	13	-	-	-	-	-	1	-	3	2	3	1
Male.....	10	-	-	-	-	-	-	-	3	1	2	1
Female.....	3	-	-	-	-	-	1	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	2	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Arkansas

All races, both sexes.....	56	4	12	-	-	2	3	3	4	1	3	5
Male.....	35	3	6	-	-	1	1	2	3	1	2	3
Female.....	21	1	6	-	-	1	2	1	1	-	1	2
White, both sexes.....	36	3	6	-	-	2	2	2	3	-	1	5
Male.....	20	2	2	-	-	1	1	1	2	-	1	3
Female.....	16	1	4	-	-	1	1	1	1	-	-	2
Black, both sexes.....	20	1	6	-	-	-	1	1	1	1	2	-
Male.....	15	1	4	-	-	-	-	1	1	1	1	-
Female.....	5	-	2	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	280	19	14	1	1	12	25	21	25	32	22	33
Male.....	209	13	7	-	-	11	19	14	21	24	15	27
Female.....	71	6	7	1	1	1	6	7	4	8	7	6
White, both sexes.....	191	15	8	-	1	12	19	10	20	21	16	22
Male.....	149	10	5	-	-	11	16	9	18	16	11	17
Female.....	42	5	3	-	1	1	3	1	2	5	5	5
Black, both sexes.....	60	4	5	1	-	-	2	6	3	5	5	7
Male.....	44	3	2	-	-	-	1	3	3	4	4	6
Female.....	16	1	3	1	-	-	1	3	-	1	1	1
American Indian 1/, both sexes.....	5	-	1	-	-	-	1	2	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	4	-	1	-	-	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	24	-	-	-	-	-	3	3	2	6	1	3
Male.....	15	-	-	-	-	-	2	2	-	4	-	3
Female.....	9	-	-	-	-	-	1	1	2	2	1	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

New Hampshire

All races, both sexes.....	2	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes.....	2	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	65	6	3	1	1	1	8	4	6	5	6	7
Male.....	43	4	1	-	-	1	5	4	2	3	4	4
Female.....	22	2	2	1	1	-	3	-	4	2	2	3
White, both sexes.....	39	3	1	-	-	1	3	3	3	3	5	5
Male.....	25	1	-	-	-	1	2	3	-	2	3	3
Female.....	14	2	1	-	-	-	1	-	3	1	2	2
Black, both sexes.....	22	3	2	1	1	-	3	1	2	2	1	2
Male.....	16	3	1	-	-	-	2	1	2	1	1	1
Female.....	6	-	1	1	1	-	1	-	-	1	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	2	-	1	-	-	-
Male.....	2	-	-	-	-	-	1	-	-	-	-	-
Female.....	2	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

New Mexico

All races, both sexes.....	47	5	2	-	2	2	3	5	3	4	1	6
Male.....	30	1	1	-	-	2	2	3	1	4	1	5
Female.....	17	4	1	-	2	-	1	2	2	-	-	1
White, both sexes.....	30	4	2	-	2	1	1	2	2	2	1	4
Male.....	16	-	1	-	-	1	1	-	1	2	1	3
Female.....	14	4	1	-	2	-	-	2	1	-	-	1
Black, both sexes.....	3	-	-	-	-	1	-	-	-	-	-	1
Male.....	3	-	-	-	-	1	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	12	1	-	-	-	-	2	3	1	2	-	1
Male.....	10	1	-	-	-	-	1	3	-	2	-	1
Female.....	2	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	139	12	11	1	1	4	10	10	9	13	13	17
Male.....	96	6	7	1	1	4	8	8	8	9	10	9
Female.....	43	6	4	-	-	-	2	2	1	4	3	8
White, both sexes.....	89	8	5	1	1	1	8	5	5	6	11	11
Male.....	62	5	3	1	1	1	6	4	5	5	8	6
Female.....	27	3	2	-	-	-	2	1	-	1	3	5
Black, both sexes.....	46	3	6	-	-	3	2	5	3	6	2	6
Male.....	33	1	4	-	-	3	2	4	3	4	2	3
Female.....	13	2	2	-	-	-	-	1	-	2	-	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	1	-	-	-	-	-	-	1	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	1	-	-	-	-	-	-	1	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Ohio

All races, both sexes.....	104	2	12	4	-	5	8	7	7	3	10	12
Male.....	72	2	7	2	-	4	7	3	6	2	6	7
Female.....	32	-	5	2	-	1	1	4	1	1	4	5
White, both sexes.....	65	1	4	2	-	3	5	4	2	2	9	7
Male.....	42	1	1	1	-	2	4	2	1	2	5	5
Female.....	23	-	3	1	-	1	1	2	1	-	4	2
Black, both sexes.....	39	1	8	2	-	2	3	3	5	1	1	5
Male.....	30	1	6	1	-	2	3	1	5	-	1	2
Female.....	9	-	2	1	-	-	-	2	-	1	-	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	46	6	2	-	-	2	2	3	7	3	5	4
Male.....	29	4	1	-	-	2	-	2	3	2	4	4
Female.....	17	2	1	-	-	-	2	1	4	1	1	-
White, both sexes.....	33	3	2	-	-	2	2	2	4	1	4	3
Male.....	22	1	1	-	-	2	-	2	3	1	3	3
Female.....	11	2	1	-	-	-	2	-	1	-	1	-
Black, both sexes.....	7	2	-	-	-	-	-	-	1	1	1	1
Male.....	5	2	-	-	-	-	-	-	-	-	1	1
Female.....	2	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes.....	5	1	-	-	-	-	-	-	2	1	-	-
Male.....	2	1	-	-	-	-	-	-	-	1	-	-
Female.....	3	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Oregon

All races, both sexes.....	21	1	2	-	-	-	3	3	-	3	1	2
Male.....	17	-	2	-	-	-	1	3	-	2	1	2
Female.....	4	1	-	-	-	-	2	-	-	1	-	-
White, both sexes.....	18	-	2	-	-	-	3	3	-	2	1	2
Male.....	16	-	2	-	-	-	1	3	-	2	1	2
Female.....	2	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes.....	3	1	-	-	-	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	84	6	8	2	1	2	6	2	7	8	11	7
Male.....	54	3	8	2	-	2	2	2	3	4	6	6
Female.....	30	3	-	-	1	-	4	-	4	4	5	1
White, both sexes.....	46	2	5	1	-	2	3	1	3	4	9	2
Male.....	26	1	5	1	-	2	1	1	1	1	4	2
Female.....	20	1	-	-	-	-	2	-	2	3	5	-
Black, both sexes.....	32	4	2	1	1	-	2	1	3	3	2	5
Male.....	23	2	2	1	-	-	1	1	1	2	2	4
Female.....	9	2	-	-	1	-	1	-	2	1	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	1	-	-	-	1	-	1	1	-	-
Male.....	5	-	1	-	-	-	-	-	1	1	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5,\*U01.8-\*U01.9,\*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Arkansas

All races, both sexes.....	6	4	3	3	1	1	-	1	-	-	-	-
Male.....	3	2	3	3	1	1	-	-	-	-	-	-
Female.....	3	2	-	-	-	-	-	1	-	-	-	-
White, both sexes.....	4	2	2	2	1	-	-	1	-	-	-	-
Male.....	2	-	2	2	1	-	-	-	-	-	-	-
Female.....	2	2	-	-	-	-	-	1	-	-	-	-
Black, both sexes.....	2	2	1	1	-	1	-	-	-	-	-	-
Male.....	1	2	1	1	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	19	22	9	10	5	2	3	1	1	-	-	3
Male.....	15	19	6	10	3	1	2	1	-	-	-	1
Female.....	4	3	3	-	2	1	1	-	1	-	-	2
White, both sexes.....	14	13	3	5	3	1	3	1	1	-	-	3
Male.....	11	12	2	5	1	1	2	1	-	-	-	1
Female.....	3	1	1	-	2	-	1	-	1	-	-	2
Black, both sexes.....	5	8	3	5	1	-	-	-	-	-	-	-
Male.....	4	6	2	5	1	-	-	-	-	-	-	-
Female.....	1	2	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	3	-	1	1	-	-	-	-	-	-
Male.....	-	1	2	-	1	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	-	-	-	-	-

























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Arkansas

All races, both sexes.....	165	-	-	-	1	5	17	24	16	18	19	28
Male.....	94	-	-	-	1	3	14	10	11	13	13	12
Female.....	71	-	-	-	-	2	3	14	5	5	6	16
White, both sexes.....	147	-	-	-	1	4	16	17	16	15	17	26
Male.....	82	-	-	-	1	3	13	7	11	10	11	11
Female.....	65	-	-	-	-	1	3	10	5	5	6	15
Black, both sexes.....	17	-	-	-	-	1	1	7	-	3	1	2
Male.....	11	-	-	-	-	-	1	3	-	3	1	1
Female.....	6	-	-	-	-	1	-	4	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	1	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	345	15	2	-	2	12	18	22	15	34	46	49
Male.....	218	9	-	-	1	8	15	11	11	26	31	23
Female.....	127	6	2	-	1	4	3	11	4	8	15	26
White, both sexes.....	284	12	1	-	1	9	16	17	11	26	38	40
Male.....	177	6	-	-	-	5	13	10	8	20	25	19
Female.....	107	6	1	-	1	4	3	7	3	6	13	21
Black, both sexes.....	33	2	-	-	-	1	-	2	3	3	5	5
Male.....	23	2	-	-	-	1	-	-	2	3	3	2
Female.....	10	-	-	-	-	-	-	2	1	-	2	3
American Indian 1/, both sexes.....	4	-	-	-	-	1	1	1	-	-	-	-
Male.....	3	-	-	-	-	1	1	1	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	24	1	1	-	1	1	1	2	1	5	3	4
Male.....	15	1	-	-	1	1	1	-	1	3	3	2
Female.....	9	-	1	-	-	-	-	2	-	2	-	2

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

## Colorado

All races, both sexes.....	126	1	1	3	-	8	4	5	7	16	18	14
Male.....	67	1	-	1	-	6	4	3	6	7	9	9
Female.....	59	-	1	2	-	2	-	2	1	9	9	5
White, both sexes.....	111	1	1	3	-	6	4	4	7	14	15	12
Male.....	61	1	-	1	-	4	4	2	6	7	8	8
Female.....	50	-	1	2	-	2	-	2	1	7	7	4
Black, both sexes.....	12	-	-	-	-	1	-	-	-	2	3	1
Male.....	3	-	-	-	-	1	-	-	-	-	1	-
Female.....	9	-	-	-	-	-	-	-	-	2	2	1
American Indian 1/, both sexes.....	3	-	-	-	-	1	-	1	-	-	-	1
Male.....	3	-	-	-	-	1	-	1	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Connecticut

All races, both sexes.....	45	-	1	-	-	2	2	4	1	2	6	9
Male.....	27	-	1	-	-	1	2	2	1	2	6	3
Female.....	18	-	-	-	-	1	-	2	-	-	-	6
White, both sexes.....	37	-	1	-	-	1	1	3	1	2	5	7
Male.....	22	-	1	-	-	1	1	1	1	2	5	2
Female.....	15	-	-	-	-	-	-	2	-	-	-	5
Black, both sexes.....	6	-	-	-	-	-	1	1	-	-	1	1
Male.....	5	-	-	-	-	-	1	1	-	-	1	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Maine

All races, both sexes.....	14	-	-	-	-	-	1	2	1	1	2	2
Male.....	10	-	-	-	-	-	1	1	-	1	2	2
Female.....	4	-	-	-	-	-	-	1	1	-	-	-
White, both sexes.....	13	-	-	-	-	-	1	1	1	1	2	2
Male.....	9	-	-	-	-	-	1	-	-	1	2	2
Female.....	4	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	778	-	2	1	-	16	54	67	55	87	144	165
Male.....	546	-	1	1	-	14	41	49	39	65	102	109
Female.....	232	-	1	-	-	2	13	18	16	22	42	56
White, both sexes.....	525	-	1	-	-	15	46	59	43	56	91	102
Male.....	367	-	-	-	-	13	34	44	30	44	65	67
Female.....	158	-	1	-	-	2	12	15	13	12	26	35
Black, both sexes.....	247	-	1	1	-	1	8	7	10	31	52	61
Male.....	175	-	1	1	-	1	7	5	7	21	36	41
Female.....	72	-	-	-	-	-	1	2	3	10	16	20
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	-	-	-	-	-	1	2	-	1	2
Male.....	4	-	-	-	-	-	-	-	2	-	1	1
Female.....	2	-	-	-	-	-	-	1	-	-	-	1













































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Oregon

All races, both sexes.....	18	11	6	1	-	1	1	-	-	-	-	-
Male.....	9	5	1	-	-	1	-	-	-	-	-	-
Female.....	9	6	5	1	-	-	1	-	-	-	-	-
White, both sexes.....	15	11	6	1	-	1	1	-	-	-	-	-
Male.....	7	5	1	-	-	1	-	-	-	-	-	-
Female.....	8	6	5	1	-	-	1	-	-	-	-	-
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	24	8	7	3	3	4	4	5	3	-	-	1
Male.....	12	5	3	3	1	2	3	3	-	-	-	1
Female.....	12	3	4	-	2	2	1	2	3	-	-	-
White, both sexes.....	21	7	5	3	2	3	3	5	2	-	-	1
Male.....	11	4	2	3	1	2	3	3	-	-	-	1
Female.....	10	3	3	-	1	1	-	2	2	-	-	-
Black, both sexes.....	3	1	2	-	1	-	1	-	1	-	-	-
Male.....	1	1	1	-	-	-	-	-	-	-	-	-
Female.....	2	-	1	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Wisconsin

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Wyoming

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

Arkansas

All races, both sexes.....	128	-	-	-	1	3	14	20	12	15	15	25
Male.....	68	-	-	-	1	3	11	7	7	10	9	12
Female.....	60	-	-	-	-	-	3	13	5	5	6	13
White, both sexes.....	119	-	-	-	1	3	14	15	12	13	15	23
Male.....	63	-	-	-	1	3	11	5	7	8	9	11
Female.....	56	-	-	-	-	-	3	10	5	5	6	12
Black, both sexes.....	9	-	-	-	-	-	-	5	-	2	-	2
Male.....	5	-	-	-	-	-	-	2	-	2	-	1
Female.....	4	-	-	-	-	-	-	3	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	160	1	2	-	-	4	7	12	8	19	24	26
Male.....	84	-	-	-	-	2	6	5	5	14	11	10
Female.....	76	1	2	-	-	2	1	7	3	5	13	16
White, both sexes.....	141	1	1	-	-	4	7	10	6	15	20	24
Male.....	73	-	-	-	-	2	6	5	4	11	9	8
Female.....	68	1	1	-	-	2	1	5	2	4	11	16
Black, both sexes.....	13	-	-	-	-	-	-	2	2	2	3	1
Male.....	8	-	-	-	-	-	-	-	1	2	1	1
Female.....	5	-	-	-	-	-	-	2	1	-	2	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	1	-	-	-	-	-	-	2	1	1
Male.....	3	-	-	-	-	-	-	-	-	1	1	1
Female.....	2	-	1	-	-	-	-	-	-	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

Colorado

All races, both sexes.....	85	-	-	2	-	3	3	5	6	11	13	9
Male.....	44	-	-	1	-	3	3	3	5	5	5	6
Female.....	41	-	-	1	-	-	-	2	1	6	8	3
White, both sexes.....	78	-	-	2	-	2	3	4	6	10	11	9
Male.....	42	-	-	1	-	2	3	2	5	5	5	6
Female.....	36	-	-	1	-	-	-	2	1	5	6	3
Black, both sexes.....	5	-	-	-	-	-	-	-	-	1	2	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	5	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes.....	2	-	-	-	-	1	-	1	-	-	-	-
Male.....	2	-	-	-	-	1	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	32	-	-	-	-	1	-	3	1	1	4	9
Male.....	15	-	-	-	-	-	-	1	1	1	4	3
Female.....	17	-	-	-	-	1	-	2	-	-	-	6
White, both sexes.....	28	-	-	-	-	-	-	2	1	1	4	7
Male.....	13	-	-	-	-	-	-	-	1	1	4	2
Female.....	15	-	-	-	-	-	-	2	-	-	-	5
Black, both sexes.....	2	-	-	-	-	-	-	1	-	-	-	1
Male.....	2	-	-	-	-	-	-	1	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

Maine

All races, both sexes.....	8	-	-	-	-	-	-	1	-	1	-	2
Male.....	5	-	-	-	-	-	-	-	-	1	-	2
Female.....	3	-	-	-	-	-	-	1	-	-	-	-
White, both sexes.....	8	-	-	-	-	-	-	1	-	1	-	2
Male.....	5	-	-	-	-	-	-	-	-	1	-	2
Female.....	3	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	677	-	-	-	-	15	47	62	49	76	132	144
Male.....	464	-	-	-	-	13	36	44	33	55	92	93
Female.....	213	-	-	-	-	2	11	18	16	21	40	51
White, both sexes.....	461	-	-	-	-	14	42	56	40	49	86	86
Male.....	314	-	-	-	-	12	31	41	27	37	60	54
Female.....	147	-	-	-	-	2	11	15	13	12	26	32
Black, both sexes.....	212	-	-	-	-	1	5	5	8	27	45	57
Male.....	147	-	-	-	-	1	5	3	5	18	31	38
Female.....	65	-	-	-	-	-	-	2	3	9	14	19
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	-	1	1	-	1	1
Male.....	3	-	-	-	-	-	-	-	1	-	1	1
Female.....	1	-	-	-	-	-	-	1	-	-	-	-



























































































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Florida

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Missouri

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

New Mexico

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for New Mexico. All values are 0.

New York

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for New York. Values are 1 or 2.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Oregon

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Rhode Island

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Washington

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Arkansas

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Florida

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Missouri

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Nebraska

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Rhode Island

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

South Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Hawaii

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

New Hampshire

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Falling, jumping or pushed from a high place, undetermined intent (Y30)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Falling, jumping or pushed from a high place, undetermined intent (Y30)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Arkansas

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Florida

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Florida. All values are dashes (-).

Georgia

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Georgia. Values are dashes (-) or 1.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Illinois

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Illinois. All cells contain dashes (-).

Indiana

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Indiana. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Minnesota

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Minnesota. All cells contain dashes (-).

Mississippi

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Mississippi. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Oregon

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Rhode Island

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Rhode Island. All cells contain dashes (-).

South Carolina

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for South Carolina. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Falling, jumping or pushed from a high place, undetermined intent (Y30)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

Arkansas

All races, both sexes.....	24	-	-	-	-	2	3	1	2	1	2	2
Male.....	14	-	-	-	-	-	3	-	2	1	2	-
Female.....	10	-	-	-	-	2	-	1	-	-	-	2
White, both sexes.....	17	-	-	-	-	1	2	-	2	-	-	2
Male.....	9	-	-	-	-	-	2	-	2	-	-	-
Female.....	8	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes.....	6	-	-	-	-	1	1	1	-	1	1	-
Male.....	4	-	-	-	-	-	1	-	-	1	1	-
Female.....	2	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	1	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	144	14	-	-	1	3	7	6	6	10	19	18
Male.....	100	9	-	-	-	1	5	2	5	9	17	8
Female.....	44	5	-	-	1	2	2	4	1	1	2	10
White, both sexes.....	109	11	-	-	1	3	6	3	5	6	15	12
Male.....	77	6	-	-	-	1	4	1	4	6	13	7
Female.....	32	5	-	-	1	2	2	2	1	-	2	5
Black, both sexes.....	17	2	-	-	-	-	-	-	-	1	2	3
Male.....	12	2	-	-	-	-	-	-	-	1	2	-
Female.....	5	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	17	1	-	-	-	-	1	2	1	3	2	3
Male.....	10	1	-	-	-	-	1	-	1	2	2	1
Female.....	7	-	-	-	-	-	-	2	-	1	-	2































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

Oregon

All races, both sexes.....	6	1	1	-	-	-	1	-	-	-	-	-
Male.....	5	-	1	-	-	-	-	-	-	-	-	-
Female.....	1	1	-	-	-	-	1	-	-	-	-	-
White, both sexes.....	3	1	1	-	-	-	1	-	-	-	-	-
Male.....	3	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	5	3	3	1	3	3	3	3	2	-	-	1
Male.....	3	3	1	1	1	2	2	2	-	-	-	1
Female.....	2	-	2	-	2	1	1	1	2	-	-	-
White, both sexes.....	5	2	3	1	2	2	2	3	1	-	-	1
Male.....	3	2	1	1	1	2	2	2	-	-	-	1
Female.....	2	-	2	-	1	-	-	1	1	-	-	-
Black, both sexes.....	-	1	-	-	1	-	1	-	1	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal intervention (Y35,Y89.0)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Legal intervention (Y35,Y89.0)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Legal intervention (Y35,Y89.0)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Massachusetts

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Legal execution (Y35.5)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Legal execution (Y35.5)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

Delaware

Table with 13 columns (Age groups) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

Nebraska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Massachusetts

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Missouri

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

New Mexico

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

North Carolina

Table with 13 columns representing age groups and 13 rows representing race and sex categories for North Carolina. All cells contain dashes (-).

North Dakota

Table with 13 columns representing age groups and 13 rows representing race and sex categories for North Dakota. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Rhode Island

Table with 13 columns representing age groups and 13 rows representing race and sex categories for Rhode Island. All cells contain dashes (-).

South Carolina

Table with 13 columns representing age groups and 13 rows representing race and sex categories for South Carolina. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Operations of war and their sequelae (Y36,Y89.1)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Alaska

All races, both sexes.....	-	-	-	2	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	2	-	-	1	-	-	-	-	-
White, both sexes.....	-	-	-	2	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	2	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	1	4	5	8	4	12	8	8	-	-	-	-
Male.....	1	3	3	6	4	5	2	3	-	-	-	-
Female.....	-	1	2	2	-	7	6	5	-	-	-	-
White, both sexes.....	1	3	4	7	4	11	6	7	-	-	-	-
Male.....	1	2	2	5	4	5	2	2	-	-	-	-
Female.....	-	1	2	2	-	6	4	5	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	1	-	-	1	1	1	-	-	-	-
Male.....	-	1	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Arkansas

All races, both sexes.....	2	4	1	-	1	1	3	4	1	1	-	-
Male.....	2	2	1	-	1	-	3	2	-	-	-	-
Female.....	-	2	-	-	-	1	-	2	1	1	-	-
White, both sexes.....	1	3	1	-	-	-	3	3	1	-	-	-
Male.....	1	2	1	-	-	-	3	2	-	-	-	-
Female.....	-	1	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	1	1	-	-	1	1	-	-	-	1	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	14	13	19	11	19	14	21	21	10	4	1	-
Male.....	8	8	7	5	6	6	8	13	2	3	1	-
Female.....	6	5	12	6	13	8	13	8	8	1	-	-
White, both sexes.....	11	10	19	9	16	13	20	18	5	4	1	-
Male.....	6	7	7	3	5	6	8	11	2	3	1	-
Female.....	5	3	12	6	11	7	12	7	3	1	-	-
Black, both sexes.....	2	1	-	2	1	-	1	1	2	-	-	-
Male.....	1	-	-	2	-	-	-	1	-	-	-	-
Female.....	1	1	-	-	1	-	1	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	-	-	2	1	-	2	3	-	-	-
Male.....	1	1	-	-	1	-	-	1	-	-	-	-
Female.....	-	1	-	-	1	1	-	1	3	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Florida

All races, both sexes.....	4	3	11	5	16	22	15	8	6	4	-	-
Male.....	1	3	3	2	10	10	10	5	-	1	-	-
Female.....	3	-	8	3	6	12	5	3	6	3	-	-
White, both sexes.....	1	2	10	5	11	22	15	8	4	4	-	-
Male.....	-	2	2	2	7	10	10	5	-	1	-	-
Female.....	1	-	8	3	4	12	5	3	4	3	-	-
Black, both sexes.....	3	1	1	-	5	-	-	-	2	-	-	-
Male.....	1	1	1	-	3	-	-	-	-	-	-	-
Female.....	2	-	-	-	2	-	-	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Georgia

All races, both sexes.....	8	7	8	12	10	15	14	10	1	1	-	-
Male.....	4	5	3	6	5	8	7	3	-	-	-	-
Female.....	4	2	5	6	5	7	7	7	1	1	-	-
White, both sexes.....	4	5	6	9	7	8	11	9	1	1	-	-
Male.....	3	3	3	5	4	5	5	3	-	-	-	-
Female.....	1	2	3	4	3	3	6	6	1	1	-	-
Black, both sexes.....	4	2	2	3	3	7	3	-	-	-	-	-
Male.....	1	2	-	1	1	3	2	-	-	-	-	-
Female.....	3	-	2	2	2	4	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

New Hampshire

All races, both sexes.....	1	-	1	-	-	2	1	-	-	1	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	1	-	-
White, both sexes.....	1	-	1	-	-	2	1	-	-	1	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	4	3	4	5	9	7	17	10	2	2	-	-
Male.....	1	1	1	5	3	2	7	5	1	-	-	-
Female.....	3	2	3	-	6	5	10	5	1	2	-	-
White, both sexes.....	2	3	2	4	7	7	13	8	2	1	-	-
Male.....	1	1	1	4	2	2	6	3	1	-	-	-
Female.....	1	2	1	-	5	5	7	5	1	1	-	-
Black, both sexes.....	2	-	2	1	2	-	3	2	-	1	-	-
Male.....	-	-	-	1	1	-	1	2	-	-	-	-
Female.....	2	-	2	-	1	-	2	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

New Mexico

All races, both sexes.....	-	-	4	3	2	2	4	3	3	-	-	-
Male.....	-	-	2	2	1	1	2	3	2	-	-	-
Female.....	-	-	2	1	1	1	2	-	1	-	-	-
White, both sexes.....	-	-	4	3	2	2	4	2	3	-	-	-
Male.....	-	-	2	2	1	1	2	2	2	-	-	-
Female.....	-	-	2	1	1	1	2	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	11	7	10	17	14	24	15	12	7	3	-	-
Male.....	5	-	5	7	8	10	6	6	2	-	-	-
Female.....	6	7	5	10	6	14	9	6	5	3	-	-
White, both sexes.....	10	6	7	14	8	19	14	12	6	2	-	-
Male.....	5	-	4	7	5	9	6	6	2	-	-	-
Female.....	5	6	3	7	3	10	8	6	4	2	-	-
Black, both sexes.....	1	1	3	3	6	2	1	-	1	1	-	-
Male.....	-	-	1	-	3	-	-	-	-	-	-	-
Female.....	1	1	2	3	3	2	1	-	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



































































































































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

New Mexico

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	1	-	1	4	4	4	4	3	1	-	-	-
Male.....	-	-	-	1	3	1	2	2	-	-	-	-
Female.....	1	-	1	3	1	3	2	1	1	-	-	-
White, both sexes.....	-	-	-	2	1	3	3	3	-	-	-	-
Male.....	-	-	-	1	1	1	2	2	-	-	-	-
Female.....	-	-	-	1	-	2	1	1	-	-	-	-
Black, both sexes.....	1	-	1	2	3	-	1	-	1	-	-	-
Male.....	-	-	-	-	2	-	-	-	-	-	-	-
Female.....	1	-	1	2	1	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



















































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Alaska

All races, both sexes.....	-	-	-	1	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes.....	-	-	-	1	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	1	3	4	4	4	10	7	8	-	-	-	-
Male.....	1	2	2	3	4	5	1	3	-	-	-	-
Female.....	-	1	2	1	-	5	6	5	-	-	-	-
White, both sexes.....	1	2	3	4	4	9	5	7	-	-	-	-
Male.....	1	1	1	3	4	5	1	2	-	-	-	-
Female.....	-	1	2	1	-	4	4	5	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	1	-	-	1	1	1	-	-	-	-
Male.....	-	1	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Arkansas

All races, both sexes.....	1	4	1	-	1	-	3	3	1	1	-	-
Male.....	1	2	1	-	1	-	3	1	-	-	-	-
Female.....	-	2	-	-	-	-	-	2	1	1	-	-
White, both sexes.....	1	3	1	-	-	-	3	2	1	-	-	-
Male.....	1	2	1	-	-	-	3	1	-	-	-	-
Female.....	-	1	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	-	1	-	-	1	-	-	-	-	1	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	10	10	17	7	15	11	17	14	10	4	1	-
Male.....	6	6	6	4	6	4	6	9	2	3	1	-
Female.....	4	4	11	3	9	7	11	5	8	1	-	-
White, both sexes.....	8	8	17	5	13	10	16	11	5	4	1	-
Male.....	5	5	6	2	5	4	6	7	2	3	1	-
Female.....	3	3	11	3	8	6	10	4	3	1	-	-
Black, both sexes.....	2	-	-	2	-	-	1	1	2	-	-	-
Male.....	1	-	-	2	-	-	-	1	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	-	-	2	1	-	2	3	-	-	-
Male.....	-	1	-	-	1	-	-	1	-	-	-	-
Female.....	-	1	-	-	1	1	-	1	3	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Florida

All races, both sexes.....	4	3	10	5	14	17	13	6	6	4	-	-
Male.....	1	3	3	2	8	8	9	4	-	1	-	-
Female.....	3	-	7	3	6	9	4	2	6	3	-	-
White, both sexes.....	1	2	9	5	10	17	13	6	4	4	-	-
Male.....	-	2	2	2	6	8	9	4	-	1	-	-
Female.....	1	-	7	3	4	9	4	2	4	3	-	-
Black, both sexes.....	3	1	1	-	4	-	-	-	2	-	-	-
Male.....	1	1	1	-	2	-	-	-	-	-	-	-
Female.....	2	-	-	-	2	-	-	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Georgia

All races, both sexes.....	7	6	8	11	9	14	14	10	1	-	-	-
Male.....	3	4	3	5	4	8	7	3	-	-	-	-
Female.....	4	2	5	6	5	6	7	7	1	-	-	-
White, both sexes.....	3	5	6	8	6	7	11	9	1	-	-	-
Male.....	2	3	3	4	3	5	5	3	-	-	-	-
Female.....	1	2	3	4	3	2	6	6	1	-	-	-
Black, both sexes.....	4	1	2	3	3	7	3	-	-	-	-	-
Male.....	1	1	-	1	1	3	2	-	-	-	-	-
Female.....	3	-	2	2	2	4	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

New Hampshire

All races, both sexes.....	1	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes.....	1	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	2	4	5	9	6	17	10	2	2	-	-
Male.....	-	1	1	5	3	1	7	5	1	-	-	-
Female.....	2	1	3	-	6	5	10	5	1	2	-	-
White, both sexes.....	-	2	2	4	7	6	13	8	2	1	-	-
Male.....	-	1	1	4	2	1	6	3	1	-	-	-
Female.....	-	1	1	-	5	5	7	5	1	1	-	-
Black, both sexes.....	2	-	2	1	2	-	3	2	-	1	-	-
Male.....	-	-	-	1	1	-	1	2	-	-	-	-
Female.....	2	-	2	-	1	-	2	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Wisconsin

All races, both sexes.....	2	-	-	5	4	2	4	1	2	-	-	-
Male.....	1	-	-	1	2	-	1	1	1	-	-	-
Female.....	1	-	-	4	2	2	3	-	1	-	-	-
White, both sexes.....	2	-	-	4	4	2	3	1	2	-	-	-
Male.....	1	-	-	1	2	-	1	1	1	-	-	-
Female.....	1	-	-	3	2	2	2	-	1	-	-	-
Black, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Wyoming

All races, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero.  
 ... Category not applicable.  
 1/ Includes Aleuts and Eskimos.  
 2/ Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.