

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### All causes

##### All origins 3/, both sexes

2007.....	803.6	684.5	28.6	15.3	79.9	104.9	184.4	420.9	877.7	2,011.3	5,011.6	12,946.5
2006.....	810.4	690.7	28.4	15.2	82.2	106.3	190.2	427.5	890.9	2,062.1	5,115.0	13,253.1
2005.....	825.9	692.5	29.4	16.3	81.4	104.4	193.3	432.0	906.9	2,137.1	5,260.0	13,798.6
2004.....	816.5	685.2	29.9	16.8	80.1	102.1	193.5	427.0	910.3	2,164.6	5,275.1	13,823.5
2003.....	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002.....	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001.....	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6

##### All origins 3/, male

2007.....	809.9	747.8	31.3	17.4	115.8	144.0	231.8	530.0	1,100.6	2,456.9	6,038.4	14,006.4
2006.....	814.8	756.3	30.5	17.6	119.3	146.8	238.7	541.0	1,110.0	2,516.2	6,177.7	14,309.1
2005.....	827.2	762.3	33.4	18.6	117.8	143.4	243.0	547.8	1,131.0	2,612.2	6,349.8	14,889.4
2004.....	817.6	753.7	32.4	19.2	114.7	139.5	243.6	543.5	1,128.8	2,644.8	6,394.3	15,031.1
2003.....	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002.....	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001.....	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000.....	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999.....	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2

##### All origins 3/, female

2007.....	797.4	618.1	25.7	13.1	42.0	64.2	136.9	315.2	670.1	1,633.0	4,304.1	12,442.3
2006.....	806.1	622.0	26.3	12.8	42.8	64.3	141.6	317.7	687.0	1,677.9	4,388.3	12,759.0
2005.....	824.6	619.4	25.1	13.9	42.7	64.1	143.6	319.9	698.5	1,736.3	4,520.0	13,297.7
2004.....	815.4	613.4	27.3	14.3	43.6	63.5	143.5	314.3	707.4	1,761.4	4,521.8	13,280.3
2003.....	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002.....	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001.....	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000.....	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999.....	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>All causes</b>												
<b>Hispanic, both sexes</b>												
2007.....	297.8	587.4	26.0	14.1	76.5	80.1	127.1	303.2	647.4	1,477.1	3,678.4	8,542.2
2006.....	300.1	590.6	26.4	14.2	80.4	81.5	134.4	310.7	657.6	1,531.0	3,745.7	9,022.8
2005.....	307.3	614.1	28.9	13.7	81.2	81.9	138.9	317.8	676.4	1,621.4	3,977.8	9,436.7
2004.....	296.2	586.9	27.3	14.4	76.5	78.2	138.7	313.5	684.0	1,607.5	3,926.7	9,483.8
2003.....	305.8	610.1	30.2	15.7	77.1	83.9	144.9	327.8	714.3	1,685.6	4,093.5	10,431.5
2002.....	302.2	592.7	29.8	15.5	77.2	79.3	145.8	321.3	719.8	1,734.1	4,143.1	10,679.0
2001.....	306.8	572.7	30.6	14.7	75.6	84.5	154.8	334.4	722.2	1,776.0	4,262.6	11,821.8
2000.....	303.8	596.3	29.6	15.7	72.8	84.6	157.6	330.2	744.4	1,803.8	4,335.9	11,822.9
1999.....	305.7	583.8	30.9	15.6	71.3	85.5	158.2	333.2	752.5	1,853.0	4,356.3	12,177.9
<b>Hispanic, male</b>												
2007.....	321.8	632.7	28.0	15.8	115.3	110.1	166.3	399.2	831.4	1,862.7	4,364.8	8,953.7
2006.....	323.9	640.7	28.8	16.4	120.7	112.7	176.5	403.8	843.6	1,910.7	4,492.6	9,435.5
2005.....	334.4	670.2	33.2	15.3	120.4	115.5	182.0	417.4	875.8	2,029.4	4,856.8	10,140.5
2004.....	321.1	636.5	30.2	16.7	115.3	109.2	184.2	417.8	874.0	1,994.3	4,791.6	9,932.8
2003.....	330.7	665.5	33.8	18.4	112.9	118.1	189.8	436.4	920.8	2,115.7	4,948.5	11,029.2
2002.....	328.7	644.0	34.2	17.4	114.4	112.5	192.5	423.4	937.4	2,193.4	5,043.5	11,674.1
2001.....	332.9	624.4	33.8	16.6	111.5	118.0	208.5	443.9	923.9	2,242.6	5,258.0	12,888.3
2000.....	331.3	637.1	31.5	17.9	107.7	120.2	211.0	439.0	965.7	2,287.9	5,395.3	13,086.2
1999.....	332.6	623.4	32.9	17.8	104.1	120.6	215.1	444.4	974.8	2,368.9	5,379.2	13,485.9
<b>Hispanic, female</b>												
2007.....	272.1	539.9	23.8	12.3	33.5	43.4	82.7	204.0	476.9	1,162.1	3,196.2	8,318.9
2006.....	274.6	538.3	24.0	11.8	35.2	43.1	87.1	215.3	486.5	1,222.7	3,222.9	8,803.5
2005.....	278.2	555.4	24.5	12.0	36.6	41.1	90.6	216.4	493.9	1,291.6	3,365.8	9,068.4
2004.....	269.7	535.1	24.3	12.0	31.8	40.9	88.2	208.3	511.3	1,297.2	3,329.6	9,253.0
2003.....	279.3	552.4	26.3	12.8	35.5	43.0	95.5	219.1	527.9	1,342.9	3,506.4	10,128.1
2002.....	274.0	539.1	25.3	13.5	34.1	40.0	94.9	219.8	524.3	1,368.7	3,526.4	10,186.0
2001.....	279.0	518.9	27.2	12.7	33.7	45.2	97.0	226.7	543.0	1,408.0	3,589.8	11,300.5
2000.....	274.6	553.6	27.5	13.4	31.7	43.4	100.5	223.8	548.4	1,423.2	3,624.5	11,202.8
1999.....	277.2	542.3	28.7	13.3	32.9	44.9	97.4	224.9	555.8	1,448.8	3,675.7	11,547.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>All causes</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	892.0	709.8	29.2	15.6	80.5	110.8	194.7	434.0	896.1	2,050.4	5,093.2	13,170.0
2006.....	897.1	714.6	28.9	15.5	82.3	112.1	199.5	439.7	908.7	2,099.0	5,192.4	13,450.1
2005.....	911.2	708.4	29.4	16.9	81.2	109.5	201.8	443.2	924.0	2,172.1	5,327.5	13,990.9
2004.....	899.4	705.6	30.5	17.2	80.6	107.3	201.7	437.4	926.1	2,200.1	5,340.3	13,997.7
2003.....	924.4	716.2	31.7	17.2	82.0	107.6	209.3	442.1	955.6	2,289.1	5,522.6	14,740.6
2002.....	928.8	713.0	31.4	17.8	81.9	108.4	210.3	438.8	967.2	2,347.0	5,610.8	14,958.7
2001.....	926.2	703.5	33.8	17.7	81.4	108.9	209.2	435.6	979.5	2,383.9	5,628.5	15,199.0
2000.....	929.6	760.4	32.9	18.4	80.9	104.1	202.9	431.8	1,006.5	2,427.9	5,707.6	15,619.8
1999.....	929.9	760.7	34.8	19.1	80.5	104.7	201.6	423.3	1,019.2	2,485.0	5,753.4	15,631.0
<b>Non-Hispanic 4/, male</b>												
2007.....	899.8	777.9	32.2	17.8	115.6	153.0	244.2	544.7	1,121.4	2,498.2	6,141.6	14,289.1
2006.....	902.8	784.5	30.8	17.8	118.6	155.7	249.7	555.3	1,129.5	2,555.7	6,274.0	14,558.8
2005.....	912.6	781.7	33.3	19.4	116.9	150.2	252.9	560.5	1,148.7	2,649.4	6,428.1	15,121.2
2004.....	900.9	778.7	32.9	19.6	114.2	146.6	252.8	554.7	1,145.4	2,683.5	6,472.4	15,260.4
2003.....	922.9	798.1	35.2	20.0	116.7	146.5	264.2	561.6	1,179.8	2,808.1	6,715.9	15,984.2
2002.....	928.0	782.6	35.4	20.4	117.3	148.6	266.0	557.0	1,198.1	2,889.1	6,825.9	16,418.0
2001.....	923.6	773.2	37.6	20.4	117.5	148.5	265.2	551.6	1,207.0	2,944.1	6,888.0	16,866.3
2000.....	928.1	836.8	36.8	21.5	115.9	141.6	259.2	548.7	1,244.2	3,009.8	7,020.7	17,639.8
1999.....	932.2	834.9	38.8	22.1	114.6	142.7	258.1	537.3	1,265.6	3,102.8	7,133.0	17,710.4
<b>Non-Hispanic 4/, female</b>												
2007.....	884.5	638.3	26.1	13.2	43.7	68.7	146.1	327.6	686.2	1,669.0	4,371.8	12,641.5
2006.....	891.7	641.4	26.9	13.0	44.3	68.8	150.3	328.5	703.2	1,711.4	4,453.8	12,934.8
2005.....	910.0	631.6	25.3	14.4	43.8	68.9	151.6	330.3	714.6	1,768.0	4,581.2	13,475.2
2004.....	898.0	628.9	28.1	14.7	45.7	68.1	151.5	324.5	722.2	1,792.8	4,579.4	13,433.1
2003.....	925.9	630.7	28.0	14.3	46.0	68.9	155.6	327.3	747.2	1,853.7	4,726.4	14,194.2
2002.....	929.5	640.1	27.2	15.0	45.3	68.5	155.7	325.2	753.0	1,894.4	4,805.4	14,329.1
2001.....	928.6	630.4	29.9	14.9	44.1	69.6	154.3	324.0	768.7	1,918.4	4,801.8	14,505.6
2000.....	931.0	680.1	28.8	15.2	45.0	66.8	147.7	319.2	786.8	1,947.5	4,852.4	14,802.9
1999.....	927.8	682.9	30.6	16.0	45.4	67.1	146.3	313.7	791.7	1,977.5	4,858.2	14,798.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### All causes

##### Non-Hispanic white, both sexes

2007 5/.....	964.1	559.7	25.5	14.2	74.5	102.4	181.5	403.8	849.6	2,006.2	5,113.1	13,413.3
2006 5/.....	968.5	564.2	25.0	14.0	76.1	102.1	184.8	406.6	861.2	2,055.0	5,205.2	13,688.2
2005 5/.....	981.8	562.6	26.2	15.2	74.8	98.4	186.7	407.1	873.5	2,126.0	5,330.0	14,222.4
2004 5/.....	967.8	564.4	26.8	15.5	75.2	95.9	185.9	400.8	875.1	2,152.5	5,337.1	14,191.1
2003 5/.....	993.6	576.5	27.6	15.6	75.8	95.8	190.9	402.3	902.4	2,237.2	5,515.4	14,944.8
2002.....	997.5	575.9	27.1	16.0	75.8	95.9	191.6	400.1	912.8	2,296.1	5,596.0	15,149.1
2001.....	991.1	555.4	30.1	16.3	74.1	96.4	188.3	396.0	924.0	2,325.5	5,604.4	15,371.2
2000.....	993.2	596.5	28.5	17.1	73.6	90.1	181.0	390.5	950.2	2,366.2	5,675.4	15,799.7
1999.....	990.7	593.1	30.1	17.3	72.4	91.3	177.8	381.5	957.8	2,420.0	5,716.0	15,795.3

##### Non-Hispanic white, male

2007 5/.....	960.4	616.8	28.1	16.1	104.6	140.8	228.4	508.7	1,057.5	2,432.7	6,152.7	14,588.3
2006 5/.....	962.0	621.9	26.7	16.2	107.6	141.1	233.1	515.1	1,064.0	2,490.3	6,278.3	14,841.1
2005 5/.....	970.6	625.7	29.9	17.4	105.8	134.1	236.1	517.2	1,079.6	2,584.5	6,420.4	15,401.3
2004 5/.....	957.4	625.2	29.0	17.9	104.8	130.9	235.7	510.5	1,076.4	2,617.9	6,461.5	15,489.2
2003 5/.....	979.1	647.2	30.2	18.1	105.9	129.9	243.8	513.2	1,110.5	2,738.5	6,692.2	16,234.4
2002.....	983.9	643.5	30.3	18.3	106.7	130.9	244.9	509.9	1,126.5	2,824.1	6,801.7	16,641.9
2001.....	975.6	611.6	33.7	18.6	105.1	131.5	241.6	502.6	1,136.3	2,869.4	6,851.5	17,055.3
2000.....	978.5	658.7	32.4	20.0	103.5	123.0	233.9	497.7	1,170.9	2,930.5	6,977.8	17,853.2
1999.....	979.6	658.1	33.4	19.9	100.8	124.5	230.0	483.7	1,187.4	3,023.2	7,088.0	17,871.2

##### Non-Hispanic white, female

2007 5/.....	967.6	499.6	22.7	12.3	42.7	63.4	134.4	300.5	651.3	1,634.9	4,385.4	12,856.7
2006 5/.....	974.7	503.7	23.2	11.8	42.9	62.5	136.3	299.8	668.0	1,677.4	4,460.7	13,150.7
2005 5/.....	992.6	496.5	22.2	12.9	42.2	62.1	137.0	298.7	677.2	1,729.6	4,579.7	13,683.1
2004 5/.....	977.7	500.7	24.4	13.0	44.3	60.3	136.0	292.9	683.8	1,752.0	4,571.1	13,609.6
2003 5/.....	1,007.6	502.5	24.9	12.9	44.3	61.2	138.0	293.1	705.0	1,807.9	4,720.5	14,377.2
2002.....	1,010.6	504.8	23.8	13.6	43.8	60.3	138.3	292.1	710.5	1,846.0	4,787.9	14,504.3
2001.....	1,006.1	496.4	26.3	13.9	41.9	60.9	134.9	291.0	723.5	1,864.1	4,777.3	14,670.6
2000.....	1,007.3	530.9	24.4	13.9	42.6	56.8	128.1	285.0	742.1	1,891.0	4,819.3	14,971.7
1999.....	1,001.3	524.6	26.7	14.7	42.9	57.8	125.6	281.0	741.6	1,915.1	4,817.7	14,967.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### All causes

##### Non-Hispanic black, both sexes

2007 5/.....	750.7	1,309.1	43.7	21.6	112.6	176.2	310.9	713.2	1,446.2	2,847.8	5,822.9	12,468.1
2006 5/.....	759.1	1,339.2	44.3	21.9	115.6	185.7	325.2	738.5	1,472.4	2,887.7	5,980.8	12,655.4
2005 5/.....	774.4	1,323.9	42.5	24.0	115.8	186.0	328.5	761.4	1,513.6	2,979.6	6,218.3	13,239.1
2004 5/.....	768.8	1,297.2	45.5	24.4	112.8	186.5	332.3	760.8	1,527.9	3,032.3	6,251.7	13,492.2
2003 5/.....	788.8	1,290.0	48.0	23.8	116.6	188.9	354.2	788.3	1,578.4	3,158.7	6,433.0	14,107.7
2002.....	792.8	1,280.3	48.4	25.3	116.9	193.9	357.8	782.0	1,592.9	3,204.8	6,584.8	14,386.7
2001.....	798.1	1,354.4	49.5	24.0	121.4	191.5	365.8	786.9	1,602.5	3,300.7	6,645.3	14,660.8
2000.....	805.5	1,477.8	51.7	25.0	123.2	194.6	365.1	802.2	1,633.9	3,378.8	6,764.3	14,837.5
1999.....	812.1	1,510.2	54.6	27.5	125.5	191.1	376.9	796.7	1,681.7	3,454.5	6,805.4	14,839.2

##### Non-Hispanic black, male

2007 5/.....	805.1	1,426.3	47.5	25.5	173.1	250.6	393.4	900.4	1,913.7	3,678.3	7,295.1	13,189.0
2006 5/.....	815.3	1,453.3	48.2	25.8	176.9	266.0	405.6	944.2	1,935.2	3,736.8	7,520.4	13,403.1
2005 5/.....	825.7	1,451.5	47.6	27.8	177.7	264.0	407.5	969.9	1,993.8	3,814.1	7,771.1	13,978.1
2004 5/.....	818.7	1,427.6	49.5	26.9	169.4	262.1	409.3	974.9	1,998.6	3,880.2	7,813.2	14,647.3
2003 5/.....	840.6	1,427.7	55.1	27.9	176.5	266.6	440.0	1,012.8	2,047.7	4,041.1	8,165.7	15,082.2
2002.....	842.3	1,369.3	55.8	29.8	177.4	274.3	445.7	1,002.9	2,074.5	4,078.9	8,246.2	15,776.9
2001.....	849.7	1,484.2	55.1	28.6	186.2	267.9	456.3	1,014.6	2,055.4	4,218.9	8,426.8	16,576.2
2000.....	859.5	1,623.2	56.1	29.1	187.2	270.2	462.7	1,036.2	2,111.4	4,303.9	8,551.0	16,944.8
1999.....	872.8	1,650.8	61.5	33.1	191.4	267.9	479.5	1,048.1	2,176.7	4,397.8	8,615.3	17,385.3

##### Non-Hispanic black, female

2007 5/.....	701.0	1,187.1	39.8	17.6	50.5	106.7	237.9	552.5	1,070.7	2,255.0	4,985.7	12,169.3
2006 5/.....	708.0	1,220.1	40.3	17.9	52.8	111.4	254.1	562.2	1,100.2	2,281.4	5,107.2	12,350.5
2005 5/.....	727.6	1,190.8	37.2	20.0	52.6	114.1	258.5	582.4	1,127.1	2,383.1	5,338.1	12,941.6
2004 5/.....	723.4	1,161.6	41.3	21.8	55.3	117.2	264.1	577.0	1,149.9	2,427.9	5,370.3	13,035.6
2003 5/.....	741.6	1,146.9	40.7	19.6	56.0	118.0	278.4	595.7	1,201.6	2,531.2	5,454.6	13,730.3
2002.....	748.0	1,187.5	40.8	20.7	56.1	121.3	280.1	592.7	1,206.3	2,584.8	5,644.9	13,855.4
2001.....	751.2	1,217.8	43.6	19.3	56.8	122.4	285.7	591.5	1,238.2	2,653.0	5,645.1	13,951.4
2000.....	756.7	1,327.7	47.1	20.7	60.1	126.5	278.8	601.2	1,249.4	2,728.0	5,758.6	14,054.9
1999.....	757.3	1,365.4	47.4	21.8	60.5	122.1	286.4	581.7	1,285.1	2,789.2	5,792.0	13,887.7





















Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

All origins 3/, both sexes

2007.....	2.2	*	*	*	*	*	0.1	0.3	1.0	4.7	18.8	52.0
2006.....	2.2	*	*	*	*	*	0.1	0.3	1.1	4.8	19.0	50.5
2005.....	1.9	*	*	*	*	*	0.1	0.2	0.8	4.4	15.9	45.8
2004.....	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.8	34.3
2003.....	1.1	*	*	*	*	*	*	0.2	0.6	2.4	9.3	24.8
2002.....	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3

All origins 3/, male

2007.....	1.7	*	*	*	*	*	*	0.2	1.1	4.6	18.3	50.4
2006.....	1.7	*	*	*	*	*	0.1	0.3	1.3	4.8	18.6	50.0
2005.....	1.5	*	*	*	*	*	*	0.3	0.9	4.4	16.4	46.1
2004.....	1.1	*	*	*	*	*	*	0.2	0.9	3.1	12.7	32.0
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.3	21.4
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5

All origins 3/, female

2007.....	2.8	*	*	*	*	*	0.1	0.3	0.9	4.9	19.1	52.7
2006.....	2.7	*	*	*	*	*	*	0.3	1.0	4.8	19.4	50.8
2005.....	2.3	*	*	*	*	*	*	0.2	0.8	4.5	15.5	45.6
2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.4	12.9	35.4
2003.....	1.3	*	*	*	*	*	*	0.2	0.6	2.6	9.3	26.3
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 4/, both sexes

2007.....	2.5	*	*	*	*	*	0.1	0.3	1.0	4.8	19.2	52.9
2006.....	2.5	*	*	*	*	*	0.1	0.3	1.1	4.9	19.2	51.8
2005.....	2.2	*	*	*	*	*	0.1	0.2	0.9	4.6	16.2	46.7
2004.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.4	13.1	35.0
2003.....	1.2	*	*	*	*	*	*	0.2	0.6	2.5	9.5	25.2
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.1	21.4
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.9
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.4

Non-Hispanic 4/, male

2007.....	1.9	*	*	*	*	*	*	0.3	1.1	4.7	18.6	51.4
2006.....	2.0	*	*	*	*	*	*	0.3	1.2	4.9	18.9	51.2
2005.....	1.7	*	*	*	*	*	*	0.3	1.0	4.6	16.9	47.4
2004.....	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.1	32.4
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.3	9.5	21.4
2002.....	0.8	*	*	*	*	*	*	0.1	0.6	2.3	7.2	21.0
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.7	12.0
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.5
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.6	9.6

Non-Hispanic 4/, female

2007.....	3.1	*	*	*	*	*	*	0.3	0.9	5.0	19.7	53.6
2006.....	3.0	*	*	*	*	*	*	0.3	1.0	4.9	19.5	52.1
2005.....	2.6	*	*	*	*	*	*	0.2	0.8	4.6	15.8	46.4
2004.....	2.0	*	*	*	*	*	*	0.2	0.8	3.5	13.1	36.1
2003.....	1.5	*	*	*	*	*	*	0.2	0.7	2.7	9.5	26.8
2002.....	1.1	*	*	*	*	*	*	0.1	0.5	1.8	7.0	21.6
2001.....	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.5	13.2
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	12.1
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic white, both sexes

2007 5/.....	3.0	*	*	*	*	*	0.1	0.3	1.0	5.0	20.4	56.1
2006 5/.....	2.9	*	*	*	*	*	*	0.3	1.1	5.2	20.6	55.1
2005 5/.....	2.5	*	*	*	*	*	0.1	0.2	0.8	4.9	17.2	49.7
2004 5/.....	2.0	*	*	*	*	*	0.1	0.2	0.9	3.6	13.8	37.3
2003 5/.....	1.4	*	*	*	*	*	*	0.1	0.6	2.7	10.0	26.5
2002.....	1.1	*	*	*	*	*	*	0.2	0.5	2.2	7.5	22.5
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.4	4.7	13.6
2000.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.3
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.4	11.0

Non-Hispanic white, male

2007 5/.....	2.3	*	*	*	*	*	*	0.3	1.1	4.7	19.5	54.7
2006 5/.....	2.3	*	*	*	*	*	*	0.3	1.2	5.1	19.9	54.4
2005 5/.....	2.0	*	*	*	*	*	*	0.3	1.0	4.8	17.5	50.9
2004 5/.....	1.5	*	*	*	*	*	*	0.2	0.9	3.3	13.8	34.5
2003 5/.....	1.0	*	*	*	*	*	*	0.2	0.6	2.4	10.1	22.7
2002.....	0.9	*	*	*	*	*	*	0.2	0.5	2.3	7.5	22.1
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.4	4.7	12.6
2000.....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.3	12.0
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.8	10.0

Non-Hispanic white, female

2007 5/.....	3.7	*	*	*	*	*	*	0.3	0.9	5.3	21.0	56.7
2006 5/.....	3.6	*	*	*	*	*	*	0.3	1.0	5.2	21.0	55.4
2005 5/.....	3.0	*	*	*	*	*	*	0.2	0.7	4.9	16.9	49.2
2004 5/.....	2.4	*	*	*	*	*	*	0.2	0.9	3.8	13.9	38.6
2003 5/.....	1.8	*	*	*	*	*	*	0.1	0.7	2.9	9.9	28.2
2002.....	1.3	*	*	*	*	*	*	0.1	0.5	2.0	7.5	22.7
2001.....	0.9	*	*	*	*	*	*	0.2	0.3	1.3	4.7	14.0
2000.....	0.7	*	*	*	*	*	*	*	0.3	1.2	3.9	12.5
1999.....	0.6	*	*	*	*	*	*	*	*	1.0	3.2	11.4



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Tuberculosis (A16-A19)

All origins 3/, both sexes

2007.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.2	1.6
2006.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.3	2.1
2005.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.3	2.2
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.4
2003.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001.....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2

All origins 3/, male

2007.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.5	2.8
2006.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	1.8	3.1
2005.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	1.7	3.3
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.4	1.1	1.6	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999.....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1

All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	0.1	0.4	0.9	1.1
2006.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.9	1.6
2005.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7
2004.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Tuberculosis (A16-A19)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.5
2006.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.0
2005.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.2
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.7	1.1	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.6	2.2
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.8	2.5
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.7
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.9	2.5
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9	3.9

Non-Hispanic 4/, male

2007.....	0.2	*	*	*	*	*	*	0.3	0.4	0.6	1.4	2.3
2006.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.8	1.8	2.8
2005.....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.6	3.2
2004.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.3	3.5
2003.....	0.3	*	*	*	*	*	*	0.4	0.5	0.8	2.1	3.0
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.2	2.3	3.8
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	4.3
2000.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.6	4.3
1999.....	0.4	*	*	*	*	*	0.2	0.4	0.8	1.3	2.6	6.8

Non-Hispanic 4/, female

2007.....	0.1	*	*	*	*	*	*	*	*	0.4	0.8	1.1
2006.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.6
2005.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7
2004.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.6
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	1.9
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.7	1.4	1.8
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Tuberculosis (A16-A19)

Non-Hispanic white, both sexes

2007 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.7	1.4
2006 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.9	1.5
2005 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.8	1.6
2004 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5
2003 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	1.2	1.7
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.0
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0

Non-Hispanic white, male

2007 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.1
2006 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.5	1.2	1.7
2005 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.3
2004 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.5	0.9	2.4
2003 5/.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.5	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.3	0.8	1.7	2.9
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.4	0.6	1.4	2.9
2000.....	0.2	*	*	*	*	*	*	*	0.2	*	0.6	1.5	3.6
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.4	0.8	1.8	4.9

Non-Hispanic white, female

2007 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.2	0.6	1.0
2006 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	0.6	1.3
2005 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.6	1.3
2004 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	0.7	1.1
2003 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	1.0	1.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	1.1	1.4
2001.....	0.2	*	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	1.1	1.4
1999.....	0.2	*	*	*	*	*	*	*	*	*	0.3	1.0	2.2



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Respiratory tuberculosis (A16)

##### All origins 3/, both sexes

2007.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.3
2006.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.4	1.0	1.6
2005.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.9
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.9	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7

##### All origins 3/, male

2007.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.2	2.4
2006.....	0.2	*	*	*	*	*	0.1	0.3	0.5	0.7	1.5	2.4
2005.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.7
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.2	3.7
2003.....	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001.....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4

##### All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
2006.....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	1.2
2005.....	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.5
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Respiratory tuberculosis (A16)

Non-Hispanic 4/, both sexes

2007.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	0.8	1.2
2006.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.5
2005.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.8	2.0	
2003.....	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.3	1.7	
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.5	2.2	
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.4	
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.6	2.2	
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.5	

Non-Hispanic 4/, male

2007.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.4	1.0	2.0
2006.....	0.2	*	*	*	*	*	*	*	0.2	0.4	0.6	1.5	2.2
2005.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.7
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.8	1.0	3.3	
2003.....	0.2	*	*	*	*	*	*	0.3	0.4	0.6	1.8	2.4	
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.3	
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	0.8	1.6	3.8	
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.2	3.8	
1999.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	2.3	6.1	

Non-Hispanic 4/, female

2007.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	0.8	
2006.....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.2	
2005.....	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.4	
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.3	
2003.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.5	
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.7	
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8	
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.5	
1999.....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4	

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Respiratory tuberculosis (A16)

Non-Hispanic white, both sexes

2007 5/.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.1
2006 5/.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.7	1.1
2005 5/.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.6	1.3
2004 5/.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4
2003 5/.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.3
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.6
2001.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
2000.....	0.1	*	*	*	*	*	*	0.1	*	0.3	1.1	1.8
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8

Non-Hispanic white, male

2007 5/.....	0.1	*	*	*	*	*	*	*	0.2	*	0.6	1.8
2006 5/.....	0.1	*	*	*	*	*	*	*	0.2	0.3	1.0	*
2005 5/.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.8
2004 5/.....	0.1	*	*	*	*	*	*	*	*	0.5	0.7	2.2
2003 5/.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.2	1.8
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.2	2.7
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.4	1.1	2.4
2000.....	0.2	*	*	*	*	*	*	0.2	*	0.5	1.3	3.2
1999.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	4.8

Non-Hispanic white, female

2007 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
2006 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.0
2005 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.0
2004 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.0
2003 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.8	1.1
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9

































Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningococcal infection (A39)

All origins 3/, both sexes

2007.....	0.0	*	*	*	0.0	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000.....	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999.....	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*

All origins 3/, male

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	*

All origins 3/, female

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*









Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Septicemia (A40-A41)

All origins 3/, both sexes

2007.....	11.5	6.6	0.5	0.2	0.4	0.7	2.1	5.5	12.9	32.8	79.9	174.4
2006.....	11.4	6.5	0.5	0.2	0.3	0.7	2.0	5.2	12.8	32.1	82.4	177.3
2005.....	11.5	7.4	0.5	0.2	0.4	0.8	1.9	5.2	12.9	32.6	81.4	187.3
2004.....	11.4	6.6	0.5	0.2	0.3	0.8	1.9	5.4	12.9	32.4	81.6	186.7
2003.....	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001.....	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7

All origins 3/, male

2007.....	10.7	7.0	0.5	0.2	0.4	0.7	2.1	6.0	14.0	36.6	89.7	189.3
2006.....	10.5	6.8	0.5	0.3	0.3	0.7	2.1	5.9	14.1	34.8	92.7	194.8
2005.....	10.5	8.1	0.6	0.2	0.4	0.7	1.9	5.6	14.2	36.3	91.4	201.9
2004.....	10.4	8.0	0.5	0.1	0.3	0.8	2.0	5.9	13.8	35.8	93.3	203.9
2003.....	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002.....	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001.....	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000.....	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999.....	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5

All origins 3/, female

2007.....	12.4	6.3	0.4	0.1	0.4	0.7	2.1	5.1	12.0	29.5	73.1	167.3
2006.....	12.3	6.2	0.6	0.2	0.3	0.7	1.9	4.6	11.5	29.7	75.3	169.2
2005.....	12.5	6.5	0.5	0.2	0.3	0.8	1.9	4.8	11.7	29.5	74.6	180.6
2004.....	12.3	5.2	0.6	0.2	0.3	0.8	1.9	4.9	12.0	29.6	73.7	179.0
2003.....	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002.....	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001.....	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000.....	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999.....	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Septicemia (A40-A41)</b>												
<b>Hispanic, both sexes</b>												
2007.....	4.2	6.3	*	*	*	0.5	1.2	3.8	9.4	26.2	62.1	124.1
2006.....	4.1	6.9	*	*	0.3	0.5	1.2	4.0	9.9	24.9	65.1	122.9
2005.....	4.0	7.1	*	*	*	0.5	1.2	3.7	9.7	25.5	61.0	134.8
2004.....	4.0	6.1	0.7	*	*	0.5	1.4	4.0	9.7	25.1	63.8	138.0
2003.....	3.8	5.8	*	*	*	0.5	1.5	3.9	9.1	22.1	62.0	151.8
2002.....	3.9	7.8	*	*	*	0.5	1.2	3.7	8.4	28.9	63.7	156.3
2001.....	3.7	6.1	*	*	*	0.3	1.3	3.9	8.2	25.4	62.4	171.3
2000.....	3.5	8.0	*	*	*	0.5	1.5	3.9	8.5	21.8	59.3	144.7
1999.....	3.6	5.7	*	*	*	0.4	1.5	3.1	8.0	25.6	61.7	180.3
<b>Hispanic, male</b>												
2007.....	4.0	6.6	*	*	*	*	1.3	4.6	11.1	31.4	73.8	113.8
2006.....	4.1	7.5	*	*	*	0.5	1.2	4.9	11.3	30.5	76.5	138.7
2005.....	3.8	7.8	*	*	*	0.5	1.0	4.3	10.6	29.6	74.2	137.0
2004.....	3.8	7.8	*	*	*	*	1.6	5.2	10.3	27.2	69.7	147.1
2003.....	3.5	6.6	*	*	*	0.5	1.5	4.9	9.6	25.3	67.1	160.1
2002.....	3.7	8.0	*	*	*	0.6	1.6	5.2	9.0	30.8	72.2	164.0
2001.....	3.6	7.0	*	*	*	*	1.5	4.7	9.9	29.8	76.0	160.0
2000.....	3.4	8.4	*	*	*	0.6	1.7	5.0	10.0	26.6	70.3	157.2
1999.....	3.5	7.0	*	*	*	*	2.0	3.8	10.2	29.4	75.1	181.2
<b>Hispanic, female</b>												
2007.....	4.3	5.9	*	*	*	0.6	1.0	3.0	7.8	22.0	53.9	129.6
2006.....	4.2	6.3	*	*	*	*	1.1	3.0	8.5	20.3	57.1	114.4
2005.....	4.2	6.4	*	*	*	*	1.3	3.0	8.8	22.2	51.9	133.6
2004.....	4.2	*	*	*	*	*	1.2	2.8	9.2	23.4	59.8	133.2
2003.....	4.1	5.0	*	*	*	*	1.3	2.8	8.7	19.6	58.5	147.7
2002.....	4.1	7.6	*	*	*	*	0.7	2.3	7.8	27.4	57.9	152.5
2001.....	3.9	5.2	*	*	*	*	1.1	3.1	6.6	21.9	53.2	176.9
2000.....	3.5	7.7	*	*	*	*	1.3	2.8	7.2	18.1	51.8	138.5
1999.....	3.7	*	*	*	*	*	1.0	2.4	6.1	22.5	52.7	179.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Septicemia (A40-A41)

##### Non-Hispanic 4/, both sexes

2007.....	12.8	6.7	0.5	0.2	0.4	0.8	2.3	5.7	13.2	33.2	80.9	176.8
2006.....	12.7	6.3	0.6	0.2	0.3	0.8	2.1	5.4	13.0	32.6	83.3	179.7
2005.....	12.8	7.4	0.6	0.2	0.4	0.8	2.0	5.4	13.1	33.0	82.5	189.6
2004.....	12.5	6.7	0.5	0.2	0.4	0.9	2.0	5.5	13.1	32.9	82.5	188.7
2003.....	12.9	7.2	0.6	0.2	0.4	0.8	2.1	5.4	13.3	33.2	86.0	204.1
2002.....	12.9	7.1	0.5	0.2	0.3	0.8	2.0	5.3	12.9	34.9	87.3	204.2
2001.....	12.4	8.1	0.7	0.1	0.4	0.8	1.9	5.1	12.5	33.1	83.0	206.8
2000.....	12.1	6.9	0.7	0.2	0.3	0.7	2.0	5.0	12.1	31.4	81.0	217.5
1999.....	12.0	7.9	0.6	0.2	0.3	0.7	1.8	4.7	11.6	31.4	79.8	221.2

##### Non-Hispanic 4/, male

2007.....	11.9	7.0	0.6	0.2	0.4	0.8	2.3	6.2	14.2	37.0	90.7	193.2
2006.....	11.7	6.5	0.6	0.3	0.3	0.8	2.2	5.9	14.3	35.1	93.7	197.5
2005.....	11.7	8.2	0.6	0.3	0.4	0.8	2.0	5.8	14.4	36.6	92.2	205.3
2004.....	11.5	7.9	0.4	0.1	0.4	0.9	2.0	5.9	14.0	36.3	94.6	206.3
2003.....	11.6	7.6	0.8	0.2	0.4	0.9	2.1	5.8	14.3	36.8	96.2	215.9
2002.....	11.6	7.3	0.5	0.3	0.3	0.9	2.0	5.7	14.0	38.2	99.8	215.6
2001.....	11.2	8.8	0.9	0.1	0.4	0.8	2.1	5.5	13.4	37.0	96.1	226.8
2000.....	10.7	6.7	0.7	0.2	0.3	0.8	2.0	5.6	13.1	34.8	92.3	224.8
1999.....	10.7	9.4	0.6	0.2	0.4	0.8	2.1	5.2	12.5	35.4	92.2	234.5

##### Non-Hispanic 4/, female

2007.....	13.7	6.4	0.4	0.1	0.4	0.8	2.3	5.3	12.3	30.0	74.2	169.0
2006.....	13.6	6.2	0.5	0.2	0.4	0.7	2.0	4.8	11.8	30.4	76.2	171.4
2005.....	13.8	6.6	0.5	0.2	0.3	0.9	2.0	5.0	11.9	29.9	75.9	182.5
2004.....	13.5	5.4	0.5	0.2	0.3	0.9	2.0	5.1	12.2	30.0	74.3	180.8
2003.....	14.2	6.8	0.4	0.2	0.4	0.8	2.1	5.0	12.4	30.1	79.1	199.0
2002.....	14.1	6.8	0.5	0.2	0.3	0.8	2.0	4.9	11.9	32.1	79.0	199.3
2001.....	13.5	7.4	0.6	0.2	0.3	0.8	1.7	4.8	11.8	29.9	74.4	198.5
2000.....	13.5	7.0	0.6	0.1	0.2	0.7	1.9	4.3	11.2	28.7	73.6	214.5
1999.....	13.2	6.2	0.5	0.2	0.3	0.7	1.6	4.2	10.8	28.2	71.8	215.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Septicemia (A40-A41)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	12.9	4.2	0.5	0.2	0.3	0.6	1.9	4.6	11.4	29.3	76.2	169.3
2006 5/.....	12.7	4.0	0.5	0.2	0.3	0.6	1.7	4.2	11.0	29.0	78.4	173.2
2005 5/.....	12.7	4.3	0.5	0.2	0.3	0.6	1.6	4.3	11.2	29.8	76.8	179.2
2004 5/.....	12.5	4.8	0.4	0.2	0.3	0.7	1.6	4.4	11.1	29.3	77.2	179.6
2003 5/.....	12.9	4.2	0.5	0.2	0.3	0.6	1.7	4.3	11.3	29.3	80.8	192.4
2002.....	12.9	5.5	0.5	0.2	0.3	0.7	1.6	4.3	10.7	31.0	81.3	192.6
2001.....	12.3	5.1	0.7	0.1	0.3	0.6	1.4	4.0	10.4	29.4	77.6	195.6
2000.....	11.9	4.8	0.5	0.2	0.2	0.5	1.5	3.8	10.0	27.7	74.2	204.3
1999.....	11.7	5.3	0.5	0.2	0.3	0.6	1.4	3.6	9.5	27.3	73.0	206.0
<b>Non-Hispanic white, male</b>												
2007 5/.....	11.8	4.4	0.6	0.2	0.4	0.7	1.9	4.8	12.0	32.3	86.5	185.4
2006 5/.....	11.7	4.1	0.5	0.2	0.2	0.6	1.8	4.8	11.9	31.5	88.5	193.3
2005 5/.....	11.5	5.3	0.6	0.2	0.3	0.6	1.6	4.6	12.3	32.7	86.2	193.6
2004 5/.....	11.4	5.7	*	0.2	0.3	0.7	1.6	4.8	11.7	32.4	88.5	198.5
2003 5/.....	11.4	3.9	0.6	0.2	0.4	0.6	1.7	4.6	12.0	32.5	90.5	204.8
2002.....	11.5	5.6	0.5	0.3	0.3	0.7	1.6	4.6	11.8	34.0	92.1	203.9
2001.....	11.0	5.8	0.9	*	0.3	0.5	1.7	4.2	11.1	32.5	89.7	213.6
2000.....	10.4	4.7	0.5	0.2	0.3	0.6	1.6	4.3	10.9	30.9	84.7	211.2
1999.....	10.2	6.7	0.5	0.2	0.3	0.7	1.5	4.0	9.9	30.5	84.6	218.1
<b>Non-Hispanic white, female</b>												
2007 5/.....	13.9	4.1	*	*	0.3	0.6	1.8	4.3	10.8	26.7	69.0	161.7
2006 5/.....	13.7	3.9	0.4	*	0.3	0.6	1.6	3.7	10.1	26.8	71.4	163.9
2005 5/.....	13.9	3.3	*	*	0.2	0.7	1.6	4.0	10.1	27.2	70.3	172.6
2004 5/.....	13.6	3.9	0.5	0.2	0.2	0.7	1.5	4.1	10.4	26.7	69.5	171.1
2003 5/.....	14.3	4.5	*	0.2	0.3	0.7	1.7	3.9	10.6	26.6	74.3	187.0
2002.....	14.2	5.4	*	0.2	0.3	0.6	1.7	3.9	9.7	28.5	74.0	187.7
2001.....	13.6	4.4	0.5	*	0.3	0.6	1.2	3.8	9.9	26.8	69.6	188.1
2000.....	13.4	4.8	0.6	*	*	0.5	1.5	3.3	9.2	24.9	67.2	201.6
1999.....	13.1	3.8	0.5	0.2	0.2	0.5	1.2	3.3	9.1	24.5	65.4	201.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Septicemia (A40-A41)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	16.4	16.7	*	*	0.7	1.6	5.1	14.0	30.5	73.5	149.6	299.4
2006 5/.....	16.0	15.6	*	0.3	0.6	1.6	5.0	13.4	31.6	70.2	153.3	290.9
2005 5/.....	16.5	19.9	0.8	0.3	0.7	2.0	4.7	13.4	31.5	68.2	159.9	340.8
2004 5/.....	16.1	14.2	*	*	0.7	1.9	4.6	13.6	32.2	71.4	157.9	324.7
2003 5/.....	16.8	18.8	1.0	*	0.7	2.0	5.1	14.0	32.8	74.0	160.0	373.9
2002.....	16.8	13.1	*	0.3	0.4	2.0	4.6	13.2	33.8	76.0	172.9	370.8
2001.....	16.4	20.3	0.9	*	0.7	1.9	4.8	13.7	32.6	72.8	161.9	368.6
2000.....	16.7	16.6	1.1	*	0.6	2.1	4.8	13.9	31.8	72.5	175.1	395.9
1999.....	16.6	18.6	*	*	0.6	1.6	5.1	13.2	30.8	75.3	169.8	414.2
<b>Non-Hispanic black, male</b>												
2007 5/.....	15.4	17.9	*	*	0.7	1.5	5.0	15.9	36.0	88.4	165.0	336.8
2006 5/.....	14.9	16.3	*	*	0.7	1.7	5.2	15.1	37.9	77.2	179.6	300.5
2005 5/.....	15.5	20.2	*	*	0.8	1.9	4.7	15.2	36.6	81.9	183.3	379.7
2004 5/.....	15.1	15.8	*	*	0.7	2.1	4.8	15.2	37.1	82.9	192.5	346.5
2003 5/.....	15.8	21.7	*	*	*	2.3	5.2	15.8	38.7	86.9	191.3	409.1
2002.....	15.5	13.9	*	*	*	1.9	5.0	15.0	36.6	87.6	213.7	399.3
2001.....	15.4	21.4	*	*	0.9	1.9	4.9	15.2	37.4	86.9	196.8	430.8
2000.....	15.3	16.4	*	*	*	2.4	4.8	16.8	35.9	81.6	208.9	432.1
1999.....	15.7	20.8	*	*	*	1.7	6.0	15.4	37.9	91.5	202.6	451.6
<b>Non-Hispanic black, female</b>												
2007 5/.....	17.2	15.5	*	*	0.8	1.8	5.1	12.3	26.1	62.9	140.8	283.9
2006 5/.....	17.0	14.8	*	*	0.6	1.6	4.8	12.0	26.5	65.3	138.4	287.0
2005 5/.....	17.4	19.5	*	*	0.6	2.1	4.7	11.8	27.3	58.4	146.7	325.2
2004 5/.....	17.0	12.6	*	*	0.8	1.8	4.5	12.3	28.3	63.1	138.3	316.0
2003 5/.....	17.7	15.8	*	*	0.7	1.7	5.1	12.5	28.2	64.9	142.3	360.3
2002.....	18.0	12.3	*	*	*	2.0	4.2	11.6	31.6	67.8	149.9	359.9
2001.....	17.2	19.1	*	*	*	1.9	4.7	12.5	28.8	62.8	142.4	345.6
2000.....	17.9	16.9	*	*	*	1.8	4.9	11.5	28.5	66.1	156.1	382.5
1999.....	17.4	16.4	*	*	*	1.6	4.3	11.3	25.1	63.9	151.4	400.2









































Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

All origins 3/, both sexes

2007.....	2.5	*	*	*	*	0.2	1.1	6.4	7.4	4.5	4.4	2.9
2006.....	2.4	*	*	*	*	0.2	1.2	6.7	6.7	4.5	4.6	3.1
2005.....	1.9	*	*	*	*	0.2	1.2	5.4	4.5	3.5	3.7	2.4
2004.....	1.8	*	*	*	*	0.2	1.4	5.3	4.3	3.4	3.6	2.7
2003.....	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001.....	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000.....	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999.....	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7

All origins 3/, male

2007.....	3.3	*	*	*	*	0.2	1.5	9.2	11.2	5.2	4.7	3.0
2006.....	3.2	*	*	*	*	0.2	1.7	9.9	9.9	5.0	4.6	3.0
2005.....	2.5	*	*	*	*	0.2	1.6	8.1	6.7	3.7	3.7	2.6
2004.....	2.4	*	*	*	*	0.2	1.8	8.0	6.1	4.0	3.6	3.0
2003.....	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002.....	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001.....	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000.....	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9

All origins 3/, female

2007.....	1.6	*	*	*	*	0.1	0.8	3.7	3.8	3.9	4.2	2.9
2006.....	1.6	*	*	*	*	0.1	0.8	3.7	3.8	4.0	4.6	3.1
2005.....	1.3	*	*	*	*	0.1	0.7	2.9	2.5	3.3	3.7	2.3
2004.....	1.3	*	*	*	*	0.1	1.0	2.7	2.5	3.0	3.5	2.5
2003.....	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002.....	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001.....	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000.....	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999.....	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Hispanic, both sexes

2007.....	2.5	*	*	*	*	*	1.5	8.3	13.3	11.0	9.0	*
2006.....	2.5	*	*	*	*	*	1.6	9.4	13.0	10.8	10.2	8.6
2005.....	1.9	*	*	*	*	*	1.8	6.9	8.5	8.6	6.6	*
2004.....	1.7	*	*	*	*	*	1.8	6.8	7.7	7.0	7.5	*
2003.....	1.8	*	*	*	*	*	1.8	7.0	7.2	8.7	9.2	*
2002.....	2.0	*	*	*	*	*	1.7	8.0	8.3	8.6	10.5	*
2001.....	2.1	*	*	*	*	*	2.3	7.8	10.4	9.9	11.0	*
2000.....	2.0	*	*	*	*	*	2.5	7.0	8.6	11.1	13.0	*
1999.....	1.9	*	*	*	*	*	2.1	6.8	7.8	12.4	13.5	*

Hispanic, male

2007.....	3.2	*	*	*	*	*	2.2	12.0	20.0	12.8	8.2	*
2006.....	3.2	*	*	*	*	*	2.5	13.9	17.6	12.0	8.2	*
2005.....	2.4	*	*	*	*	*	2.5	10.3	11.4	9.0	*	*
2004.....	2.2	*	*	*	*	*	2.6	10.2	10.4	8.0	*	*
2003.....	2.3	*	*	*	*	*	2.6	10.8	10.4	7.8	7.7	*
2002.....	2.5	*	*	*	*	*	2.5	12.0	11.6	7.6	10.3	*
2001.....	2.6	*	*	*	*	*	3.1	12.2	14.2	7.0	10.3	*
2000.....	2.5	*	*	*	*	*	3.8	10.6	11.7	11.0	12.3	*
1999.....	2.3	*	*	*	*	*	3.3	10.5	9.8	13.0	*	*

Hispanic, female

2007.....	1.7	*	*	*	*	*	0.7	4.4	7.1	9.5	9.6	*
2006.....	1.8	*	*	*	*	*	*	4.7	8.7	9.7	11.5	*
2005.....	1.3	*	*	*	*	*	1.0	3.4	5.8	8.2	6.7	*
2004.....	1.2	*	*	*	*	*	0.9	3.4	5.2	6.2	9.1	*
2003.....	1.3	*	*	*	*	*	1.0	3.1	4.4	9.4	10.3	*
2002.....	1.4	*	*	*	*	*	0.9	4.0	5.4	9.4	10.7	*
2001.....	1.7	*	*	*	*	*	1.4	3.5	7.1	12.3	11.5	*
2000.....	1.5	*	*	*	*	*	1.0	3.3	5.8	11.3	13.5	*
1999.....	1.5	*	*	*	*	*	0.8	3.3	6.0	12.0	16.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Non-Hispanic 4/, both sexes

2007.....	2.4	*	*	*	*	0.2	1.1	6.2	6.8	4.0	4.1	2.8
2006.....	2.4	*	*	*	*	0.1	1.2	6.4	6.2	3.9	4.2	2.8
2005.....	1.9	*	*	*	*	0.2	1.0	5.3	4.2	3.1	3.5	2.2
2004.....	1.8	*	*	*	*	0.2	1.3	5.2	4.0	3.2	3.3	2.6
2003.....	1.9	*	*	*	*	0.2	1.4	5.4	3.6	3.4	4.0	2.3
2002.....	2.0	*	*	*	*	0.2	1.6	5.6	3.6	3.8	4.3	3.7
2001.....	1.9	*	*	*	*	0.2	1.7	5.0	3.5	4.2	4.6	3.6
2000.....	1.9	*	*	*	*	0.2	1.6	4.8	3.3	4.2	5.0	3.8
1999.....	1.7	*	*	*	*	0.2	1.5	3.8	3.0	4.3	5.0	4.7

Non-Hispanic 4/, male

2007.....	3.3	*	*	*	*	0.2	1.4	8.9	10.3	4.6	4.5	2.9
2006.....	3.2	*	*	*	*	0.2	1.5	9.3	9.2	4.4	4.4	3.0
2005.....	2.5	*	*	*	*	0.2	1.4	7.8	6.2	3.3	3.5	2.4
2004.....	2.5	*	*	*	*	0.2	1.7	7.8	5.8	3.7	3.5	3.0
2003.....	2.5	*	*	*	*	0.2	2.0	8.0	5.0	3.9	4.7	2.7
2002.....	2.7	*	*	*	*	0.3	2.3	8.6	4.9	4.3	5.1	4.8
2001.....	2.5	*	*	*	*	0.2	2.4	7.7	4.6	4.6	5.0	4.3
2000.....	2.4	*	*	*	*	0.2	2.2	7.4	4.4	4.4	6.1	5.4
1999.....	2.2	*	*	*	*	0.2	2.2	5.9	3.9	4.7	6.4	6.8

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	0.8	3.6	3.6	3.4	3.4	3.8	2.7
2006.....	1.6	*	*	*	*	0.8	3.6	3.6	3.4	3.5	4.2	2.7
2005.....	1.3	*	*	*	*	0.1	0.7	2.8	2.3	2.9	3.5	2.2
2004.....	1.3	*	*	*	*	0.2	1.0	2.7	2.3	2.7	3.2	2.4
2003.....	1.3	*	*	*	*	0.1	0.8	2.8	2.3	3.0	3.5	2.2
2002.....	1.4	*	*	*	*	0.2	1.0	2.7	2.4	3.5	3.8	3.2
2001.....	1.4	*	*	*	*	0.2	1.0	2.3	2.5	3.8	4.3	3.3
2000.....	1.4	*	*	*	*	0.2	1.0	2.4	2.3	4.0	4.3	3.2
1999.....	1.2	*	*	*	*	0.2	0.9	1.8	2.1	4.0	4.0	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Non-Hispanic white, both sexes

2007 5/.....	2.4	*	*	*	*	0.2	1.0	6.2	6.1	3.4	3.6	2.5
2006 5/.....	2.3	*	*	*	*	0.1	1.1	6.3	5.3	3.3	3.6	2.5
2005 5/.....	1.8	*	*	*	*	0.1	1.0	5.1	3.6	2.4	3.2	2.0
2004 5/.....	1.8	*	*	*	*	0.1	1.3	4.9	3.4	2.7	3.0	2.4
2003 5/.....	1.8	*	*	*	*	0.2	1.4	5.2	2.9	2.9	3.6	2.2
2002.....	1.9	*	*	*	*	0.2	1.6	5.3	3.0	3.3	4.1	3.5
2001.....	1.8	*	*	*	*	0.2	1.7	4.6	2.9	3.7	4.3	3.3
2000.....	1.8	*	*	*	*	0.1	1.6	4.3	2.7	3.7	4.7	3.7
1999.....	1.6	*	*	*	*	0.2	1.4	3.4	2.5	3.7	4.7	4.5

Non-Hispanic white, male

2007 5/.....	3.3	*	*	*	*	0.2	1.3	8.9	9.3	4.0	4.1	2.8
2006 5/.....	3.1	*	*	*	*	0.2	1.4	9.3	7.9	3.7	3.9	3.1
2005 5/.....	2.5	*	*	*	*	*	1.4	7.7	5.5	2.8	3.4	2.2
2004 5/.....	2.4	*	*	*	*	*	1.7	7.4	4.8	3.2	3.2	2.7
2003 5/.....	2.4	*	*	*	*	0.2	2.0	7.7	4.2	3.4	4.2	2.7
2002.....	2.6	*	*	*	*	0.3	2.3	8.2	4.1	3.9	4.9	4.6
2001.....	2.5	*	*	*	*	0.2	2.3	7.2	4.0	4.3	5.1	4.4
2000.....	2.3	*	*	*	*	*	2.1	6.7	3.6	3.8	5.9	5.4
1999.....	2.1	*	*	*	*	0.2	2.1	5.2	3.4	4.3	6.4	6.5

Non-Hispanic white, female

2007 5/.....	1.6	*	*	*	*	*	0.8	3.5	3.1	2.9	3.2	2.4
2006 5/.....	1.5	*	*	*	*	*	0.8	3.4	2.9	2.9	3.4	2.2
2005 5/.....	1.2	*	*	*	*	*	0.7	2.6	1.9	2.2	3.1	1.9
2004 5/.....	1.2	*	*	*	*	*	0.9	2.5	2.0	2.3	2.8	2.3
2003 5/.....	1.2	*	*	*	*	*	0.8	2.6	1.8	2.5	3.1	2.0
2002.....	1.3	*	*	*	*	*	0.9	2.5	2.0	2.8	3.5	3.1
2001.....	1.2	*	*	*	*	*	1.0	2.0	2.0	3.1	3.8	2.9
2000.....	1.3	*	*	*	*	*	1.0	2.0	1.8	3.6	4.0	3.1
1999.....	1.1	*	*	*	*	0.2	0.8	1.7	1.6	3.3	3.6	3.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Non-Hispanic black, both sexes

2007 5/.....	2.7	*	*	*	*	*	1.1	7.3	12.2	5.8	5.6	*
2006 5/.....	2.9	*	*	*	*	*	1.3	7.8	12.5	7.3	6.7	*
2005 5/.....	2.3	*	*	*	*	*	1.2	7.1	8.5	6.2	5.1	*
2004 5/.....	2.3	*	*	*	*	*	1.4	7.6	8.8	4.6	5.0	*
2003 5/.....	2.3	*	*	*	*	*	1.8	7.6	8.2	5.3	5.9	*
2002.....	2.3	*	*	*	*	0.4	2.0	8.2	7.2	6.0	5.2	*
2001.....	2.3	*	*	*	*	*	2.2	8.1	6.8	5.6	5.7	*
2000.....	2.2	*	*	*	*	0.4	2.1	8.0	5.8	5.8	6.3	*
1999.....	2.1	*	*	*	*	*	2.2	7.0	5.1	6.8	5.9	*

Non-Hispanic black, male

2007 5/.....	3.7	*	*	*	*	*	1.6	10.5	19.7	8.1	6.3	*
2006 5/.....	3.8	*	*	*	*	*	1.7	11.1	20.0	9.3	7.5	*
2005 5/.....	2.9	*	*	*	*	*	1.5	10.3	12.9	7.3	*	*
2004 5/.....	3.1	*	*	*	*	*	1.5	11.9	14.1	6.7	*	*
2003 5/.....	3.1	*	*	*	*	*	2.4	11.7	12.1	5.9	6.1	*
2002.....	3.1	*	*	*	*	*	2.6	12.6	10.7	7.0	*	*
2001.....	2.8	*	*	*	*	*	3.0	12.0	9.0	5.0	*	*
2000.....	2.9	*	*	*	*	*	3.0	12.3	8.1	7.2	7.7	*
1999.....	2.6	*	*	*	*	*	3.1	11.8	6.3	5.7	*	*

Non-Hispanic black, female

2007 5/.....	1.8	*	*	*	*	*	0.8	4.6	6.2	4.2	5.2	*
2006 5/.....	2.0	*	*	*	*	*	1.0	5.0	6.5	5.9	6.3	*
2005 5/.....	1.7	*	*	*	*	*	0.9	4.4	5.0	5.4	5.9	*
2004 5/.....	1.5	*	*	*	*	*	1.4	3.9	4.5	3.1	5.3	*
2003 5/.....	1.6	*	*	*	*	*	1.2	4.1	5.1	4.9	5.9	*
2002.....	1.7	*	*	*	*	*	1.5	4.4	4.4	5.2	5.3	*
2001.....	1.8	*	*	*	*	*	1.4	4.7	5.0	6.0	6.7	*
2000.....	1.6	*	*	*	*	*	1.4	4.3	3.9	4.7	5.6	*
1999.....	1.6	*	*	*	*	*	1.4	2.9	4.1	7.5	6.7	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes

2007.....	3.7	*	*	*	0.4	2.7	8.3	9.5	5.3	2.3	0.8	*
2006.....	4.0	*	*	*	0.5	2.9	9.2	10.1	5.5	2.5	0.8	*
2005.....	4.2	*	*	*	0.4	3.3	9.9	10.6	5.3	2.3	0.8	*
2004.....	4.4	*	*	0.1	0.5	3.7	10.9	10.6	5.4	2.4	0.8	*
2003.....	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002.....	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001.....	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000.....	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999.....	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*

All origins 3/, male

2007.....	5.4	*	*	*	0.4	3.2	11.6	14.0	8.5	4.2	1.6	*
2006.....	5.9	*	*	*	0.6	3.5	12.9	15.3	8.9	4.2	1.6	*
2005.....	6.3	*	*	*	0.4	4.0	14.3	16.4	8.8	4.1	1.4	*
2004.....	6.6	*	*	*	0.5	4.5	15.7	16.3	9.0	4.0	1.4	*
2003.....	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002.....	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001.....	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000.....	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999.....	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*

All origins 3/, female

2007.....	2.1	*	*	*	0.3	2.2	4.9	5.1	2.2	0.8	0.3	*
2006.....	2.2	*	*	*	0.4	2.3	5.4	5.1	2.3	1.1	0.3	*
2005.....	2.2	*	*	*	0.3	2.6	5.6	5.1	2.0	0.9	0.4	*
2004.....	2.4	*	*	*	0.4	2.8	6.2	5.2	2.0	1.0	0.5	*
2003.....	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002.....	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001.....	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000.....	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999.....	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, both sexes

2007.....	3.3	*	*	*	0.4	2.2	7.4	10.8	7.1	4.8	*	*
2006.....	3.6	*	*	*	0.4	2.4	8.8	11.7	7.6	4.7	*	*
2005.....	3.9	*	*	*	0.4	2.5	9.6	13.0	7.2	4.3	*	*
2004.....	4.3	*	*	*	0.4	2.8	11.0	14.1	8.4	5.0	*	*
2003.....	4.7	*	*	*	0.4	3.3	12.3	15.9	9.8	5.2	*	*
2002.....	4.8	*	*	*	0.4	3.6	13.4	14.7	9.5	3.4	*	*
2001.....	5.0	*	*	*	0.5	4.4	13.9	15.5	9.2	4.5	*	*
2000.....	5.4	*	*	*	0.3	5.6	14.9	16.9	9.5	3.8	*	*
1999.....	5.8	*	*	*	0.5	6.4	17.3	15.4	8.4	4.7	*	*

Hispanic, male

2007.....	5.0	*	*	*	0.6	3.0	10.9	16.9	11.2	8.9	*	*
2006.....	5.6	*	*	*	*	3.4	12.9	18.6	12.9	7.9	*	*
2005.....	5.9	*	*	*	0.6	3.3	14.7	20.8	12.6	7.3	*	*
2004.....	6.4	*	*	*	*	4.1	16.1	22.0	14.4	8.0	*	*
2003.....	7.1	*	*	*	0.5	4.6	17.7	26.1	15.4	9.2	*	*
2002.....	7.2	*	*	*	*	5.1	20.1	23.1	15.0	6.9	*	*
2001.....	7.6	*	*	*	0.6	6.0	20.8	24.9	14.8	8.6	*	*
2000.....	8.2	*	*	*	*	7.8	22.7	27.5	15.3	7.2	*	*
1999.....	8.7	*	*	*	0.6	9.3	26.2	24.2	15.4	9.0	*	*

Hispanic, female

2007.....	1.5	*	*	*	*	1.3	3.4	4.6	3.4	*	*	*
2006.....	1.6	*	*	*	*	1.2	4.1	4.5	2.8	*	*	*
2005.....	1.6	*	*	*	*	1.4	4.0	5.2	2.3	*	*	*
2004.....	1.9	*	*	*	*	1.2	5.3	6.2	2.9	*	*	*
2003.....	2.2	*	*	*	*	1.8	6.3	5.6	4.6	*	*	*
2002.....	2.2	*	*	*	*	1.8	6.2	6.3	4.6	*	*	*
2001.....	2.3	*	*	*	*	2.5	6.5	6.2	4.1	*	*	*
2000.....	2.4	*	*	*	*	3.0	6.4	6.6	4.4	*	*	*
1999.....	2.6	*	*	*	*	3.2	7.9	6.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 4/, both sexes

2007.....	3.8	*	*	*	0.4	2.8	8.4	9.2	5.1	2.1	0.8	*
2006.....	4.1	*	*	*	0.5	3.0	9.2	9.9	5.2	2.3	0.7	*
2005.....	4.3	*	*	*	0.4	3.5	9.9	10.3	5.1	2.2	0.7	*
2004.....	4.4	*	*	0.1	0.5	3.9	10.9	10.1	5.0	2.1	0.7	*
2003.....	4.6	*	*	0.1	0.4	4.1	11.9	10.2	5.0	2.1	0.7	*
2002.....	4.8	*	*	0.1	0.4	4.8	12.5	10.6	4.7	2.1	0.7	*
2001.....	4.9	*	*	0.1	0.6	5.4	12.7	9.9	4.8	1.9	0.6	*
2000.....	5.0	*	*	0.1	0.5	6.2	12.7	10.3	4.7	2.0	0.6	*
1999.....	5.2	*	*	0.2	0.5	6.8	13.2	10.0	4.5	2.0	0.6	*

Non-Hispanic 4/, male

2007.....	5.5	*	*	*	0.4	3.2	11.7	13.6	8.2	3.7	1.5	*
2006.....	6.0	*	*	*	0.6	3.6	12.9	14.8	8.5	3.9	1.4	*
2005.....	6.3	*	*	*	0.4	4.1	14.1	15.7	8.5	3.8	1.2	*
2004.....	6.5	*	*	*	0.5	4.6	15.5	15.5	8.5	3.6	1.2	*
2003.....	7.0	*	*	*	0.4	5.2	17.3	16.0	8.4	3.6	1.4	*
2002.....	7.3	*	*	*	0.4	6.0	18.4	16.9	7.9	3.7	1.3	*
2001.....	7.5	*	*	0.1	0.5	7.3	19.0	15.8	8.0	3.1	1.3	*
2000.....	7.7	*	*	*	0.5	7.9	19.1	16.7	8.0	3.4	1.2	*
1999.....	8.0	*	*	0.2	0.5	9.4	20.1	16.5	7.6	3.4	1.0	*

Non-Hispanic 4/, female

2007.....	2.2	*	*	*	0.4	2.4	5.2	5.1	2.1	0.7	0.3	*
2006.....	2.3	*	*	*	0.4	2.5	5.6	5.1	2.2	1.0	0.3	*
2005.....	2.3	*	*	*	0.4	2.8	5.8	5.0	2.0	0.8	0.4	*
2004.....	2.4	*	*	*	0.5	3.1	6.3	5.0	1.9	0.9	0.4	*
2003.....	2.4	*	*	*	0.4	3.0	6.5	4.7	1.9	0.9	*	*
2002.....	2.5	*	*	*	0.5	3.6	6.7	4.7	1.7	0.8	0.3	*
2001.....	2.5	*	*	*	0.7	3.6	6.6	4.2	1.8	0.9	*	*
2000.....	2.5	*	*	0.1	0.5	4.4	6.4	4.2	1.6	0.8	*	*
1999.....	2.5	*	*	0.1	0.6	4.2	6.5	3.9	1.6	0.7	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic white, both sexes

2007 5/.....	1.6	*	*	*	*	0.8	3.6	3.9	2.1	1.0	0.4	*
2006 5/.....	1.8	*	*	*	0.1	0.9	4.1	4.3	2.2	1.0	0.4	*
2005 5/.....	1.9	*	*	*	0.1	1.1	4.6	4.5	2.2	0.9	0.4	*
2004 5/.....	1.9	*	*	*	0.1	1.2	5.1	4.3	2.2	0.9	0.4	*
2003 5/.....	2.1	*	*	*	0.1	1.5	5.6	4.5	2.2	0.8	0.3	*
2002.....	2.1	*	*	*	0.1	1.7	5.8	4.6	2.2	0.8	0.3	*
2001.....	2.2	*	*	*	0.1	2.1	6.0	4.2	2.1	0.7	0.3	*
2000.....	2.3	*	*	*	0.1	2.4	6.2	4.6	2.2	0.8	0.3	*
1999.....	2.4	*	*	*	0.1	3.0	6.4	4.6	2.0	0.8	0.3	*

Non-Hispanic white, male

2007 5/.....	2.7	*	*	*	*	1.1	6.1	6.5	3.6	1.9	0.8	*
2006 5/.....	3.0	*	*	*	*	1.3	6.8	7.4	4.0	1.8	0.7	*
2005 5/.....	3.1	*	*	*	*	1.6	7.6	7.6	3.8	1.7	0.7	*
2004 5/.....	3.3	*	*	*	*	1.8	8.4	7.4	4.0	1.7	0.7	*
2003 5/.....	3.6	*	*	*	*	2.4	9.4	7.8	4.0	1.5	0.8	*
2002.....	3.7	*	*	*	*	2.7	9.8	8.0	4.0	1.5	0.5	*
2001.....	3.8	*	*	*	*	3.2	10.4	7.5	3.8	1.4	0.6	*
2000.....	4.0	*	*	*	0.2	3.7	10.4	8.2	4.0	1.5	0.6	*
1999.....	4.2	*	*	*	0.2	4.9	10.9	8.2	3.6	1.5	0.6	*

Non-Hispanic white, female

2007 5/.....	0.5	*	*	*	*	0.5	1.2	1.4	0.6	*	*	*
2006 5/.....	0.6	*	*	*	*	0.5	1.5	1.3	0.5	0.2	*	*
2005 5/.....	0.6	*	*	*	*	0.7	1.6	1.5	0.6	*	*	*
2004 5/.....	0.6	*	*	*	*	0.6	1.8	1.3	0.4	*	*	*
2003 5/.....	0.6	*	*	*	*	0.7	1.8	1.2	0.5	0.3	*	*
2002.....	0.6	*	*	*	*	0.8	1.8	1.2	0.5	*	*	*
2001.....	0.6	*	*	*	*	0.9	1.7	1.0	0.5	*	*	*
2000.....	0.7	*	*	*	*	1.1	2.0	1.1	0.5	*	*	*
1999.....	0.7	*	*	*	*	1.2	1.9	1.1	0.4	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic black, both sexes

2007 5/.....	16.8	*	*	*	1.7	13.0	36.8	45.3	30.0	12.6	5.3	*
2006 5/.....	17.9	*	*	*	2.2	14.3	39.7	48.3	30.8	14.3	4.8	*
2005 5/.....	18.5	*	*	*	1.7	15.6	42.1	50.5	30.4	14.0	5.0	*
2004 5/.....	19.4	*	*	*	2.3	17.9	46.0	50.9	29.9	13.6	4.6	*
2003 5/.....	20.1	*	*	0.5	1.9	18.1	50.5	51.3	29.9	14.4	4.7	*
2002.....	21.3	*	*	*	2.1	21.5	54.0	54.8	26.3	14.6	5.7	*
2001.....	21.6	*	*	0.4	2.8	24.0	55.0	51.9	28.0	12.8	5.3	*
2000.....	22.0	*	*	0.4	2.2	27.2	54.5	53.9	26.3	13.2	4.6	*
1999.....	22.3	*	*	0.6	2.3	28.3	57.2	52.2	26.4	13.2	3.7	*

Non-Hispanic black, male

2007 5/.....	22.7	*	*	*	1.7	14.3	48.0	64.9	50.3	22.8	10.2	*
2006 5/.....	24.4	*	*	*	2.6	16.2	52.3	70.3	50.3	24.1	10.3	*
2005 5/.....	25.8	*	*	*	1.8	17.9	56.7	76.4	52.0	25.0	8.7	*
2004 5/.....	26.7	*	*	*	2.4	20.0	62.1	76.3	50.8	23.4	7.7	*
2003 5/.....	28.5	*	*	*	1.9	21.7	70.0	79.0	51.4	25.3	8.9	*
2002.....	30.3	*	*	*	2.0	25.5	75.6	86.2	44.8	26.6	12.3	*
2001.....	30.8	*	*	*	2.4	30.8	77.8	81.4	47.7	21.4	11.4	*
2000.....	31.7	*	*	*	2.2	33.4	79.0	86.6	46.4	23.8	8.9	*
1999.....	32.6	*	*	0.7	1.9	37.1	83.5	85.5	45.6	24.0	6.6	*

Non-Hispanic black, female

2007 5/.....	11.4	*	*	*	1.7	11.8	27.0	28.5	13.8	5.4	*	*
2006 5/.....	12.1	*	*	*	1.9	12.6	28.5	29.4	15.1	7.2	*	*
2005 5/.....	11.8	*	*	*	1.6	13.4	29.2	28.3	12.9	6.1	*	*
2004 5/.....	12.8	*	*	*	2.2	15.9	31.6	29.0	13.1	6.6	*	*
2003 5/.....	12.6	*	*	*	2.0	14.8	33.2	27.6	12.6	6.6	*	*
2002.....	13.2	*	*	*	2.2	17.9	34.9	28.0	11.5	6.0	*	*
2001.....	13.3	*	*	*	3.2	17.8	34.8	26.6	12.1	6.7	*	*
2000.....	13.1	*	*	*	2.3	21.7	32.8	25.8	10.1	5.8	*	*
1999.....	13.0	*	*	*	2.8	20.3	34.1	23.8	11.0	5.6	*	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)												
All origins 3/, both sexes												
2007.....	1.9	3.6	0.4	0.1	0.2	0.3	0.5	1.2	2.8	5.7	11.2	20.0
2006.....	2.0	4.0	0.4	0.2	0.2	0.3	0.6	1.3	2.7	5.8	11.5	20.4
2005.....	2.6	4.1	0.4	0.2	0.2	0.3	0.9	3.2	4.2	7.0	13.7	21.2
2004.....	2.6	4.1	0.4	0.1	0.2	0.4	1.0	3.0	4.3	7.1	13.2	22.5
2003.....	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
2002.....	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001.....	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000.....	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999.....	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
All origins 3/, male												
2007.....	2.0	3.9	0.4	0.2	0.3	0.4	0.6	1.4	3.2	6.6	13.5	25.3
2006.....	2.1	4.5	0.5	0.2	0.2	0.3	0.8	1.6	3.2	6.7	13.9	25.1
2005.....	2.9	4.8	0.5	0.1	0.2	0.4	1.2	4.5	5.3	8.3	16.4	23.6
2004.....	2.8	4.8	0.4	0.1	0.3	0.4	1.2	4.1	5.4	8.4	15.1	29.6
2003.....	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
2002.....	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001.....	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000.....	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999.....	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9
All origins 3/, female												
2007.....	1.9	3.4	0.4	*	0.2	0.3	0.5	0.9	2.5	4.9	9.7	17.5
2006.....	1.9	3.6	0.3	0.2	0.2	0.3	0.5	1.0	2.3	5.1	9.9	18.2
2005.....	2.3	3.4	0.3	0.2	0.2	0.3	0.6	2.0	3.2	5.9	11.8	20.1
2004.....	2.3	3.4	0.4	0.1	0.2	0.3	0.8	1.9	3.2	6.0	11.9	19.3
2003.....	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
2002.....	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001.....	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000.....	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999.....	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, both sexes

2007.....	1.1	4.0	*	*	0.3	0.5	0.5	1.3	2.9	5.1	10.1	15.4
2006.....	0.9	2.8	*	*	0.3	0.3	0.7	1.2	2.1	5.2	7.9	10.2
2005.....	1.8	3.5	*	*	*	0.4	1.3	4.8	5.5	8.6	13.3	15.4
2004.....	1.8	3.4	0.6	*	*	0.3	1.3	4.3	6.0	7.6	14.6	18.6
2003.....	1.7	2.7	*	*	*	0.3	1.3	4.4	5.6	8.4	15.3	13.0
2002.....	1.5	4.9	*	*	*	0.7	1.2	3.5	4.3	6.4	11.3	17.7
2001.....	1.1	5.0	*	*	*	0.5	0.8	2.0	3.0	5.9	10.3	15.0
2000.....	1.2	3.0	*	*	*	0.4	1.0	1.9	3.2	5.8	11.3	19.2
1999.....	1.5	4.8	*	*	*	0.6	1.4	2.3	3.8	7.7	14.2	26.0

Hispanic, male

2007.....	1.2	4.2	*	*	*	0.6	0.7	1.8	3.2	4.8	10.9	*
2006.....	1.1	*	*	*	*	*	0.8	1.8	2.5	6.2	11.1	*
2005.....	2.0	*	*	*	*	0.5	1.7	7.0	7.2	8.8	13.0	*
2004.....	2.1	*	*	*	*	*	1.7	6.0	8.3	8.8	17.5	*
2003.....	2.0	*	*	*	*	*	1.8	6.8	7.7	9.8	16.5	*
2002.....	1.8	6.3	*	*	*	0.8	1.6	5.2	5.6	7.6	12.3	*
2001.....	1.3	5.0	*	*	*	0.6	1.1	2.8	4.4	7.6	12.1	*
2000.....	1.3	*	*	*	*	0.6	1.4	2.1	4.4	5.9	13.3	*
1999.....	1.5	5.7	*	*	*	0.6	1.9	2.7	3.8	7.9	15.1	*

Hispanic, female

2007.....	1.1	*	*	*	*	*	*	0.9	2.6	5.3	9.6	13.2
2006.....	0.8	*	*	*	*	*	*	*	1.8	4.5	5.8	*
2005.....	1.6	*	*	*	*	*	0.8	2.6	4.0	8.3	13.5	17.6
2004.....	1.4	*	*	*	*	*	0.9	2.6	4.0	6.7	12.6	19.4
2003.....	1.3	*	*	*	*	*	0.7	2.0	3.6	7.4	14.5	*
2002.....	1.2	*	*	*	*	0.6	*	1.7	3.2	5.3	10.7	*
2001.....	1.0	5.0	*	*	*	*	*	1.3	*	4.6	9.1	*
2000.....	1.0	*	*	*	*	*	*	1.8	*	5.6	9.9	*
1999.....	1.4	*	*	*	*	*	*	1.9	3.8	7.6	13.5	28.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Non-Hispanic 4/, both sexes												
2007.....	2.1	3.5	0.4	0.1	0.2	0.3	0.5	1.1	2.8	5.7	11.3	20.3
2006.....	2.1	4.4	0.4	0.2	0.2	0.3	0.6	1.3	2.8	5.9	11.7	20.9
2005.....	2.7	4.3	0.4	0.2	0.2	0.3	0.9	3.0	4.1	6.9	13.7	21.5
2004.....	2.7	4.3	0.4	0.1	0.2	0.4	0.9	2.8	4.1	7.0	13.1	22.6
2003.....	2.7	4.8	0.5	0.1	0.2	0.4	1.0	2.7	3.9	7.2	13.2	22.9
2002.....	2.4	5.7	0.4	0.2	0.2	0.4	1.0	2.2	3.3	6.7	12.5	21.8
2001.....	2.1	4.6	0.5	0.2	0.2	0.3	0.8	1.6	3.0	6.2	10.6	20.3
2000.....	2.1	4.7	0.4	0.2	0.2	0.3	0.7	1.5	2.9	5.8	11.1	21.2
1999.....	2.8	5.5	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
Non-Hispanic 4/, male												
2007.....	2.2	3.8	0.3	0.2	0.2	0.3	0.6	1.4	3.2	6.7	13.6	25.6
2006.....	2.2	4.7	0.5	0.2	0.2	0.3	0.8	1.6	3.3	6.7	14.1	25.8
2005.....	3.0	5.1	0.5	0.1	0.2	0.4	1.1	4.2	5.0	8.3	16.5	24.2
2004.....	3.0	5.2	0.4	*	0.2	0.4	1.1	3.9	5.2	8.4	15.0	30.2
2003.....	2.9	4.8	0.5	*	0.2	0.4	1.2	3.6	4.9	8.4	15.8	27.0
2002.....	2.7	5.9	0.5	0.2	0.3	0.4	1.3	3.0	4.0	8.0	15.6	26.8
2001.....	2.2	4.8	0.5	0.1	0.2	0.4	1.0	2.1	3.6	6.7	13.2	24.0
2000.....	2.2	5.1	0.5	0.2	0.3	0.3	0.9	1.9	3.2	7.3	13.5	27.8
1999.....	2.8	6.2	0.6	0.2	0.3	0.6	1.4	2.6	4.2	9.2	17.2	32.3
Non-Hispanic 4/, female												
2007.....	2.0	3.2	0.4	*	0.2	0.2	0.5	0.9	2.4	4.9	9.6	17.7
2006.....	2.1	4.1	*	0.2	0.2	0.3	0.4	1.0	2.3	5.2	10.1	18.6
2005.....	2.5	3.4	*	0.2	0.2	0.2	0.6	1.9	3.1	5.7	11.7	20.2
2004.....	2.4	3.3	0.3	*	0.2	0.3	0.8	1.8	3.1	5.9	11.9	19.2
2003.....	2.5	4.8	0.5	*	0.2	0.3	0.7	1.9	3.0	6.2	11.5	21.2
2002.....	2.2	5.4	*	0.2	0.2	0.3	0.7	1.4	2.5	5.6	10.5	19.7
2001.....	2.1	4.5	0.4	0.2	0.2	0.3	0.6	1.2	2.5	5.8	8.9	18.8
2000.....	2.0	4.3	0.4	0.2	0.2	0.4	0.6	1.1	2.6	4.7	9.5	18.6
1999.....	2.7	4.7	0.4	0.2	0.2	0.6	1.0	1.9	3.3	6.7	12.2	24.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Non-Hispanic white, both sexes												
2007 5/.....	2.2	2.9	0.4	0.1	0.2	0.2	0.4	1.0	2.7	5.6	11.2	20.8
2006 5/.....	2.2	2.8	0.4	0.2	0.2	0.2	0.5	1.1	2.6	5.8	11.7	21.1
2005 5/.....	2.8	3.7	0.4	0.2	0.2	0.2	0.7	2.9	3.8	6.7	13.4	21.5
2004 5/.....	2.8	3.7	0.3	0.1	0.2	0.3	0.8	2.8	3.8	6.7	12.8	22.6
2003 5/.....	2.8	3.8	0.4	0.1	0.2	0.3	0.8	2.6	3.6	7.2	12.8	23.4
2002.....	2.5	4.8	0.4	0.2	0.2	0.3	0.8	1.9	2.9	6.6	12.4	22.2
2001.....	2.1	3.3	0.4	0.1	0.2	0.3	0.7	1.5	2.8	6.0	10.4	20.2
2000.....	2.1	3.5	0.4	0.2	0.2	0.2	0.6	1.3	2.6	5.6	10.9	21.0
1999.....	2.7	4.4	0.5	0.2	0.2	0.4	0.9	1.8	3.3	7.4	13.6	26.4
Non-Hispanic white, male												
2007 5/.....	2.2	3.1	*	0.2	0.2	0.2	0.4	1.2	3.0	6.4	13.6	26.1
2006 5/.....	2.3	3.1	0.4	*	0.2	0.2	0.6	1.4	3.0	6.7	14.0	26.5
2005 5/.....	3.1	4.4	0.4	*	*	0.2	1.0	4.1	4.7	8.2	16.2	24.4
2004 5/.....	3.0	4.6	*	*	0.2	0.3	0.9	3.8	4.8	8.0	14.6	30.8
2003 5/.....	2.9	3.6	*	*	0.2	0.3	1.1	3.4	4.5	8.2	15.4	27.6
2002.....	2.7	4.8	0.4	0.2	0.2	0.3	1.0	2.7	3.5	7.9	15.3	27.2
2001.....	2.2	3.3	0.4	*	0.2	0.3	0.8	1.9	3.3	6.5	13.1	24.2
2000.....	2.2	3.9	*	*	0.2	0.2	0.7	1.7	3.0	7.0	13.2	27.0
1999.....	2.7	5.5	0.6	0.2	0.2	0.4	1.2	2.2	3.8	8.8	16.3	30.8
Non-Hispanic white, female												
2007 5/.....	2.1	2.7	*	*	0.2	0.2	0.3	0.7	2.3	4.9	9.5	18.3
2006 5/.....	2.2	2.4	*	0.2	0.2	0.2	0.4	0.9	2.3	5.1	10.2	18.6
2005 5/.....	2.5	2.9	*	0.2	0.2	0.2	0.5	1.7	3.0	5.5	11.4	20.1
2004 5/.....	2.5	2.7	*	*	0.2	0.3	0.7	1.8	2.9	5.5	11.6	18.9
2003 5/.....	2.6	4.0	*	*	0.2	0.3	0.6	1.7	2.8	6.3	11.1	21.6
2002.....	2.3	4.9	*	0.2	0.2	0.3	0.5	1.2	2.3	5.5	10.5	20.0
2001.....	2.1	3.4	*	0.2	0.2	0.2	0.5	1.0	2.3	5.7	8.7	18.6
2000.....	2.0	3.1	*	0.2	0.2	0.2	0.5	1.0	2.3	4.5	9.5	18.6
1999.....	2.6	3.3	*	*	0.2	0.4	0.6	1.5	2.9	6.3	11.8	24.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic black, both sexes

2007 5/.....	2.0	5.9	*	*	0.4	0.7	1.2	2.2	4.4	6.8	12.8	17.4
2006 5/.....	2.2	10.2	*	*	*	0.7	1.3	2.6	4.5	7.4	12.4	22.6
2005 5/.....	2.6	6.8	*	*	0.4	0.8	1.5	4.2	6.0	7.8	15.2	22.7
2004 5/.....	2.6	6.8	*	*	*	0.8	1.7	3.4	6.3	8.8	16.8	24.8
2003 5/.....	2.6	8.4	0.9	*	*	0.6	1.8	4.4	6.1	7.0	16.4	19.9
2002.....	2.6	9.0	*	*	0.4	0.9	2.2	4.0	5.9	7.4	13.8	21.8
2001.....	2.3	10.0	*	*	0.4	0.9	1.6	3.0	5.2	8.3	11.3	20.9
2000.....	2.3	10.4	*	*	0.4	1.0	1.7	3.0	4.9	8.0	13.3	24.2
1999.....	3.6	10.0	1.0	*	0.4	1.7	3.1	5.6	7.3	11.9	20.9	34.3

Non-Hispanic black, male

2007 5/.....	2.2	6.5	*	*	*	1.0	1.2	2.8	4.7	9.8	14.0	24.4
2006 5/.....	2.3	10.5	*	*	*	0.8	1.8	3.5	6.1	7.9	15.6	21.1
2005 5/.....	2.9	8.4	*	*	*	0.9	2.1	5.2	8.2	9.4	19.6	23.8
2004 5/.....	3.0	7.4	*	*	*	0.9	1.9	5.0	9.1	10.7	20.3	30.5
2003 5/.....	3.0	9.3	*	*	*	0.8	2.2	5.8	8.1	10.6	19.0	23.4
2002.....	3.0	10.0	*	*	*	1.1	2.6	5.6	8.7	9.1	17.0	28.4
2001.....	2.5	11.5	*	*	*	*	2.1	4.0	6.4	9.8	12.6	*
2000.....	2.6	10.6	*	*	*	0.9	2.2	4.0	5.8	10.5	17.9	31.7
1999.....	3.7	10.0	*	*	*	1.9	3.1	6.3	7.9	14.1	29.4	46.4

Non-Hispanic black, female

2007 5/.....	1.8	*	*	*	*	*	1.3	1.7	4.0	4.6	12.0	14.5
2006 5/.....	2.1	10.0	*	*	*	*	0.9	1.9	3.1	7.0	10.6	23.2
2005 5/.....	2.3	*	*	*	*	*	1.0	3.4	4.2	6.7	12.7	22.2
2004 5/.....	2.3	*	*	*	*	*	1.5	2.1	4.1	7.4	14.8	22.5
2003 5/.....	2.2	7.4	*	*	*	*	1.4	3.1	4.6	4.4	14.8	18.6
2002.....	2.2	7.9	*	*	*	*	1.8	2.6	3.7	6.2	11.9	19.2
2001.....	2.1	8.4	*	*	*	1.0	1.2	2.1	4.3	7.2	10.6	20.5
2000.....	2.1	10.2	*	*	*	1.0	1.3	2.2	4.1	6.2	10.8	21.4
1999.....	3.5	10.0	*	*	*	1.5	3.0	4.9	6.7	10.4	16.1	29.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms (C00-C97)

All origins 3/, both sexes

2007.....	186.6	1.7	2.2	2.4	3.9	8.5	30.8	114.3	315.4	715.5	1,256.3	1,590.2
2006.....	187.0	1.8	2.3	2.2	3.9	9.0	31.9	116.3	321.2	727.2	1,263.8	1,606.1
2005.....	188.7	1.8	2.3	2.5	4.1	9.0	33.2	118.6	326.9	742.7	1,274.8	1,637.7
2004.....	188.6	1.8	2.5	2.5	4.1	9.1	33.4	119.0	333.4	755.1	1,280.4	1,653.3
2003.....	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002.....	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001.....	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8

All origins 3/, male

2007.....	197.0	1.8	2.3	2.4	4.5	8.2	26.4	117.5	358.5	854.3	1,617.4	2,249.2
2006.....	196.6	1.8	2.5	2.5	4.6	8.6	27.4	119.0	363.6	870.4	1,631.3	2,248.7
2005.....	198.9	2.1	2.6	2.7	4.8	8.8	28.9	121.6	369.5	899.1	1,649.7	2,319.3
2004.....	198.4	1.8	2.6	2.7	4.8	8.6	29.1	124.3	376.7	907.6	1,662.1	2,349.5
2003.....	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002.....	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001.....	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000.....	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999.....	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2

All origins 3/, female

2007.....	176.5	1.6	2.2	2.3	3.2	8.9	35.2	111.3	275.2	597.6	1,007.4	1,276.7
2006.....	177.6	1.8	2.1	2.0	3.1	9.5	36.4	113.7	281.8	605.9	1,012.5	1,305.5
2005.....	178.8	1.5	2.0	2.2	3.3	9.1	37.5	115.8	287.4	610.9	1,020.3	1,324.6
2004.....	179.1	1.9	2.4	2.2	3.4	9.6	37.7	113.8	293.2	627.1	1,023.5	1,340.1
2003.....	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002.....	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001.....	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000.....	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999.....	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms (C00-C97)

Hispanic, both sexes

2007.....	60.8	*	1.8	2.8	4.6	7.5	21.7	74.8	197.9	452.5	833.6	1,026.6
2006.....	60.1	*	2.2	2.3	4.4	7.2	21.6	74.7	198.6	461.9	844.0	1,077.5
2005.....	61.3	*	2.2	2.5	4.5	6.8	22.1	77.9	203.3	481.3	882.8	1,128.1
2004.....	59.3	*	2.0	2.8	4.3	6.9	21.8	75.5	209.7	472.0	872.2	1,131.2
2003.....	60.3	*	3.0	2.9	4.1	7.1	22.9	80.7	214.7	485.2	890.1	1,234.9
2002.....	59.7	*	2.3	2.9	4.5	6.8	23.2	78.2	215.1	496.0	894.9	1,308.8
2001.....	60.5	*	2.8	2.5	4.3	7.2	25.6	82.8	223.4	504.0	913.9	1,363.7
2000.....	59.9	*	2.5	2.9	4.2	7.3	25.2	82.1	221.0	516.2	953.5	1,383.5
1999.....	59.6	*	2.5	2.7	4.1	7.4	24.4	84.9	225.1	520.5	926.8	1,399.6

Hispanic, male

2007.....	61.6	*	1.9	2.7	5.3	6.3	17.4	74.2	221.9	560.3	1,072.6	1,417.9
2006.....	60.4	*	2.6	2.3	5.1	6.1	16.0	71.4	224.8	574.8	1,098.4	1,440.1
2005.....	63.0	*	2.8	2.5	5.3	6.5	17.8	75.9	236.9	603.5	1,161.8	1,601.5
2004.....	60.9	*	1.6	3.0	5.5	6.4	18.6	77.4	239.0	585.8	1,174.2	1,508.8
2003.....	61.5	*	3.6	3.2	4.9	6.8	18.2	81.1	246.5	617.6	1,163.9	1,668.6
2002.....	61.2	*	2.2	2.9	5.2	6.3	18.4	78.4	254.3	622.3	1,190.8	1,869.0
2001.....	62.2	*	2.9	2.5	5.5	6.3	21.7	81.5	253.5	642.8	1,258.3	1,967.4
2000.....	61.3	*	2.9	3.5	4.5	6.9	20.1	79.4	253.1	651.2	1,306.4	2,049.7
1999.....	61.2	*	2.3	2.8	4.6	6.6	19.8	84.6	268.1	668.9	1,252.9	1,945.3

Hispanic, female

2007.....	59.9	*	1.6	2.8	3.9	8.9	26.5	75.5	175.6	364.4	665.7	814.2
2006.....	59.7	*	1.8	2.2	3.5	8.5	27.9	78.1	174.4	370.2	665.9	884.9
2005.....	59.5	*	1.6	2.5	3.6	7.1	27.0	79.9	172.5	382.5	688.5	880.4
2004.....	57.7	*	2.4	2.5	3.0	7.5	25.4	73.5	183.0	380.7	663.6	937.0
2003.....	59.1	*	2.4	2.6	3.2	7.4	28.0	80.2	185.9	379.7	702.1	1,014.8
2002.....	58.1	*	2.4	2.9	3.6	7.5	28.4	78.0	179.8	395.6	692.2	1,031.2
2001.....	58.7	*	2.7	2.5	2.9	8.3	29.8	84.2	196.7	394.5	681.2	1,068.6
2000.....	58.5	*	2.2	2.3	3.7	7.8	30.7	84.7	192.5	410.0	716.5	1,056.5
1999.....	58.0	*	2.7	2.6	3.5	8.3	29.4	85.1	187.2	404.3	709.8	1,136.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms (C00-C97)

Non-Hispanic 4/, both sexes

2007.....	208.7	1.6	2.4	2.3	3.7	8.8	32.4	119.0	325.5	735.6	1,282.8	1,619.0
2006.....	208.7	1.8	2.3	2.2	3.8	9.5	33.7	120.9	331.3	746.7	1,288.4	1,630.6
2005.....	209.8	1.8	2.3	2.5	4.0	9.5	35.0	122.9	336.9	761.5	1,296.6	1,660.5
2004.....	209.4	1.7	2.6	2.4	4.0	9.6	35.2	123.4	342.9	774.5	1,301.1	1,674.3
2003.....	211.9	2.0	2.3	2.6	4.0	9.9	36.7	126.0	352.4	788.9	1,321.6	1,714.7
2002.....	213.4	1.7	2.6	2.5	4.2	10.3	37.6	128.0	360.9	810.5	1,329.2	1,736.8
2001.....	213.9	1.6	2.7	2.4	4.2	10.6	38.2	130.3	365.9	820.5	1,331.4	1,776.6
2000.....	215.6	2.5	2.7	2.4	4.4	10.3	37.9	131.2	376.6	832.8	1,348.9	1,831.1
1999.....	215.5	1.7	2.7	2.4	4.5	10.4	38.6	131.0	384.6	843.2	1,344.5	1,816.1

Non-Hispanic 4/, male

2007.....	222.2	1.9	2.4	2.4	4.3	8.7	28.1	122.7	370.2	876.0	1,652.2	2,296.0
2006.....	221.3	1.7	2.5	2.5	4.5	9.3	29.5	124.4	374.8	891.3	1,662.8	2,289.8
2005.....	222.8	2.4	2.6	2.7	4.7	9.4	30.8	126.5	379.9	919.5	1,677.1	2,354.9
2004.....	221.8	1.8	2.8	2.7	4.6	9.1	30.8	129.2	387.0	928.8	1,687.2	2,387.3
2003.....	224.3	1.9	2.6	2.7	4.5	9.4	32.8	131.9	396.9	951.1	1,720.2	2,444.0
2002.....	226.7	1.9	2.8	2.8	4.8	9.9	33.5	132.6	410.0	985.1	1,733.2	2,513.2
2001.....	227.2	1.6	2.9	2.5	4.9	9.9	34.0	134.6	415.3	1,003.8	1,745.1	2,634.8
2000.....	228.7	2.6	2.9	2.5	5.2	9.7	34.3	135.1	426.6	1,020.2	1,776.0	2,732.4
1999.....	229.9	1.6	3.0	2.6	5.3	10.3	35.0	135.4	438.8	1,036.8	1,783.7	2,747.5

Non-Hispanic 4/, female

2007.....	195.9	1.3	2.3	2.2	3.1	8.9	36.7	115.4	283.8	615.9	1,028.7	1,299.3
2006.....	196.7	1.9	2.2	2.0	3.0	9.7	37.7	117.5	290.8	624.0	1,032.8	1,324.2
2005.....	197.5	*	2.1	2.2	3.3	9.5	39.2	119.5	296.8	627.7	1,038.6	1,343.6
2004.....	197.5	1.6	2.4	2.2	3.4	10.0	39.5	117.8	301.9	644.5	1,041.6	1,355.6
2003.....	200.0	2.1	2.1	2.4	3.4	10.4	40.6	120.5	311.1	652.7	1,055.6	1,394.3
2002.....	200.8	1.5	2.4	2.2	3.6	10.6	41.6	123.5	315.4	664.7	1,061.5	1,401.8
2001.....	201.2	1.6	2.5	2.3	3.6	11.3	42.4	126.0	320.0	668.1	1,059.8	1,419.6
2000.....	203.1	2.4	2.6	2.2	3.6	10.9	41.4	127.3	330.3	678.1	1,070.6	1,466.6
1999.....	201.9	1.8	2.5	2.2	3.7	10.5	42.1	126.8	334.5	684.2	1,059.4	1,443.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms (C00-C97)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	226.9	1.8	2.5	2.4	3.6	8.5	31.4	113.8	318.2	736.3	1,292.4	1,631.9
2006 5/.....	227.1	2.1	2.3	2.3	3.7	9.2	32.4	115.8	325.1	748.2	1,297.8	1,643.1
2005 5/.....	227.7	2.0	2.3	2.5	4.0	9.0	34.0	117.1	329.0	762.3	1,304.0	1,668.6
2004 5/.....	227.0	1.9	2.7	2.4	3.9	9.3	33.7	117.3	336.2	773.6	1,305.4	1,680.9
2003 5/.....	229.5	2.2	2.4	2.6	4.0	9.6	35.2	119.4	345.9	788.8	1,322.6	1,717.8
2002.....	230.9	1.6	2.6	2.5	4.3	10.0	36.2	122.3	353.0	809.2	1,328.4	1,737.9
2001.....	230.6	1.8	2.9	2.4	4.3	10.3	36.5	124.6	358.3	815.9	1,326.5	1,775.2
2000.....	232.2	2.6	2.8	2.4	4.3	10.1	35.6	125.1	369.3	827.7	1,342.9	1,826.6
1999.....	231.5	1.6	3.0	2.3	4.3	10.2	36.6	124.8	375.0	837.1	1,336.4	1,808.3
<b>Non-Hispanic white, male</b>												
2007 5/.....	240.6	2.0	2.4	2.4	4.3	8.6	27.7	116.8	357.6	867.3	1,652.8	2,300.4
2006 5/.....	239.9	1.9	2.5	2.5	4.5	9.3	28.9	118.7	363.4	883.0	1,662.9	2,300.2
2005 5/.....	240.7	2.7	2.5	2.7	4.7	9.0	30.5	120.3	366.1	910.4	1,673.7	2,358.3
2004 5/.....	239.2	2.0	2.8	2.7	4.4	9.3	29.9	121.9	374.6	917.5	1,677.3	2,387.1
2003 5/.....	241.8	2.1	2.6	2.7	4.6	9.4	31.9	123.8	384.8	942.0	1,707.8	2,441.7
2002.....	243.8	2.0	2.8	2.7	4.9	9.8	32.5	125.9	395.5	975.3	1,716.5	2,507.7
2001.....	243.4	1.7	3.1	2.4	4.9	10.0	32.6	127.3	401.7	988.2	1,721.8	2,616.8
2000.....	244.4	2.9	3.1	2.6	5.0	9.7	32.3	127.2	412.0	1,002.1	1,750.2	2,714.1
1999.....	245.1	1.7	3.3	2.6	5.0	10.1	33.5	126.7	422.3	1,019.3	1,757.7	2,721.3
<b>Non-Hispanic white, female</b>												
2007 5/.....	213.7	*	2.5	2.3	3.0	8.4	35.2	110.9	280.6	622.2	1,040.1	1,315.2
2006 5/.....	214.7	2.2	2.1	2.0	2.9	9.2	35.9	113.0	288.5	631.3	1,044.4	1,336.7
2005 5/.....	215.1	*	2.2	2.2	3.3	8.9	37.5	113.9	293.6	634.4	1,049.5	1,353.2
2004 5/.....	215.3	1.8	2.5	2.1	3.3	9.3	37.5	112.9	299.8	649.8	1,052.0	1,364.5
2003 5/.....	217.6	2.3	2.2	2.4	3.4	9.8	38.6	115.1	308.9	657.6	1,062.4	1,399.1
2002.....	218.5	*	2.3	2.3	3.6	10.3	39.9	118.7	312.8	667.7	1,068.3	1,405.4
2001.....	218.4	1.8	2.8	2.4	3.6	10.7	40.4	121.9	317.3	669.7	1,064.4	1,425.1
2000.....	220.6	2.4	2.4	2.3	3.5	10.5	38.9	123.0	328.9	681.0	1,075.3	1,468.7
1999.....	218.4	*	2.7	2.1	3.6	10.3	39.7	123.0	330.5	684.6	1,060.5	1,444.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms (C00-C97)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	166.3	*	2.0	2.2	4.0	11.3	43.2	171.2	448.0	889.0	1,410.1	1,696.8
2006 5/.....	165.5	*	2.5	2.2	4.1	11.9	45.8	174.0	450.1	890.6	1,420.3	1,700.4
2005 5/.....	167.4	*	2.5	2.6	4.1	12.7	46.2	179.7	466.0	906.1	1,435.5	1,764.4
2004 5/.....	167.5	*	2.5	2.6	5.0	12.4	48.3	181.8	467.2	931.9	1,465.9	1,794.3
2003 5/.....	170.0	*	2.3	2.7	4.1	12.6	50.5	189.7	480.9	941.1	1,508.2	1,867.5
2002.....	171.5	*	3.0	2.6	4.1	12.8	51.5	186.0	494.6	977.2	1,537.4	1,904.3
2001.....	172.8	*	2.0	2.7	4.4	13.4	52.8	188.7	494.5	1,005.6	1,559.5	1,947.5
2000.....	174.9	*	2.8	2.3	5.3	12.3	55.3	196.7	505.2	1,019.3	1,582.8	2,015.1
1999.....	176.9	*	2.0	2.8	5.6	12.6	55.3	196.8	530.5	1,033.6	1,592.9	2,024.0
<b>Non-Hispanic black, male</b>												
2007 5/.....	180.1	*	2.4	2.2	4.4	10.0	35.3	183.6	558.0	1,162.2	1,974.3	2,691.9
2006 5/.....	179.2	*	2.7	2.5	4.7	10.6	38.1	187.0	557.2	1,178.9	2,015.7	2,577.4
2005 5/.....	181.8	*	2.8	3.1	4.8	12.3	37.4	190.4	581.2	1,205.6	2,047.4	2,720.4
2004 5/.....	182.4	*	2.8	2.5	5.6	10.1	39.8	201.8	580.5	1,231.5	2,116.9	2,781.1
2003 5/.....	184.8	*	2.4	2.9	4.7	10.6	42.9	211.5	597.1	1,242.1	2,175.8	2,859.6
2002.....	187.8	*	2.7	3.1	4.6	11.7	44.3	202.3	621.8	1,293.5	2,247.6	3,008.4
2001.....	190.8	*	2.4	3.0	5.1	10.9	45.8	209.6	614.1	1,354.7	2,310.5	3,238.5
2000.....	194.9	*	2.6	2.6	6.4	10.5	49.6	219.0	637.4	1,382.7	2,372.3	3,290.1
1999.....	197.4	*	2.0	2.9	6.6	11.9	48.6	225.8	670.9	1,387.0	2,377.4	3,361.1
<b>Non-Hispanic black, female</b>												
2007 5/.....	153.7	*	1.6	2.1	3.7	12.6	50.2	160.6	359.7	694.0	1,089.3	1,284.3
2006 5/.....	153.0	*	2.4	2.0	3.4	13.2	52.7	162.8	363.9	684.6	1,082.5	1,342.7
2005 5/.....	154.3	*	2.1	2.2	3.5	13.1	54.0	170.6	373.3	692.1	1,088.7	1,379.5
2004 5/.....	153.9	*	2.2	2.7	4.4	14.5	55.8	164.6	376.3	718.3	1,098.4	1,404.3
2003 5/.....	156.5	*	2.1	2.4	3.5	14.5	57.2	170.9	387.6	727.0	1,131.2	1,483.2
2002.....	156.7	*	3.2	2.0	3.6	13.9	57.8	172.1	392.5	752.8	1,135.6	1,482.4
2001.....	156.5	*	*	2.3	3.8	15.7	59.0	170.7	398.3	759.4	1,137.8	1,469.3
2000.....	156.9	*	3.0	1.9	4.2	14.0	60.4	177.5	398.8	763.6	1,138.4	1,541.6
1999.....	158.3	*	2.0	2.7	4.8	13.3	61.3	172.0	417.9	784.2	1,153.6	1,524.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/, both sexes

2007.....	2.7	*	*	*	*	0.1	0.6	2.6	6.2	9.9	13.4	17.1
2006.....	2.6	*	*	*	*	0.1	0.6	2.6	5.8	9.6	13.5	16.3
2005.....	2.6	*	*	*	0.1	0.1	0.6	2.6	6.4	9.8	12.8	17.0
2004.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.1	10.1	13.7	18.0
2003.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3

All origins 3/, male

2007.....	3.7	*	*	*	*	0.1	0.8	4.0	10.0	15.5	20.1	24.0
2006.....	3.6	*	*	*	*	0.1	0.9	4.1	9.3	14.8	20.4	21.9
2005.....	3.6	*	*	*	*	0.2	0.9	4.2	10.3	15.3	18.9	22.0
2004.....	3.7	*	*	*	*	0.2	0.8	4.6	10.0	15.6	20.3	24.9
2003.....	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002.....	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001.....	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000.....	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999.....	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1

All origins 3/, female

2007.....	1.7	*	*	*	*	0.1	0.4	1.3	2.6	5.2	8.7	13.8
2006.....	1.6	*	*	*	*	*	0.4	1.1	2.4	5.2	8.8	13.7
2005.....	1.7	*	*	*	*	0.1	0.3	1.2	2.8	5.2	8.7	14.7
2004.....	1.7	*	*	*	*	*	0.4	1.2	2.5	5.6	9.2	14.9
2003.....	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002.....	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001.....	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000.....	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999.....	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 4/, both sexes

2007.....	3.0	*	*	*	*	0.1	0.6	2.8	6.4	10.1	13.6	17.4
2006.....	2.9	*	*	*	*	0.1	0.7	2.8	6.0	9.7	13.7	16.5
2005.....	2.9	*	*	*	0.1	0.2	0.7	2.8	6.6	10.2	12.9	17.6
2004.....	3.0	*	*	*	0.1	0.1	0.6	3.0	6.3	10.4	13.9	18.2
2003.....	3.0	*	*	*	0.1	0.2	0.7	2.9	6.8	10.5	13.3	18.7
2002.....	3.0	*	*	*	0.1	0.2	0.6	2.9	7.1	10.7	13.7	18.7
2001.....	3.0	*	*	*	0.1	0.2	0.7	2.8	7.1	10.6	14.3	18.4
2000.....	2.9	*	*	*	0.1	0.1	0.7	3.0	6.9	10.8	13.3	19.5
1999.....	2.9	*	*	*	0.1	0.1	0.7	3.0	7.0	10.5	14.3	19.4

Non-Hispanic 4/, male

2007.....	4.2	*	*	*	*	0.1	0.9	4.2	10.5	15.8	20.6	24.4
2006.....	4.0	*	*	*	*	0.2	0.9	4.4	9.7	15.0	20.6	22.3
2005.....	4.1	*	*	*	*	0.2	1.0	4.4	10.6	15.7	19.1	22.8
2004.....	4.1	*	*	*	*	0.2	0.9	4.8	10.3	15.8	20.7	24.9
2003.....	4.1	*	*	*	*	0.2	1.0	4.8	10.9	16.0	19.5	27.1
2002.....	4.1	*	*	*	*	0.2	0.8	4.6	11.3	16.2	19.8	26.7
2001.....	4.1	*	*	*	*	0.2	1.0	4.6	11.6	16.2	20.6	25.8
2000.....	3.9	*	*	*	*	0.1	1.0	4.6	11.1	16.1	19.4	26.5
1999.....	4.0	*	*	*	0.1	0.2	1.0	4.8	11.0	15.8	21.0	27.3

Non-Hispanic 4/, female

2007.....	1.9	*	*	*	*	*	0.4	1.4	2.7	5.3	8.9	14.2
2006.....	1.8	*	*	*	*	*	0.4	1.2	2.6	5.3	9.1	13.8
2005.....	1.9	*	*	*	*	0.1	0.4	1.2	2.9	5.5	8.8	15.2
2004.....	1.9	*	*	*	*	*	0.4	1.3	2.6	5.8	9.4	15.1
2003.....	1.9	*	*	*	*	*	0.4	1.1	2.9	5.9	9.2	15.0
2002.....	1.9	*	*	*	*	*	0.4	1.2	3.2	6.1	9.7	15.3
2001.....	1.9	*	*	*	*	0.1	0.4	1.2	2.9	6.0	10.2	15.3
2000.....	2.0	*	*	*	*	0.1	0.4	1.3	3.0	6.5	9.3	16.6
1999.....	2.0	*	*	*	*	*	0.4	1.4	3.2	6.0	9.9	16.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic white, both sexes

2007 5/.....	3.2	*	*	*	*	0.1	0.6	2.5	6.1	9.9	13.8	17.9
2006 5/.....	3.0	*	*	*	*	0.1	0.6	2.6	5.7	9.4	13.8	17.1
2005 5/.....	3.0	*	*	*	*	0.1	0.6	2.5	6.2	9.8	12.9	17.9
2004 5/.....	3.1	*	*	*	*	0.1	0.5	2.6	5.8	10.2	13.9	18.7
2003 5/.....	3.1	*	*	*	*	0.1	0.6	2.6	6.3	10.1	13.0	19.1
2002.....	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.6	13.9	18.9
2001.....	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.2	14.5	18.9
2000.....	3.0	*	*	*	*	0.1	0.6	2.5	6.1	10.4	13.3	19.8
1999.....	3.0	*	*	*	*	0.1	0.5	2.4	6.3	10.1	14.2	19.9

Non-Hispanic white, male

2007 5/.....	4.3	*	*	*	*	*	0.7	3.9	10.0	15.0	20.5	24.8
2006 5/.....	4.1	*	*	*	*	*	0.8	4.1	9.0	14.1	20.4	23.0
2005 5/.....	4.1	*	*	*	*	*	0.8	3.9	9.7	14.8	18.5	22.9
2004 5/.....	4.1	*	*	*	*	0.2	0.7	4.0	9.4	15.0	20.4	24.8
2003 5/.....	4.1	*	*	*	*	0.2	1.0	4.2	9.9	14.9	18.7	27.4
2002.....	4.1	*	*	*	*	0.2	0.7	4.0	10.2	15.7	19.9	26.8
2001.....	4.1	*	*	*	*	0.2	0.8	4.0	10.4	15.3	20.5	26.4
2000.....	3.9	*	*	*	*	*	0.8	3.8	9.7	15.0	18.9	27.4
1999.....	3.9	*	*	*	*	*	0.8	3.8	9.7	15.1	20.3	27.5

Non-Hispanic white, female

2007 5/.....	2.0	*	*	*	*	*	0.4	1.2	2.5	5.4	9.0	14.7
2006 5/.....	2.0	*	*	*	*	*	0.4	1.0	2.5	5.3	9.3	14.4
2005 5/.....	2.0	*	*	*	*	*	0.4	1.1	2.8	5.6	9.1	15.6
2004 5/.....	2.0	*	*	*	*	*	0.3	1.1	2.4	6.0	9.6	16.0
2003 5/.....	2.0	*	*	*	*	*	0.2	0.9	2.9	5.9	9.2	15.4
2002.....	2.1	*	*	*	*	*	0.3	1.0	2.9	6.2	9.8	15.5
2001.....	2.1	*	*	*	*	*	0.3	1.0	2.8	5.9	10.5	15.7
2000.....	2.1	*	*	*	*	*	0.4	1.2	2.8	6.5	9.6	16.8
1999.....	2.1	*	*	*	*	*	0.3	1.1	3.0	5.9	10.1	16.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic black, both sexes

2007 5/.....	2.8	*	*	*	*	*	1.0	4.2	10.0	13.9	13.8	14.6
2006 5/.....	2.7	*	*	*	*	*	0.8	4.4	9.7	13.9	13.8	10.1
2005 5/.....	2.9	*	*	*	*	*	0.9	5.1	10.9	14.2	13.7	12.8
2004 5/.....	3.1	*	*	*	*	*	1.3	6.2	11.4	14.1	15.3	12.7
2003 5/.....	3.0	*	*	*	*	*	1.1	5.2	11.5	15.7	15.6	15.5
2002.....	2.9	*	*	*	*	*	1.1	5.6	12.4	12.9	12.1	16.0
2001.....	3.0	*	*	*	*	*	1.3	5.3	12.9	15.7	13.9	13.1
2000.....	3.1	*	*	*	*	*	1.1	6.1	13.7	15.7	15.1	14.9
1999.....	3.3	*	*	*	*	*	1.2	7.3	13.6	15.4	16.8	14.5

Non-Hispanic black, male

2007 5/.....	4.2	*	*	*	*	*	1.4	6.8	16.8	26.1	24.2	24.4
2006 5/.....	4.1	*	*	*	*	*	1.3	7.1	17.4	25.5	25.9	*
2005 5/.....	4.7	*	*	*	*	*	1.6	8.8	20.0	26.7	27.8	21.9
2004 5/.....	4.9	*	*	*	*	*	1.9	10.7	20.3	27.1	26.2	27.4
2003 5/.....	4.8	*	*	*	*	*	1.2	9.2	21.8	29.1	28.6	25.6
2002.....	4.4	*	*	*	*	*	1.6	9.8	22.5	23.0	18.8	26.3
2001.....	4.6	*	*	*	*	*	1.7	8.9	23.9	27.3	23.4	*
2000.....	4.9	*	*	*	*	*	1.8	10.7	24.7	28.6	29.0	*
1999.....	5.0	*	*	*	*	*	1.8	12.3	23.5	25.4	31.9	25.0

Non-Hispanic black, female

2007 5/.....	1.5	*	*	*	*	*	*	2.0	4.5	5.2	7.8	10.5
2006 5/.....	1.3	*	*	*	*	*	*	2.0	3.6	5.7	6.9	8.6
2005 5/.....	1.2	*	*	*	*	*	*	1.9	3.5	5.2	5.7	9.2
2004 5/.....	1.4	*	*	*	*	*	0.9	2.3	4.3	4.8	9.2	*
2003 5/.....	1.4	*	*	*	*	*	1.1	1.8	3.3	6.2	8.3	11.6
2002.....	1.5	*	*	*	*	*	0.7	2.1	4.3	5.8	8.3	12.1
2001.....	1.5	*	*	*	*	*	0.9	2.1	4.0	7.5	8.6	10.7
2000.....	1.5	*	*	*	*	*	*	2.1	4.9	6.6	7.3	14.4
1999.....	1.7	*	*	*	*	*	0.7	3.0	5.7	8.4	8.4	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

All origins 3/, both sexes

2007.....	4.5	*	*	*	*	0.1	0.6	3.3	10.3	19.3	25.9	25.0
2006.....	4.6	*	*	*	*	0.1	0.6	3.5	10.4	19.5	26.9	25.6
2005.....	4.6	*	*	*	*	0.1	0.7	3.6	10.2	19.8	27.4	24.9
2004.....	4.4	*	*	*	*	0.1	0.7	3.4	10.3	19.2	26.7	25.8
2003.....	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002.....	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001.....	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000.....	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999.....	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8

All origins 3/, male

2007.....	7.2	*	*	*	*	0.1	1.0	5.8	18.2	33.8	47.8	48.4
2006.....	7.3	*	*	*	*	0.1	1.1	6.0	18.4	34.3	48.8	48.5
2005.....	7.3	*	*	*	*	0.1	1.2	6.2	17.9	34.9	50.7	46.0
2004.....	7.0	*	*	*	*	0.1	1.1	5.9	17.9	33.4	48.8	48.5
2003.....	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002.....	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001.....	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000.....	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999.....	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0

All origins 3/, female

2007.....	1.9	*	*	*	*	*	0.2	0.9	3.0	6.9	10.9	13.9
2006.....	2.0	*	*	*	*	*	0.2	1.1	3.0	7.0	12.0	14.9
2005.....	1.9	*	*	*	*	*	0.2	1.0	3.1	7.0	11.6	15.3
2004.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.3	11.9	15.7
2003.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000.....	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999.....	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

Hispanic, both sexes

2007.....	1.1	*	*	*	*	*	*	1.7	4.3	8.5	14.1	12.9
2006.....	1.2	*	*	*	*	*	0.3	1.5	4.9	9.9	15.9	14.9
2005.....	1.1	*	*	*	*	*	*	1.6	4.3	11.2	14.5	18.8
2004.....	1.1	*	*	*	*	*	*	1.0	4.7	10.3	15.0	18.6
2003.....	1.1	*	*	*	*	*	*	1.3	5.0	10.0	14.4	16.1
2002.....	1.1	*	*	*	*	*	0.4	1.5	4.1	10.2	15.2	23.3
2001.....	1.1	*	*	*	*	*	*	1.9	5.2	8.4	18.2	20.0
2000.....	1.1	*	*	*	*	*	*	1.2	5.0	11.1	19.4	19.9
1999.....	1.1	*	*	*	*	*	*	1.5	5.2	9.2	18.1	26.7

Hispanic, male

2007.....	1.7	*	*	*	*	*	*	3.1	8.2	15.1	26.0	24.4
2006.....	1.8	*	*	*	*	*	*	2.7	8.6	17.5	28.8	25.9
2005.....	1.8	*	*	*	*	*	*	2.7	7.8	20.7	26.7	37.4
2004.....	1.6	*	*	*	*	*	*	1.9	8.6	18.2	27.2	29.4
2003.....	1.6	*	*	*	*	*	*	2.4	8.0	17.2	28.4	*
2002.....	1.6	*	*	*	*	*	*	2.7	8.0	18.3	27.1	36.8
2001.....	1.7	*	*	*	*	*	*	3.6	8.5	16.0	30.4	41.9
2000.....	1.6	*	*	*	*	*	*	2.2	8.0	19.4	35.4	40.3
1999.....	1.7	*	*	*	*	*	*	2.7	10.2	16.3	32.8	49.6

Hispanic, female

2007.....	0.4	*	*	*	*	*	*	*	*	3.0	5.7	*
2006.....	0.5	*	*	*	*	*	*	*	1.5	3.7	6.9	*
2005.....	0.4	*	*	*	*	*	*	*	*	3.6	6.0	*
2004.....	0.5	*	*	*	*	*	*	*	*	4.0	6.7	*
2003.....	0.5	*	*	*	*	*	*	*	2.4	4.3	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	3.6	7.0	16.6
2001.....	0.5	*	*	*	*	*	*	*	2.2	*	10.0	*
2000.....	0.5	*	*	*	*	*	*	*	2.4	4.6	8.6	*
1999.....	0.4	*	*	*	*	*	*	*	*	3.6	8.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic 4/, both sexes

2007.....	5.1	*	*	*	*	0.1	0.6	3.5	10.9	20.1	26.7	25.7
2006.....	5.2	*	*	*	*	0.1	0.7	3.7	10.9	20.2	27.6	26.1
2005.....	5.1	*	*	*	*	0.1	0.8	3.8	10.7	20.4	28.2	25.2
2004.....	5.0	*	*	*	*	0.1	0.7	3.7	10.7	19.9	27.3	26.2
2003.....	4.9	*	*	*	*	0.1	0.7	3.7	10.9	20.4	27.0	25.4
2002.....	4.9	*	*	*	*	*	0.7	3.7	10.8	20.8	27.2	24.5
2001.....	4.9	*	*	*	*	0.1	0.7	3.7	10.4	21.3	26.6	26.0
2000.....	4.8	*	*	*	*	0.1	0.7	3.8	11.2	20.4	25.8	26.1
1999.....	4.7	*	*	*	*	0.1	0.7	3.6	11.0	20.6	24.6	26.8

Non-Hispanic 4/, male

2007.....	8.3	*	*	*	*	0.1	1.1	6.1	19.0	35.3	49.2	49.7
2006.....	8.3	*	*	*	*	0.1	1.2	6.4	19.2	35.5	50.0	49.8
2005.....	8.2	*	*	*	*	0.2	1.4	6.6	18.7	35.9	52.2	46.3
2004.....	7.9	*	*	*	*	0.2	1.2	6.3	18.6	34.5	49.9	49.4
2003.....	7.8	*	*	*	*	0.1	1.2	6.4	19.3	35.3	48.8	46.3
2002.....	7.8	*	*	*	*	*	1.2	6.3	18.9	36.6	49.3	43.5
2001.....	7.7	*	*	*	*	0.1	1.3	6.5	18.2	37.1	48.1	48.4
2000.....	7.5	*	*	*	*	0.2	1.3	6.6	19.4	34.8	46.1	47.8
1999.....	7.4	*	*	*	*	0.2	1.2	6.4	18.8	36.2	44.7	48.0

Non-Hispanic 4/, female

2007.....	2.1	*	*	*	*	*	0.2	1.0	3.2	7.2	11.2	14.3
2006.....	2.2	*	*	*	*	*	0.2	1.2	3.1	7.3	12.3	15.1
2005.....	2.2	*	*	*	*	*	0.3	1.0	3.2	7.2	11.9	15.5
2004.....	2.2	*	*	*	*	*	0.2	1.1	3.4	7.6	12.2	15.8
2003.....	2.2	*	*	*	*	*	0.2	1.1	3.2	7.9	12.4	16.1
2002.....	2.2	*	*	*	*	*	0.3	1.1	3.4	7.6	12.5	16.3
2001.....	2.2	*	*	*	*	*	0.3	1.1	3.2	8.2	12.5	16.7
2000.....	2.3	*	*	*	*	*	0.1	1.0	3.6	8.5	12.6	17.3
1999.....	2.2	*	*	*	*	*	0.2	0.9	3.8	7.8	11.6	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic white, both sexes

2007 5/.....	5.7	*	*	*	*	0.1	0.7	3.5	10.8	20.6	27.3	26.5
2006 5/.....	5.7	*	*	*	*	0.1	0.8	3.7	10.9	20.6	28.3	27.0
2005 5/.....	5.6	*	*	*	*	0.1	0.8	3.8	10.6	20.7	28.8	26.2
2004 5/.....	5.4	*	*	*	*	0.1	0.8	3.6	10.5	19.9	27.8	26.5
2003 5/.....	5.3	*	*	*	*	0.1	0.7	3.4	10.6	20.3	27.3	26.1
2002.....	5.3	*	*	*	*	*	0.7	3.6	10.4	20.5	27.3	25.1
2001.....	5.1	*	*	*	*	0.1	0.8	3.5	9.7	20.5	26.8	26.5
2000.....	5.1	*	*	*	*	0.1	0.7	3.5	10.4	20.0	25.5	26.7
1999.....	4.9	*	*	*	*	0.1	0.7	3.3	10.2	20.0	24.4	27.2

Non-Hispanic white, male

2007 5/.....	9.2	*	*	*	*	0.2	1.2	6.2	19.1	36.2	50.2	51.3
2006 5/.....	9.2	*	*	*	*	*	1.4	6.4	19.4	36.0	51.7	51.3
2005 5/.....	9.1	*	*	*	*	0.2	1.4	6.7	18.7	36.4	53.4	49.0
2004 5/.....	8.7	*	*	*	*	*	1.3	6.3	18.4	34.6	51.0	50.1
2003 5/.....	8.5	*	*	*	*	*	1.2	6.0	18.8	35.1	49.9	47.9
2002.....	8.4	*	*	*	*	*	1.3	6.3	18.2	36.3	49.9	44.7
2001.....	8.1	*	*	*	*	*	1.3	6.1	17.0	36.0	48.6	49.0
2000.....	8.0	*	*	*	*	0.2	1.4	6.3	18.3	34.3	45.9	49.4
1999.....	7.7	*	*	*	*	0.2	1.2	6.0	17.5	35.4	44.4	48.0

Non-Hispanic white, female

2007 5/.....	2.3	*	*	*	*	*	0.2	0.9	3.0	7.1	11.3	14.7
2006 5/.....	2.3	*	*	*	*	*	0.2	1.1	2.8	7.2	12.2	15.7
2005 5/.....	2.3	*	*	*	*	*	0.2	0.8	3.0	7.1	11.8	15.8
2004 5/.....	2.3	*	*	*	*	*	0.2	0.9	3.0	7.2	12.0	16.0
2003 5/.....	2.3	*	*	*	*	*	0.2	0.9	2.8	7.7	12.0	16.5
2002.....	2.3	*	*	*	*	*	0.2	0.9	3.0	7.1	12.2	16.6
2001.....	2.3	*	*	*	*	*	0.2	1.0	2.7	7.5	12.4	17.1
2000.....	2.3	*	*	*	*	*	*	0.8	3.0	8.0	12.0	17.5
1999.....	2.2	*	*	*	*	*	0.2	0.8	3.3	7.1	11.3	18.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	3.8	*	*	*	*	*	0.6	4.3	14.3	21.8	25.7	19.2
2006 5/.....	3.8	*	*	*	*	*	0.6	5.2	13.9	22.5	24.9	18.3
2005 5/.....	3.9	*	*	*	*	*	0.9	4.9	14.5	22.9	27.7	17.5
2004 5/.....	4.1	*	*	*	*	*	0.7	5.3	15.8	25.2	27.4	21.9
2003 5/.....	4.3	*	*	*	*	*	0.9	6.1	17.3	26.1	28.4	21.4
2002.....	4.3	*	*	*	*	*	1.0	5.2	17.7	28.9	29.9	20.6
2001.....	4.7	*	*	*	*	*	1.0	6.1	18.9	32.8	30.0	24.0
2000.....	4.7	*	*	*	*	*	0.8	6.8	20.2	28.6	34.1	22.6
1999.....	4.7	*	*	*	*	*	1.0	6.6	20.3	30.2	31.4	23.6
<b>Non-Hispanic black, male</b>												
2007 5/.....	5.8	*	*	*	*	*	0.8	7.0	24.6	39.1	47.6	41.0
2006 5/.....	5.8	*	*	*	*	*	1.0	8.4	23.2	40.6	43.1	35.7
2005 5/.....	6.0	*	*	*	*	*	1.5	8.0	25.2	41.1	52.1	26.6
2004 5/.....	6.0	*	*	*	*	*	1.0	8.5	26.0	42.2	47.6	39.6
2003 5/.....	6.5	*	*	*	*	*	1.4	10.4	29.4	45.2	46.7	38.4
2002.....	6.5	*	*	*	*	*	1.5	8.4	30.7	49.8	52.2	38.3
2001.....	7.2	*	*	*	*	*	1.6	10.5	33.0	56.2	55.5	47.4
2000.....	7.0	*	*	*	*	*	1.2	11.5	34.8	47.6	58.9	38.7
1999.....	7.2	*	*	*	*	*	1.7	11.5	35.3	51.1	57.2	49.9
<b>Non-Hispanic black, female</b>												
2007 5/.....	2.0	*	*	*	*	*	*	2.1	6.0	9.4	13.3	10.1
2006 5/.....	2.0	*	*	*	*	*	*	2.4	6.4	9.6	14.5	11.2
2005 5/.....	2.0	*	*	*	*	*	*	2.3	5.9	9.9	13.8	13.8
2004 5/.....	2.4	*	*	*	*	*	*	2.5	7.6	13.1	15.9	14.9
2003 5/.....	2.4	*	*	*	*	*	*	2.5	7.5	12.5	18.1	14.9
2002.....	2.4	*	*	*	*	*	*	2.4	7.2	14.0	17.2	13.8
2001.....	2.4	*	*	*	*	*	*	2.3	7.5	16.2	15.7	15.4
2000.....	2.6	*	*	*	*	*	*	2.7	8.5	15.2	20.1	16.6
1999.....	2.4	*	*	*	*	*	*	2.4	8.3	15.5	17.0	13.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

All origins 3/, both sexes

2007.....	3.8	*	*	*	*	0.3	1.1	2.6	5.7	13.2	25.0	35.4
2006.....	3.8	*	*	*	*	0.3	1.1	2.7	5.9	13.0	25.6	35.8
2005.....	3.9	*	*	*	0.0	0.3	1.0	2.6	5.9	14.0	26.6	38.5
2004.....	4.0	*	*	*	*	0.3	1.1	2.7	6.5	14.9	26.3	42.2
2003.....	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002.....	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001.....	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000.....	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999.....	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1

All origins 3/, male

2007.....	4.5	*	*	*	*	0.3	1.1	3.3	8.0	19.1	35.2	51.1
2006.....	4.5	*	*	*	*	0.3	1.1	3.4	8.3	18.0	36.8	50.2
2005.....	4.6	*	*	*	*	0.3	1.2	3.3	8.2	19.7	37.4	54.2
2004.....	4.8	*	*	*	*	0.3	1.2	3.5	9.1	20.6	37.2	62.9
2003.....	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002.....	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001.....	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000.....	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999.....	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3

All origins 3/, female

2007.....	3.0	*	*	*	*	0.3	1.0	2.0	3.7	8.2	17.9	27.9
2006.....	3.1	*	*	*	*	0.3	1.1	2.0	3.7	8.8	18.0	29.1
2005.....	3.2	*	*	*	*	0.3	0.8	1.9	3.7	9.2	19.3	31.2
2004.....	3.3	*	*	*	*	0.3	0.9	2.0	4.1	10.1	19.0	32.9
2003.....	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002.....	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001.....	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000.....	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999.....	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of stomach (C16)</b>												
<b>Hispanic, both sexes</b>												
2007.....	2.9	*	*	*	*	0.6	1.7	3.8	7.7	19.6	37.9	59.7
2006.....	2.9	*	*	*	*	0.6	1.7	3.9	10.2	20.4	37.2	47.0
2005.....	3.0	*	*	*	*	0.5	1.5	4.6	9.3	20.7	44.0	50.9
2004.....	3.3	*	*	*	*	0.5	2.0	4.6	11.7	24.4	44.0	62.8
2003.....	3.1	*	*	*	*	0.5	1.6	5.1	11.4	23.8	43.6	62.7
2002.....	3.1	*	*	*	*	0.5	1.7	4.2	11.0	21.9	47.2	72.1
2001.....	3.2	*	*	*	*	0.6	1.8	5.4	11.3	26.4	45.6	63.8
2000.....	3.1	*	*	*	*	0.4	2.0	4.8	9.8	28.4	50.2	65.7
1999.....	3.1	*	*	*	*	0.6	1.7	4.1	11.7	26.2	49.8	73.7
<b>Hispanic, male</b>												
2007.....	3.2	*	*	*	*	0.6	1.6	4.0	10.8	28.3	52.3	84.4
2006.....	3.1	*	*	*	*	0.5	1.4	4.5	13.1	27.1	49.3	55.3
2005.....	3.4	*	*	*	*	0.5	1.5	5.0	12.1	27.6	65.8	76.0
2004.....	3.6	*	*	*	*	0.5	2.0	4.7	14.8	32.0	66.9	84.1
2003.....	3.4	*	*	*	*	*	1.5	5.7	14.1	33.2	63.7	73.9
2002.....	3.4	*	*	*	*	0.7	1.5	4.7	13.7	32.9	61.6	107.1
2001.....	3.6	*	*	*	*	*	1.9	6.2	15.0	37.1	60.4	95.2
2000.....	3.6	*	*	*	*	*	2.0	5.0	13.7	43.3	70.8	98.8
1999.....	3.5	*	*	*	*	0.7	1.6	5.2	16.4	35.6	73.5	75.5
<b>Hispanic, female</b>												
2007.....	2.5	*	*	*	*	0.6	1.8	3.6	4.7	12.6	27.8	46.3
2006.....	2.7	*	*	*	*	0.7	2.1	3.3	7.5	14.9	28.8	42.5
2005.....	2.6	*	*	*	*	*	1.4	4.2	6.8	15.2	28.8	37.8
2004.....	2.9	*	*	*	*	*	2.1	4.5	8.8	18.3	28.2	51.9
2003.....	2.8	*	*	*	*	*	1.6	4.5	9.0	16.3	29.8	57.0
2002.....	2.7	*	*	*	*	*	1.9	3.8	8.7	13.2	37.4	54.7
2001.....	2.8	*	*	*	*	0.7	1.7	4.6	8.1	18.0	35.7	48.4
2000.....	2.7	*	*	*	*	*	2.1	4.6	6.3	16.8	36.3	49.5
1999.....	2.7	*	*	*	*	*	1.8	3.0	7.6	18.8	34.0	72.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of stomach (C16)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	3.9	*	*	*	*	0.2	0.9	2.5	5.6	12.7	24.1	34.1
2006.....	3.9	*	*	*	*	0.2	0.9	2.5	5.5	12.4	24.9	35.2
2005.....	4.0	*	*	*	*	0.2	0.9	2.4	5.6	13.5	25.5	37.8
2004.....	4.2	*	*	*	*	0.3	0.9	2.5	6.0	14.2	25.3	41.2
2003.....	4.3	*	*	*	*	0.3	1.0	2.7	6.3	14.6	26.7	42.7
2002.....	4.4	*	*	*	0.1	0.3	0.9	2.6	6.5	14.9	27.7	46.3
2001.....	4.5	*	*	*	*	0.2	0.9	2.6	6.6	15.8	29.4	44.7
2000.....	4.7	*	*	*	*	0.3	0.9	2.7	7.2	15.8	30.1	52.0
1999.....	4.7	*	*	*	*	0.3	1.0	2.9	7.5	16.5	30.8	49.2
<b>Non-Hispanic 4/, male</b>												
2007.....	4.8	*	*	*	*	0.2	1.1	3.2	7.7	18.4	34.1	49.2
2006.....	4.8	*	*	*	*	0.2	1.0	3.3	7.9	17.3	36.0	49.9
2005.....	4.8	*	*	*	*	0.2	1.1	3.1	7.9	19.1	35.6	53.0
2004.....	5.0	*	*	*	*	0.2	1.1	3.4	8.6	19.7	35.5	61.7
2003.....	5.2	*	*	*	*	0.3	1.1	3.6	9.2	21.0	36.9	59.2
2002.....	5.3	*	*	*	*	0.3	1.0	3.6	9.1	21.7	39.4	65.0
2001.....	5.4	*	*	*	*	0.3	1.0	3.6	9.8	22.3	42.1	61.0
2000.....	5.5	*	*	*	*	0.3	1.2	3.7	10.3	22.4	42.6	74.5
1999.....	5.7	*	*	*	*	0.3	1.2	4.0	10.6	23.4	45.3	71.0
<b>Non-Hispanic 4/, female</b>												
2007.....	3.1	*	*	*	*	0.3	0.8	1.8	3.6	7.8	17.3	27.0
2006.....	3.1	*	*	*	*	0.3	0.9	1.8	3.4	8.3	17.3	28.4
2005.....	3.3	*	*	*	*	0.3	0.7	1.6	3.4	8.7	18.7	30.9
2004.....	3.3	*	*	*	*	0.3	0.8	1.7	3.7	9.5	18.4	32.1
2003.....	3.5	*	*	*	*	0.3	0.9	1.9	3.7	9.2	19.8	35.5
2002.....	3.5	*	*	*	*	0.3	0.7	1.7	4.1	9.2	19.9	38.3
2001.....	3.6	*	*	*	*	0.2	0.8	1.7	3.7	10.4	21.0	37.9
2000.....	3.8	*	*	*	*	0.3	0.7	1.8	4.4	10.4	21.9	42.9
1999.....	3.8	*	*	*	*	0.3	0.9	1.8	4.7	10.8	21.3	40.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

Non-Hispanic white, both sexes

2007 5/.....	3.6	*	*	*	*	0.2	0.8	1.9	4.4	10.6	20.7	30.3
2006 5/.....	3.6	*	*	*	*	0.2	0.7	2.1	4.6	10.4	21.4	31.5
2005 5/.....	3.7	*	*	*	*	0.1	0.7	1.9	4.6	11.3	22.0	33.8
2004 5/.....	3.9	*	*	*	*	0.2	0.7	2.0	4.9	11.8	22.3	37.6
2003 5/.....	4.0	*	*	*	*	0.2	0.8	2.1	5.4	12.6	23.3	37.5
2002.....	4.1	*	*	*	*	0.2	0.7	2.1	5.4	12.6	24.1	40.7
2001.....	4.2	*	*	*	*	0.2	0.7	2.1	5.6	13.5	25.8	40.0
2000.....	4.4	*	*	*	*	0.2	0.7	2.2	6.1	13.5	26.1	46.3
1999.....	4.4	*	*	*	*	0.2	0.7	2.3	6.1	14.4	27.0	43.5

Non-Hispanic white, male

2007 5/.....	4.4	*	*	*	*	0.2	0.9	2.5	6.2	15.6	28.8	44.0
2006 5/.....	4.5	*	*	*	*	*	0.7	2.8	6.6	14.7	31.3	45.2
2005 5/.....	4.4	*	*	*	*	*	0.8	2.5	6.5	15.9	30.8	47.2
2004 5/.....	4.7	*	*	*	*	0.2	0.9	2.7	7.1	16.3	31.1	56.9
2003 5/.....	4.8	*	*	*	*	0.2	0.9	2.8	7.7	18.0	32.4	52.3
2002.....	5.0	*	*	*	*	0.2	0.8	2.9	7.8	18.6	35.0	58.0
2001.....	5.0	*	*	*	*	0.2	0.8	2.8	8.2	18.8	37.4	55.7
2000.....	5.2	*	*	*	*	0.2	0.8	3.0	8.9	19.3	37.4	67.3
1999.....	5.4	*	*	*	*	0.2	0.9	3.3	8.7	20.6	39.9	62.6

Non-Hispanic white, female

2007 5/.....	2.8	*	*	*	*	0.2	0.6	1.4	2.8	6.3	14.9	23.9
2006 5/.....	2.8	*	*	*	*	0.2	0.6	1.4	2.7	6.7	14.5	25.2
2005 5/.....	3.0	*	*	*	*	*	0.6	1.3	2.8	7.3	15.9	27.7
2004 5/.....	3.1	*	*	*	*	0.2	0.5	1.2	2.9	8.0	16.3	28.9
2003 5/.....	3.3	*	*	*	*	0.2	0.6	1.4	3.1	8.0	17.2	31.0
2002.....	3.2	*	*	*	*	0.2	0.5	1.2	3.2	7.5	16.8	33.3
2001.....	3.4	*	*	*	*	*	0.6	1.3	3.1	9.0	18.1	33.5
2000.....	3.5	*	*	*	*	0.2	0.5	1.4	3.6	8.6	18.7	37.9
1999.....	3.5	*	*	*	*	0.2	0.6	1.4	3.7	9.1	18.6	35.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of stomach (C16)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	5.2	*	*	*	*	*	1.5	5.1	12.1	25.3	50.6	65.5
2006 5/.....	5.0	*	*	*	*	0.4	1.8	4.5	11.1	24.8	52.0	68.2
2005 5/.....	5.2	*	*	*	*	0.4	1.8	4.6	11.5	27.0	52.6	75.4
2004 5/.....	5.4	*	*	*	*	0.5	1.7	5.4	12.9	30.8	50.9	71.4
2003 5/.....	5.5	*	*	*	*	0.4	1.8	5.8	12.9	27.5	55.0	86.0
2002.....	5.8	*	*	*	*	0.6	1.6	5.5	13.2	31.5	57.7	100.1
2001.....	5.8	*	*	*	*	0.5	2.2	5.5	13.1	31.0	59.1	89.3
2000.....	6.1	*	*	*	*	0.7	2.1	5.8	14.1	32.3	64.3	102.7
1999.....	6.3	*	*	*	*	0.7	2.3	6.0	16.1	31.5	65.1	103.1
<b>Non-Hispanic black, male</b>												
2007 5/.....	6.5	*	*	*	*	*	1.7	7.4	17.8	39.0	79.5	96.0
2006 5/.....	6.0	*	*	*	*	*	2.0	6.2	16.7	34.8	82.0	85.2
2005 5/.....	6.4	*	*	*	*	*	2.6	6.5	18.3	41.6	72.0	117.0
2004 5/.....	6.7	*	*	*	*	*	1.9	7.3	19.6	48.2	75.9	99.6
2003 5/.....	6.8	*	*	*	*	*	2.0	8.5	20.0	44.8	80.5	107.6
2002.....	6.9	*	*	*	*	0.8	1.9	7.9	19.5	47.3	80.9	136.7
2001.....	6.9	*	*	*	*	*	2.4	8.0	20.0	49.2	87.9	106.3
2000.....	7.2	*	*	*	*	0.8	3.0	8.1	20.9	47.0	91.0	138.6
1999.....	7.6	*	*	*	*	*	2.8	8.7	23.3	46.4	101.6	149.8
<b>Non-Hispanic black, female</b>												
2007 5/.....	4.0	*	*	*	*	*	1.4	3.1	7.5	15.6	34.1	52.8
2006 5/.....	4.1	*	*	*	*	*	1.6	2.9	6.7	17.6	35.0	61.3
2005 5/.....	4.1	*	*	*	*	*	1.1	3.0	6.0	16.5	41.6	58.6
2004 5/.....	4.3	*	*	*	*	*	1.5	3.7	7.6	18.4	36.9	60.2
2003 5/.....	4.4	*	*	*	*	*	1.6	3.5	7.3	15.2	40.6	77.6
2002.....	4.9	*	*	*	*	*	1.4	3.4	8.1	20.3	44.6	86.1
2001.....	4.7	*	*	*	*	*	2.0	3.3	7.6	18.1	43.0	83.0
2000.....	5.2	*	*	*	*	*	1.4	3.9	8.6	22.0	49.3	89.4
1999.....	5.1	*	*	*	*	*	1.9	3.7	10.2	21.0	44.6	85.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes

2007.....	17.8	*	*	*	0.1	0.7	3.0	10.9	27.7	60.1	118.4	200.8
2006.....	17.9	*	*	*	0.1	0.8	3.0	10.4	27.3	61.9	122.8	207.5
2005.....	18.0	*	*	*	0.1	0.8	3.0	10.2	26.9	63.3	124.5	216.9
2004.....	18.3	*	*	*	0.1	0.7	3.0	10.3	28.2	64.7	128.9	225.6
2003.....	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002.....	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001.....	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000.....	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999.....	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7

All origins 3/, male

2007.....	18.2	*	*	*	0.1	0.7	3.3	12.3	33.8	74.5	144.0	223.5
2006.....	18.3	*	*	*	0.1	0.8	3.2	11.6	33.0	76.8	149.5	234.6
2005.....	18.4	*	*	*	0.1	0.8	3.2	11.5	32.9	79.6	151.6	245.9
2004.....	18.6	*	*	*	0.2	0.7	3.2	11.9	34.2	80.6	156.9	254.7
2003.....	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002.....	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001.....	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000.....	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999.....	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0

All origins 3/, female

2007.....	17.3	*	*	*	*	0.6	2.8	9.6	22.0	47.9	100.7	190.0
2006.....	17.5	*	*	*	0.1	0.8	2.9	9.2	22.1	49.3	104.6	194.9
2005.....	17.6	*	*	*	0.1	0.7	2.8	9.0	21.2	49.5	106.1	203.6
2004.....	18.0	*	*	*	0.1	0.6	2.9	8.7	22.7	51.4	110.0	212.5
2003.....	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002.....	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001.....	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000.....	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999.....	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Hispanic, both sexes

2007.....	6.2	*	*	*	*	0.5	2.1	8.0	21.3	42.0	85.9	133.4
2006.....	6.2	*	*	*	*	0.5	2.0	7.5	20.4	45.7	93.7	142.4
2005.....	6.0	*	*	*	*	0.5	1.9	7.1	18.4	46.5	92.4	146.7
2004.....	5.9	*	*	*	*	0.6	1.9	7.2	21.2	47.5	87.5	147.5
2003.....	6.2	*	*	*	*	0.5	2.1	8.2	23.6	49.7	95.5	144.6
2002.....	6.1	*	*	*	*	0.4	2.1	7.7	23.0	51.2	95.6	165.7
2001.....	6.1	*	*	*	*	0.3	2.3	7.2	22.5	52.1	96.5	193.8
2000.....	6.0	*	*	*	*	0.5	2.3	8.4	23.1	49.2	98.2	181.1
1999.....	6.0	*	*	*	*	0.5	1.7	8.3	21.0	56.1	101.0	176.8

Hispanic, male

2007.....	6.4	*	*	*	*	0.5	2.4	9.1	25.7	56.6	101.3	145.3
2006.....	6.3	*	*	*	*	0.5	1.7	7.8	25.5	60.3	112.9	157.9
2005.....	6.1	*	*	*	*	*	1.8	8.0	23.3	60.3	111.6	176.8
2004.....	6.2	*	*	*	*	0.5	2.2	8.5	26.6	59.4	114.1	172.3
2003.....	6.6	*	*	*	*	0.5	2.3	10.0	29.8	64.8	120.4	172.4
2002.....	6.4	*	*	*	*	*	2.0	9.3	31.1	67.1	121.1	174.0
2001.....	6.2	*	*	*	*	*	2.7	7.4	28.2	66.5	116.7	257.1
2000.....	6.2	*	*	*	*	*	2.0	9.0	28.2	63.5	131.3	235.8
1999.....	6.1	*	*	*	*	*	1.5	9.1	24.1	72.7	134.0	211.4

Hispanic, female

2007.....	5.9	*	*	*	*	*	1.8	6.9	17.2	30.1	75.2	126.9
2006.....	6.1	*	*	*	*	*	2.3	7.2	15.8	33.7	80.3	134.2
2005.....	5.8	*	*	*	*	0.7	2.0	6.2	13.9	35.3	79.1	131.0
2004.....	5.6	*	*	*	*	0.6	1.4	5.8	16.3	38.0	69.1	134.7
2003.....	5.7	*	*	*	*	*	1.8	6.4	18.0	37.6	78.5	130.5
2002.....	5.7	*	*	*	*	*	2.1	6.1	15.7	38.6	78.1	161.6
2001.....	5.9	*	*	*	*	*	1.9	7.1	17.4	40.8	82.8	162.9
2000.....	5.8	*	*	*	*	*	2.7	7.8	18.5	38.0	75.9	154.3
1999.....	5.9	*	*	*	*	*	2.0	7.5	18.3	43.0	79.1	160.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 4/, both sexes

2007.....	19.8	*	*	*	0.1	0.7	3.2	11.3	28.2	61.5	120.4	204.3
2006.....	19.9	*	*	*	0.1	0.9	3.2	10.7	27.8	63.1	124.5	210.6
2005.....	20.0	*	*	*	0.1	0.8	3.1	10.5	27.5	64.4	126.2	220.1
2004.....	20.3	*	*	*	0.1	0.7	3.2	10.6	28.8	65.9	130.9	228.8
2003.....	21.3	*	*	*	0.1	0.8	3.1	11.1	30.2	71.9	136.5	243.2
2002.....	21.7	*	*	*	0.1	0.8	3.3	11.8	32.3	74.6	139.1	244.0
2001.....	22.0	*	*	*	0.1	0.9	3.3	12.0	32.2	76.7	141.1	257.4
2000.....	22.4	*	*	*	0.1	0.7	3.1	11.5	33.3	78.2	147.8	273.1
1999.....	22.4	*	*	*	0.1	0.7	3.1	11.3	34.3	78.5	148.1	274.3

Non-Hispanic 4/, male

2007.....	20.4	*	*	*	0.1	0.8	3.4	12.7	34.5	75.7	146.8	228.0
2006.....	20.4	*	*	*	*	0.9	3.5	12.0	33.5	77.9	151.7	238.4
2005.....	20.5	*	*	*	0.1	0.9	3.4	11.9	33.7	80.9	153.7	249.2
2004.....	20.7	*	*	*	0.2	0.8	3.3	12.3	34.8	81.9	159.1	258.5
2003.....	21.7	*	*	*	*	0.9	3.4	12.7	36.4	89.4	166.3	277.3
2002.....	22.3	*	*	*	0.2	0.9	3.8	13.4	39.9	92.7	169.3	279.2
2001.....	22.3	*	*	*	0.1	0.9	3.6	13.6	38.9	95.2	171.8	299.6
2000.....	22.8	*	*	*	0.1	0.8	3.6	13.1	40.3	97.2	181.3	318.6
1999.....	22.7	*	*	*	0.1	0.9	3.2	12.8	41.6	96.2	183.2	327.5

Non-Hispanic 4/, female

2007.....	19.2	*	*	*	*	0.6	2.9	9.9	22.4	49.3	102.3	193.0
2006.....	19.4	*	*	*	*	0.9	3.0	9.4	22.6	50.4	105.9	197.7
2005.....	19.4	*	*	*	*	0.7	2.9	9.3	21.8	50.5	107.6	206.8
2004.....	19.9	*	*	*	*	0.7	3.1	8.9	23.2	52.3	112.0	215.6
2003.....	20.9	*	*	*	*	0.7	2.8	9.6	24.4	57.2	116.6	228.1
2002.....	21.2	*	*	*	*	0.7	2.9	10.2	25.2	59.6	119.1	228.8
2001.....	21.7	*	*	*	*	0.8	3.0	10.5	25.9	61.3	121.0	239.9
2000.....	22.1	*	*	*	*	0.6	2.7	9.9	26.9	62.6	126.1	254.7
1999.....	22.2	*	*	*	*	0.6	3.0	9.9	27.6	63.9	125.3	253.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic white, both sexes

2007 5/.....	21.1	*	*	*	*	0.7	3.0	10.4	26.3	59.5	119.0	204.3
2006 5/.....	21.2	*	*	*	0.1	0.8	3.0	9.9	26.3	61.3	122.0	210.5
2005 5/.....	21.2	*	*	*	0.1	0.8	3.0	9.6	25.7	62.5	124.0	218.8
2004 5/.....	21.7	*	*	*	0.1	0.7	3.0	9.8	27.3	63.8	129.4	228.5
2003 5/.....	22.7	*	*	*	*	0.7	2.8	10.1	28.5	69.9	135.0	242.1
2002.....	23.2	*	*	*	0.1	0.8	3.0	10.9	30.6	73.0	136.8	243.2
2001.....	23.5	*	*	*	0.1	0.8	3.1	11.1	30.6	74.7	138.5	256.9
2000.....	24.0	*	*	*	0.1	0.6	2.8	10.6	31.7	76.5	145.7	271.9
1999.....	24.0	*	*	*	0.1	0.7	2.9	10.6	32.8	76.7	145.9	273.4

Non-Hispanic white, male

2007 5/.....	21.7	*	*	*	*	0.8	3.2	11.7	32.2	73.3	145.5	227.6
2006 5/.....	21.6	*	*	*	*	0.8	3.2	11.1	31.8	75.5	148.1	236.7
2005 5/.....	21.8	*	*	*	*	0.8	3.3	10.9	31.5	78.5	151.2	248.8
2004 5/.....	22.2	*	*	*	*	0.7	3.1	11.5	33.3	79.4	156.5	259.6
2003 5/.....	23.2	*	*	*	*	0.8	3.1	11.6	34.5	87.4	164.5	276.8
2002.....	23.8	*	*	*	0.2	0.9	3.4	12.5	38.0	91.1	165.8	277.6
2001.....	23.9	*	*	*	*	0.9	3.3	12.8	37.4	93.2	169.3	298.2
2000.....	24.2	*	*	*	*	0.8	3.2	12.2	38.2	95.1	178.4	315.8
1999.....	24.3	*	*	*	*	0.8	3.1	12.1	39.7	94.8	180.6	328.2

Non-Hispanic white, female

2007 5/.....	20.5	*	*	*	*	0.6	2.9	9.2	20.6	47.6	100.5	193.3
2006 5/.....	20.7	*	*	*	*	0.7	2.7	8.7	21.0	49.0	103.9	198.2
2005 5/.....	20.6	*	*	*	*	0.7	2.6	8.3	20.2	48.6	105.2	205.1
2004 5/.....	21.3	*	*	*	*	0.6	2.8	8.1	21.7	50.3	110.9	214.6
2003 5/.....	22.2	*	*	*	*	0.7	2.5	8.5	22.8	54.9	115.0	226.8
2002.....	22.7	*	*	*	*	0.7	2.7	9.4	23.7	57.6	117.3	228.3
2001.....	23.1	*	*	*	*	0.7	2.8	9.5	24.2	59.0	118.1	239.8
2000.....	23.8	*	*	*	*	0.5	2.4	9.0	25.5	60.9	124.1	254.1
1999.....	23.7	*	*	*	*	0.6	2.7	9.0	26.2	61.5	123.2	251.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic black, both sexes

2007 5/.....	18.0	*	*	*	*	1.0	4.3	18.4	47.7	88.6	157.3	229.9
2006 5/.....	18.1	*	*	*	*	1.3	5.2	17.5	44.2	90.5	171.2	244.7
2005 5/.....	18.2	*	*	*	*	1.2	4.4	18.1	46.1	92.4	170.7	264.1
2004 5/.....	17.8	*	*	*	*	1.0	4.6	16.9	45.8	96.4	167.6	262.5
2003 5/.....	18.8	*	*	*	*	1.0	5.2	19.2	48.9	102.2	175.0	282.9
2002.....	18.9	*	*	*	*	0.7	5.1	19.0	50.5	100.6	185.9	278.3
2001.....	19.2	*	*	*	*	1.1	5.0	19.5	50.6	105.9	189.7	289.0
2000.....	19.4	*	*	*	*	1.0	5.2	20.0	51.1	106.3	192.5	312.9
1999.....	19.3	*	*	*	*	1.1	4.9	18.1	53.7	107.2	194.5	308.8

Non-Hispanic black, male

2007 5/.....	18.8	*	*	*	*	1.1	5.0	21.5	59.4	114.3	196.1	273.1
2006 5/.....	18.9	*	*	*	*	1.3	5.7	20.5	53.0	118.6	223.6	314.3
2005 5/.....	18.5	*	*	*	*	1.5	4.5	20.2	58.5	116.4	213.6	297.8
2004 5/.....	18.0	*	*	*	*	1.1	4.7	19.5	54.7	122.8	218.8	277.4
2003 5/.....	19.0	*	*	*	*	1.0	5.4	22.4	59.9	128.2	217.0	321.8
2002.....	19.3	*	*	*	*	0.9	6.0	21.8	62.7	123.0	240.9	332.6
2001.....	19.0	*	*	*	*	1.2	5.8	21.4	59.3	130.7	229.9	352.3
2000.....	19.8	*	*	*	*	1.1	6.1	22.7	63.7	133.7	244.4	381.6
1999.....	19.2	*	*	*	*	1.2	4.8	20.6	67.0	127.8	245.8	351.8

Non-Hispanic black, female

2007 5/.....	17.2	*	*	*	*	0.9	3.7	15.7	38.3	70.2	135.3	211.9
2006 5/.....	17.3	*	*	*	*	1.3	4.7	14.9	37.2	70.5	141.5	216.4
2005 5/.....	17.9	*	*	*	*	0.8	4.3	16.3	36.2	75.3	146.3	250.5
2004 5/.....	17.6	*	*	*	*	1.0	4.4	14.8	38.7	77.6	138.8	256.6
2003 5/.....	18.7	*	*	*	*	0.9	5.0	16.4	40.1	83.8	151.2	267.8
2002.....	18.4	*	*	*	*	*	4.3	16.7	40.7	84.7	154.9	257.5
2001.....	19.4	*	*	*	*	1.0	4.3	17.9	43.6	88.4	167.1	265.6
2000.....	19.1	*	*	*	*	1.0	4.4	17.6	40.9	87.1	163.2	287.4
1999.....	19.5	*	*	*	*	0.9	5.0	15.9	43.1	92.7	165.8	292.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes

2007.....	5.7	*	*	0.1	0.1	0.2	0.9	5.7	12.8	20.1	32.8	32.0
2006.....	5.5	*	*	*	0.1	0.2	0.8	5.6	12.2	20.2	31.9	33.5
2005.....	5.4	*	0.1	*	0.1	0.2	0.9	5.9	11.3	20.1	32.0	32.5
2004.....	5.2	*	0.1	*	0.1	0.2	0.9	5.6	10.7	19.9	31.3	32.4
2003.....	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002.....	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001.....	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000.....	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999.....	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6

All origins 3/, male

2007.....	7.6	*	*	*	*	0.3	1.2	9.1	20.5	28.7	46.0	43.3
2006.....	7.3	*	*	*	*	0.3	1.1	8.9	19.0	29.3	44.6	43.8
2005.....	7.2	*	*	*	*	0.3	1.2	9.6	17.9	29.0	45.0	44.8
2004.....	6.8	*	*	*	0.1	0.3	1.3	9.2	16.6	28.3	42.7	41.5
2003.....	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002.....	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001.....	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000.....	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999.....	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1

All origins 3/, female

2007.....	3.8	*	*	*	0.1	0.2	0.5	2.4	5.5	12.7	23.6	26.7
2006.....	3.8	*	*	*	*	0.2	0.5	2.3	5.8	12.6	23.2	28.7
2005.....	3.7	*	*	*	*	0.1	0.6	2.4	5.2	12.6	23.2	26.8
2004.....	3.7	*	*	*	*	0.1	0.4	2.2	5.3	12.8	23.6	28.3
2003.....	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002.....	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001.....	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000.....	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999.....	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Hispanic, both sexes

2007.....	4.2	*	*	*	*	*	0.9	6.7	17.6	32.4	53.8	41.5
2006.....	4.1	*	*	*	*	*	0.8	7.3	17.2	31.5	51.6	55.2
2005.....	4.2	*	*	*	*	*	0.9	8.0	18.2	33.8	52.1	54.8
2004.....	3.8	*	*	*	*	*	1.0	7.2	16.6	30.8	48.6	50.9
2003.....	3.8	*	*	*	*	*	0.8	7.5	16.1	32.5	49.8	51.8
2002.....	3.5	*	*	*	*	*	0.8	5.9	15.7	34.1	44.2	59.9
2001.....	3.4	*	*	*	*	*	0.8	6.7	14.1	28.8	52.1	52.5
2000.....	3.3	*	*	*	*	*	1.2	5.6	13.7	31.1	51.8	58.4
1999.....	3.2	*	*	*	*	*	0.9	5.7	13.1	31.3	49.8	53.3

Hispanic, male

2007.....	5.3	*	*	*	*	*	1.1	10.3	28.2	44.8	72.3	52.9
2006.....	5.2	*	*	*	*	*	1.0	11.5	27.5	45.7	69.2	62.0
2005.....	5.4	*	*	*	*	*	1.1	12.6	28.8	46.9	68.5	71.0
2004.....	4.9	*	*	*	*	*	1.6	11.2	27.5	42.3	62.9	71.4
2003.....	4.8	*	*	*	*	*	1.2	12.5	24.3	45.2	66.0	55.4
2002.....	4.5	*	*	*	*	*	1.1	9.8	24.1	48.9	59.9	75.3
2001.....	4.3	*	*	*	*	*	1.3	11.1	21.5	41.1	68.0	64.8
2000.....	4.2	*	*	*	*	*	1.7	9.1	20.5	43.9	74.7	72.6
1999.....	3.9	*	*	*	*	*	1.1	9.6	20.7	42.8	63.1	66.9

Hispanic, female

2007.....	2.9	*	*	*	*	*	*	3.0	7.9	22.4	40.8	35.3
2006.....	2.9	*	*	*	*	*	*	3.1	7.7	20.0	39.2	51.5
2005.....	3.0	*	*	*	*	*	0.7	3.3	8.6	23.1	40.7	46.3
2004.....	2.6	*	*	*	*	*	*	3.1	6.7	21.6	38.8	40.3
2003.....	2.7	*	*	*	*	*	*	2.5	8.8	22.4	38.7	50.0
2002.....	2.5	*	*	*	*	*	*	2.1	8.2	22.4	33.4	52.2
2001.....	2.5	*	*	*	*	*	*	2.3	7.6	19.1	41.4	46.5
2000.....	2.4	*	*	*	*	*	*	2.2	7.6	21.1	36.3	51.4
1999.....	2.4	*	*	*	*	*	*	1.9	6.3	22.2	40.9	46.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 4/, both sexes

2007.....	5.9	*	*	*	0.1	0.2	0.9	5.6	12.3	19.0	31.3	31.5
2006.....	5.8	*	*	*	0.1	0.2	0.8	5.3	11.7	19.3	30.6	32.3
2005.....	5.6	*	*	*	0.1	0.3	0.9	5.7	10.7	19.0	30.8	31.3
2004.....	5.4	*	0.2	*	0.1	0.2	0.8	5.5	10.2	19.0	30.3	31.5
2003.....	5.2	*	*	*	0.1	0.2	0.9	5.2	9.4	18.8	29.7	32.6
2002.....	5.1	*	0.2	*	0.1	0.3	0.9	4.8	9.1	18.6	28.4	33.9
2001.....	4.9	*	*	*	0.1	0.2	0.9	4.6	8.9	18.1	27.9	33.2
2000.....	4.8	*	0.2	*	0.1	0.2	0.9	4.3	8.9	17.9	27.7	31.7
1999.....	4.6	*	0.2	*	0.1	0.2	1.0	3.9	8.4	18.2	27.1	29.7

Non-Hispanic 4/, male

2007.....	8.0	*	*	*	*	0.3	1.2	9.0	19.8	27.4	44.3	42.6
2006.....	7.7	*	*	*	*	0.3	1.2	8.5	18.2	27.9	43.0	42.6
2005.....	7.5	*	*	*	*	0.4	1.2	9.2	16.9	27.6	43.5	43.4
2004.....	7.1	*	*	*	0.1	0.3	1.3	9.0	15.6	27.2	41.4	40.0
2003.....	6.9	*	*	*	*	0.3	1.3	8.4	14.1	26.6	43.5	47.1
2002.....	6.6	*	*	*	*	0.4	1.4	8.0	13.7	27.1	40.5	44.8
2001.....	6.3	*	*	*	*	0.2	1.3	7.4	13.2	25.8	39.7	49.7
2000.....	6.2	*	*	*	*	0.3	1.3	6.9	13.3	25.9	40.5	43.4
1999.....	5.8	*	*	*	0.1	0.3	1.5	6.2	12.2	26.5	37.9	38.7

Non-Hispanic 4/, female

2007.....	3.9	*	*	*	*	0.2	0.5	2.3	5.3	11.9	22.4	26.2
2006.....	3.9	*	*	*	*	0.2	0.5	2.3	5.6	12.0	22.2	27.6
2005.....	3.8	*	*	*	*	0.1	0.5	2.3	4.9	11.8	22.1	25.8
2004.....	3.8	*	*	*	*	0.2	0.4	2.1	5.2	12.1	22.7	27.7
2003.....	3.6	*	*	*	*	0.2	0.5	2.1	5.0	12.3	20.4	26.2
2002.....	3.6	*	*	*	*	0.2	0.5	1.7	4.8	11.4	20.5	29.2
2001.....	3.5	*	*	*	*	0.2	0.4	1.9	4.8	11.7	20.2	26.4
2000.....	3.4	*	*	*	*	0.1	0.6	1.9	4.9	11.2	19.4	27.0
1999.....	3.4	*	*	*	*	0.2	0.5	1.8	4.9	11.3	20.0	26.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic white, both sexes

2007 5/.....	5.8	*	*	*	0.1	0.2	0.7	4.8	10.4	17.4	29.9	30.4
2006 5/.....	5.6	*	*	*	0.1	0.2	0.6	4.4	9.9	17.7	29.1	31.7
2005 5/.....	5.5	*	*	*	*	0.2	0.6	4.8	9.0	17.2	29.2	30.4
2004 5/.....	5.3	*	*	*	0.1	0.1	0.6	4.5	8.6	17.1	28.8	30.1
2003 5/.....	5.1	*	*	*	0.1	0.1	0.7	4.2	8.0	17.0	28.2	31.1
2002.....	5.0	*	*	*	*	0.2	0.7	3.9	7.7	17.0	27.4	32.6
2001.....	4.8	*	*	*	0.1	0.1	0.6	3.7	7.6	16.6	26.5	32.4
2000.....	4.6	*	*	*	*	0.1	0.6	3.4	7.5	16.2	26.5	30.8
1999.....	4.5	*	0.2	*	0.1	0.1	0.7	3.1	7.1	16.8	26.1	28.9

Non-Hispanic white, male

2007 5/.....	7.7	*	*	*	*	0.2	0.9	7.7	16.4	24.8	42.4	41.0
2006 5/.....	7.3	*	*	*	*	0.2	0.7	7.1	14.9	25.2	41.2	42.3
2005 5/.....	7.2	*	*	*	*	0.2	0.8	7.6	14.0	24.8	41.5	43.4
2004 5/.....	6.8	*	*	*	*	0.2	0.9	7.4	12.8	24.4	40.0	38.0
2003 5/.....	6.7	*	*	*	*	*	1.0	6.7	11.9	24.3	41.6	45.0
2002.....	6.4	*	*	*	*	0.2	1.1	6.4	11.5	24.6	38.7	43.6
2001.....	6.1	*	*	*	*	*	0.9	5.9	11.2	24.2	38.0	48.9
2000.....	5.9	*	*	*	*	0.2	0.8	5.3	11.0	23.6	39.2	41.6
1999.....	5.7	*	*	*	*	*	1.1	4.8	10.2	24.6	36.7	39.0

Non-Hispanic white, female

2007 5/.....	4.0	*	*	*	*	*	0.5	2.0	4.7	11.1	21.2	25.4
2006 5/.....	4.0	*	*	*	*	0.2	0.5	1.9	5.1	11.1	20.6	26.8
2005 5/.....	3.8	*	*	*	*	*	0.5	2.0	4.3	10.6	20.8	24.5
2004 5/.....	3.8	*	*	*	*	*	0.4	1.7	4.7	10.9	21.2	26.5
2003 5/.....	3.6	*	*	*	*	*	0.5	1.7	4.2	10.7	19.2	25.0
2002.....	3.6	*	*	*	*	0.2	0.4	1.4	4.2	10.4	19.9	27.8
2001.....	3.4	*	*	*	*	*	0.3	1.5	4.1	10.2	18.9	25.5
2000.....	3.4	*	*	*	*	*	0.4	1.6	4.3	10.1	18.2	26.5
1999.....	3.4	*	*	*	*	*	0.4	1.6	4.2	10.2	19.1	24.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic black, both sexes

2007 5/.....	5.9	*	*	*	*	0.5	1.4	8.7	23.3	24.8	34.5	34.8
2006 5/.....	5.7	*	*	*	*	0.4	1.3	9.6	22.3	24.6	33.4	31.6
2005 5/.....	5.6	*	*	*	*	0.5	1.5	9.9	20.5	25.4	33.6	28.9
2004 5/.....	5.4	*	*	*	*	0.4	1.4	9.8	17.7	26.1	34.7	36.6
2003 5/.....	5.1	*	*	*	*	0.5	1.4	10.0	16.5	23.1	31.7	41.9
2002.....	4.7	*	*	*	*	0.5	1.5	8.9	15.0	23.3	30.9	38.1
2001.....	4.6	*	*	*	*	*	1.7	8.9	15.1	21.9	32.3	35.3
2000.....	4.5	*	*	*	*	0.4	1.8	8.8	14.5	23.7	29.1	31.8
1999.....	4.3	*	*	*	*	0.4	1.7	7.4	13.4	24.6	29.7	34.9

Non-Hispanic black, male

2007 5/.....	8.7	*	*	*	*	0.8	2.3	14.7	42.0	39.6	49.5	50.6
2006 5/.....	8.3	*	*	*	*	*	2.2	15.7	40.6	40.0	45.9	34.8
2005 5/.....	8.2	*	*	*	*	*	2.2	17.3	36.4	40.2	51.0	31.4
2004 5/.....	7.6	*	*	*	*	*	2.3	16.9	30.5	41.3	44.2	47.8
2003 5/.....	7.0	*	*	*	*	*	2.2	17.3	26.4	33.8	44.7	63.9
2002.....	6.6	*	*	*	*	*	2.3	15.4	24.4	38.2	46.3	41.6
2001.....	6.1	*	*	*	*	*	2.7	15.1	24.2	30.3	42.6	42.7
2000.....	6.3	*	*	*	*	*	2.5	15.0	24.1	37.3	40.4	48.1
1999.....	5.7	*	*	*	*	*	2.5	13.1	20.3	36.8	41.6	30.9

Non-Hispanic black, female

2007 5/.....	3.3	*	*	*	*	*	*	3.6	8.3	14.2	26.0	28.2
2006 5/.....	3.3	*	*	*	*	*	*	4.4	7.5	13.5	26.3	30.3
2005 5/.....	3.2	*	*	*	*	*	0.9	3.5	7.8	14.8	23.7	28.0
2004 5/.....	3.4	*	*	*	*	*	*	3.7	7.4	15.2	29.3	32.1
2003 5/.....	3.3	*	*	*	*	*	*	3.6	8.6	15.5	24.3	33.4
2002.....	3.0	*	*	*	*	*	0.8	3.3	7.4	12.8	22.2	36.8
2001.....	3.3	*	*	*	*	*	0.8	3.6	7.8	16.0	26.4	32.5
2000.....	2.9	*	*	*	*	*	1.2	3.4	6.7	14.1	22.8	25.7
1999.....	3.1	*	*	*	*	*	0.9	2.5	7.9	15.9	23.1	36.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

All origins 3/, both sexes

2007.....	11.3	*	*	*	*	0.1	1.2	6.4	19.9	44.8	79.2	94.6
2006.....	11.2	*	*	*	*	0.1	1.2	6.3	19.9	44.8	78.6	96.9
2005.....	11.1	*	*	*	*	0.2	1.3	6.5	19.7	44.5	77.7	97.4
2004.....	10.8	*	*	*	*	0.2	1.2	6.3	19.8	44.3	76.0	97.7
2003.....	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000.....	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999.....	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7

All origins 3/, male

2007.....	11.5	*	*	*	*	0.2	1.5	7.8	24.5	52.5	88.1	104.9
2006.....	11.2	*	*	*	*	0.1	1.4	7.8	24.5	52.3	86.2	102.1
2005.....	11.1	*	*	*	*	0.2	1.5	7.7	24.3	51.8	86.0	102.8
2004.....	10.9	*	*	*	*	0.2	1.5	8.1	24.1	52.6	84.6	102.0
2003.....	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002.....	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001.....	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000.....	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999.....	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5

All origins 3/, female

2007.....	11.1	*	*	*	*	0.1	1.0	5.0	15.6	38.3	73.1	89.8
2006.....	11.1	*	*	*	*	0.1	0.9	4.9	15.7	38.4	73.4	94.4
2005.....	11.0	*	*	*	*	0.1	1.1	5.2	15.5	38.2	72.1	94.9
2004.....	10.7	*	*	*	*	0.1	0.9	4.7	15.8	37.3	70.1	95.8
2003.....	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002.....	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001.....	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000.....	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999.....	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

Hispanic, both sexes

2007.....	3.9	*	*	*	*	*	0.7	4.2	13.7	32.5	60.9	71.5
2006.....	3.9	*	*	*	*	*	0.7	4.3	13.8	33.7	59.9	77.1
2005.....	4.1	*	*	*	*	*	0.9	4.4	15.5	36.0	63.8	78.7
2004.....	3.8	*	*	*	*	*	0.7	4.1	14.4	35.3	62.8	69.9
2003.....	3.7	*	*	*	*	*	0.8	4.1	14.4	35.1	60.3	79.8
2002.....	3.5	*	*	*	*	*	0.9	4.0	13.8	34.7	55.5	79.3
2001.....	3.6	*	*	*	*	*	0.8	3.7	14.2	36.9	61.3	83.8
2000.....	3.5	*	*	*	*	*	0.7	4.1	13.7	36.9	61.8	82.9
1999.....	3.3	*	*	*	*	*	0.9	4.6	13.3	32.6	56.7	84.2

Hispanic, male

2007.....	3.7	*	*	*	*	*	1.0	5.0	16.6	37.0	53.8	77.2
2006.....	3.8	*	*	*	*	*	0.8	5.0	15.4	39.7	68.9	65.4
2005.....	4.0	*	*	*	*	*	1.1	5.2	19.2	40.5	65.8	77.2
2004.....	3.7	*	*	*	*	*	1.0	4.7	17.4	42.4	68.3	57.4
2003.....	3.6	*	*	*	*	*	1.0	5.0	17.8	40.0	67.1	70.8
2002.....	3.3	*	*	*	*	*	1.2	5.3	15.7	40.9	55.8	60.2
2001.....	3.4	*	*	*	*	*	0.9	4.3	16.2	42.3	68.9	62.9
2000.....	3.4	*	*	*	*	*	0.8	5.2	16.6	45.2	62.9	80.6
1999.....	3.2	*	*	*	*	*	1.2	5.7	15.2	40.0	56.3	86.3

Hispanic, female

2007.....	4.1	*	*	*	*	*	*	3.3	11.0	28.8	65.8	68.4
2006.....	4.1	*	*	*	*	*	*	3.7	12.3	28.9	53.6	83.3
2005.....	4.2	*	*	*	*	*	0.8	3.5	12.1	32.3	62.3	79.5
2004.....	3.9	*	*	*	*	*	*	3.5	11.8	29.5	59.0	76.3
2003.....	3.8	*	*	*	*	*	*	3.2	11.3	31.2	55.6	84.4
2002.....	3.7	*	*	*	*	*	*	2.8	12.1	29.8	55.4	88.7
2001.....	3.9	*	*	*	*	*	0.8	3.2	12.5	32.6	56.2	94.0
2000.....	3.6	*	*	*	*	*	*	3.0	11.1	30.4	61.1	84.1
1999.....	3.5	*	*	*	*	*	*	3.5	11.6	26.8	56.9	83.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic 4/, both sexes

2007.....	12.6	*	*	*	*	0.1	1.3	6.7	20.4	45.8	80.3	95.7
2006.....	12.4	*	*	*	*	0.1	1.2	6.5	20.4	45.6	79.7	97.7
2005.....	12.2	*	*	*	*	0.2	1.3	6.7	20.0	45.0	78.5	98.1
2004.....	11.9	*	*	*	*	0.2	1.3	6.6	20.2	44.9	76.5	98.9
2003.....	11.7	*	*	*	*	0.2	1.3	6.4	19.4	44.8	76.1	95.7
2002.....	11.6	*	*	*	*	0.1	1.4	6.3	19.6	45.4	75.8	94.9
2001.....	11.5	*	*	*	*	0.1	1.3	6.4	19.6	44.9	74.6	100.6
2000.....	11.4	*	*	*	*	0.1	1.3	6.5	19.8	44.4	77.0	96.6
1999.....	11.4	*	*	*	*	0.1	1.4	6.6	20.2	45.6	73.5	100.1

Non-Hispanic 4/, male

2007.....	13.0	*	*	*	*	0.2	1.6	8.2	25.2	53.6	90.3	106.2
2006.....	12.6	*	*	*	*	*	1.5	8.1	25.3	53.2	87.1	104.0
2005.....	12.3	*	*	*	*	0.2	1.6	8.0	24.7	52.6	87.2	104.1
2004.....	12.1	*	*	*	*	0.2	1.6	8.4	24.6	53.2	85.4	104.1
2003.....	11.7	*	*	*	*	0.2	1.6	8.0	23.5	51.6	86.2	96.5
2002.....	11.7	*	*	*	*	0.1	1.7	8.0	24.1	52.1	85.3	102.4
2001.....	11.4	*	*	*	*	0.2	1.5	7.9	24.2	51.2	84.6	106.4
2000.....	11.3	*	*	*	*	0.2	1.5	8.3	24.1	50.7	86.7	103.5
1999.....	11.4	*	*	*	*	0.2	1.7	8.4	24.4	52.9	82.7	108.2

Non-Hispanic 4/, female

2007.....	12.3	*	*	*	*	*	1.1	5.2	16.0	39.0	73.4	90.8
2006.....	12.3	*	*	*	*	0.1	1.0	5.0	16.0	39.1	74.6	94.8
2005.....	12.1	*	*	*	*	0.1	1.1	5.4	15.7	38.7	72.6	95.4
2004.....	11.8	*	*	*	*	*	1.0	4.8	16.1	37.9	70.6	96.5
2003.....	11.6	*	*	*	*	0.1	1.0	4.9	15.6	39.1	69.3	95.3
2002.....	11.5	*	*	*	*	*	1.1	4.6	15.4	39.8	69.5	91.7
2001.....	11.5	*	*	*	*	0.1	1.2	4.9	15.4	39.8	68.0	98.2
2000.....	11.4	*	*	*	*	*	1.0	4.7	15.8	39.2	70.6	93.8
1999.....	11.4	*	*	*	*	*	1.0	4.8	16.4	39.6	67.6	96.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic white, both sexes

2007 5/.....	13.6	*	*	*	*	0.1	1.3	6.3	20.1	45.1	79.7	94.6
2006 5/.....	13.4	*	*	*	*	0.1	1.2	6.3	20.0	44.9	79.5	96.9
2005 5/.....	13.2	*	*	*	*	0.2	1.3	6.4	19.6	44.5	77.7	97.6
2004 5/.....	12.8	*	*	*	*	0.2	1.2	6.3	19.5	44.0	76.1	98.1
2003 5/.....	12.5	*	*	*	*	0.2	1.2	6.0	18.7	44.1	75.4	94.9
2002.....	12.4	*	*	*	*	0.1	1.4	6.0	19.0	44.3	75.1	94.6
2001.....	12.3	*	*	*	*	0.1	1.3	6.1	19.3	44.1	73.7	99.4
2000.....	12.2	*	*	*	*	0.2	1.2	6.3	19.2	43.4	75.7	94.7
1999.....	12.1	*	*	*	*	0.1	1.3	6.3	19.5	44.7	72.1	98.9

Non-Hispanic white, male

2007 5/.....	14.1	*	*	*	*	*	1.6	7.8	24.6	53.5	90.6	106.6
2006 5/.....	13.7	*	*	*	*	*	1.5	7.9	24.8	53.0	88.0	105.3
2005 5/.....	13.3	*	*	*	*	0.2	1.6	7.6	24.2	52.5	86.7	104.3
2004 5/.....	13.1	*	*	*	*	0.2	1.5	8.1	23.8	52.0	85.8	105.0
2003 5/.....	12.6	*	*	*	*	0.2	1.5	7.5	22.8	51.4	86.0	96.8
2002.....	12.6	*	*	*	*	*	1.7	7.7	23.5	51.2	85.4	102.9
2001.....	12.4	*	*	*	*	0.2	1.5	7.7	24.0	50.3	84.5	108.0
2000.....	12.2	*	*	*	*	0.2	1.5	8.0	23.4	50.0	86.1	103.0
1999.....	12.1	*	*	*	*	0.2	1.6	8.0	23.4	52.1	81.6	108.1

Non-Hispanic white, female

2007 5/.....	13.1	*	*	*	*	*	1.1	4.9	15.7	37.8	72.0	88.9
2006 5/.....	13.1	*	*	*	*	*	0.9	4.7	15.4	38.0	73.6	93.0
2005 5/.....	13.0	*	*	*	*	*	1.1	5.1	15.2	37.7	71.5	94.5
2004 5/.....	12.6	*	*	*	*	*	0.9	4.5	15.4	37.1	69.5	95.1
2003 5/.....	12.4	*	*	*	*	*	1.0	4.6	14.8	38.0	68.1	94.0
2002.....	12.2	*	*	*	*	*	1.1	4.3	14.7	38.3	68.1	91.0
2001.....	12.3	*	*	*	*	*	1.2	4.6	14.9	38.8	66.6	95.8
2000.....	12.2	*	*	*	*	*	1.0	4.5	15.3	37.9	68.9	91.4
1999.....	12.1	*	*	*	*	*	1.0	4.7	15.7	38.5	65.9	95.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	10.6	*	*	*	*	*	1.6	9.9	27.8	59.9	99.3	116.1
2006 5/.....	10.2	*	*	*	*	*	1.6	9.4	28.5	58.4	93.1	114.7
2005 5/.....	10.0	*	*	*	*	*	1.6	9.8	27.5	57.7	95.9	111.7
2004 5/.....	9.8	*	*	*	*	*	1.7	9.5	28.8	59.0	89.1	115.1
2003 5/.....	9.8	*	*	*	*	*	1.9	10.3	27.7	57.5	94.0	113.3
2002.....	9.6	*	*	*	*	*	1.8	8.9	28.3	61.7	90.9	108.3
2001.....	9.4	*	*	*	*	*	1.6	9.2	25.9	59.0	93.3	122.0
2000.....	9.7	*	*	*	*	*	1.7	9.2	27.9	60.0	98.9	121.2
1999.....	9.7	*	*	*	*	*	2.0	9.9	28.8	60.4	95.0	119.0
<b>Non-Hispanic black, male</b>												
2007 5/.....	10.3	*	*	*	*	*	1.9	12.4	36.9	66.9	101.5	109.1
2006 5/.....	9.7	*	*	*	*	*	2.0	11.5	36.2	67.3	93.4	92.5
2005 5/.....	9.8	*	*	*	*	*	1.7	12.1	34.7	65.7	106.0	101.8
2004 5/.....	9.8	*	*	*	*	*	2.1	11.6	36.5	73.8	91.3	102.6
2003 5/.....	9.6	*	*	*	*	*	2.5	13.6	33.7	64.8	99.0	106.5
2002.....	9.3	*	*	*	*	*	2.2	11.6	34.9	69.5	97.3	102.8
2001.....	8.9	*	*	*	*	*	1.8	11.4	30.8	68.2	100.5	100.5
2000.....	9.3	*	*	*	*	*	2.1	12.1	34.6	66.9	108.0	110.4
1999.....	9.7	*	*	*	*	*	2.4	13.6	37.4	68.7	107.6	115.3
<b>Non-Hispanic black, female</b>												
2007 5/.....	10.8	*	*	*	*	*	1.3	7.7	20.4	54.9	98.1	119.0
2006 5/.....	10.6	*	*	*	*	*	1.3	7.5	22.4	52.1	92.9	123.7
2005 5/.....	10.3	*	*	*	*	*	1.5	7.9	21.7	52.0	90.2	115.7
2004 5/.....	9.9	*	*	*	*	*	1.2	7.7	22.6	48.5	87.9	120.1
2003 5/.....	10.0	*	*	*	*	*	1.3	7.4	22.8	52.4	91.3	116.0
2002.....	9.8	*	*	*	*	*	1.5	6.5	23.0	56.2	87.2	110.4
2001.....	9.9	*	*	*	*	*	1.4	7.3	21.9	52.4	89.3	130.0
2000.....	10.0	*	*	*	*	*	1.5	6.8	22.5	55.1	93.8	125.2
1999.....	9.7	*	*	*	*	*	1.7	6.7	21.9	54.6	87.9	120.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of larynx (C32)</b>												
All origins 3/, both sexes												
2007.....	1.2	*	*	*	*	*	0.1	1.0	2.8	5.1	6.7	6.0
2006.....	1.3	*	*	*	*	*	0.1	1.0	3.1	5.7	7.1	6.3
2005.....	1.3	*	*	*	*	*	0.1	1.0	3.0	6.2	7.0	6.3
2004.....	1.2	*	*	*	*	*	0.1	1.1	3.2	5.7	6.6	6.5
2003.....	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002.....	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001.....	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000.....	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999.....	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All origins 3/, male												
2007.....	1.9	*	*	*	*	*	0.2	1.7	4.9	8.6	12.9	13.9
2006.....	2.0	*	*	*	*	*	0.2	1.7	5.1	9.8	13.4	14.8
2005.....	2.0	*	*	*	*	*	0.2	1.5	5.1	10.7	13.1	15.7
2004.....	2.0	*	*	*	*	*	0.2	1.8	5.4	10.1	13.0	15.6
2003.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001.....	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000.....	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999.....	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All origins 3/, female												
2007.....	0.5	*	*	*	*	*	*	0.4	0.9	2.1	2.4	2.3
2006.....	0.5	*	*	*	*	*	*	0.4	1.2	2.2	2.8	2.3
2005.....	0.5	*	*	*	*	*	*	0.5	1.0	2.4	2.8	1.9
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	2.0	2.3	2.4
2003.....	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002.....	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001.....	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000.....	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Malignant neoplasm of larynx (C32)

##### Non-Hispanic 4/, both sexes

2007.....	1.4	*	*	*	*	*	0.1	1.1	3.0	5.2	6.8	6.1
2006.....	1.4	*	*	*	*	*	0.2	1.1	3.2	5.8	7.1	6.3
2005.....	1.4	*	*	*	*	*	0.2	1.0	3.2	6.4	7.0	6.2
2004.....	1.4	*	*	*	*	*	0.2	1.1	3.3	5.7	6.6	6.5
2003.....	1.4	*	*	*	*	*	0.2	1.1	3.5	6.3	6.9	6.5
2002.....	1.4	*	*	*	*	*	0.2	1.1	3.5	6.0	7.7	5.4
2001.....	1.5	*	*	*	*	*	0.2	1.1	3.7	6.4	7.3	6.7
2000.....	1.5	*	*	*	*	*	0.1	1.2	3.7	6.7	8.1	6.5
1999.....	1.5	*	*	*	*	*	0.2	1.1	3.9	6.8	7.2	6.8

##### Non-Hispanic 4/, male

2007.....	2.2	*	*	*	*	*	0.2	1.8	5.1	8.7	13.1	14.0
2006.....	2.3	*	*	*	*	*	0.2	1.8	5.3	10.0	13.3	14.7
2005.....	2.3	*	*	*	*	*	0.2	1.6	5.4	10.9	13.0	15.7
2004.....	2.2	*	*	*	*	*	0.2	1.9	5.5	10.0	12.8	15.6
2003.....	2.3	*	*	*	*	*	0.3	1.8	5.9	10.9	13.7	15.0
2002.....	2.3	*	*	*	*	*	0.3	1.9	5.9	10.1	15.0	13.2
2001.....	2.4	*	*	*	*	*	0.3	1.9	6.1	11.1	14.3	18.4
2000.....	2.4	*	*	*	*	*	0.2	1.9	6.4	11.5	15.7	17.3
1999.....	2.4	*	*	*	*	*	0.3	1.9	6.6	11.4	14.0	17.3

##### Non-Hispanic 4/, female

2007.....	0.6	*	*	*	*	*	*	0.4	0.9	2.3	2.5	2.3
2006.....	0.6	*	*	*	*	*	*	0.4	1.3	2.3	2.9	2.4
2005.....	0.6	*	*	*	*	*	*	0.5	1.1	2.5	2.9	1.9
2004.....	0.6	*	*	*	*	*	*	0.4	1.3	2.1	2.4	2.5
2003.....	0.6	*	*	*	*	*	*	0.3	1.3	2.5	2.4	2.7
2002.....	0.6	*	*	*	*	*	*	0.4	1.3	2.6	2.9	2.1
2001.....	0.6	*	*	*	*	*	0.1	0.4	1.4	2.6	2.7	1.8
2000.....	0.6	*	*	*	*	*	*	0.5	1.2	2.8	3.1	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.4	3.0	2.8	2.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of larynx (C32)

Non-Hispanic white, both sexes

2007 5/.....	1.4	*	*	*	*	*	0.1	1.0	2.7	4.9	6.5	5.9
2006 5/.....	1.5	*	*	*	*	*	0.1	1.0	3.0	5.5	7.0	6.2
2005 5/.....	1.4	*	*	*	*	*	0.1	0.9	2.8	6.0	6.9	5.9
2004 5/.....	1.4	*	*	*	*	*	0.1	1.0	3.0	5.4	6.3	6.4
2003 5/.....	1.4	*	*	*	*	*	0.2	0.9	3.2	6.0	6.6	6.1
2002.....	1.4	*	*	*	*	*	0.1	0.9	3.1	5.8	7.3	5.5
2001.....	1.5	*	*	*	*	*	0.2	0.9	3.3	6.0	6.9	6.5
2000.....	1.5	*	*	*	*	*	0.1	1.0	3.2	6.2	7.7	6.5
1999.....	1.5	*	*	*	*	*	0.1	0.9	3.5	6.4	7.1	6.5

Non-Hispanic white, male

2007 5/.....	2.2	*	*	*	*	*	0.2	1.6	4.5	7.9	12.3	13.4
2006 5/.....	2.3	*	*	*	*	*	0.2	1.5	4.9	9.1	12.8	14.5
2005 5/.....	2.3	*	*	*	*	*	0.2	1.4	4.6	10.0	12.7	14.8
2004 5/.....	2.2	*	*	*	*	*	0.2	1.6	4.9	9.4	12.1	15.0
2003 5/.....	2.3	*	*	*	*	*	0.3	1.5	5.2	9.9	12.8	14.1
2002.....	2.3	*	*	*	*	*	0.2	1.5	5.1	9.5	14.2	13.3
2001.....	2.3	*	*	*	*	*	0.3	1.5	5.5	10.2	13.3	18.3
2000.....	2.4	*	*	*	*	*	0.2	1.5	5.4	10.5	15.0	17.4
1999.....	2.3	*	*	*	*	*	0.2	1.6	5.8	10.7	13.6	16.0

Non-Hispanic white, female

2007 5/.....	0.6	*	*	*	*	*	*	0.4	0.9	2.2	2.5	2.3
2006 5/.....	0.7	*	*	*	*	*	*	0.4	1.2	2.4	3.0	2.3
2005 5/.....	0.6	*	*	*	*	*	*	0.4	1.1	2.5	3.0	1.8
2004 5/.....	0.6	*	*	*	*	*	*	0.4	1.3	2.0	2.4	2.5
2003 5/.....	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.4	2.6
2002.....	0.6	*	*	*	*	*	*	0.3	1.3	2.6	2.7	2.2
2001.....	0.6	*	*	*	*	*	*	0.3	1.2	2.4	2.7	1.6
2000.....	0.6	*	*	*	*	*	*	0.4	1.1	2.6	3.0	2.1
1999.....	0.7	*	*	*	*	*	*	0.3	1.3	2.9	2.9	2.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of larynx (C32)

Non-Hispanic black, both sexes

2007 5/.....	1.7	*	*	*	*	*	*	2.2	6.1	9.8	11.7	8.4
2006 5/.....	1.7	*	*	*	*	*	*	2.2	6.0	10.6	10.2	8.2
2005 5/.....	1.8	*	*	*	*	*	*	2.3	7.2	11.6	9.8	10.4
2004 5/.....	1.7	*	*	*	*	*	*	2.3	6.6	10.3	11.4	8.1
2003 5/.....	1.9	*	*	*	*	*	*	2.6	7.7	11.6	11.9	11.3
2002.....	1.9	*	*	*	*	*	0.5	2.7	7.7	10.6	13.9	*
2001.....	1.9	*	*	*	*	*	0.4	2.9	7.9	11.5	12.0	8.4
2000.....	2.0	*	*	*	*	*	*	2.9	9.2	12.9	14.3	7.0
1999.....	1.9	*	*	*	*	*	0.6	3.0	8.7	11.7	10.0	8.4

Non-Hispanic black, male

2007 5/.....	3.0	*	*	*	*	*	*	4.2	12.0	18.3	26.4	23.6
2006 5/.....	2.8	*	*	*	*	*	*	4.0	10.8	21.2	22.5	19.2
2005 5/.....	3.0	*	*	*	*	*	*	3.5	14.0	22.5	21.0	29.5
2004 5/.....	2.9	*	*	*	*	*	*	4.3	12.6	20.2	25.4	23.4
2003 5/.....	3.3	*	*	*	*	*	*	4.6	14.0	24.0	28.3	27.7
2002.....	3.1	*	*	*	*	*	0.8	4.6	15.1	19.8	29.6	*
2001.....	3.1	*	*	*	*	*	*	5.2	14.2	21.4	28.2	*
2000.....	3.4	*	*	*	*	*	*	5.2	17.5	24.6	29.0	*
1999.....	3.2	*	*	*	*	*	1.0	5.3	16.3	22.1	22.5	25.0

Non-Hispanic black, female

2007 5/.....	0.6	*	*	*	*	*	*	*	1.4	3.8	3.3	*
2006 5/.....	0.6	*	*	*	*	*	*	*	2.1	3.0	3.2	*
2005 5/.....	0.7	*	*	*	*	*	*	1.2	1.7	3.9	3.5	*
2004 5/.....	0.6	*	*	*	*	*	*	*	1.7	3.1	3.5	*
2003 5/.....	0.6	*	*	*	*	*	*	0.8	2.6	2.8	*	*
2002.....	0.7	*	*	*	*	*	*	1.1	1.8	4.1	5.0	*
2001.....	0.7	*	*	*	*	*	*	*	2.9	4.6	*	*
2000.....	0.8	*	*	*	*	*	*	1.0	2.4	4.7	6.1	*
1999.....	0.7	*	*	*	*	*	*	0.9	2.7	4.3	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/, both sexes

2007.....	52.6	*	*	*	0.1	0.3	4.3	28.4	95.4	248.8	371.3	299.8
2006.....	53.0	*	*	*	0.1	0.4	4.6	29.1	99.1	253.1	373.5	300.5
2005.....	53.7	*	*	*	*	0.3	5.3	29.7	103.3	259.6	375.6	302.3
2004.....	53.8	*	*	*	0.1	0.4	5.5	29.9	106.9	265.5	373.6	297.5
2003.....	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002.....	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001.....	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000.....	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999.....	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0

All origins 3/, male

2007.....	59.4	*	*	*	*	0.4	4.2	32.1	116.2	310.2	498.3	453.0
2006.....	60.5	*	*	*	*	0.4	4.7	33.8	121.6	319.4	509.9	457.7
2005.....	61.8	*	*	*	*	0.4	5.5	35.1	127.6	330.7	515.1	468.0
2004.....	62.0	*	*	*	*	0.4	5.8	35.8	131.7	337.6	517.3	466.8
2003.....	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002.....	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001.....	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000.....	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999.....	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2

All origins 3/, female

2007.....	46.0	*	*	*	*	0.3	4.4	24.9	76.1	196.7	283.8	227.0
2006.....	45.7	*	*	*	*	0.4	4.5	24.6	78.2	197.0	280.3	226.9
2005.....	45.9	*	*	*	*	0.3	5.1	24.5	80.7	199.6	280.9	226.2
2004.....	45.9	*	*	*	*	0.3	5.2	24.2	83.9	205.0	277.0	221.3
2003.....	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002.....	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001.....	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000.....	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999.....	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Hispanic, both sexes

2007.....	10.2	*	*	*	*	0.2	1.4	8.5	31.8	100.0	167.1	169.8
2006.....	9.8	*	*	*	*	*	1.5	8.4	31.3	98.8	167.9	160.8
2005.....	10.5	*	*	*	*	*	1.7	9.5	35.7	102.6	183.3	177.1
2004.....	10.2	*	*	*	*	0.3	1.4	9.4	35.4	102.6	178.7	193.1
2003.....	10.4	*	*	*	*	0.3	1.6	10.2	38.1	105.8	183.8	189.2
2002.....	10.4	*	*	*	*	0.4	1.5	9.9	40.9	110.4	183.7	189.0
2001.....	10.3	*	*	*	*	*	1.9	10.7	41.0	108.3	182.3	200.7
2000.....	10.3	*	*	*	*	*	1.8	10.9	39.3	110.6	198.7	213.7
1999.....	10.4	*	*	*	*	*	2.0	10.4	44.2	112.9	187.7	227.3

Hispanic, male

2007.....	12.0	*	*	*	*	*	1.2	10.3	42.1	140.7	246.1	256.1
2006.....	12.0	*	*	*	*	*	1.5	10.5	44.8	140.1	254.2	263.9
2005.....	13.0	*	*	*	*	*	1.9	11.9	52.3	151.4	281.6	269.0
2004.....	12.7	*	*	*	*	*	1.7	12.6	50.0	151.7	279.7	287.2
2003.....	12.8	*	*	*	*	*	1.7	12.6	55.4	161.8	277.3	290.9
2002.....	13.1	*	*	*	*	*	1.5	12.6	60.7	161.7	299.1	307.9
2001.....	13.1	*	*	*	*	*	2.4	14.2	59.5	163.7	293.8	335.2
2000.....	13.3	*	*	*	*	*	2.1	14.8	58.6	167.3	327.5	368.8
1999.....	13.6	*	*	*	*	*	2.3	14.2	68.1	175.3	316.0	362.3

Hispanic, female

2007.....	8.2	*	*	*	*	*	1.5	6.7	22.2	66.7	111.7	123.0
2006.....	7.5	*	*	*	*	*	1.5	6.3	19.0	65.2	107.6	106.0
2005.....	7.8	*	*	*	*	*	1.6	7.1	20.4	63.1	114.9	129.0
2004.....	7.6	*	*	*	*	*	1.0	6.2	22.1	63.1	108.9	144.8
2003.....	7.8	*	*	*	*	*	1.4	7.9	22.4	61.2	119.6	137.5
2002.....	7.6	*	*	*	*	*	1.6	7.3	23.2	69.5	104.6	130.1
2001.....	7.4	*	*	*	*	*	1.4	7.3	24.6	64.6	107.0	135.0
2000.....	7.2	*	*	*	*	*	1.4	7.1	22.2	66.0	112.3	137.5
1999.....	7.1	*	*	*	*	*	1.7	6.7	23.1	64.0	102.3	162.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 4/, both sexes

2007.....	60.1	*	*	*	0.1	0.4	4.8	30.8	101.0	260.4	384.4	306.5
2006.....	60.4	*	*	*	0.1	0.4	5.2	31.5	104.9	264.8	385.9	306.9
2005.....	60.9	*	*	*	*	0.4	5.9	31.9	108.9	271.1	386.5	307.9
2004.....	60.9	*	*	*	0.1	0.4	6.2	32.0	112.6	277.0	383.9	301.6
2003.....	61.2	*	*	*	0.1	0.4	6.2	32.3	116.6	280.3	387.2	303.0
2002.....	61.4	*	*	*	*	0.4	6.7	32.3	121.0	285.7	386.2	301.0
2001.....	61.3	*	*	*	*	0.4	6.8	32.5	123.3	290.4	379.4	305.8
2000.....	61.5	*	*	*	*	0.5	6.6	33.4	128.4	294.2	377.4	304.5
1999.....	60.5	*	*	*	*	0.5	6.5	33.0	128.9	290.7	368.2	298.8

Non-Hispanic 4/, male

2007.....	68.3	*	*	*	*	0.4	4.8	34.8	122.7	322.9	514.7	464.2
2006.....	69.3	*	*	*	*	0.5	5.3	36.5	128.0	332.5	525.5	467.4
2005.....	70.3	*	*	*	*	0.4	6.1	37.6	133.7	343.4	528.5	477.9
2004.....	70.4	*	*	*	*	0.4	6.5	38.2	138.1	350.1	529.8	474.8
2003.....	71.1	*	*	*	*	0.4	6.8	38.9	142.9	358.3	537.2	483.1
2002.....	71.8	*	*	*	*	0.5	6.9	38.9	150.3	368.1	538.0	488.9
2001.....	72.6	*	*	*	*	0.4	7.3	39.5	153.2	382.4	539.5	518.4
2000.....	73.2	*	*	*	*	0.6	7.5	40.5	160.6	389.7	539.9	526.2
1999.....	72.7	*	*	*	*	0.5	7.1	40.1	164.0	387.8	534.9	534.6

Non-Hispanic 4/, female

2007.....	52.3	*	*	*	*	0.3	4.9	27.0	80.8	207.1	294.8	232.1
2006.....	51.9	*	*	*	*	0.4	5.0	26.7	83.3	207.4	290.6	232.3
2005.....	51.9	*	*	*	*	0.3	5.7	26.4	85.8	210.0	290.3	230.3
2004.....	51.8	*	*	*	*	0.3	5.8	26.1	88.9	215.4	285.8	224.2
2003.....	51.8	*	*	*	*	0.4	5.6	26.0	92.2	214.9	287.2	223.9
2002.....	51.6	*	*	*	*	0.4	6.4	25.9	93.9	216.9	285.5	220.0
2001.....	50.5	*	*	*	*	0.5	6.2	25.8	95.6	213.9	274.3	217.4
2000.....	50.4	*	*	*	*	0.5	5.8	26.5	98.6	215.4	271.6	214.8
1999.....	48.8	*	*	*	*	0.4	6.0	26.1	96.5	211.0	260.0	204.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic white, both sexes

2007 5/.....	66.7	*	*	*	*	0.4	5.0	30.0	101.4	266.7	393.8	310.7
2006 5/.....	66.9	*	*	*	*	0.4	5.2	30.4	105.1	271.5	395.0	313.0
2005 5/.....	67.3	*	*	*	*	0.3	5.9	31.0	108.6	277.8	394.7	311.3
2004 5/.....	67.1	*	*	*	*	0.3	6.1	31.0	112.5	283.2	389.6	306.3
2003 5/.....	67.5	*	*	*	*	0.4	6.1	31.0	117.0	286.5	392.6	306.7
2002.....	67.7	*	*	*	*	0.4	6.5	31.2	120.6	292.0	392.2	304.8
2001.....	67.3	*	*	*	*	0.5	6.6	31.5	124.0	295.2	383.0	308.6
2000.....	67.4	*	*	*	*	0.5	6.2	32.0	129.1	298.0	381.7	306.8
1999.....	65.9	*	*	*	*	0.5	6.2	31.5	128.9	293.5	370.3	300.5

Non-Hispanic white, male

2007 5/.....	74.6	*	*	*	*	0.4	4.8	33.0	119.8	324.2	520.4	464.7
2006 5/.....	75.8	*	*	*	*	0.5	5.3	34.5	124.8	334.0	531.2	474.2
2005 5/.....	76.8	*	*	*	*	0.4	6.0	36.0	129.7	345.6	534.1	480.2
2004 5/.....	76.5	*	*	*	*	0.4	6.3	35.8	134.2	351.4	530.6	480.3
2003 5/.....	77.3	*	*	*	*	0.4	6.5	36.5	139.4	359.6	537.0	486.8
2002.....	77.8	*	*	*	*	0.5	6.5	36.5	145.7	369.5	538.3	493.3
2001.....	78.4	*	*	*	*	0.4	6.9	37.0	150.3	381.1	536.5	520.0
2000.....	78.9	*	*	*	*	0.6	6.7	37.7	157.7	387.3	537.7	527.3
1999.....	78.1	*	*	*	*	0.5	6.6	37.1	160.1	384.4	531.1	536.9

Non-Hispanic white, female

2007 5/.....	59.0	*	*	*	*	0.3	5.1	27.1	83.8	216.7	305.2	237.7
2006 5/.....	58.4	*	*	*	*	0.4	5.1	26.3	86.4	217.3	300.5	237.8
2005 5/.....	58.2	*	*	*	*	0.3	5.9	26.0	88.5	219.3	298.7	234.1
2004 5/.....	58.1	*	*	*	*	0.3	5.9	26.2	92.0	224.6	293.5	228.4
2003 5/.....	58.1	*	*	*	*	0.4	5.7	25.6	95.7	223.9	295.1	227.4
2002.....	57.9	*	*	*	*	0.3	6.5	26.0	96.8	225.9	294.3	223.4
2001.....	56.6	*	*	*	*	0.5	6.4	26.0	99.2	222.3	281.3	220.6
2000.....	56.4	*	*	*	*	0.5	5.7	26.4	102.2	222.9	279.2	218.0
1999.....	54.3	*	*	*	*	0.5	5.8	26.1	99.5	217.5	265.0	206.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic black, both sexes

2007 5/.....	42.9	*	*	*	*	*	5.3	43.6	126.0	276.0	363.2	287.9
2006 5/.....	43.3	*	*	*	*	0.6	6.3	46.2	131.3	277.8	370.8	268.4
2005 5/.....	44.0	*	*	*	*	*	7.0	45.5	140.7	283.5	373.4	294.1
2004 5/.....	44.2	*	*	*	*	0.5	8.0	46.5	144.2	285.3	390.1	281.5
2003 5/.....	44.5	*	*	*	*	*	8.0	48.5	147.4	293.9	389.4	283.5
2002.....	44.7	*	*	*	*	0.4	9.0	48.1	155.6	299.5	385.6	274.9
2001.....	44.6	*	*	*	*	*	9.2	47.5	149.4	311.4	390.9	281.6
2000.....	45.4	*	*	*	*	0.6	10.4	51.5	155.0	320.2	379.4	291.0
1999.....	45.6	*	*	*	*	0.4	9.6	51.5	161.1	317.7	390.9	293.3

Non-Hispanic black, male

2007 5/.....	53.6	*	*	*	*	*	5.7	55.5	182.5	403.9	554.8	514.8
2006 5/.....	54.1	*	*	*	*	*	6.8	58.5	189.5	410.3	574.5	448.1
2005 5/.....	55.0	*	*	*	*	*	7.9	58.2	203.5	417.3	575.5	503.4
2004 5/.....	56.3	*	*	*	*	*	9.0	64.3	211.1	419.0	619.8	488.8
2003 5/.....	57.0	*	*	*	*	*	9.7	65.3	215.1	434.9	632.5	509.3
2002.....	58.1	*	*	*	*	*	11.0	66.3	228.0	451.4	633.9	490.1
2001.....	59.2	*	*	*	*	*	11.4	67.5	218.3	484.4	668.0	550.9
2000.....	60.3	*	*	*	*	*	13.8	72.4	227.1	495.4	649.5	567.1
1999.....	60.9	*	*	*	*	*	11.7	73.7	241.9	491.0	665.1	580.0

Non-Hispanic black, female

2007 5/.....	33.2	*	*	*	*	*	5.0	33.4	80.6	184.8	254.3	193.9
2006 5/.....	33.5	*	*	*	*	*	5.9	35.7	84.5	183.1	255.3	195.1
2005 5/.....	34.0	*	*	*	*	*	6.3	34.6	90.1	187.9	258.8	209.9
2004 5/.....	33.2	*	*	*	*	*	7.0	31.3	90.5	190.0	260.5	199.6
2003 5/.....	33.1	*	*	*	*	*	6.5	34.1	93.1	193.5	252.1	196.0
2002.....	32.6	*	*	*	*	*	7.3	32.5	97.5	191.8	245.1	192.7
2001.....	31.3	*	*	*	*	*	7.3	30.4	94.0	189.4	235.3	181.8
2000.....	31.9	*	*	*	*	*	7.4	33.6	97.0	196.9	227.3	188.4
1999.....	31.8	*	*	*	*	*	7.9	32.5	96.4	195.5	237.4	186.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant melanoma of skin (C43)

All origins 3/, both sexes

2007.....	2.8	*	*	*	0.1	0.4	1.1	2.7	5.3	9.0	15.6	19.9
2006.....	2.8	*	*	*	0.1	0.5	1.2	2.7	5.2	9.4	15.6	20.1
2005.....	2.8	*	*	*	0.1	0.5	1.2	2.8	5.2	9.4	15.6	19.8
2004.....	2.7	*	*	*	0.1	0.5	1.2	2.8	5.3	9.3	14.7	17.7
2003.....	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002.....	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000.....	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4

All origins 3/, male

2007.....	3.7	*	*	*	0.1	0.5	1.2	3.5	7.7	13.5	24.8	35.8
2006.....	3.7	*	*	*	0.1	0.5	1.4	3.5	7.4	13.7	25.6	36.7
2005.....	3.6	*	*	*	0.1	0.7	1.4	3.6	7.0	13.8	24.5	34.9
2004.....	3.5	*	*	*	*	0.6	1.4	3.7	7.6	13.7	23.7	31.4
2003.....	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002.....	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001.....	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000.....	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999.....	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5

All origins 3/, female

2007.....	1.9	*	*	*	*	0.4	0.9	1.9	3.1	5.2	9.3	12.4
2006.....	2.0	*	*	*	*	0.4	0.9	1.9	3.2	5.8	8.9	12.4
2005.....	2.0	*	*	*	*	0.4	1.0	2.1	3.5	5.6	9.5	12.9
2004.....	1.9	*	*	*	0.1	0.4	1.0	2.0	3.2	5.5	8.7	11.5
2003.....	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002.....	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001.....	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000.....	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999.....	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant melanoma of skin (C43)

Non-Hispanic 4/, both sexes

2007.....	3.2	*	*	*	0.1	0.5	1.2	2.9	5.7	9.5	16.4	20.6
2006.....	3.2	*	*	*	0.1	0.5	1.3	2.9	5.5	9.9	16.3	20.8
2005.....	3.2	*	*	*	0.1	0.6	1.4	3.0	5.5	9.9	16.3	20.5
2004.....	3.1	*	*	*	0.1	0.6	1.3	3.0	5.6	9.8	15.3	18.3
2003.....	3.0	*	*	*	0.1	0.6	1.4	3.1	5.9	9.1	15.1	18.1
2002.....	2.9	*	*	*	0.1	0.6	1.5	3.0	5.4	9.3	14.1	18.2
2001.....	3.0	*	*	*	0.1	0.5	1.5	3.3	5.6	9.7	13.7	18.3
2000.....	3.0	*	*	*	0.1	0.5	1.5	3.4	5.7	9.3	13.5	19.3
1999.....	2.9	*	*	*	0.1	0.6	1.5	3.3	5.6	9.3	13.7	16.7

Non-Hispanic 4/, male

2007.....	4.3	*	*	*	*	0.6	1.4	3.8	8.3	14.2	26.0	37.2
2006.....	4.3	*	*	*	0.2	0.6	1.6	3.8	7.9	14.5	26.8	38.0
2005.....	4.2	*	*	*	0.1	0.8	1.6	3.9	7.4	14.5	25.6	36.2
2004.....	4.1	*	*	*	*	0.7	1.6	3.9	8.0	14.5	24.8	32.7
2003.....	4.0	*	*	*	0.1	0.7	1.7	4.0	8.2	13.2	24.2	30.9
2002.....	3.8	*	*	*	*	0.7	1.8	4.1	7.6	13.7	22.0	30.2
2001.....	3.9	*	*	*	0.1	0.6	1.9	4.2	7.8	14.6	22.0	31.0
2000.....	3.7	*	*	*	0.1	0.6	1.7	4.4	7.8	13.4	21.3	32.6
1999.....	3.7	*	*	*	0.1	0.6	2.0	4.2	7.8	13.7	21.5	28.2

Non-Hispanic 4/, female

2007.....	2.2	*	*	*	*	0.5	1.0	2.0	3.3	5.4	9.7	12.7
2006.....	2.2	*	*	*	*	0.4	1.0	2.1	3.4	6.0	9.1	12.8
2005.....	2.3	*	*	*	*	0.4	1.1	2.2	3.7	5.9	9.9	13.4
2004.....	2.1	*	*	*	*	0.5	1.1	2.1	3.4	5.8	9.0	11.9
2003.....	2.2	*	*	*	*	0.4	1.1	2.2	3.6	5.7	9.0	12.4
2002.....	2.1	*	*	*	*	0.4	1.2	2.0	3.3	5.7	8.9	12.9
2001.....	2.1	*	*	*	*	0.5	1.2	2.4	3.5	5.6	8.3	12.9
2000.....	2.2	*	*	*	*	0.4	1.3	2.4	3.7	5.9	8.4	13.9
1999.....	2.1	*	*	*	*	0.5	1.1	2.3	3.5	5.7	8.7	12.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant melanoma of skin (C43)

Non-Hispanic white, both sexes

2007 5/.....	4.0	*	*	*	0.1	0.7	1.5	3.5	6.7	10.9	18.2	22.4
2006 5/.....	4.0	*	*	*	0.2	0.7	1.6	3.5	6.4	11.5	18.0	22.5
2005 5/.....	4.0	*	*	*	0.1	0.8	1.7	3.7	6.3	11.3	18.0	22.2
2004 5/.....	3.8	*	*	*	0.1	0.8	1.7	3.6	6.6	11.2	16.9	19.9
2003 5/.....	3.8	*	*	*	0.1	0.7	1.8	3.7	6.8	10.4	16.6	19.6
2002.....	3.6	*	*	*	0.1	0.7	1.8	3.6	6.2	10.5	15.6	19.6
2001.....	3.7	*	*	*	0.2	0.7	1.9	4.0	6.5	10.8	15.0	19.9
2000.....	3.6	*	*	*	0.2	0.7	1.8	4.0	6.5	10.4	14.7	20.8
1999.....	3.5	*	*	*	0.1	0.7	1.9	3.9	6.4	10.5	14.9	17.9

Non-Hispanic white, male

2007 5/.....	5.4	*	*	*	*	0.7	1.8	4.6	9.6	16.3	28.9	40.6
2006 5/.....	5.3	*	*	*	0.2	0.8	2.0	4.6	9.0	16.6	29.5	41.3
2005 5/.....	5.2	*	*	*	*	1.0	2.0	4.6	8.5	16.6	28.3	39.7
2004 5/.....	5.1	*	*	*	*	0.9	2.0	4.7	9.3	16.6	27.2	35.8
2003 5/.....	4.9	*	*	*	0.2	0.9	2.1	4.8	9.4	14.9	26.5	33.8
2002.....	4.7	*	*	*	*	0.9	2.2	4.8	8.6	15.3	24.1	32.9
2001.....	4.8	*	*	*	0.2	0.8	2.3	5.0	9.0	16.3	24.1	34.1
2000.....	4.6	*	*	*	0.2	0.8	2.0	5.2	8.9	15.0	23.2	35.7
1999.....	4.5	*	*	*	*	0.8	2.4	5.0	8.9	15.4	23.3	30.5

Non-Hispanic white, female

2007 5/.....	2.7	*	*	*	*	0.6	1.3	2.4	3.9	6.1	10.7	13.8
2006 5/.....	2.7	*	*	*	*	0.6	1.2	2.5	3.9	7.0	10.0	13.8
2005 5/.....	2.8	*	*	*	*	0.6	1.4	2.7	4.3	6.7	11.0	14.3
2004 5/.....	2.6	*	*	*	*	0.6	1.4	2.5	4.0	6.6	9.9	12.8
2003 5/.....	2.6	*	*	*	*	0.5	1.4	2.6	4.2	6.4	9.8	13.3
2002.....	2.6	*	*	*	*	0.6	1.4	2.4	3.9	6.5	9.8	13.8
2001.....	2.6	*	*	*	*	0.6	1.5	2.9	4.1	6.2	9.0	13.9
2000.....	2.7	*	*	*	*	0.6	1.6	2.8	4.2	6.6	9.1	14.7
1999.....	2.5	*	*	*	*	0.6	1.4	2.8	4.1	6.4	9.4	12.9



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of breast (C50)

All origins 3/, both sexes

2007.....	13.6	*	*	*	*	0.8	5.1	13.7	26.8	42.3	69.7	116.6
2006.....	13.8	*	*	*	*	0.9	5.4	14.1	28.1	42.1	71.6	116.7
2005.....	14.0	*	*	*	*	0.9	5.7	14.7	28.5	43.5	71.9	122.7
2004.....	14.1	*	*	*	*	1.0	5.7	15.0	29.1	44.9	72.2	124.2
2003.....	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002.....	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001.....	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000.....	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999.....	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4

All origins 3/, male

2007.....	0.2	*	*	*	*	*	*	0.2	0.4	1.0	2.0	3.4
2006.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.0
2005.....	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.4
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.3
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5

All origins 3/, female

2007.....	26.5	*	*	*	*	1.7	10.1	26.7	51.3	77.3	116.3	170.4
2006.....	26.9	*	*	*	*	1.8	10.8	27.6	53.7	76.9	119.2	169.9
2005.....	27.3	*	*	*	*	1.8	11.3	28.7	54.5	79.2	119.2	177.9
2004.....	27.5	*	*	*	*	2.0	11.3	29.3	55.8	81.6	119.5	178.6
2003.....	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002.....	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001.....	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000.....	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999.....	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of breast (C50)</b>												
<b>Hispanic, both sexes</b>												
2007.....	4.6	*	*	*	*	0.6	3.5	9.6	17.8	26.8	39.7	57.2
2006.....	4.7	*	*	*	*	0.8	3.7	9.7	19.0	26.3	41.0	58.7
2005.....	4.5	*	*	*	*	0.5	3.8	9.9	18.2	26.0	43.5	62.5
2004.....	4.7	*	*	*	*	0.6	3.7	9.9	20.2	29.4	41.2	72.3
2003.....	4.7	*	*	*	*	0.7	4.3	10.5	20.5	27.1	42.0	77.7
2002.....	4.5	*	*	*	*	0.7	3.8	11.0	17.6	27.4	43.7	71.0
2001.....	4.8	*	*	*	*	0.8	4.5	12.0	22.0	28.0	39.5	68.8
2000.....	4.8	*	*	*	*	0.8	4.3	12.1	21.0	31.3	45.6	71.7
1999.....	4.6	*	*	*	*	0.7	4.0	12.2	19.6	28.0	43.5	86.3
<b>Hispanic, male</b>												
2007.....	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Hispanic, female</b>												
2007.....	9.4	*	*	*	*	1.4	7.4	19.4	33.9	48.3	66.5	88.3
2006.....	9.6	*	*	*	*	1.8	7.9	19.6	36.4	47.3	68.8	89.9
2005.....	9.4	*	*	*	*	1.0	8.0	20.0	34.7	46.9	73.3	95.1
2004.....	9.6	*	*	*	*	1.4	7.7	19.5	38.2	52.1	69.1	108.8
2003.....	9.8	*	*	*	*	1.6	9.1	21.0	38.8	48.4	70.8	114.8
2002.....	9.2	*	*	*	*	1.6	7.8	21.6	33.5	48.7	73.1	105.3
2001.....	9.7	*	*	*	*	1.7	9.2	23.8	41.5	49.3	64.7	102.4
2000.....	9.7	*	*	*	*	1.8	8.7	23.9	39.1	54.9	74.9	105.8
1999.....	9.3	*	*	*	*	1.6	8.3	23.9	36.6	49.1	71.8	127.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of breast (C50)

Non-Hispanic 4/, both sexes

2007.....	15.2	*	*	*	*	0.9	5.3	14.1	27.5	43.5	71.6	119.6
2006.....	15.3	*	*	*	*	0.9	5.7	14.6	28.8	43.2	73.5	119.4
2005.....	15.6	*	*	*	*	1.0	6.0	15.2	29.3	44.8	73.4	125.4
2004.....	15.6	*	*	*	*	1.1	6.0	15.5	29.9	45.9	73.9	126.3
2003.....	15.9	*	*	*	*	1.1	6.4	16.1	30.3	46.7	76.4	134.7
2002.....	16.0	*	*	*	*	1.1	6.4	16.6	30.3	47.7	78.0	136.8
2001.....	16.1	*	*	*	*	1.2	6.5	17.2	30.7	48.8	78.4	136.4
2000.....	16.5	*	*	*	*	1.2	6.5	17.3	31.9	50.1	80.1	149.7
1999.....	16.3	*	*	*	*	1.2	6.3	17.3	31.9	50.6	80.2	146.2

Non-Hispanic 4/, male

2007.....	0.3	*	*	*	*	*	*	0.2	0.4	1.1	2.0	3.6
2006.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.2
2005.....	0.3	*	*	*	*	*	*	0.2	0.5	1.4	2.2	2.6
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.3
2003.....	0.3	*	*	*	*	*	*	0.3	0.6	1.1	2.1	3.8
2002.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.3	3.5
2001.....	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.3	3.7
2000.....	0.3	*	*	*	*	*	*	0.3	0.7	1.4	2.2	4.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.1	4.7

Non-Hispanic 4/, female

2007.....	29.4	*	*	*	*	1.8	10.5	27.5	52.8	79.6	119.5	174.4
2006.....	29.7	*	*	*	*	1.9	11.2	28.5	55.1	79.1	122.2	173.5
2005.....	30.1	*	*	*	*	2.0	11.9	29.6	56.1	81.5	121.7	181.5
2004.....	30.2	*	*	*	*	2.1	11.9	30.3	57.2	83.7	122.1	181.3
2003.....	30.8	*	*	*	*	2.1	12.6	31.3	57.9	84.9	126.0	192.2
2002.....	31.0	*	*	*	*	2.2	12.6	32.3	57.9	86.7	128.2	194.4
2001.....	31.1	*	*	*	*	2.5	12.8	33.6	58.6	88.1	128.3	191.6
2000.....	31.8	*	*	*	*	2.4	12.8	33.8	60.7	90.3	130.8	208.6
1999.....	31.4	*	*	*	*	2.3	12.4	33.6	60.7	91.2	130.9	202.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of breast (C50)

Non-Hispanic white, both sexes

2007 5/.....	16.0	*	*	*	*	0.7	4.7	12.8	26.2	43.1	71.8	120.8
2006 5/.....	16.2	*	*	*	*	0.8	5.0	13.3	27.6	43.1	73.8	119.8
2005 5/.....	16.3	*	*	*	*	0.8	5.3	13.6	27.9	44.5	74.0	126.3
2004 5/.....	16.4	*	*	*	*	0.9	5.2	14.0	28.7	45.8	74.7	126.8
2003 5/.....	16.8	*	*	*	*	0.9	5.7	14.8	29.0	46.3	76.7	134.7
2002.....	16.9	*	*	*	*	0.9	5.6	15.2	29.4	47.4	78.0	137.7
2001.....	17.0	*	*	*	*	1.0	5.8	16.0	29.6	48.2	78.7	137.1
2000.....	17.4	*	*	*	*	1.1	5.8	16.1	30.8	50.3	80.6	149.6
1999.....	17.1	*	*	*	*	1.0	5.6	16.0	30.6	50.2	80.3	147.8

Non-Hispanic white, male

2007 5/.....	0.3	*	*	*	*	*	*	0.2	0.4	1.0	1.9	3.6
2006 5/.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.3
2005 5/.....	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.6
2004 5/.....	0.3	*	*	*	*	*	*	0.1	0.5	1.0	2.2	2.9
2003 5/.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.5
2000.....	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.0
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.1	2.0	4.5

Non-Hispanic white, female

2007 5/.....	31.1	*	*	*	*	1.4	9.3	25.3	50.8	79.8	120.7	176.3
2006 5/.....	31.4	*	*	*	*	1.5	10.0	26.3	53.4	79.6	123.6	174.1
2005 5/.....	31.8	*	*	*	*	1.7	10.6	26.8	53.9	81.9	123.4	182.9
2004 5/.....	32.0	*	*	*	*	1.8	10.4	27.7	55.6	84.4	124.2	182.4
2003 5/.....	32.6	*	*	*	*	1.8	11.3	29.1	56.0	85.1	127.1	192.3
2002.....	32.9	*	*	*	*	1.9	11.1	30.0	56.7	87.0	128.9	195.8
2001.....	32.9	*	*	*	*	2.0	11.5	31.6	57.0	88.0	129.4	192.6
2000.....	33.8	*	*	*	*	2.1	11.6	31.7	59.2	91.4	132.2	208.3
1999.....	33.3	*	*	*	*	2.0	11.1	31.5	58.9	91.2	131.5	204.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of breast (C50)

Non-Hispanic black, both sexes

2007 5/.....	15.2	*	*	*	*	2.0	10.0	24.7	43.4	57.8	88.6	138.8
2006 5/.....	15.1	*	*	*	*	2.0	10.5	24.9	43.3	54.6	90.8	144.7
2005 5/.....	15.5	*	*	*	*	1.9	11.2	27.8	46.3	58.1	83.2	146.1
2004 5/.....	15.0	*	*	*	*	2.2	11.3	26.7	44.7	57.4	83.1	145.6
2003 5/.....	15.5	*	*	*	*	2.3	11.4	27.1	46.1	60.1	91.3	163.3
2002.....	15.4	*	*	*	*	2.4	12.0	27.7	43.7	61.1	95.3	154.0
2001.....	15.5	*	*	*	*	2.7	11.5	27.9	45.6	64.6	93.0	153.9
2000.....	15.3	*	*	*	*	2.3	11.5	28.4	46.4	59.4	91.7	178.7
1999.....	15.5	*	*	*	*	2.4	12.0	29.2	47.1	64.5	95.6	149.1

Non-Hispanic black, male

2007 5/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/.....	0.3	*	*	*	*	*	*	*	*	2.9	*	*
2004 5/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic black, female

2007 5/.....	28.8	*	*	*	*	3.8	18.9	45.6	77.5	97.8	137.3	194.6
2006 5/.....	28.5	*	*	*	*	3.9	19.8	45.6	77.3	92.3	140.7	202.5
2005 5/.....	29.4	*	*	*	*	3.7	21.0	51.4	82.8	97.6	128.2	204.1
2004 5/.....	28.4	*	*	*	*	4.2	21.2	49.3	79.8	96.8	128.5	199.6
2003 5/.....	29.4	*	*	*	*	4.3	21.4	50.1	82.4	101.6	141.8	224.5
2002.....	29.1	*	*	*	*	4.5	22.5	51.0	77.8	102.7	147.1	210.7
2001.....	29.2	*	*	*	*	5.1	21.5	51.3	81.6	108.4	143.0	208.3
2000.....	28.9	*	*	*	*	4.3	21.5	52.5	82.5	100.1	141.5	241.6
1999.....	29.2	*	*	*	*	4.5	22.3	53.7	83.7	108.2	147.2	201.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes

2007.....	1.3	*	*	*	*	0.5	1.5	2.1	2.7	3.2	3.9	4.7
2006.....	1.3	*	*	*	*	0.4	1.4	2.2	2.6	3.2	3.8	5.2
2005.....	1.3	*	*	*	*	0.4	1.4	2.3	2.6	3.1	4.0	5.4
2004.....	1.3	*	*	*	*	0.5	1.4	2.0	2.8	3.3	3.7	5.1
2003.....	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
2002.....	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001.....	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000.....	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999.....	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9

All origins 3/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins 3/, female

2007.....	2.6	*	*	*	*	0.9	3.0	4.1	5.3	5.8	6.6	7.0
2006.....	2.6	*	*	*	*	0.9	2.9	4.4	5.0	5.9	6.4	7.7
2005.....	2.6	*	*	*	*	0.9	2.8	4.4	5.1	5.7	6.7	7.8
2004.....	2.6	*	*	*	*	1.1	2.8	4.0	5.4	6.0	6.2	7.3
2003.....	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
2002.....	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001.....	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000.....	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999.....	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Hispanic, both sexes

2007.....	1.0	*	*	*	*	0.5	1.3	2.7	4.0	3.6	5.2	*
2006.....	1.0	*	*	*	*	0.4	1.4	2.8	3.3	4.4	6.6	*
2005.....	1.0	*	*	*	*	0.3	1.6	2.8	3.8	4.8	4.5	*
2004.....	1.0	*	*	*	*	0.4	1.6	2.4	4.3	4.3	5.7	*
2003.....	1.0	*	*	*	*	0.4	1.4	2.9	3.1	3.5	6.6	*
2002.....	1.0	*	*	*	*	0.4	1.5	2.4	3.6	4.7	6.7	*
2001.....	1.1	*	*	*	*	0.5	1.9	3.0	4.1	3.4	7.0	*
2000.....	1.2	*	*	*	*	0.5	2.1	2.8	4.6	5.6	8.5	13.9
1999.....	1.1	*	*	*	*	0.5	1.8	3.3	3.9	4.9	6.5	*

Hispanic, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, female

2007.....	2.2	*	*	*	*	1.1	2.8	5.4	7.7	6.5	8.9	*
2006.....	2.2	*	*	*	*	1.0	2.9	5.7	6.3	7.9	11.3	*
2005.....	2.2	*	*	*	*	0.6	3.4	5.7	7.2	8.8	7.7	*
2004.....	2.2	*	*	*	*	0.9	3.3	4.8	8.2	7.7	9.6	*
2003.....	2.0	*	*	*	*	0.9	3.0	5.7	5.9	6.3	11.1	*
2002.....	2.1	*	*	*	*	0.9	3.1	4.7	6.8	8.4	11.2	*
2001.....	2.3	*	*	*	*	1.1	3.9	6.0	7.7	6.1	11.8	*
2000.....	2.5	*	*	*	*	1.1	4.4	5.5	8.6	10.0	14.2	20.8
1999.....	2.3	*	*	*	*	1.1	3.6	6.6	7.4	8.8	10.8	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 4/, both sexes

2007.....	1.4	*	*	*	*	0.4	1.5	2.0	2.6	3.1	3.8	4.7
2006.....	1.4	*	*	*	*	0.5	1.4	2.2	2.5	3.1	3.6	5.2
2005.....	1.4	*	*	*	*	0.5	1.4	2.2	2.5	2.9	3.9	5.3
2004.....	1.4	*	*	*	*	0.6	1.4	2.0	2.7	3.2	3.6	5.0
2003.....	1.4	*	*	*	*	0.5	1.4	2.1	2.8	3.0	4.2	5.7
2002.....	1.4	*	*	*	*	0.5	1.5	2.2	2.6	3.4	4.0	6.1
2001.....	1.5	*	*	*	*	0.5	1.5	2.4	2.8	3.2	4.5	7.0
2000.....	1.5	*	*	*	*	0.6	1.5	2.4	3.0	3.6	4.7	7.1
1999.....	1.6	*	*	*	*	0.5	1.5	2.4	3.2	3.7	4.6	7.9

Non-Hispanic 4/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 4/, female

2007.....	2.7	*	*	*	*	0.9	3.0	4.0	5.0	5.8	6.4	6.9
2006.....	2.7	*	*	*	*	0.9	2.9	4.2	4.9	5.7	6.1	7.6
2005.....	2.7	*	*	*	*	0.9	2.7	4.3	4.9	5.4	6.6	7.7
2004.....	2.6	*	*	*	*	1.1	2.7	3.9	5.2	5.9	6.0	7.2
2003.....	2.7	*	*	*	*	1.0	2.8	4.1	5.4	5.6	7.0	8.2
2002.....	2.8	*	*	*	*	1.1	2.9	4.3	5.1	6.2	6.7	8.7
2001.....	2.9	*	*	*	*	1.0	3.0	4.6	5.3	5.8	7.5	10.0
2000.....	3.0	*	*	*	*	1.2	2.9	4.7	5.7	6.6	7.7	9.9
1999.....	3.0	*	*	*	*	1.1	3.0	4.7	6.1	6.7	7.5	11.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic white, both sexes

2007 5/.....	1.3	*	*	*	*	0.4	1.4	1.8	2.3	2.7	3.2	3.9
2006 5/.....	1.3	*	*	*	*	0.4	1.3	1.9	2.4	2.6	3.1	4.1
2005 5/.....	1.3	*	*	*	*	0.5	1.3	1.9	2.1	2.6	3.4	4.7
2004 5/.....	1.2	*	*	*	*	0.5	1.2	1.8	2.4	2.6	3.1	3.9
2003 5/.....	1.3	*	*	*	*	0.4	1.3	1.9	2.5	2.5	3.4	4.8
2002.....	1.3	*	*	*	*	0.5	1.3	1.9	2.3	2.7	3.4	4.9
2001.....	1.4	*	*	*	*	0.5	1.4	2.1	2.4	2.6	3.8	6.1
2000.....	1.4	*	*	*	*	0.5	1.3	2.1	2.5	2.9	4.1	6.0
1999.....	1.4	*	*	*	*	0.6	1.4	2.3	2.7	3.0	3.9	6.7

Non-Hispanic white, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic white, female

2007 5/.....	2.5	*	*	*	*	0.8	2.8	3.5	4.5	5.0	5.5	5.8
2006 5/.....	2.5	*	*	*	*	0.8	2.6	3.8	4.6	4.8	5.2	6.1
2005 5/.....	2.5	*	*	*	*	0.9	2.5	3.8	4.2	4.8	5.7	6.8
2004 5/.....	2.4	*	*	*	*	1.1	2.5	3.5	4.6	4.9	5.2	5.7
2003 5/.....	2.5	*	*	*	*	0.9	2.6	3.7	4.9	4.7	5.7	7.0
2002.....	2.5	*	*	*	*	1.0	2.7	3.8	4.5	5.0	5.7	7.0
2001.....	2.7	*	*	*	*	1.0	2.8	4.2	4.7	4.9	6.3	8.6
2000.....	2.7	*	*	*	*	1.1	2.5	4.2	4.8	5.3	6.8	8.4
1999.....	2.8	*	*	*	*	1.1	2.8	4.5	5.2	5.5	6.5	9.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic black, both sexes

2007 5/.....	2.1	*	*	*	*	0.9	2.5	3.6	4.8	6.9	9.5	13.0
2006 5/.....	2.1	*	*	*	*	0.8	2.5	3.7	4.1	7.7	8.8	17.5
2005 5/.....	2.1	*	*	*	*	0.5	2.4	3.8	5.3	6.5	10.7	13.4
2004 5/.....	2.1	*	*	*	*	0.8	2.3	3.8	5.4	7.7	9.1	16.4
2003 5/.....	2.2	*	*	*	*	0.9	2.3	3.7	5.7	7.5	11.6	14.9
2002.....	2.3	*	*	*	*	0.8	2.6	4.2	5.1	9.2	10.7	17.5
2001.....	2.2	*	*	*	*	0.8	2.5	3.8	5.3	7.9	12.0	19.0
2000.....	2.5	*	*	*	*	1.0	2.6	4.3	6.4	9.8	12.0	19.7
1999.....	2.4	*	*	*	*	0.7	2.5	3.6	7.2	10.8	11.3	21.7

Non-Hispanic black, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic black, female

2007 5/.....	4.0	*	*	*	*	1.8	4.7	6.6	8.7	11.9	14.9	18.4
2006 5/.....	3.9	*	*	*	*	1.5	4.7	6.8	7.3	13.2	13.7	24.7
2005 5/.....	4.0	*	*	*	*	1.1	4.6	7.1	9.5	11.2	16.8	18.8
2004 5/.....	4.0	*	*	*	*	1.6	4.3	7.1	9.8	13.2	14.2	22.9
2003 5/.....	4.1	*	*	*	*	1.8	4.3	6.9	10.2	12.8	18.1	20.6
2002.....	4.3	*	*	*	*	1.6	4.9	7.8	9.2	15.8	16.7	24.2
2001.....	4.2	*	*	*	*	1.6	4.6	7.1	9.6	13.5	18.7	26.1
2000.....	4.7	*	*	*	*	1.9	5.0	8.0	11.5	16.6	18.8	27.0
1999.....	4.5	*	*	*	*	1.2	4.8	6.6	13.0	18.3	17.7	29.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins 3/, both sexes

2007.....	2.5	*	*	*	*	0.1	0.4	1.3	4.8	10.4	15.0	20.1
2006.....	2.5	*	*	*	*	0.1	0.3	1.4	4.8	10.0	15.7	21.6
2005.....	2.4	*	*	*	*	0.1	0.3	1.3	4.8	9.7	15.0	22.1
2004.....	2.4	*	*	*	*	0.0	0.4	1.3	4.4	10.1	15.5	22.3
2003.....	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
2002.....	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000.....	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999.....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0

All origins 3/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins 3/, female

2007.....	4.9	*	*	*	*	0.2	0.8	2.6	9.3	19.3	25.4	29.6
2006.....	4.9	*	*	*	*	0.1	0.7	2.7	9.3	18.5	26.4	31.6
2005.....	4.7	*	*	*	*	0.1	0.7	2.6	9.3	17.9	25.1	32.2
2004.....	4.7	*	*	*	*	0.1	0.8	2.6	8.6	18.5	25.9	32.3
2003.....	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
2002.....	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001.....	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999.....	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Hispanic, both sexes

2007.....	0.9	*	*	*	*	*	0.4	1.1	3.6	8.3	8.4	8.2
2006.....	0.9	*	*	*	*	*	0.3	1.2	3.5	6.5	11.6	15.3
2005.....	0.9	*	*	*	*	*	0.4	1.5	3.4	7.8	11.9	11.1
2004.....	0.8	*	*	*	*	*	*	0.9	3.5	7.9	10.5	9.5
2003.....	0.8	*	*	*	*	*	0.4	1.2	2.9	7.0	9.8	20.7
2002.....	0.9	*	*	*	*	*	0.3	1.1	3.2	9.0	11.0	20.5
2001.....	0.9	*	*	*	*	*	*	1.4	3.8	7.5	11.7	18.8
2000.....	0.8	*	*	*	*	*	*	1.1	3.6	7.0	10.7	19.9
1999.....	0.8	*	*	*	*	*	*	1.1	3.7	8.3	10.6	15.4

Hispanic, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, female

2007.....	1.8	*	*	*	*	*	0.9	2.2	6.9	15.1	14.2	12.7
2006.....	1.8	*	*	*	*	*	0.7	2.4	6.7	11.9	19.7	23.4
2005.....	1.9	*	*	*	*	*	0.8	3.1	6.5	14.1	20.2	16.9
2004.....	1.6	*	*	*	*	*	*	1.7	6.7	14.2	17.8	14.4
2003.....	1.7	*	*	*	*	*	0.9	2.3	5.6	12.5	16.6	31.3
2002.....	1.8	*	*	*	*	*	0.7	2.1	6.2	16.1	18.5	30.7
2001.....	1.8	*	*	*	*	*	*	2.8	7.3	13.4	19.6	27.9
2000.....	1.6	*	*	*	*	*	*	2.2	6.8	12.4	17.8	29.7
1999.....	1.6	*	*	*	*	*	*	2.2	7.0	14.8	17.7	22.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 4/, both sexes

2007.....	2.8	*	*	*	*	*	0.4	1.4	4.9	10.6	15.5	20.7
2006.....	2.7	*	*	*	*	0.1	0.3	1.4	4.9	10.3	15.9	21.9
2005.....	2.6	*	*	*	*	0.1	0.3	1.3	4.9	9.8	15.1	22.6
2004.....	2.6	*	*	*	*	*	0.4	1.4	4.5	10.2	15.8	22.8
2003.....	2.6	*	*	*	*	0.1	0.3	1.3	4.8	10.3	15.4	22.9
2002.....	2.6	*	*	*	*	*	0.3	1.4	4.6	10.1	15.5	24.6
2001.....	2.6	*	*	*	*	0.1	0.3	1.4	4.4	9.9	16.6	24.8
2000.....	2.6	*	*	*	*	0.1	0.3	1.4	4.7	9.8	16.0	24.9
1999.....	2.5	*	*	*	*	0.1	0.3	1.3	4.3	9.9	16.6	24.3

Non-Hispanic 4/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 4/, female

2007.....	5.4	*	*	*	*	*	0.8	2.7	9.6	19.6	26.1	30.5
2006.....	5.4	*	*	*	*	0.1	0.7	2.8	9.5	19.0	26.8	32.0
2005.....	5.2	*	*	*	*	0.1	0.6	2.6	9.5	18.2	25.4	32.9
2004.....	5.2	*	*	*	*	*	0.8	2.7	8.7	18.8	26.4	33.0
2003.....	5.1	*	*	*	*	0.1	0.7	2.6	9.2	18.9	25.6	33.0
2002.....	5.1	*	*	*	*	*	0.6	2.8	8.8	18.5	25.8	35.3
2001.....	5.1	*	*	*	*	0.2	0.6	2.8	8.5	18.1	27.5	35.1
2000.....	5.0	*	*	*	*	0.1	0.6	2.7	9.0	17.9	26.5	34.9
1999.....	4.9	*	*	*	*	0.1	0.6	2.5	8.3	18.0	27.4	34.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic white, both sexes

2007 5/.....	2.8	*	*	*	*	*	0.3	1.3	4.6	9.2	14.3	20.6
2006 5/.....	2.8	*	*	*	*	*	0.3	1.3	4.4	9.0	15.1	22.0
2005 5/.....	2.7	*	*	*	*	*	0.3	1.2	4.4	8.9	14.3	22.2
2004 5/.....	2.7	*	*	*	*	*	0.4	1.3	4.1	8.9	15.2	23.0
2003 5/.....	2.7	*	*	*	*	*	0.3	1.3	4.4	9.3	14.5	22.8
2002.....	2.6	*	*	*	*	*	0.3	1.3	4.1	8.9	14.6	23.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	4.0	8.9	15.9	24.5
2000.....	2.6	*	*	*	*	*	0.3	1.3	4.3	8.6	15.0	24.5
1999.....	2.6	*	*	*	*	*	0.3	1.3	4.0	8.9	16.0	23.6

Non-Hispanic white, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic white, female

2007 5/.....	5.4	*	*	*	*	*	0.7	2.6	8.9	17.1	24.3	30.3
2006 5/.....	5.5	*	*	*	*	*	0.7	2.6	8.6	16.8	25.5	32.2
2005 5/.....	5.3	*	*	*	*	*	0.6	2.5	8.6	16.5	24.1	32.4
2004 5/.....	5.3	*	*	*	*	*	0.8	2.6	8.0	16.6	25.5	33.3
2003 5/.....	5.2	*	*	*	*	*	0.6	2.5	8.5	17.2	24.4	32.8
2002.....	5.1	*	*	*	*	*	0.6	2.7	8.0	16.5	24.4	34.1
2001.....	5.2	*	*	*	*	*	0.6	2.7	7.8	16.4	26.5	34.7
2000.....	5.1	*	*	*	*	*	0.5	2.7	8.4	15.9	24.8	34.3
1999.....	5.1	*	*	*	*	*	0.6	2.5	7.7	16.3	26.4	33.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic black, both sexes

2007 5/.....	3.4	*	*	*	*	*	0.6	1.9	8.7	25.1	30.6	27.4
2006 5/.....	3.2	*	*	*	*	*	*	1.8	10.0	23.2	29.2	26.3
2005 5/.....	3.1	*	*	*	*	*	0.4	1.9	10.0	20.6	27.5	32.0
2004 5/.....	3.0	*	*	*	*	*	0.5	2.0	8.1	23.4	26.6	26.5
2003 5/.....	3.0	*	*	*	*	*	0.5	1.7	9.1	21.4	28.0	30.3
2002.....	3.1	*	*	*	*	*	0.4	2.0	8.6	22.7	28.4	39.6
2001.....	2.9	*	*	*	*	*	0.4	2.1	7.7	21.6	28.6	32.8
2000.....	3.0	*	*	*	*	*	0.4	1.7	8.4	22.9	31.1	33.7
1999.....	2.8	*	*	*	*	*	0.5	1.4	8.1	21.5	27.5	35.9

Non-Hispanic black, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic black, female

2007 5/.....	6.4	*	*	*	*	*	1.2	3.5	15.7	43.1	48.0	38.7
2006 5/.....	6.2	*	*	*	*	*	*	3.3	18.0	39.7	45.8	37.0
2005 5/.....	6.0	*	*	*	*	*	0.8	3.5	18.1	35.3	43.1	44.8
2004 5/.....	5.7	*	*	*	*	*	1.0	3.6	14.7	40.1	41.5	36.9
2003 5/.....	5.7	*	*	*	*	*	1.0	3.2	16.5	36.7	43.8	42.1
2002.....	5.8	*	*	*	*	*	0.7	3.6	15.5	38.7	44.4	54.8
2001.....	5.5	*	*	*	*	*	0.7	4.0	13.9	36.8	44.6	44.9
2000.....	5.7	*	*	*	*	*	0.7	3.1	15.3	39.1	48.6	46.2
1999.....	5.4	*	*	*	*	*	0.9	2.7	14.7	36.7	42.9	49.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

All origins 3/, both sexes

2007.....	4.8	*	*	*	0.1	0.2	0.8	3.5	9.2	18.7	30.3	37.6
2006.....	5.0	*	*	*	0.1	0.2	0.9	3.8	9.6	18.9	31.1	37.8
2005.....	5.0	*	*	*	*	0.2	0.9	4.0	10.0	19.1	30.9	38.1
2004.....	5.0	*	*	*	0.1	0.2	1.0	4.0	9.6	20.0	32.1	37.7
2003.....	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
2002.....	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001.....	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000.....	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999.....	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6

All origins 3/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins 3/, female

2007.....	9.6	*	*	*	0.1	0.4	1.6	6.9	17.7	34.6	51.2	55.4
2006.....	9.8	*	*	*	0.1	0.4	1.8	7.6	18.6	34.9	52.4	55.5
2005.....	9.8	*	*	*	*	0.4	1.9	7.8	19.4	35.3	51.8	55.6
2004.....	9.9	*	*	*	0.1	0.5	2.1	7.8	18.5	36.8	53.7	54.6
2003.....	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
2002.....	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001.....	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000.....	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999.....	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

Hispanic, both sexes

2007.....	1.7	*	*	*	*	*	0.7	2.7	6.1	12.1	21.6	22.9
2006.....	1.7	*	*	*	*	*	0.7	2.6	6.1	12.8	24.0	27.8
2005.....	1.7	*	*	*	*	*	0.6	3.1	7.1	13.1	19.9	27.4
2004.....	1.7	*	*	*	*	*	0.8	2.8	6.5	12.9	20.9	28.1
2003.....	1.6	*	*	*	*	*	0.8	2.7	6.4	12.7	22.0	28.5
2002.....	1.6	*	*	*	*	*	0.8	3.2	5.6	12.3	24.2	23.8
2001.....	1.6	*	*	*	*	*	0.8	2.7	7.1	14.4	20.0	30.6
2000.....	1.6	*	*	*	*	*	0.6	3.0	6.5	14.3	24.9	32.5
1999.....	1.4	*	*	*	*	*	0.6	2.7	6.1	12.2	20.8	23.9

Hispanic, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, female

2007.....	3.5	*	*	*	*	*	1.5	5.5	11.8	22.0	36.7	35.3
2006.....	3.5	*	*	*	*	*	1.4	5.2	11.6	23.2	40.7	42.5
2005.....	3.6	*	*	*	*	*	1.4	6.2	13.6	23.7	33.7	41.7
2004.....	3.4	*	*	*	*	*	1.7	5.5	12.5	23.3	35.3	42.5
2003.....	3.3	*	*	*	*	*	1.7	5.5	12.2	22.8	37.1	43.0
2002.....	3.3	*	*	*	*	*	1.7	6.3	10.7	22.1	40.8	35.6
2001.....	3.4	*	*	*	*	*	1.6	5.3	13.4	25.8	33.5	45.6
2000.....	3.4	*	*	*	*	*	1.2	5.9	12.2	25.5	41.6	48.5
1999.....	2.9	*	*	*	*	*	1.2	5.3	11.5	21.7	34.7	35.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

Non-Hispanic 4/, both sexes

2007.....	5.4	*	*	*	0.1	0.2	0.8	3.6	9.4	19.2	30.8	38.3
2006.....	5.5	*	*	*	*	0.2	1.0	4.0	10.0	19.3	31.5	38.3
2005.....	5.5	*	*	*	*	0.2	1.0	4.1	10.3	19.6	31.5	38.6
2004.....	5.6	*	*	*	0.1	0.3	1.1	4.1	9.8	20.5	32.7	38.1
2003.....	5.6	*	*	*	0.1	0.2	1.0	4.1	10.7	19.9	33.1	38.8
2002.....	5.6	*	*	*	0.1	0.3	1.1	4.4	10.7	20.2	33.8	38.7
2001.....	5.6	*	*	*	0.1	0.3	1.1	4.3	10.5	20.3	33.7	39.8
2000.....	5.5	*	*	*	*	0.3	1.1	4.3	10.3	19.9	33.4	41.2
1999.....	5.3	*	*	*	0.1	0.2	1.2	4.4	10.5	20.0	32.0	38.1

Non-Hispanic 4/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 4/, female

2007.....	10.6	*	*	*	0.1	0.4	1.7	7.0	18.2	35.6	52.1	56.4
2006.....	10.8	*	*	*	*	0.5	1.9	7.8	19.2	35.7	53.1	56.0
2005.....	10.8	*	*	*	*	0.4	2.0	8.0	19.9	36.2	52.9	56.2
2004.....	10.8	*	*	*	0.1	0.5	2.1	8.0	19.0	37.7	54.7	55.1
2003.....	10.9	*	*	*	0.1	0.5	2.1	8.1	20.5	36.6	55.2	55.9
2002.....	11.0	*	*	*	0.2	0.5	2.1	8.6	20.6	37.0	56.3	55.3
2001.....	10.9	*	*	*	0.1	0.5	2.1	8.4	20.3	37.3	55.8	56.4
2000.....	10.7	*	*	*	*	0.5	2.2	8.4	19.9	36.3	55.2	57.8
1999.....	10.4	*	*	*	0.2	0.4	2.3	8.6	20.1	36.4	52.7	53.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

Non-Hispanic white, both sexes

2007 5/.....	6.1	*	*	*	*	0.2	0.9	3.7	9.8	20.1	32.0	39.7
2006 5/.....	6.2	*	*	*	*	0.3	1.0	4.1	10.4	20.3	32.5	39.3
2005 5/.....	6.2	*	*	*	*	0.2	1.0	4.2	10.7	20.1	32.4	40.0
2004 5/.....	6.2	*	*	*	*	0.2	1.1	4.2	10.2	21.0	33.9	39.1
2003 5/.....	6.3	*	*	*	*	0.3	1.1	4.2	11.1	20.5	34.2	39.5
2002.....	6.3	*	*	*	0.1	0.3	1.1	4.6	11.1	20.9	34.8	39.6
2001.....	6.2	*	*	*	*	0.3	1.1	4.5	10.8	21.0	34.8	40.6
2000.....	6.2	*	*	*	*	0.3	1.1	4.5	10.8	20.5	34.5	42.2
1999.....	6.0	*	*	*	0.1	0.2	1.2	4.5	10.9	20.5	32.7	39.2

Non-Hispanic white, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic white, female

2007 5/.....	12.0	*	*	*	*	0.4	1.8	7.4	19.2	37.6	54.4	58.5
2006 5/.....	12.3	*	*	*	*	0.5	2.0	8.2	20.3	37.9	55.1	57.6
2005 5/.....	12.2	*	*	*	*	0.5	2.0	8.4	20.9	37.5	54.7	58.2
2004 5/.....	12.2	*	*	*	*	0.5	2.2	8.3	19.8	39.1	57.0	56.5
2003 5/.....	12.3	*	*	*	*	0.5	2.2	8.4	21.6	38.1	57.2	56.8
2002.....	12.4	*	*	*	0.2	0.5	2.2	9.1	21.7	38.7	58.0	56.7
2001.....	12.2	*	*	*	*	0.5	2.2	9.0	21.0	38.8	57.8	57.5
2000.....	12.0	*	*	*	*	0.6	2.3	8.9	21.0	37.7	57.2	59.2
1999.....	11.7	*	*	*	0.2	0.4	2.4	9.0	21.2	37.7	54.1	54.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

Non-Hispanic black, both sexes

2007 5/.....	3.0	*	*	*	*	*	0.6	3.0	8.4	16.7	25.7	27.9
2006 5/.....	3.1	*	*	*	*	*	0.9	3.5	8.6	16.1	25.6	33.7
2005 5/.....	3.2	*	*	*	*	*	1.0	3.2	8.8	18.9	28.2	31.1
2004 5/.....	3.4	*	*	*	*	*	0.9	4.0	8.8	20.5	28.0	32.5
2003 5/.....	3.2	*	*	*	*	*	1.0	3.9	8.8	17.8	27.8	37.2
2002.....	3.2	*	*	*	*	*	1.1	3.5	9.2	17.4	29.3	35.4
2001.....	3.2	*	*	*	*	*	1.2	3.1	10.2	18.5	27.2	36.5
2000.....	3.0	*	*	*	*	*	1.1	3.4	8.6	18.2	27.1	34.7
1999.....	3.1	*	*	*	*	*	0.9	3.6	9.1	18.7	28.7	29.1

Non-Hispanic black, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic black, female

2007 5/.....	5.8	*	*	*	*	*	1.2	5.6	15.2	28.7	40.4	39.4
2006 5/.....	5.9	*	*	*	*	*	1.7	6.6	15.5	27.6	40.1	47.5
2005 5/.....	6.2	*	*	*	*	*	1.9	6.0	15.9	32.4	44.2	43.7
2004 5/.....	6.5	*	*	*	*	*	1.8	7.5	15.9	35.2	43.8	45.4
2003 5/.....	6.2	*	*	*	*	*	1.9	7.2	15.9	30.5	43.5	51.6
2002.....	6.1	*	*	*	*	*	2.0	6.5	16.6	29.8	45.9	48.9
2001.....	6.1	*	*	*	*	*	2.2	5.7	18.4	31.5	42.5	50.0
2000.....	5.8	*	*	*	*	*	2.0	6.3	15.6	31.1	42.4	47.5
1999.....	5.8	*	*	*	*	*	1.6	6.7	16.4	31.9	44.8	40.0











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins 3/, both sexes

2007.....	4.2	*	*	0.1	0.1	0.1	0.6	2.8	7.9	16.3	27.2	33.0
2006.....	4.1	*	*	0.1	0.0	0.1	0.6	2.8	7.8	16.3	26.2	34.3
2005.....	4.2	*	*	0.1	*	0.1	0.7	3.0	8.3	16.9	26.9	32.6
2004.....	4.2	*	*	0.1	*	0.1	0.6	3.0	8.7	16.7	26.8	32.7
2003.....	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002.....	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001.....	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000.....	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999.....	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4

All origins 3/, male

2007.....	5.4	*	*	*	0.1	0.2	0.7	4.1	11.5	23.0	39.2	50.9
2006.....	5.2	*	*	*	*	0.2	0.9	4.0	11.4	23.1	37.1	52.1
2005.....	5.3	*	*	*	*	0.2	0.8	4.3	12.0	24.4	37.9	50.6
2004.....	5.3	*	*	*	*	0.2	0.8	4.3	12.8	23.9	38.2	47.0
2003.....	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002.....	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001.....	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000.....	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999.....	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4

All origins 3/, female

2007.....	3.1	*	*	*	*	*	0.4	1.6	4.5	10.6	18.9	24.5
2006.....	3.1	*	*	*	*	0.1	0.4	1.6	4.5	10.5	18.7	25.9
2005.....	3.1	*	*	*	*	*	0.5	1.8	4.8	10.5	19.4	24.3
2004.....	3.1	*	*	*	*	*	0.4	1.7	5.0	10.6	19.1	26.3
2003.....	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002.....	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001.....	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000.....	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999.....	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Hispanic, both sexes

2007.....	1.9	*	*	*	*	*	0.5	2.7	7.2	14.1	27.8	25.4
2006.....	1.7	*	*	*	*	*	0.4	2.3	6.9	13.2	22.1	23.5
2005.....	1.8	*	*	*	*	*	0.7	2.8	6.3	15.7	23.4	32.9
2004.....	1.8	*	*	*	*	*	0.7	3.1	8.2	14.0	23.4	28.1
2003.....	1.7	*	*	*	*	*	0.5	2.1	8.1	16.2	23.6	26.9
2002.....	1.7	*	*	*	*	*	0.6	2.8	8.1	12.9	24.4	26.1
2001.....	1.7	*	*	*	*	*	0.7	3.3	6.9	14.4	23.3	31.9
2000.....	1.6	*	*	*	*	*	0.6	2.8	7.9	12.3	24.7	25.9
1999.....	1.6	*	*	*	*	*	0.6	2.8	7.9	13.9	22.5	28.8

Hispanic, male

2007.....	2.5	*	*	*	*	*	0.6	4.1	10.7	20.2	44.1	38.6
2006.....	2.0	*	*	*	*	*	*	3.3	9.7	19.8	32.3	32.7
2005.....	2.2	*	*	*	*	*	0.7	3.4	9.3	22.5	33.1	46.1
2004.....	2.4	*	*	*	*	*	0.8	4.4	11.2	23.3	34.7	43.4
2003.....	2.1	*	*	*	*	*	0.7	2.8	11.0	22.3	32.2	38.5
2002.....	2.1	*	*	*	*	*	0.9	3.9	11.3	18.5	39.0	40.2
2001.....	2.2	*	*	*	*	*	1.2	4.7	9.9	21.4	35.3	57.1
2000.....	2.0	*	*	*	*	*	1.0	3.9	11.4	17.9	38.4	40.3
1999.....	2.0	*	*	*	*	*	0.9	4.3	11.8	21.1	29.7	*

Hispanic, female

2007.....	1.3	*	*	*	*	*	*	1.2	4.0	9.0	16.4	18.2
2006.....	1.2	*	*	*	*	*	*	1.3	4.3	7.8	15.0	18.6
2005.....	1.4	*	*	*	*	*	*	2.2	3.5	10.1	16.7	26.1
2004.....	1.3	*	*	*	*	*	*	1.7	5.5	6.5	15.6	20.2
2003.....	1.4	*	*	*	*	*	*	1.5	5.6	11.3	17.6	21.1
2002.....	1.2	*	*	*	*	*	*	1.7	5.2	8.4	14.3	19.1
2001.....	1.2	*	*	*	*	*	*	1.9	4.2	8.8	15.1	19.5
2000.....	1.1	*	*	*	*	*	*	1.8	4.7	7.8	15.5	*
1999.....	1.2	*	*	*	*	*	*	1.4	4.4	8.3	17.7	23.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 4/, both sexes

2007.....	4.6	*	*	0.1	0.1	0.1	0.6	2.8	7.9	16.5	27.1	33.4
2006.....	4.6	*	*	0.1	0.1	0.2	0.6	2.9	7.9	16.5	26.4	34.8
2005.....	4.6	*	*	0.1	*	0.1	0.7	3.0	8.4	16.9	27.1	32.5
2004.....	4.6	*	*	0.1	*	0.1	0.6	3.0	8.7	16.8	26.9	32.9
2003.....	4.6	*	*	0.1	*	0.2	0.8	3.0	9.0	16.9	27.2	33.4
2002.....	4.6	*	*	0.1	*	0.1	0.7	3.3	8.8	18.0	26.4	32.7
2001.....	4.6	*	*	0.1	0.1	0.1	0.8	3.3	9.4	17.8	26.4	32.3
2000.....	4.5	*	*	0.1	*	0.1	0.8	3.3	9.1	17.3	26.6	33.9
1999.....	4.3	*	*	0.1	*	0.1	0.8	3.2	8.9	16.8	24.6	31.4

Non-Hispanic 4/, male

2007.....	5.9	*	*	*	*	0.2	0.8	4.1	11.6	23.2	38.8	51.6
2006.....	5.8	*	*	*	*	0.2	0.9	4.1	11.5	23.3	37.3	53.2
2005.....	5.9	*	*	*	*	0.2	0.8	4.3	12.2	24.4	38.1	50.7
2004.....	5.8	*	*	*	*	0.2	0.8	4.3	12.9	23.9	38.4	47.1
2003.....	5.9	*	*	*	*	0.2	1.1	4.4	13.5	23.7	39.8	49.0
2002.....	5.9	*	*	*	*	0.2	1.0	4.9	13.0	25.5	37.7	50.4
2001.....	5.9	*	*	*	*	0.1	1.0	4.9	13.7	25.4	37.5	52.6
2000.....	5.7	*	*	*	*	0.1	1.0	4.8	13.2	24.4	38.1	55.4
1999.....	5.4	*	*	*	*	0.2	1.1	4.6	13.0	23.7	35.9	49.8

Non-Hispanic 4/, female

2007.....	3.4	*	*	*	*	*	0.4	1.6	4.5	10.8	19.0	24.8
2006.....	3.4	*	*	*	*	*	0.4	1.7	4.4	10.8	18.9	26.2
2005.....	3.4	*	*	*	*	*	0.5	1.7	5.0	10.6	19.6	24.2
2004.....	3.4	*	*	*	*	*	0.4	1.7	4.9	10.9	19.3	26.5
2003.....	3.4	*	*	*	*	0.1	0.4	1.7	4.9	11.1	18.7	26.6
2002.....	3.4	*	*	*	*	*	0.4	1.8	4.9	11.8	18.8	25.1
2001.....	3.4	*	*	0.1	*	0.1	0.6	1.8	5.4	11.4	19.1	23.8
2000.....	3.4	*	*	*	*	*	0.5	1.9	5.3	11.4	19.1	25.2
1999.....	3.2	*	*	*	*	*	0.5	1.8	5.2	11.2	17.3	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic white, both sexes

2007 5/.....	5.1	*	*	*	*	0.1	0.5	2.8	8.1	17.0	27.6	33.9
2006 5/.....	5.1	*	*	*	*	0.1	0.6	2.9	8.2	17.1	27.2	35.5
2005 5/.....	5.1	*	*	*	*	*	0.7	3.1	8.6	17.5	27.8	33.0
2004 5/.....	5.1	*	*	0.1	*	0.1	0.6	3.1	9.1	17.3	27.5	33.8
2003 5/.....	5.1	*	*	0.1	*	0.1	0.7	3.1	9.2	17.4	27.8	34.0
2002.....	5.1	*	*	0.1	*	0.1	0.7	3.4	9.1	18.7	26.8	33.5
2001.....	5.1	*	*	*	*	0.1	0.7	3.4	9.6	18.4	26.8	32.8
2000.....	5.0	*	*	*	*	0.1	0.7	3.4	9.3	17.7	26.9	34.6
1999.....	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5

Non-Hispanic white, male

2007 5/.....	6.5	*	*	*	*	*	0.7	4.0	11.7	23.5	39.7	53.0
2006 5/.....	6.4	*	*	*	*	*	0.9	4.1	11.7	23.9	38.5	54.6
2005 5/.....	6.5	*	*	*	*	*	0.8	4.3	12.3	25.1	38.9	52.1
2004 5/.....	6.5	*	*	*	*	*	0.8	4.5	13.3	24.4	39.1	48.6
2003 5/.....	6.6	*	*	*	*	*	1.0	4.5	13.7	24.3	40.7	50.4
2002.....	6.5	*	*	*	*	*	1.0	5.0	13.3	26.3	38.4	52.4
2001.....	6.5	*	*	*	*	*	1.0	4.9	13.8	26.2	37.7	54.3
2000.....	6.3	*	*	*	*	*	1.0	4.9	13.5	24.6	38.5	56.9
1999.....	6.0	*	*	*	*	*	1.0	4.5	13.3	24.0	36.5	50.2

Non-Hispanic white, female

2007 5/.....	3.8	*	*	*	*	*	0.3	1.6	4.7	11.3	19.1	24.8
2006 5/.....	3.8	*	*	*	*	*	0.3	1.6	4.8	11.2	19.4	26.6
2005 5/.....	3.8	*	*	*	*	*	0.5	1.8	5.1	11.0	20.1	24.3
2004 5/.....	3.8	*	*	*	*	*	0.3	1.8	5.1	11.2	19.6	27.2
2003 5/.....	3.8	*	*	*	*	*	0.4	1.7	4.8	11.6	19.1	26.8
2002.....	3.8	*	*	*	*	*	0.4	1.8	5.1	12.2	19.0	25.3
2001.....	3.8	*	*	*	*	*	0.5	1.8	5.5	11.8	19.5	23.8
2000.....	3.8	*	*	*	*	*	0.5	1.9	5.4	11.8	19.3	25.7
1999.....	3.5	*	*	*	*	*	0.5	1.8	5.3	11.5	17.3	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic black, both sexes

2007 5/.....	3.2	*	*	*	*	*	0.8	3.5	8.7	15.3	27.3	30.9
2006 5/.....	3.0	*	*	*	*	0.4	1.0	3.6	7.5	15.1	22.5	33.7
2005 5/.....	3.1	*	*	*	*	0.4	0.9	3.3	9.1	15.5	24.4	30.9
2004 5/.....	2.9	*	*	*	*	*	0.8	3.2	8.2	16.0	26.3	28.8
2003 5/.....	3.0	*	*	*	*	*	1.1	3.3	9.7	14.8	24.2	34.2
2002.....	2.9	*	*	*	*	*	0.8	3.6	8.8	15.8	26.5	28.1
2001.....	3.1	*	*	*	*	*	1.3	3.8	9.7	15.3	26.0	30.0
2000.....	2.9	*	*	*	*	*	0.9	3.5	9.0	16.8	25.9	29.6
1999.....	3.0	*	*	*	*	0.4	1.1	3.9	8.7	16.4	23.9	31.7

Non-Hispanic black, male

2007 5/.....	4.1	*	*	*	*	*	1.2	5.3	13.6	23.6	37.1	42.8
2006 5/.....	3.9	*	*	*	*	0.8	1.6	5.3	12.3	22.9	32.8	49.5
2005 5/.....	4.0	*	*	*	*	*	1.4	5.3	14.2	24.3	35.9	40.0
2004 5/.....	3.7	*	*	*	*	*	1.1	4.6	12.4	24.1	38.2	39.6
2003 5/.....	3.8	*	*	*	*	*	1.7	5.0	14.1	22.3	37.2	44.7
2002.....	3.6	*	*	*	*	*	1.2	5.0	14.4	22.1	36.6	39.4
2001.....	4.0	*	*	*	*	*	1.5	5.5	15.3	22.7	40.8	46.2
2000.....	3.7	*	*	*	*	*	1.0	5.0	13.4	26.8	38.3	49.3
1999.....	3.7	*	*	*	*	*	1.6	6.0	13.3	23.9	32.5	49.9

Non-Hispanic black, female

2007 5/.....	2.4	*	*	*	*	*	*	1.9	4.8	9.4	21.7	26.0
2006 5/.....	2.1	*	*	*	*	*	*	2.2	3.6	9.5	16.7	27.3
2005 5/.....	2.2	*	*	*	*	*	*	1.7	5.0	9.2	18.0	27.2
2004 5/.....	2.3	*	*	*	*	*	*	1.9	4.9	10.3	19.6	24.5
2003 5/.....	2.3	*	*	*	*	*	0.7	1.8	6.2	9.5	16.8	30.1
2002.....	2.3	*	*	*	*	*	*	2.5	4.4	11.3	20.7	23.8
2001.....	2.3	*	*	*	*	*	1.1	2.2	5.3	10.0	17.7	24.0
2000.....	2.2	*	*	*	*	*	0.8	2.2	5.4	9.7	18.9	22.2
1999.....	2.2	*	*	*	*	*	*	2.1	5.0	11.0	19.1	24.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

All origins 3/, both sexes

2007.....	4.6	*	*	*	*	*	0.2	1.3	4.8	14.6	38.5	68.6
2006.....	4.5	*	*	*	*	*	0.2	1.2	5.0	14.9	37.4	67.4
2005.....	4.5	*	*	*	*	*	0.3	1.2	4.9	14.9	37.2	68.3
2004.....	4.4	*	*	*	*	*	0.3	1.3	5.0	14.9	37.4	67.9
2003.....	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002.....	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001.....	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000.....	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999.....	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5

All origins 3/, male

2007.....	6.5	*	*	*	*	*	0.3	1.9	7.5	23.7	66.6	132.7
2006.....	6.4	*	*	*	*	*	0.3	1.8	7.8	23.7	65.5	131.6
2005.....	6.3	*	*	*	*	*	0.4	1.7	7.6	24.3	65.1	132.1
2004.....	6.2	*	*	*	*	*	0.3	2.0	7.9	24.0	64.7	133.0
2003.....	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002.....	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001.....	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000.....	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999.....	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8

All origins 3/, female

2007.....	2.7	*	*	*	*	*	0.2	0.7	2.3	6.8	19.1	38.1
2006.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	18.1	37.4
2005.....	2.7	*	*	*	*	*	0.2	0.8	2.3	6.9	18.3	39.0
2004.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	19.0	38.6
2003.....	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002.....	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001.....	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000.....	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999.....	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

Non-Hispanic 4/, both sexes

2007.....	5.2	*	*	*	*	*	0.2	1.4	5.0	15.1	39.5	70.5
2006.....	5.1	*	*	*	*	*	0.3	1.3	5.2	15.4	38.4	69.0
2005.....	5.0	*	*	*	*	*	0.3	1.3	5.1	15.4	38.2	69.6
2004.....	5.0	*	*	*	*	*	0.3	1.4	5.2	15.4	38.2	69.4
2003.....	4.8	*	*	*	*	*	0.2	1.3	5.2	15.4	37.1	65.6
2002.....	4.9	*	*	*	*	*	0.3	1.3	5.1	16.4	37.8	66.9
2001.....	4.8	*	*	*	*	*	0.3	1.3	5.2	15.8	36.4	69.6
2000.....	4.7	*	*	*	*	*	0.3	1.2	5.2	16.1	36.3	70.3
1999.....	4.7	*	*	*	*	*	0.2	1.3	5.4	16.3	36.0	69.3

Non-Hispanic 4/, male

2007.....	7.4	*	*	*	*	*	0.3	2.0	7.7	24.5	68.3	137.1
2006.....	7.3	*	*	*	*	*	0.4	2.0	8.2	24.7	67.4	135.5
2005.....	7.1	*	*	*	*	*	0.4	1.8	7.9	25.1	66.9	134.6
2004.....	7.1	*	*	*	*	*	0.4	2.1	8.1	24.9	66.6	136.5
2003.....	6.8	*	*	*	*	*	0.3	2.1	8.1	24.7	64.5	127.2
2002.....	6.8	*	*	*	*	*	0.4	1.9	7.9	26.0	64.7	129.8
2001.....	6.7	*	*	*	*	*	0.4	1.9	8.1	26.0	63.8	136.4
2000.....	6.6	*	*	*	*	*	0.3	1.8	8.1	26.1	63.6	141.3
1999.....	6.6	*	*	*	*	*	0.3	2.0	8.4	26.3	63.8	136.6

Non-Hispanic 4/, female

2007.....	3.1	*	*	*	*	*	0.2	0.8	2.4	7.1	19.7	39.1
2006.....	3.0	*	*	*	*	*	0.2	0.7	2.4	7.4	18.6	38.1
2005.....	3.0	*	*	*	*	*	0.2	0.8	2.5	7.2	18.7	40.0
2004.....	3.0	*	*	*	*	*	0.2	0.7	2.4	7.5	19.1	39.4
2003.....	2.9	*	*	*	*	*	0.2	0.6	2.5	7.6	18.8	38.6
2002.....	3.1	*	*	*	*	*	0.2	0.8	2.4	8.5	20.0	39.8
2001.....	3.0	*	*	*	*	*	0.3	0.6	2.5	7.4	18.3	41.9
2000.....	3.0	*	*	*	*	*	0.2	0.6	2.5	7.8	18.5	41.6
1999.....	3.0	*	*	*	*	*	0.2	0.6	2.6	8.1	17.9	42.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

Non-Hispanic white, both sexes

2007 5/.....	6.0	*	*	*	*	*	0.2	1.4	5.2	16.1	41.3	73.3
2006 5/.....	5.9	*	*	*	*	*	0.3	1.4	5.5	16.1	40.1	71.5
2005 5/.....	5.8	*	*	*	*	*	0.3	1.3	5.4	16.2	39.8	71.5
2004 5/.....	5.8	*	*	*	*	*	0.3	1.4	5.5	16.4	40.1	71.1
2003 5/.....	5.5	*	*	*	*	*	0.2	1.3	5.5	16.3	38.3	67.6
2002.....	5.6	*	*	*	*	*	0.3	1.4	5.3	17.1	39.1	68.0
2001.....	5.5	*	*	*	*	*	0.3	1.3	5.4	16.6	37.4	71.5
2000.....	5.4	*	*	*	*	*	0.3	1.3	5.5	16.8	37.1	72.8
1999.....	5.4	*	*	*	*	*	0.2	1.3	5.5	16.8	37.1	70.7

Non-Hispanic white, male

2007 5/.....	8.7	*	*	*	*	*	0.3	2.1	8.0	26.1	72.1	145.0
2006 5/.....	8.6	*	*	*	*	*	0.4	2.1	8.6	26.3	71.1	142.3
2005 5/.....	8.3	*	*	*	*	*	0.5	1.9	8.4	26.6	70.2	140.2
2004 5/.....	8.3	*	*	*	*	*	0.3	2.2	8.7	26.7	70.5	143.1
2003 5/.....	7.9	*	*	*	*	*	0.3	2.2	8.6	26.1	67.2	133.6
2002.....	7.9	*	*	*	*	*	0.4	2.0	8.5	27.4	67.2	135.2
2001.....	7.8	*	*	*	*	*	0.4	2.0	8.6	27.6	66.6	143.0
2000.....	7.7	*	*	*	*	*	0.3	2.0	8.6	27.5	65.9	148.0
1999.....	7.6	*	*	*	*	*	0.3	2.0	8.7	27.6	66.2	143.0

Non-Hispanic white, female

2007 5/.....	3.5	*	*	*	*	*	0.2	0.7	2.4	7.3	19.7	39.3
2006 5/.....	3.3	*	*	*	*	*	0.2	0.7	2.5	7.3	18.5	38.5
2005 5/.....	3.4	*	*	*	*	*	0.2	0.8	2.5	7.3	18.9	40.1
2004 5/.....	3.3	*	*	*	*	*	0.2	0.7	2.5	7.5	19.4	38.9
2003 5/.....	3.3	*	*	*	*	*	0.2	0.5	2.5	7.8	18.7	38.6
2002.....	3.4	*	*	*	*	*	0.2	0.8	2.3	8.4	20.2	39.0
2001.....	3.2	*	*	*	*	*	0.3	0.6	2.4	7.4	18.1	41.7
2000.....	3.3	*	*	*	*	*	0.2	0.7	2.6	7.8	18.2	42.5
1999.....	3.2	*	*	*	*	*	0.2	0.7	2.5	7.8	18.0	41.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

Non-Hispanic black, both sexes

2007 5/.....	2.7	*	*	*	*	*	*	1.6	5.0	12.1	32.9	55.2
2006 5/.....	2.7	*	*	*	*	*	*	1.3	4.9	13.6	31.7	55.7
2005 5/.....	2.5	*	*	*	*	*	*	1.5	4.7	12.5	29.8	57.6
2004 5/.....	2.4	*	*	*	*	*	0.5	1.6	4.2	11.9	27.6	61.3
2003 5/.....	2.5	*	*	*	*	*	0.4	1.7	4.1	12.3	32.8	53.8
2002.....	2.6	*	*	*	*	*	*	1.1	4.5	14.7	31.2	64.1
2001.....	2.4	*	*	*	*	*	0.4	1.3	5.1	12.1	31.4	55.6
2000.....	2.4	*	*	*	*	*	*	1.1	4.2	13.5	34.9	49.6
1999.....	2.6	*	*	*	*	*	*	1.5	5.6	15.2	29.4	60.8

Non-Hispanic black, male

2007 5/.....	3.1	*	*	*	*	*	*	1.9	7.4	18.4	46.2	81.1
2006 5/.....	3.1	*	*	*	*	*	*	1.9	7.7	18.2	42.8	95.3
2005 5/.....	3.0	*	*	*	*	*	*	1.7	6.6	18.3	45.7	101.8
2004 5/.....	2.7	*	*	*	*	*	*	2.3	6.0	15.0	40.2	83.3
2003 5/.....	2.9	*	*	*	*	*	*	2.2	5.5	17.8	49.6	79.9
2002.....	2.8	*	*	*	*	*	*	1.3	5.5	18.7	48.1	91.9
2001.....	2.6	*	*	*	*	*	*	1.8	6.5	16.5	43.2	78.5
2000.....	2.8	*	*	*	*	*	*	1.4	5.9	17.8	52.8	88.1
1999.....	3.0	*	*	*	*	*	*	2.6	7.4	19.2	46.6	86.8

Non-Hispanic black, female

2007 5/.....	2.3	*	*	*	*	*	*	1.4	3.1	7.7	25.3	44.5
2006 5/.....	2.3	*	*	*	*	*	*	0.7	2.7	10.4	25.4	39.6
2005 5/.....	2.1	*	*	*	*	*	*	1.3	3.1	8.4	20.8	39.8
2004 5/.....	2.2	*	*	*	*	*	*	1.0	2.7	9.7	20.4	52.6
2003 5/.....	2.1	*	*	*	*	*	*	1.2	2.9	8.3	23.3	43.7
2002.....	2.4	*	*	*	*	*	*	0.9	3.7	11.8	21.7	53.5
2001.....	2.3	*	*	*	*	*	*	*	3.9	9.0	24.8	47.1
2000.....	2.1	*	*	*	*	*	*	*	2.7	10.4	24.8	35.3
1999.....	2.3	*	*	*	*	*	*	*	4.2	12.3	19.8	51.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes

2007.....	4.4	*	0.7	0.8	0.5	0.9	2.0	4.4	9.1	15.5	20.0	16.1
2006.....	4.3	*	0.6	0.8	0.5	0.9	2.0	4.5	9.3	14.7	19.0	16.7
2005.....	4.4	*	0.5	0.8	0.6	0.9	2.2	4.6	9.4	15.8	20.2	15.8
2004.....	4.4	*	0.6	0.8	0.5	0.9	2.0	4.6	9.5	15.7	19.6	16.6
2003.....	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002.....	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001.....	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000.....	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999.....	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0

All origins 3/, male

2007.....	4.9	*	0.7	0.7	0.5	1.1	2.4	5.3	11.3	18.5	24.1	21.1
2006.....	4.9	*	0.7	0.8	0.5	1.0	2.4	5.5	11.4	18.2	24.0	21.2
2005.....	5.0	*	0.5	0.8	0.7	1.2	2.8	5.6	11.3	19.3	25.3	19.3
2004.....	4.9	*	0.7	0.8	0.6	1.1	2.5	5.8	11.8	18.4	23.7	19.4
2003.....	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002.....	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001.....	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000.....	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999.....	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7

All origins 3/, female

2007.....	3.9	*	0.6	0.8	0.4	0.7	1.5	3.4	7.0	13.0	17.2	13.8
2006.....	3.7	*	0.6	0.7	0.4	0.8	1.5	3.6	7.3	11.8	15.5	14.6
2005.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.6	7.5	12.9	16.7	14.3
2004.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.4	7.3	13.4	16.8	15.4
2003.....	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002.....	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001.....	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000.....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999.....	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Hispanic, both sexes

2007.....	1.8	*	0.5	0.7	0.6	0.5	1.2	3.0	5.8	8.9	14.3	11.1
2006.....	1.6	*	0.7	0.7	0.3	0.6	1.1	2.9	5.4	7.5	13.0	12.1
2005.....	1.8	*	0.6	0.7	0.5	0.5	1.3	2.9	5.9	10.2	13.4	15.4
2004.....	1.6	*	*	0.7	0.5	0.4	1.1	2.4	5.9	9.0	12.4	14.3
2003.....	1.7	*	0.6	0.8	0.3	0.6	1.1	2.5	5.9	9.5	15.3	*
2002.....	1.7	*	0.7	0.7	0.4	0.5	0.9	3.4	5.8	10.0	13.7	13.3
2001.....	1.7	*	0.7	0.5	0.4	0.6	1.4	2.6	5.9	10.3	14.8	*
2000.....	1.8	*	*	0.9	0.4	0.7	1.5	2.9	6.2	9.8	14.2	15.9
1999.....	1.7	*	0.9	0.7	*	0.5	1.3	3.0	6.0	10.0	14.0	*

Hispanic, male

2007.....	1.9	*	*	0.6	0.6	0.6	1.3	3.3	6.6	9.5	17.8	*
2006.....	1.7	*	*	0.8	*	0.5	1.5	3.3	6.0	9.2	16.1	*
2005.....	2.0	*	*	0.5	0.6	0.6	1.6	3.6	7.4	11.5	18.0	*
2004.....	1.7	*	*	0.6	0.6	*	1.4	2.9	7.2	10.1	15.4	*
2003.....	1.8	*	*	0.9	*	0.6	1.2	3.2	6.3	11.8	16.5	*
2002.....	1.8	*	*	0.7	*	*	1.0	4.4	6.6	12.4	16.0	*
2001.....	1.8	*	*	*	*	0.6	1.4	3.0	6.8	14.4	17.9	*
2000.....	1.9	*	*	1.2	*	0.8	1.5	2.6	7.7	12.5	18.7	*
1999.....	1.7	*	*	0.7	*	*	1.8	2.9	7.8	10.5	16.2	*

Hispanic, female

2007.....	1.7	*	*	0.7	*	*	1.0	2.7	4.9	8.5	11.9	11.0
2006.....	1.5	*	*	0.7	*	0.6	0.7	2.4	4.8	6.1	10.8	14.4
2005.....	1.6	*	*	0.9	*	*	1.0	2.2	4.6	9.2	10.2	15.0
2004.....	1.5	*	*	0.9	*	*	0.8	1.8	4.7	8.1	10.4	15.1
2003.....	1.6	*	*	0.7	*	*	0.9	1.9	5.6	7.6	14.5	*
2002.....	1.5	*	*	0.7	*	*	0.9	2.3	5.1	8.1	12.1	*
2001.....	1.6	*	*	0.6	*	*	1.4	2.2	5.1	7.1	12.7	*
2000.....	1.7	*	*	0.7	*	*	1.5	3.2	4.9	7.6	11.2	*
1999.....	1.7	*	*	0.8	*	*	0.8	3.1	4.5	9.6	12.5	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 4/, both sexes

2007.....	4.8	*	0.7	0.8	0.5	1.0	2.1	4.5	9.4	16.0	20.4	16.4
2006.....	4.8	*	0.6	0.8	0.5	1.0	2.1	4.7	9.6	15.2	19.3	16.9
2005.....	4.9	*	0.5	0.8	0.6	1.1	2.3	4.8	9.6	16.2	20.6	15.9
2004.....	4.8	*	0.7	0.8	0.5	1.0	2.2	4.8	9.7	16.2	20.0	16.7
2003.....	4.9	*	0.7	0.8	0.6	1.1	2.6	5.0	10.0	15.9	19.7	15.4
2002.....	4.9	*	0.7	0.8	0.5	1.1	2.4	5.3	9.9	16.6	19.4	16.5
2001.....	4.8	*	0.6	0.8	0.6	1.2	2.4	5.2	10.0	15.7	20.6	16.5
2000.....	4.9	*	0.7	0.7	0.6	1.1	2.5	5.3	9.8	16.9	20.2	16.9
1999.....	5.0	*	0.7	0.8	0.6	1.2	2.5	5.3	10.6	17.3	20.6	16.1

Non-Hispanic 4/, male

2007.....	5.5	*	0.7	0.7	0.5	1.3	2.7	5.6	11.7	19.2	24.5	21.7
2006.....	5.4	*	0.6	0.8	0.6	1.2	2.6	5.7	11.9	18.8	24.4	21.9
2005.....	5.5	*	0.5	0.9	0.7	1.4	3.0	5.8	11.6	19.8	25.7	19.4
2004.....	5.4	*	0.7	0.9	0.6	1.2	2.7	6.1	12.2	19.0	24.2	19.7
2003.....	5.5	*	0.7	0.8	0.7	1.3	3.1	6.5	11.8	19.1	25.0	20.1
2002.....	5.5	*	0.8	0.9	0.7	1.3	3.0	6.4	12.0	20.7	24.1	20.2
2001.....	5.5	*	0.6	0.7	0.7	1.5	3.0	6.4	12.5	19.4	26.1	20.5
2000.....	5.5	*	0.7	0.8	0.7	1.4	3.3	6.5	12.1	20.6	24.5	22.3
1999.....	5.6	*	0.7	0.8	0.7	1.5	3.0	6.5	12.7	21.4	25.8	21.3

Non-Hispanic 4/, female

2007.....	4.2	*	0.7	0.8	0.4	0.8	1.6	3.5	7.2	13.3	17.5	13.9
2006.....	4.1	*	0.6	0.7	0.5	0.9	1.6	3.7	7.5	12.2	15.8	14.6
2005.....	4.2	*	0.5	0.8	0.4	0.7	1.7	3.8	7.8	13.1	17.0	14.2
2004.....	4.2	*	0.6	0.7	0.4	0.8	1.7	3.5	7.5	13.8	17.2	15.4
2003.....	4.3	*	0.6	0.8	0.5	1.0	2.1	3.7	8.3	13.2	16.2	13.4
2002.....	4.2	*	0.7	0.7	0.4	0.9	1.8	4.2	7.9	13.1	16.2	14.9
2001.....	4.2	*	0.6	0.8	0.5	0.8	1.8	4.1	7.7	12.7	17.0	14.9
2000.....	4.2	*	0.7	0.7	0.5	0.7	1.8	4.1	7.8	13.8	17.5	14.7
1999.....	4.4	*	0.7	0.8	0.5	0.9	2.0	4.2	8.6	14.0	17.2	14.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic white, both sexes

2007 5/.....	5.7	*	0.8	0.8	0.5	1.1	2.4	5.0	10.3	17.4	22.0	17.2
2006 5/.....	5.5	*	0.6	0.8	0.5	1.2	2.4	5.3	10.5	16.6	20.7	17.7
2005 5/.....	5.6	*	0.5	0.9	0.6	1.2	2.6	5.3	10.5	17.6	21.8	16.5
2004 5/.....	5.6	*	0.7	0.8	0.6	1.1	2.5	5.3	10.7	17.6	21.3	17.4
2003 5/.....	5.6	*	0.7	0.8	0.7	1.3	3.0	5.5	10.8	17.1	20.9	15.9
2002.....	5.6	*	0.6	0.8	0.6	1.2	2.8	5.9	10.7	17.9	20.4	17.3
2001.....	5.6	*	0.7	0.8	0.6	1.3	2.7	5.7	10.8	16.8	21.9	17.2
2000.....	5.6	*	0.7	0.8	0.6	1.2	2.8	5.9	10.7	18.3	21.2	17.5
1999.....	5.7	*	0.7	0.8	0.6	1.4	2.8	5.8	11.5	18.7	21.7	16.7

Non-Hispanic white, male

2007 5/.....	6.4	*	0.8	0.8	0.6	1.4	3.0	6.1	12.7	20.8	26.5	23.2
2006 5/.....	6.3	*	0.7	0.9	0.6	1.4	3.0	6.3	12.9	20.3	26.0	22.9
2005 5/.....	6.4	*	0.4	0.9	0.7	1.5	3.4	6.3	12.5	21.4	26.9	20.4
2004 5/.....	6.3	*	0.8	0.9	0.7	1.4	3.0	6.6	13.2	20.8	25.4	21.0
2003 5/.....	6.3	*	0.8	0.8	0.7	1.5	3.5	7.0	12.8	20.4	26.3	21.0
2002.....	6.4	*	0.7	0.9	0.8	1.4	3.5	7.1	12.8	22.3	25.2	21.4
2001.....	6.3	*	0.7	0.7	0.7	1.7	3.5	7.0	13.4	20.9	27.5	20.9
2000.....	6.3	*	0.7	0.8	0.7	1.5	3.7	7.1	13.1	22.2	25.7	23.5
1999.....	6.4	*	0.8	0.8	0.7	1.8	3.3	7.1	13.7	23.0	27.2	22.4

Non-Hispanic white, female

2007 5/.....	4.9	*	0.8	0.9	0.4	0.9	1.8	3.8	8.0	14.5	18.9	14.4
2006 5/.....	4.8	*	0.5	0.7	0.5	0.9	1.8	4.2	8.3	13.3	17.1	15.3
2005 5/.....	4.9	*	0.6	0.8	0.5	0.9	1.9	4.2	8.6	14.3	18.2	14.8
2004 5/.....	4.9	*	0.7	0.7	0.4	0.9	1.9	3.9	8.3	14.9	18.5	15.8
2003 5/.....	4.9	*	0.6	0.9	0.6	1.1	2.5	4.1	9.0	14.3	17.3	13.7
2002.....	4.9	*	0.6	0.8	0.4	1.1	2.1	4.7	8.7	14.2	17.1	15.5
2001.....	4.8	*	0.6	0.9	0.5	0.9	2.0	4.5	8.4	13.4	18.2	15.7
2000.....	4.9	*	0.7	0.7	0.5	0.9	1.9	4.6	8.4	15.0	18.3	15.1
1999.....	5.0	*	0.7	0.7	0.5	1.1	2.3	4.6	9.5	15.1	18.1	14.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic black, both sexes

2007 5/.....	2.0	*	*	0.6	0.3	0.6	1.0	3.0	5.3	9.6	8.8	8.9
2006 5/.....	2.0	*	*	0.7	0.5	0.5	1.2	2.6	5.4	8.2	9.6	9.6
2005 5/.....	2.1	*	*	0.8	0.4	0.5	1.2	2.8	5.5	8.5	12.3	9.6
2004 5/.....	2.0	*	*	0.8	0.5	0.7	1.3	2.9	5.0	7.9	10.1	10.6
2003 5/.....	2.2	*	*	0.7	0.6	0.6	1.5	3.1	5.6	9.0	10.6	10.7
2002.....	2.1	*	1.1	0.8	0.4	0.6	1.2	2.8	5.6	8.6	11.9	9.7
2001.....	2.1	*	*	0.7	0.5	0.5	1.1	3.1	5.9	9.4	9.9	11.9
2000.....	2.2	*	*	0.8	0.6	0.7	1.6	2.9	5.8	8.2	12.1	10.5
1999.....	2.2	*	*	0.8	0.7	0.7	1.4	3.3	6.1	8.2	11.7	10.0

Non-Hispanic black, male

2007 5/.....	2.2	*	*	*	*	0.8	1.4	3.6	7.1	11.4	9.6	*
2006 5/.....	2.3	*	*	0.6	0.6	*	1.5	3.2	7.2	10.4	12.5	*
2005 5/.....	2.4	*	*	0.8	*	0.8	1.4	3.7	7.2	10.6	17.9	*
2004 5/.....	2.3	*	*	0.7	*	*	1.9	4.0	6.9	8.1	14.5	*
2003 5/.....	2.5	*	*	0.8	0.7	*	2.1	4.1	6.8	11.9	15.6	*
2002.....	2.3	*	*	1.1	*	0.9	1.3	2.7	8.1	10.2	16.1	*
2001.....	2.2	*	*	0.8	*	*	1.3	3.7	7.9	10.4	12.0	*
2000.....	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.2	9.9	13.3	*
1999.....	2.4	*	*	0.7	0.9	0.9	1.8	4.5	7.4	10.2	12.5	*

Non-Hispanic black, female

2007 5/.....	1.9	*	*	0.8	*	*	*	2.4	3.9	8.3	8.3	9.0
2006 5/.....	1.8	*	*	0.8	*	*	0.9	2.0	4.0	6.6	7.9	9.0
2005 5/.....	1.8	*	*	0.7	*	*	1.0	2.0	4.2	7.0	9.1	10.0
2004 5/.....	1.8	*	*	0.8	*	*	0.8	1.9	3.5	7.8	7.6	11.6
2003 5/.....	1.9	*	*	0.7	*	*	0.9	2.3	4.7	6.9	7.8	12.0
2002.....	1.9	*	*	*	*	*	1.1	2.8	3.7	7.5	9.5	10.5
2001.....	1.9	*	*	*	*	*	1.0	2.5	4.2	8.7	8.8	9.8
2000.....	1.9	*	*	0.7	*	*	1.2	2.2	4.7	7.1	11.5	10.5
1999.....	1.9	*	*	0.9	*	*	1.0	2.2	5.1	6.8	11.2	9.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes

2007.....	18.2	0.6	0.7	0.9	1.5	1.9	3.4	8.2	23.5	63.2	137.3	185.5
2006.....	18.4	0.8	0.8	0.7	1.6	2.0	3.5	8.5	24.4	65.2	137.8	186.1
2005.....	18.6	0.6	0.8	0.8	1.6	2.0	3.4	8.7	25.1	67.0	138.7	189.8
2004.....	18.6	0.6	0.8	0.9	1.5	2.1	3.4	9.0	25.8	68.2	139.8	189.4
2003.....	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002.....	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001.....	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000.....	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999.....	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4

All origins 3/, male

2007.....	20.3	*	0.7	1.0	1.7	2.2	4.1	10.2	29.4	80.7	182.2	253.4
2006.....	20.3	*	0.9	0.9	2.0	2.4	4.1	10.5	30.8	82.4	183.1	244.7
2005.....	20.6	*	1.0	1.0	1.9	2.3	4.0	10.4	31.1	86.6	183.0	262.4
2004.....	20.4	*	0.9	1.0	1.9	2.4	4.2	11.0	32.1	87.4	181.3	254.9
2003.....	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002.....	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001.....	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000.....	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999.....	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0

All origins 3/, female

2007.....	16.2	*	0.7	0.7	1.2	1.6	2.7	6.3	18.1	48.2	106.4	153.3
2006.....	16.5	*	0.8	0.6	1.1	1.7	2.9	6.6	18.3	50.7	106.8	158.7
2005.....	16.6	*	0.6	0.7	1.2	1.8	2.8	7.0	19.6	50.5	108.6	156.4
2004.....	16.8	*	0.6	0.7	1.2	1.7	2.7	7.0	19.9	52.0	111.8	160.0
2003.....	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002.....	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001.....	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000.....	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999.....	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Hispanic, both sexes

2007.....	7.2	*	0.5	1.3	2.0	2.0	3.3	7.3	20.9	48.9	92.3	95.1
2006.....	7.0	*	0.7	0.8	2.0	2.0	3.1	7.1	18.6	51.3	93.6	100.9
2005.....	7.3	*	0.8	1.1	2.2	1.9	2.8	8.1	20.9	52.7	97.6	107.8
2004.....	6.9	*	0.8	1.4	2.0	2.0	3.2	7.7	19.8	48.3	95.4	103.2
2003.....	7.1	*	1.3	1.3	1.9	2.0	3.4	8.5	20.4	47.1	99.1	129.6
2002.....	7.3	*	0.9	1.3	2.1	1.9	3.9	7.7	22.1	53.7	100.8	132.5
2001.....	7.4	*	1.3	1.2	2.2	2.2	3.9	8.2	23.9	52.6	103.3	134.4
2000.....	7.5	*	1.0	1.1	2.0	1.9	3.9	8.9	25.0	58.8	103.3	143.3
1999.....	7.5	*	1.0	1.2	2.0	2.1	3.9	10.1	24.2	58.5	101.0	131.9

Hispanic, male

2007.....	7.5	*	*	1.3	2.2	2.0	3.6	8.7	22.9	58.9	110.9	133.2
2006.....	7.4	*	*	0.8	2.5	2.1	3.5	8.3	22.4	60.5	115.7	129.7
2005.....	7.8	*	1.1	1.3	2.7	2.2	3.3	9.0	23.7	64.7	124.9	144.5
2004.....	7.6	*	*	1.8	2.6	2.2	4.1	9.9	23.9	60.8	117.0	119.1
2003.....	7.4	*	1.5	1.5	2.4	2.4	4.2	9.5	24.3	57.4	114.7	149.3
2002.....	7.6	*	*	1.3	2.7	2.3	4.4	8.9	26.2	63.3	119.0	180.7
2001.....	8.3	*	1.4	1.3	2.7	2.7	5.0	9.7	28.9	67.5	142.2	180.9
2000.....	8.0	*	*	1.4	2.3	2.4	4.3	11.1	30.7	71.3	126.9	197.5
1999.....	7.6	*	*	1.3	2.1	2.2	4.2	12.0	29.4	68.5	117.8	135.9

Hispanic, female

2007.....	7.0	*	*	1.3	1.8	2.1	2.9	5.9	19.0	40.6	79.2	74.5
2006.....	6.6	*	*	0.8	1.4	1.8	2.8	6.0	15.1	43.9	78.1	85.7
2005.....	6.7	*	*	0.9	1.5	1.6	2.4	7.0	18.4	43.0	78.6	88.6
2004.....	6.2	*	*	1.1	1.2	1.6	2.2	5.6	16.1	38.2	80.5	95.1
2003.....	6.8	*	*	1.1	1.3	1.5	2.6	7.4	16.8	39.0	88.5	119.5
2002.....	7.0	*	*	1.3	1.5	1.4	3.3	6.5	18.5	46.1	88.3	108.6
2001.....	6.5	*	*	1.1	1.5	1.7	2.6	6.7	19.5	40.8	77.1	111.7
2000.....	6.9	*	*	0.8	1.5	1.4	3.4	6.8	19.8	48.9	87.5	116.7
1999.....	7.4	*	*	1.0	1.9	2.1	3.4	8.3	19.6	50.6	89.9	130.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 4/, both sexes

2007.....	20.2	0.7	0.7	0.8	1.4	1.9	3.4	8.3	23.7	64.2	140.1	190.2
2006.....	20.3	0.8	0.8	0.7	1.5	2.1	3.6	8.7	24.8	66.2	140.4	190.1
2005.....	20.4	*	0.8	0.8	1.4	2.0	3.5	8.7	25.4	68.0	141.0	193.6
2004.....	20.5	*	0.8	0.7	1.4	2.1	3.5	9.1	26.2	69.6	142.1	193.0
2003.....	21.0	0.8	0.8	0.9	1.4	2.2	4.0	9.3	27.2	71.8	145.4	200.5
2002.....	21.3	0.8	0.7	0.8	1.6	2.4	3.9	10.1	28.6	75.1	145.3	199.8
2001.....	21.6	0.6	0.9	0.8	1.7	2.5	4.1	10.5	29.8	75.8	146.8	202.1
2000.....	21.8	1.0	0.9	0.9	1.8	2.4	4.1	10.8	30.3	77.3	150.9	206.5
1999.....	21.9	0.7	0.9	0.8	1.8	2.4	4.3	10.8	32.3	78.8	149.5	201.4

Non-Hispanic 4/, male

2007.....	22.7	*	0.7	0.9	1.6	2.3	4.2	10.3	29.9	82.4	186.8	260.3
2006.....	22.7	*	0.9	0.9	1.9	2.5	4.3	10.7	31.5	83.9	187.1	250.7
2005.....	22.8	*	1.0	0.9	1.7	2.3	4.1	10.5	31.6	88.2	186.3	268.5
2004.....	22.6	*	0.9	0.8	1.7	2.4	4.2	11.1	32.7	89.3	184.8	261.1
2003.....	23.3	*	0.8	1.0	1.6	2.7	4.8	11.6	34.6	92.6	188.2	273.6
2002.....	23.6	*	0.8	1.0	1.8	2.8	4.7	12.1	35.6	97.0	190.9	272.8
2001.....	23.5	*	1.0	1.0	1.9	2.9	5.0	12.9	36.8	96.3	188.6	278.8
2000.....	23.7	*	1.0	1.0	2.2	2.7	4.8	13.1	36.4	98.1	195.2	285.0
1999.....	23.7	*	0.9	0.9	2.2	2.9	5.2	13.3	39.5	98.7	191.5	284.3

Non-Hispanic 4/, female

2007.....	17.7	*	0.7	0.6	1.1	1.4	2.6	6.4	18.0	48.8	108.0	157.1
2006.....	18.1	*	0.8	0.6	1.0	1.6	2.9	6.7	18.6	51.2	108.5	162.0
2005.....	18.2	*	0.6	0.6	1.1	1.8	2.9	7.0	19.6	51.0	110.3	159.5
2004.....	18.5	*	0.6	0.6	1.2	1.7	2.8	7.2	20.1	52.9	113.3	162.6
2003.....	18.9	*	0.7	0.8	1.1	1.7	3.2	7.1	20.4	54.3	116.8	168.4
2002.....	19.2	*	0.7	0.7	1.4	2.0	3.1	8.0	22.2	56.8	115.1	168.3
2001.....	19.7	*	0.9	0.7	1.4	2.2	3.3	8.2	23.3	58.9	119.3	170.2
2000.....	20.1	*	0.9	0.7	1.3	2.0	3.4	8.6	24.7	60.1	122.1	174.8
1999.....	20.1	*	0.8	0.7	1.4	2.0	3.4	8.5	25.6	62.5	122.3	168.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic white, both sexes

2007 5/.....	22.4	*	0.7	0.8	1.3	1.8	3.2	7.9	23.3	65.4	144.7	198.0
2006 5/.....	22.6	*	0.9	0.7	1.4	2.0	3.4	8.4	24.5	67.6	145.3	196.6
2005 5/.....	22.8	*	0.7	0.8	1.4	1.8	3.4	8.3	25.3	69.4	146.2	201.5
2004 5/.....	22.7	*	0.8	0.7	1.4	2.0	3.4	8.8	26.1	70.4	145.9	198.8
2003 5/.....	23.4	1.0	0.8	0.9	1.4	2.1	3.8	9.0	27.4	73.2	149.3	207.2
2002.....	23.7	*	0.8	0.8	1.7	2.3	3.8	9.7	28.4	76.7	149.5	206.5
2001.....	23.8	*	1.1	0.8	1.7	2.4	4.0	10.3	29.5	77.2	150.5	208.2
2000.....	24.2	1.3	1.0	0.9	1.7	2.3	4.0	10.5	30.3	79.0	155.1	210.7
1999.....	24.1	*	1.0	0.8	1.8	2.4	4.2	10.6	32.0	80.4	153.0	206.4

Non-Hispanic white, male

2007 5/.....	25.3	*	0.6	0.9	1.6	2.3	4.0	9.9	29.7	84.1	193.2	271.4
2006 5/.....	25.2	*	0.9	0.8	1.9	2.5	4.0	10.3	31.0	85.5	193.4	258.6
2005 5/.....	25.4	*	0.8	0.9	1.7	2.0	4.0	9.9	31.5	90.1	192.5	281.9
2004 5/.....	25.1	*	0.9	0.9	1.6	2.4	4.1	10.7	32.8	90.3	190.1	270.9
2003 5/.....	26.0	*	0.9	1.0	1.6	2.6	4.7	11.2	34.9	95.2	194.8	284.6
2002.....	26.2	*	0.9	1.0	1.9	2.7	4.5	11.8	35.4	99.3	196.8	283.5
2001.....	25.9	*	1.0	0.8	2.0	2.8	4.7	12.8	36.4	97.9	193.6	288.7
2000.....	26.2	*	1.1	1.0	2.1	2.6	4.8	12.7	36.3	100.1	201.0	293.8
1999.....	26.1	*	1.0	0.9	2.2	2.7	5.0	13.1	39.2	100.7	196.1	291.8

Non-Hispanic white, female

2007 5/.....	19.6	*	0.8	0.6	1.0	1.3	2.4	6.0	17.2	49.0	110.8	163.2
2006 5/.....	20.2	*	0.9	0.6	1.0	1.6	2.8	6.5	18.2	52.1	111.9	167.7
2005 5/.....	20.2	*	0.6	0.6	1.1	1.6	2.7	6.6	19.4	51.4	114.3	164.7
2004 5/.....	20.4	*	0.6	0.6	1.2	1.5	2.6	7.0	19.8	53.4	115.8	166.5
2003 5/.....	20.8	*	0.7	0.8	1.1	1.6	3.0	6.8	20.3	54.3	118.5	173.1
2002.....	21.2	*	0.7	0.7	1.4	1.9	3.1	7.6	21.7	57.5	117.8	173.2
2001.....	21.8	*	1.1	0.7	1.4	2.0	3.2	7.9	23.0	59.5	122.0	174.6
2000.....	22.3	*	0.9	0.8	1.2	1.9	3.2	8.4	24.6	61.2	125.0	177.2
1999.....	22.2	*	0.9	0.6	1.3	2.0	3.3	8.3	25.2	63.4	124.7	172.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic black, both sexes

2007 5/.....	13.6	*	0.8	0.8	1.5	2.4	5.3	12.0	31.3	69.3	117.7	140.4
2006 5/.....	13.8	*	*	0.8	1.6	2.5	5.3	12.4	33.5	69.9	118.7	145.5
2005 5/.....	13.4	*	1.0	0.7	1.6	3.4	4.7	12.6	31.0	69.8	114.7	139.3
2004 5/.....	14.0	*	*	0.7	1.7	3.1	4.6	12.8	32.9	76.0	125.2	156.3
2003 5/.....	13.9	*	*	0.9	1.5	3.0	5.4	13.4	31.8	74.3	127.4	152.3
2002.....	14.1	*	*	0.8	1.5	3.0	4.8	13.9	36.4	77.0	123.5	153.7
2001.....	14.5	*	*	1.0	1.8	3.5	5.8	13.5	37.1	79.0	127.6	157.3
2000.....	14.3	*	*	0.7	2.3	3.0	5.4	15.0	36.5	76.2	122.3	177.1
1999.....	14.5	*	*	0.9	2.4	3.0	5.6	14.1	39.5	77.5	128.2	162.3

Non-Hispanic black, male

2007 5/.....	14.5	*	*	1.0	1.7	2.8	6.5	15.0	37.5	86.1	150.5	185.0
2006 5/.....	15.5	*	*	1.0	2.0	2.9	6.7	15.8	44.0	92.6	156.0	198.8
2005 5/.....	14.6	*	*	1.0	1.9	3.9	5.5	16.0	39.3	90.0	154.5	168.4
2004 5/.....	14.8	*	*	0.7	2.0	3.4	5.4	16.0	40.3	99.5	152.9	197.1
2003 5/.....	14.5	*	*	0.9	1.7	3.5	6.6	17.1	41.1	88.6	150.3	188.6
2002.....	15.1	*	*	1.0	1.5	3.8	6.2	16.5	45.2	99.4	155.3	203.5
2001.....	15.6	*	*	1.3	2.0	4.0	7.6	16.3	46.6	100.7	161.4	206.7
2000.....	15.2	*	*	0.8	2.8	3.7	5.7	18.2	45.3	97.8	157.4	226.6
1999.....	15.6	*	*	1.2	2.6	3.9	7.0	17.5	48.4	96.9	162.9	221.1

Non-Hispanic black, female

2007 5/.....	12.8	*	*	*	1.3	2.1	4.3	9.4	26.3	57.3	99.0	121.9
2006 5/.....	12.3	*	*	*	1.3	2.1	3.9	9.5	25.1	53.6	97.5	123.7
2005 5/.....	12.2	*	*	*	1.3	2.9	4.0	9.7	24.4	55.4	92.2	127.5
2004 5/.....	13.2	*	*	0.7	1.4	2.8	3.9	10.0	27.0	59.2	109.6	140.1
2003 5/.....	13.3	*	*	0.8	1.2	2.5	4.4	10.2	24.3	64.1	114.4	138.3
2002.....	13.1	*	*	*	1.5	2.3	3.5	11.6	29.4	61.2	105.4	134.6
2001.....	13.5	*	*	*	1.7	3.1	4.2	11.1	29.5	63.7	108.7	139.0
2000.....	13.5	*	*	*	1.8	2.5	5.2	12.2	29.4	61.0	102.6	158.7
1999.....	13.6	*	*	0.7	2.1	2.2	4.3	11.1	32.3	63.8	108.7	140.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All origins 3/, both sexes												
2007.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	1.1	2.2	2.4
2006.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.6	1.1	2.2	2.6
2005.....	0.4	*	*	*	0.1	0.3	0.3	0.4	0.6	1.1	2.3	2.1
2004.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	1.9	2.2
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All origins 3/, male												
2007.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.6	2.8
2006.....	0.5	*	*	*	0.2	0.4	0.5	0.5	0.8	1.4	2.6	3.7
2005.....	0.5	*	*	*	0.1	0.3	0.3	0.4	0.8	1.4	2.5	3.1
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	0.9	1.6	2.5	3.2
2003.....	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All origins 3/, female												
2007.....	0.4	*	*	*	0.1	0.2	0.2	0.2	0.5	0.9	1.9	2.2
2006.....	0.4	*	*	*	0.1	0.3	0.2	0.2	0.4	0.8	2.0	2.1
2005.....	0.4	*	*	*	0.2	0.3	0.2	0.3	0.4	0.9	2.2	1.6
2004.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.5	1.7
2003.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic 4/, both sexes												
2007.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	1.0	2.2	2.4
2006.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.1	2.2	2.6
2005.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.6	1.0	2.3	2.0
2004.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	1.9	2.1
2003.....	0.5	*	*	*	0.2	0.5	0.4	0.4	0.7	1.2	2.2	2.3
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.8
2001.....	0.5	*	*	*	0.2	0.5	0.4	0.5	0.5	1.1	2.2	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.4
1999.....	0.5	*	*	*	0.3	0.4	0.4	0.4	0.7	1.5	2.1	2.3
Non-Hispanic 4/, male												
2007.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.7	2.7
2006.....	0.6	*	*	*	0.2	0.5	0.5	0.5	0.8	1.3	2.5	3.8
2005.....	0.5	*	*	*	0.1	0.4	0.3	0.4	0.8	1.3	2.4	3.0
2004.....	0.5	*	*	*	0.2	0.4	0.3	0.5	1.0	1.6	2.5	3.1
2003.....	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.5	2.7
2002.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.4	3.7
2001.....	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.5	2.6	2.5
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.3	3.7
1999.....	0.6	*	*	*	0.3	0.5	0.5	0.5	1.0	1.9	2.3	3.7
Non-Hispanic 4/, female												
2007.....	0.4	*	*	*	0.1	0.2	0.2	0.2	0.4	0.9	1.9	2.2
2006.....	0.4	*	*	*	0.1	0.3	0.3	0.2	0.4	0.8	2.0	2.1
2005.....	0.4	*	*	*	0.2	0.3	0.2	0.4	0.4	0.8	2.2	1.6
2004.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.5	1.6
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	0.9	1.9	2.2
2002.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	1.5	2.4
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	2.0	2.0
2000.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.7	2.0
1999.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic white, both sexes												
2007 5/.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.5	1.1	2.4	2.5
2006 5/.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.1	2.3	2.7
2005 5/.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.6	1.1	2.4	2.1
2004 5/.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.0	2.1
2003 5/.....	0.6	*	*	*	0.2	0.5	0.4	0.4	0.7	1.2	2.3	2.5
2002.....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.6	1.4	2.0	2.9
2001.....	0.5	*	*	*	0.2	0.5	0.4	0.5	0.6	1.2	2.3	2.3
2000.....	0.5	*	*	*	0.2	0.5	0.4	0.4	0.6	1.3	2.1	2.6
1999.....	0.6	*	*	*	0.3	0.4	0.4	0.5	0.8	1.5	2.2	2.3
Non-Hispanic white, male												
2007 5/.....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.6	1.3	2.9	2.9
2006 5/.....	0.6	*	*	*	0.2	0.5	0.5	0.5	0.9	1.4	2.6	3.9
2005 5/.....	0.5	*	*	*	0.1	0.3	0.3	0.4	0.8	1.3	2.5	3.2
2004 5/.....	0.6	*	*	*	0.2	0.4	0.3	0.5	1.0	1.6	2.6	3.3
2003 5/.....	0.6	*	*	*	*	0.5	0.5	0.5	0.9	1.5	2.6	2.7
2002.....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.8	1.9	2.6	4.0
2001.....	0.6	*	*	*	0.3	0.5	0.5	0.6	0.7	1.6	2.7	2.6
2000.....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.7	1.6	2.5	3.9
1999.....	0.7	*	*	*	0.3	0.5	0.5	0.5	1.1	2.0	2.5	3.8
Non-Hispanic white, female												
2007 5/.....	0.4	*	*	*	*	0.2	0.2	0.2	0.4	0.9	2.1	2.4
2006 5/.....	0.4	*	*	*	*	0.3	0.3	0.3	0.4	0.9	2.1	2.1
2005 5/.....	0.5	*	*	*	0.2	0.2	0.2	0.4	0.5	0.9	2.4	1.5
2004 5/.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	0.9	1.6	1.6
2003 5/.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	2.0	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.6	2.5
2001.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	2.0	2.3
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.6	1.1	1.9	2.1
1999.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	1.2	2.0	1.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's Lymphoma (C82-C85)

All origins 3/, both sexes

2007.....	6.8	*	*	0.1	0.3	0.5	1.2	3.2	8.9	23.1	52.7	72.1
2006.....	6.9	*	*	0.1	0.3	0.5	1.2	3.3	9.1	24.0	53.7	72.3
2005.....	7.0	*	*	0.1	0.3	0.6	1.3	3.3	9.6	24.8	54.5	75.2
2004.....	7.1	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.1	54.6	75.0
2003.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002.....	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001.....	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000.....	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999.....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7

All origins 3/, male

2007.....	7.4	*	*	0.1	0.4	0.6	1.6	4.1	11.3	29.6	65.7	91.6
2006.....	7.4	*	*	0.1	0.4	0.7	1.5	4.3	11.8	29.7	66.9	89.6
2005.....	7.6	*	*	0.2	0.4	0.7	1.7	4.4	11.9	31.4	67.4	100.5
2004.....	7.6	*	*	0.1	0.4	0.8	1.7	4.8	12.3	32.5	67.3	95.8
2003.....	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002.....	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001.....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000.....	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999.....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4

All origins 3/, female

2007.....	6.2	*	*	*	0.2	0.4	0.8	2.3	6.7	17.6	43.8	62.8
2006.....	6.4	*	*	*	0.2	0.4	0.9	2.3	6.7	19.2	44.7	64.2
2005.....	6.5	*	*	*	0.2	0.4	0.9	2.3	7.5	19.3	45.7	63.5
2004.....	6.6	*	*	*	0.3	0.5	0.8	2.6	7.6	20.7	46.0	65.7
2003.....	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002.....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001.....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000.....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999.....	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's Lymphoma (C82-C85)												
Hispanic, both sexes												
2007.....	2.7	*	*	*	0.4	0.5	1.1	2.9	8.1	18.2	40.0	47.6
2006.....	2.6	*	*	*	0.4	0.5	0.9	2.7	8.2	18.9	40.1	40.3
2005.....	2.6	*	*	*	0.3	0.4	1.0	2.9	7.7	19.5	44.7	44.5
2004.....	2.6	*	*	*	0.4	0.6	1.2	3.3	7.5	18.7	40.6	43.8
2003.....	2.6	*	*	*	0.4	0.6	1.4	3.0	8.4	19.1	40.4	54.4
2002.....	2.8	*	*	*	0.4	0.6	1.3	3.0	9.4	22.7	43.7	50.4
2001.....	2.7	*	*	*	0.3	0.5	1.4	3.6	10.9	18.3	43.3	59.4
2000.....	2.9	*	*	*	0.3	0.6	1.6	3.6	11.1	25.7	41.5	59.7
1999.....	2.8	*	*	*	*	0.6	1.4	4.4	9.3	25.2	42.7	56.1
Hispanic, male												
2007.....	2.8	*	*	*	*	*	1.3	3.7	8.8	23.3	49.0	52.9
2006.....	2.7	*	*	*	0.6	0.5	1.2	3.1	10.2	22.1	47.7	54.1
2005.....	2.7	*	*	*	*	0.5	1.2	3.2	8.9	23.9	52.8	52.3
2004.....	2.8	*	*	*	*	0.7	1.8	4.9	8.8	24.0	47.2	50.4
2003.....	2.8	*	*	*	*	0.8	1.8	3.4	10.5	23.8	45.3	57.0
2002.....	2.7	*	*	*	*	0.6	1.5	3.6	11.7	26.0	45.5	60.2
2001.....	3.0	*	*	*	*	0.6	2.0	4.4	13.2	24.4	58.6	68.6
2000.....	3.1	*	*	*	*	0.7	1.9	4.6	13.9	30.2	48.2	82.6
1999.....	2.8	*	*	*	*	*	1.7	5.7	11.8	27.5	48.0	60.4
Hispanic, female												
2007.....	2.6	*	*	*	*	*	1.0	2.1	7.5	14.0	33.8	44.7
2006.....	2.4	*	*	*	*	*	0.6	2.2	6.4	16.3	34.7	33.0
2005.....	2.5	*	*	*	*	*	0.8	2.5	6.5	15.9	39.1	40.4
2004.....	2.3	*	*	*	*	*	*	1.6	6.4	14.5	36.1	40.3
2003.....	2.5	*	*	*	*	*	0.9	2.7	6.6	15.3	37.1	53.1
2002.....	2.8	*	*	*	*	*	1.2	2.3	7.4	20.1	42.4	45.6
2001.....	2.4	*	*	*	*	*	0.8	2.8	8.8	13.5	32.9	54.9
2000.....	2.7	*	*	*	*	*	1.3	2.6	8.6	22.2	37.0	48.5
1999.....	2.8	*	*	*	*	*	1.0	3.1	7.1	23.4	39.2	54.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's Lymphoma (C82-C85)												
Non-Hispanic 4/, both sexes												
2007.....	7.5	*	*	0.1	0.3	0.5	1.2	3.2	9.0	23.5	53.5	73.4
2006.....	7.6	*	*	0.1	0.3	0.6	1.2	3.3	9.2	24.4	54.5	73.9
2005.....	7.8	*	*	0.1	0.3	0.6	1.3	3.4	9.8	25.2	55.0	76.6
2004.....	7.9	*	*	0.1	0.3	0.6	1.2	3.7	10.0	26.6	55.3	76.3
2003.....	8.1	*	*	0.1	0.3	0.6	1.4	3.9	10.9	27.3	56.8	79.6
2002.....	8.3	*	*	0.1	0.3	0.7	1.5	4.0	11.6	29.2	57.7	78.2
2001.....	8.6	*	*	0.1	0.4	0.8	1.7	4.5	12.3	29.9	59.2	80.3
2000.....	8.8	*	*	0.1	0.5	0.8	1.7	4.7	13.2	30.7	61.7	80.9
1999.....	8.9	*	*	0.1	0.4	0.7	1.8	4.8	13.5	32.1	62.4	78.3
Non-Hispanic 4/, male												
2007.....	8.3	*	*	*	0.4	0.7	1.6	4.1	11.5	30.1	66.7	93.9
2006.....	8.3	*	*	0.1	0.4	0.7	1.5	4.4	11.9	30.3	68.1	91.5
2005.....	8.5	*	*	0.2	0.4	0.7	1.8	4.5	12.2	31.9	68.2	103.0
2004.....	8.5	*	*	0.1	0.4	0.8	1.6	4.8	12.6	33.1	68.4	97.8
2003.....	8.8	*	*	0.1	0.4	0.8	1.8	5.0	14.0	34.5	70.3	105.6
2002.....	9.0	*	*	0.2	0.4	0.9	2.0	5.1	14.6	36.5	71.8	101.4
2001.....	9.2	*	*	0.1	0.5	0.9	2.3	5.9	15.1	37.2	71.7	108.1
2000.....	9.4	*	*	0.2	0.7	0.9	2.2	6.1	15.9	37.7	75.6	106.7
1999.....	9.4	*	*	0.2	0.5	0.9	2.3	6.2	16.5	38.8	75.5	106.0
Non-Hispanic 4/, female												
2007.....	6.8	*	*	*	0.2	0.3	0.8	2.3	6.6	17.9	44.4	63.8
2006.....	7.0	*	*	*	0.2	0.4	0.9	2.3	6.7	19.4	45.3	65.7
2005.....	7.1	*	*	*	0.2	0.5	0.9	2.3	7.6	19.5	46.0	64.6
2004.....	7.3	*	*	*	0.3	0.4	0.8	2.7	7.7	21.2	46.5	66.7
2003.....	7.4	*	*	*	0.3	0.5	1.0	2.8	8.1	21.3	47.8	68.2
2002.....	7.7	*	*	*	0.3	0.5	1.1	3.0	8.9	23.0	48.3	68.2
2001.....	8.0	*	*	*	0.3	0.6	1.2	3.1	9.7	23.8	50.9	68.7
2000.....	8.3	*	*	*	0.3	0.6	1.2	3.4	10.6	24.8	52.6	70.4
1999.....	8.4	*	*	*	0.3	0.5	1.3	3.4	10.8	26.5	53.9	67.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's Lymphoma (C82-C85)												
Non-Hispanic white, both sexes												
2007 5/.....	8.6	*	*	*	0.3	0.4	1.1	3.1	9.1	24.9	56.6	78.2
2006 5/.....	8.7	*	*	0.1	0.2	0.5	1.1	3.3	9.3	25.8	57.9	78.2
2005 5/.....	8.9	*	*	0.1	0.3	0.5	1.3	3.3	10.0	26.6	58.4	81.3
2004 5/.....	9.0	*	*	0.1	0.3	0.6	1.2	3.7	10.4	28.1	58.5	80.2
2003 5/.....	9.3	*	*	0.1	0.3	0.5	1.4	3.8	11.4	28.8	60.0	84.3
2002.....	9.5	*	*	0.2	0.3	0.7	1.5	4.0	12.0	30.6	60.9	83.1
2001.....	9.7	*	*	0.1	0.3	0.7	1.7	4.5	12.7	31.4	62.3	84.1
2000.....	10.1	*	*	0.1	0.4	0.7	1.7	4.8	13.8	32.3	65.1	85.1
1999.....	10.1	*	*	0.1	0.4	0.7	1.7	4.9	14.1	33.9	65.6	82.0
Non-Hispanic white, male												
2007 5/.....	9.4	*	*	*	0.4	0.6	1.5	4.0	11.8	31.4	70.4	99.6
2006 5/.....	9.4	*	*	*	0.3	0.6	1.3	4.3	12.0	31.8	72.2	96.2
2005 5/.....	9.7	*	*	0.2	0.4	0.6	1.7	4.3	12.4	33.6	72.4	109.8
2004 5/.....	9.6	*	*	*	0.4	0.8	1.7	4.7	12.9	34.6	72.0	103.5
2003 5/.....	10.1	*	*	*	0.4	0.7	1.8	4.9	14.6	36.6	74.4	112.0
2002.....	10.2	*	*	0.2	0.4	0.8	1.9	5.1	15.0	38.3	75.6	107.6
2001.....	10.4	*	*	*	0.4	0.9	2.1	6.0	15.5	38.8	75.2	113.1
2000.....	10.6	*	*	0.2	0.6	0.8	2.2	6.1	16.5	39.5	79.6	113.0
1999.....	10.6	*	*	0.2	0.5	0.8	2.2	6.3	17.0	40.8	78.9	110.0
Non-Hispanic white, female												
2007 5/.....	7.9	*	*	*	0.2	0.2	0.7	2.2	6.6	19.2	47.0	68.0
2006 5/.....	8.1	*	*	*	*	0.4	0.8	2.3	6.8	20.5	48.0	69.8
2005 5/.....	8.2	*	*	*	*	0.4	0.8	2.3	7.8	20.6	48.8	68.3
2004 5/.....	8.4	*	*	*	0.3	0.4	0.8	2.7	8.0	22.5	49.2	69.8
2003 5/.....	8.5	*	*	*	0.3	0.4	1.0	2.8	8.4	22.1	50.3	72.2
2002.....	8.8	*	*	*	0.2	0.5	1.1	2.9	9.2	24.1	51.0	72.6
2001.....	9.2	*	*	*	0.3	0.5	1.2	3.1	10.1	25.2	53.7	72.0
2000.....	9.5	*	*	*	0.3	0.6	1.1	3.4	11.3	26.3	55.6	73.8
1999.....	9.6	*	*	*	0.2	0.5	1.2	3.5	11.4	28.1	56.9	70.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's Lymphoma (C82-C85)												
Non-Hispanic black, both sexes												
2007 5/.....	3.6	*	*	*	0.4	0.9	1.9	4.0	9.4	16.7	26.1	31.7
2006 5/.....	3.9	*	*	*	0.5	0.9	2.2	4.6	10.2	17.3	28.3	34.5
2005 5/.....	3.7	*	*	*	0.4	1.1	2.0	4.4	9.2	17.2	27.0	34.4
2004 5/.....	3.8	*	*	*	0.5	1.1	1.5	4.8	9.1	18.4	28.2	39.4
2003 5/.....	3.7	*	*	*	0.4	1.1	2.0	4.9	9.1	18.4	28.6	31.2
2002.....	4.0	*	*	*	0.4	1.2	2.0	4.8	10.8	21.6	30.1	31.2
2001.....	4.1	*	*	*	0.6	1.2	2.3	5.1	10.9	21.3	30.6	40.0
2000.....	3.9	*	*	*	0.6	1.1	2.1	5.0	10.8	19.7	27.1	36.6
1999.....	4.1	*	*	*	0.6	1.1	2.4	4.8	11.2	20.5	32.1	40.4
Non-Hispanic black, male												
2007 5/.....	4.1	*	*	*	*	1.1	2.6	5.2	12.0	23.6	30.8	37.5
2006 5/.....	4.4	*	*	*	*	1.0	2.9	6.6	13.9	22.1	32.8	43.1
2005 5/.....	4.2	*	*	*	*	1.3	2.3	6.3	12.3	23.1	31.1	40.9
2004 5/.....	4.1	*	*	*	0.6	1.4	1.9	6.6	12.0	23.5	33.4	42.7
2003 5/.....	3.9	*	*	*	*	1.4	2.4	6.7	11.5	20.7	31.2	35.2
2002.....	4.5	*	*	*	*	1.6	2.9	5.7	13.9	28.3	35.8	43.8
2001.....	4.7	*	*	*	0.8	1.3	3.4	6.7	13.7	28.2	37.2	55.4
2000.....	4.3	*	*	*	*	1.2	2.6	7.0	14.8	25.3	31.8	43.4
1999.....	4.7	*	*	*	*	1.6	3.3	6.6	14.8	26.2	41.0	59.4
Non-Hispanic black, female												
2007 5/.....	3.2	*	*	*	*	0.8	1.3	2.9	7.3	11.8	23.5	29.3
2006 5/.....	3.4	*	*	*	*	0.7	1.6	2.8	7.2	13.9	25.7	31.0
2005 5/.....	3.3	*	*	*	*	0.8	1.7	2.7	6.7	13.0	24.6	31.8
2004 5/.....	3.4	*	*	*	*	0.8	1.2	3.4	6.9	14.8	25.3	38.1
2003 5/.....	3.5	*	*	*	*	0.9	1.5	3.4	7.1	16.9	27.2	29.7
2002.....	3.6	*	*	*	*	0.8	1.2	4.0	8.3	16.9	26.9	26.3
2001.....	3.7	*	*	*	*	1.1	1.4	3.7	8.6	16.5	27.0	34.2
2000.....	3.5	*	*	*	*	0.9	1.7	3.3	7.5	15.7	24.5	34.0
1999.....	3.6	*	*	*	*	*	1.7	3.2	8.2	16.5	27.1	33.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Leukemia (C91-C95)

All origins 3/, both sexes

2007.....	7.2	0.5	0.6	0.8	1.0	1.1	1.5	3.1	8.6	23.8	52.7	76.7
2006.....	7.3	0.8	0.8	0.6	1.1	1.1	1.6	3.2	8.9	25.0	52.6	76.9
2005.....	7.3	0.5	0.7	0.7	1.1	1.1	1.6	3.3	8.8	25.1	52.3	78.7
2004.....	7.3	0.6	0.8	0.8	1.0	1.0	1.6	3.3	9.2	25.4	52.4	78.8
2003.....	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002.....	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001.....	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000.....	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999.....	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4

All origins 3/, male

2007.....	8.3	*	0.6	0.9	1.1	1.2	1.7	3.6	10.9	31.6	75.7	109.4
2006.....	8.4	*	0.8	0.7	1.4	1.3	1.8	3.7	11.4	33.1	75.9	105.1
2005.....	8.4	*	1.0	0.8	1.4	1.3	1.6	3.6	11.3	34.0	75.0	112.4
2004.....	8.3	*	0.9	0.9	1.3	1.2	1.8	3.8	11.9	34.4	73.2	110.7
2003.....	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002.....	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001.....	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000.....	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999.....	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2

All origins 3/, female

2007.....	6.2	*	0.7	0.7	0.9	0.9	1.4	2.6	6.4	17.2	36.8	61.2
2006.....	6.3	*	0.8	0.6	0.8	0.9	1.5	2.7	6.7	18.2	36.7	63.7
2005.....	6.2	*	0.5	0.6	0.8	1.0	1.5	3.0	6.4	17.5	37.0	63.3
2004.....	6.3	*	0.6	0.7	0.8	0.9	1.4	2.8	6.8	17.8	38.4	64.5
2003.....	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002.....	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001.....	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000.....	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999.....	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Hispanic, both sexes												
2007.....	2.8	*	*	1.2	1.5	1.3	1.6	2.9	6.7	14.2	28.1	30.4
2006.....	2.8	*	0.7	0.7	1.5	1.3	1.7	2.7	5.9	16.9	29.4	35.2
2005.....	3.0	*	0.7	1.0	1.7	1.3	1.5	3.2	7.0	17.6	28.2	40.6
2004.....	2.8	*	0.8	1.3	1.5	1.1	1.5	3.0	6.3	15.7	28.6	36.2
2003.....	2.9	*	1.2	1.1	1.4	1.1	1.7	3.6	6.3	13.9	32.9	47.2
2002.....	2.8	*	0.8	1.1	1.6	1.1	2.1	2.5	7.2	15.5	27.4	47.1
2001.....	3.0	*	1.3	1.1	1.7	1.5	1.9	2.9	6.5	17.3	33.4	42.5
2000.....	2.9	*	0.9	1.1	1.5	1.1	1.9	3.5	7.8	16.9	32.6	49.1
1999.....	2.9	*	0.9	1.1	1.6	1.4	2.0	3.4	7.0	17.3	31.0	44.2
Hispanic, male												
2007.....	3.0	*	*	1.2	1.7	1.4	1.7	3.0	7.6	17.0	34.8	51.8
2006.....	3.0	*	*	0.7	1.8	1.4	1.6	3.0	7.1	20.9	40.5	42.9
2005.....	3.4	*	*	1.1	2.2	1.5	1.7	3.4	8.4	22.9	41.8	57.3
2004.....	3.1	*	*	1.6	2.0	1.3	1.8	3.3	8.1	20.2	37.9	43.4
2003.....	3.2	*	1.4	1.3	1.7	1.3	1.9	4.1	7.6	18.3	40.6	61.6
2002.....	3.0	*	*	1.1	2.0	1.4	2.3	2.8	7.2	18.5	38.6	65.3
2001.....	3.4	*	1.4	1.1	2.2	1.8	2.3	3.3	8.4	19.6	45.6	68.6
2000.....	3.3	*	*	1.3	1.7	1.5	2.0	4.1	10.2	20.7	42.8	76.6
1999.....	3.1	*	*	1.1	1.7	1.4	1.9	3.6	8.6	23.5	39.1	49.6
Hispanic, female												
2007.....	2.7	*	*	1.2	1.3	1.3	1.5	2.7	5.7	11.9	23.4	18.8
2006.....	2.6	*	*	0.7	1.2	1.2	1.7	2.5	4.8	13.7	21.7	31.2
2005.....	2.5	*	*	0.9	1.2	1.1	1.3	3.0	5.7	13.4	18.8	31.9
2004.....	2.4	*	*	1.0	0.9	0.9	1.3	2.6	4.6	12.1	22.2	32.4
2003.....	2.6	*	*	1.0	1.0	0.9	1.5	3.2	5.2	10.4	27.7	39.8
2002.....	2.6	*	*	1.2	1.1	0.7	1.9	2.3	7.3	13.1	19.7	38.1
2001.....	2.6	*	*	1.1	1.1	1.1	1.4	2.6	4.8	15.6	25.1	29.8
2000.....	2.6	*	*	0.8	1.3	*	1.8	2.9	5.6	13.9	25.8	35.6
1999.....	2.8	*	*	1.0	1.5	1.3	2.1	3.2	5.6	12.4	25.7	41.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic 4/, both sexes												
2007.....	8.0	*	0.7	0.7	0.9	1.0	1.5	3.1	8.7	24.6	54.2	79.0
2006.....	8.1	0.7	0.8	0.6	1.0	1.1	1.6	3.3	9.2	25.6	54.0	78.9
2005.....	8.0	*	0.8	0.6	1.0	1.1	1.5	3.3	8.9	25.6	53.7	80.6
2004.....	8.0	*	0.8	0.6	0.9	1.0	1.6	3.3	9.5	26.1	53.7	80.6
2003.....	8.1	0.8	0.7	0.8	0.9	1.1	1.8	3.3	9.6	26.4	54.9	81.6
2002.....	8.2	*	0.6	0.7	1.1	1.2	1.6	3.7	10.0	27.0	54.3	81.7
2001.....	8.2	*	0.9	0.7	1.1	1.2	1.7	3.8	10.3	27.2	54.1	82.7
2000.....	8.2	1.0	0.9	0.7	1.1	1.2	1.7	3.8	10.1	27.8	55.3	84.7
1999.....	8.2	*	0.7	0.7	1.2	1.2	1.7	3.8	10.8	27.8	53.6	83.7
Non-Hispanic 4/, male												
2007.....	9.3	*	0.7	0.8	1.0	1.2	1.7	3.7	11.1	32.7	78.3	112.6
2006.....	9.4	*	0.8	0.7	1.3	1.3	1.8	3.8	11.7	34.0	78.1	108.3
2005.....	9.3	*	1.0	0.7	1.2	1.2	1.6	3.7	11.5	34.8	76.9	115.3
2004.....	9.2	*	0.9	0.7	1.1	1.2	1.8	3.9	12.2	35.4	75.1	113.9
2003.....	9.3	*	0.8	0.8	1.1	1.4	2.0	4.0	12.2	36.2	77.0	113.7
2002.....	9.4	*	0.7	0.8	1.2	1.4	1.9	4.3	12.6	37.3	76.5	116.5
2001.....	9.3	*	0.9	0.8	1.2	1.4	1.9	4.3	13.1	36.6	75.4	119.3
2000.....	9.3	*	0.9	0.8	1.4	1.4	1.9	4.3	12.6	38.2	77.0	121.5
1999.....	9.2	*	0.8	0.8	1.4	1.3	1.9	4.4	13.4	37.5	74.6	122.1
Non-Hispanic 4/, female												
2007.....	6.8	*	0.7	0.6	0.8	0.9	1.3	2.6	6.5	17.6	37.6	63.2
2006.....	6.9	*	0.8	0.5	0.7	0.9	1.4	2.8	6.8	18.6	37.6	65.2
2005.....	6.8	*	0.5	0.5	0.8	1.0	1.5	3.0	6.5	17.8	38.0	64.8
2004.....	6.9	*	0.6	0.6	0.8	0.9	1.4	2.8	7.0	18.2	39.3	65.7
2003.....	7.0	*	0.7	0.7	0.7	0.8	1.6	2.7	7.1	18.3	40.2	67.4
2002.....	7.0	*	0.5	0.6	0.9	1.0	1.4	3.2	7.6	18.4	39.5	66.6
2001.....	7.2	*	0.9	0.6	0.9	1.1	1.5	3.4	7.7	19.4	40.1	67.5
2000.....	7.2	*	0.9	0.6	0.8	1.0	1.6	3.3	7.7	19.3	41.1	69.8
1999.....	7.2	*	0.7	0.6	0.9	1.1	1.5	3.2	8.5	19.8	40.0	68.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Leukemia (C91-C95)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	9.1	*	0.7	0.7	0.9	1.0	1.4	3.1	8.9	25.4	56.8	82.9
2006 5/.....	9.2	*	0.9	0.6	1.0	1.1	1.6	3.4	9.4	26.8	56.3	82.0
2005 5/.....	9.1	*	0.7	0.6	1.0	1.0	1.6	3.3	9.2	26.7	56.1	84.5
2004 5/.....	9.0	*	0.7	0.7	0.9	1.0	1.6	3.3	9.7	26.8	55.7	83.6
2003 5/.....	9.1	1.0	0.8	0.8	0.9	1.1	1.8	3.3	9.8	27.5	57.1	84.7
2002.....	9.1	*	0.7	0.7	1.1	1.2	1.7	3.7	10.0	28.2	56.2	84.4
2001.....	9.1	*	1.0	0.7	1.1	1.2	1.7	3.8	10.5	28.1	56.0	86.1
2000.....	9.2	1.3	0.9	0.7	1.1	1.1	1.7	3.7	10.2	29.0	57.5	87.3
1999.....	9.1	*	0.8	0.7	1.1	1.2	1.7	3.7	11.0	28.6	55.3	86.8
<b>Non-Hispanic white, male</b>												
2007 5/.....	10.6	*	0.6	0.8	1.0	1.1	1.6	3.7	11.5	33.8	82.2	118.1
2006 5/.....	10.6	*	0.8	0.7	1.3	1.4	1.8	3.9	12.0	35.3	81.0	112.1
2005 5/.....	10.5	*	0.8	0.7	1.1	1.1	1.6	3.6	11.9	36.3	79.7	121.9
2004 5/.....	10.3	*	0.8	0.7	1.1	1.2	1.7	3.8	12.5	36.1	78.0	118.8
2003 5/.....	10.5	*	0.8	0.9	1.1	1.3	1.9	4.0	12.5	37.7	80.6	119.0
2002.....	10.5	*	0.8	0.7	1.3	1.3	1.9	4.2	12.7	38.7	79.2	121.3
2001.....	10.3	*	1.0	0.7	1.3	1.4	1.8	4.3	13.3	37.7	78.0	125.3
2000.....	10.4	*	1.0	0.8	1.3	1.3	1.9	4.2	12.7	39.5	79.9	125.5
1999.....	10.2	*	0.8	0.7	1.4	1.3	1.9	4.3	13.6	38.6	77.0	127.2
<b>Non-Hispanic white, female</b>												
2007 5/.....	7.6	*	0.8	0.5	0.7	0.8	1.2	2.6	6.5	18.1	39.1	66.2
2006 5/.....	7.8	*	0.9	0.5	0.7	0.9	1.5	2.9	6.9	19.4	39.1	68.0
2005 5/.....	7.7	*	0.5	0.5	0.8	0.9	1.6	3.0	6.6	18.5	39.9	67.3
2004 5/.....	7.7	*	0.6	0.6	0.8	0.8	1.4	2.8	7.0	18.8	40.4	67.8
2003 5/.....	7.8	*	0.7	0.6	0.7	0.8	1.6	2.6	7.3	18.8	41.3	69.6
2002.....	7.8	*	0.5	0.6	0.9	1.0	1.5	3.2	7.5	19.3	40.7	68.4
2001.....	8.0	*	1.1	0.6	0.9	1.0	1.6	3.3	7.8	19.9	41.4	69.8
2000.....	8.1	*	0.9	0.6	0.8	0.9	1.5	3.2	7.8	20.1	42.7	72.0
1999.....	8.0	*	0.9	0.6	0.8	1.1	1.5	3.2	8.5	20.3	41.1	70.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic black, both sexes												
2007 5/.....	4.7	*	*	0.7	1.0	1.2	2.1	3.7	9.3	22.9	39.8	52.9
2006 5/.....	4.7	*	*	0.6	1.0	1.1	1.9	3.4	9.7	23.1	41.0	56.8
2005 5/.....	4.5	*	1.0	0.6	1.0	1.6	1.6	4.0	8.6	21.5	39.0	53.0
2004 5/.....	5.0	*	*	0.6	1.0	1.3	2.1	4.0	10.4	25.4	41.7	60.7
2003 5/.....	4.8	*	*	0.7	0.9	1.2	2.4	3.8	9.8	23.4	42.1	60.4
2002.....	4.8	*	*	0.6	0.9	1.3	1.6	4.4	11.5	22.6	42.9	65.3
2001.....	4.9	*	*	0.9	1.0	1.5	2.1	4.3	10.1	25.0	42.1	57.1
2000.....	4.9	*	*	0.6	1.4	1.7	2.1	5.0	10.8	23.1	38.8	66.1
1999.....	4.9	*	*	0.8	1.5	1.3	1.9	4.4	10.8	25.2	41.6	55.9
Non-Hispanic black, male												
2007 5/.....	5.2	*	*	0.8	1.1	1.3	2.4	4.6	10.8	30.1	53.1	78.5
2006 5/.....	5.5	*	*	0.9	1.2	1.2	2.5	3.9	12.8	32.5	60.6	87.0
2005 5/.....	5.1	*	*	0.7	1.3	1.6	1.9	4.5	10.8	29.2	59.1	63.8
2004 5/.....	5.5	*	*	0.7	1.1	1.4	2.3	4.9	12.8	37.3	52.8	81.3
2003 5/.....	5.2	*	*	0.7	1.2	1.6	2.8	4.6	12.6	31.5	54.0	78.8
2002.....	5.3	*	*	0.7	1.0	1.5	1.8	5.2	14.1	32.2	58.0	87.5
2001.....	5.4	*	*	1.2	1.1	1.8	2.6	4.6	13.4	34.4	56.1	75.1
2000.....	5.5	*	*	0.7	1.9	2.0	2.0	5.6	14.0	32.3	55.8	98.6
1999.....	5.4	*	*	1.0	1.6	1.5	2.1	5.7	13.7	33.4	55.7	78.4
Non-Hispanic black, female												
2007 5/.....	4.3	*	*	*	0.9	1.0	1.8	2.9	8.0	17.8	32.2	42.3
2006 5/.....	4.0	*	*	*	0.8	1.1	1.4	2.9	7.3	16.3	29.8	44.5
2005 5/.....	4.1	*	*	*	0.7	1.6	1.3	3.7	6.7	16.0	27.6	48.6
2004 5/.....	4.5	*	*	*	0.9	1.2	1.8	3.2	8.5	17.0	35.4	52.6
2003 5/.....	4.4	*	*	0.8	0.7	0.9	1.9	3.1	7.5	17.6	35.4	53.2
2002.....	4.4	*	*	*	0.8	1.1	1.4	3.6	9.4	15.8	34.3	56.9
2001.....	4.4	*	*	*	1.0	1.1	1.7	4.0	7.4	18.4	34.2	50.5
2000.....	4.4	*	*	*	0.9	1.4	2.2	4.4	8.3	16.5	29.2	54.1
1999.....	4.4	*	*	0.6	1.3	1.1	1.7	3.3	8.5	19.4	33.8	47.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All origins 3/, both sexes

2007.....	3.7	*	*	*	*	*	0.4	1.6	5.5	15.1	29.6	34.1
2006.....	3.7	*	*	*	*	*	0.3	1.7	5.7	15.0	29.1	33.8
2005.....	3.8	*	*	*	*	*	0.3	1.6	6.1	15.9	29.5	33.6
2004.....	3.7	*	*	*	*	*	0.3	1.6	5.9	15.4	30.7	33.1
2003.....	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
2002.....	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001.....	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000.....	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999.....	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6

All origins 3/, male

2007.....	4.1	*	*	*	*	*	0.4	2.0	6.5	18.2	38.0	49.4
2006.....	4.0	*	*	*	*	*	0.4	2.0	6.8	18.1	37.5	45.8
2005.....	4.1	*	*	*	*	*	0.4	2.0	7.0	19.7	38.0	46.1
2004.....	3.9	*	*	*	*	*	0.4	1.9	6.9	18.8	38.1	44.6
2003.....	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
2002.....	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001.....	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000.....	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999.....	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3

All origins 3/, female

2007.....	3.4	*	*	*	*	*	0.3	1.3	4.5	12.4	23.8	26.8
2006.....	3.4	*	*	*	*	*	0.3	1.4	4.6	12.4	23.3	28.2
2005.....	3.5	*	*	*	*	*	0.2	1.3	5.2	12.8	23.7	27.9
2004.....	3.5	*	*	*	*	*	0.3	1.3	5.0	12.5	25.7	28.0
2003.....	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
2002.....	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001.....	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000.....	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999.....	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Hispanic, both sexes

2007.....	1.5	*	*	*	*	*	0.3	1.3	5.3	15.0	22.0	14.3
2006.....	1.4	*	*	*	*	*	0.3	1.5	4.0	14.2	21.6	22.7
2005.....	1.4	*	*	*	*	*	*	1.5	5.4	13.4	21.4	20.1
2004.....	1.4	*	*	*	*	*	*	1.3	5.5	12.4	24.0	18.6
2003.....	1.3	*	*	*	*	*	*	1.4	5.2	12.6	23.9	24.9
2002.....	1.5	*	*	*	*	*	*	1.8	5.0	14.3	26.7	33.3
2001.....	1.4	*	*	*	*	*	*	1.3	5.8	14.9	24.7	29.4
2000.....	1.4	*	*	*	*	*	*	1.4	5.3	15.0	24.5	31.2
1999.....	1.4	*	*	*	*	*	*	1.7	6.8	13.4	23.1	28.1

Hispanic, male

2007.....	1.5	*	*	*	*	*	*	1.8	5.5	16.8	25.1	24.4
2006.....	1.4	*	*	*	*	*	*	1.8	4.6	15.9	24.7	30.4
2005.....	1.3	*	*	*	*	*	*	1.8	4.9	15.2	26.4	29.9
2004.....	1.3	*	*	*	*	*	*	1.3	6.4	14.1	30.0	*
2003.....	1.2	*	*	*	*	*	*	1.6	5.8	13.3	25.7	*
2002.....	1.6	*	*	*	*	*	*	2.1	6.6	17.6	31.6	53.5
2001.....	1.5	*	*	*	*	*	*	1.4	6.0	20.4	35.8	38.1
2000.....	1.3	*	*	*	*	*	*	1.5	5.5	18.4	30.0	*
1999.....	1.3	*	*	*	*	*	*	1.9	7.8	14.7	24.5	*

Hispanic, female

2007.....	1.4	*	*	*	*	*	*	0.9	5.2	13.5	19.7	*
2006.....	1.4	*	*	*	*	*	*	1.2	3.4	12.8	19.5	18.6
2005.....	1.4	*	*	*	*	*	*	1.3	5.8	11.9	17.9	15.0
2004.....	1.4	*	*	*	*	*	*	1.2	4.7	11.1	19.8	18.7
2003.....	1.4	*	*	*	*	*	*	1.2	4.7	12.1	22.6	22.7
2002.....	1.4	*	*	*	*	*	*	1.5	3.5	11.7	23.3	23.2
2001.....	1.3	*	*	*	*	*	*	1.2	5.6	10.6	17.2	25.1
2000.....	1.4	*	*	*	*	*	*	*	5.1	12.3	20.8	30.7
1999.....	1.5	*	*	*	*	*	*	1.5	5.9	12.4	22.2	31.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 4/, both sexes

2007.....	4.2	*	*	*	*	*	0.4	1.6	5.5	15.1	30.1	35.1
2006.....	4.1	*	*	*	*	*	0.4	1.7	5.8	15.1	29.5	34.4
2005.....	4.2	*	*	*	*	*	0.3	1.6	6.1	16.1	29.9	34.2
2004.....	4.1	*	*	*	*	*	0.3	1.6	5.9	15.6	31.0	33.7
2003.....	4.3	*	*	*	*	*	0.3	1.8	6.0	16.8	31.4	36.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	6.4	17.5	31.3	36.7
2001.....	4.3	*	*	*	*	*	0.3	1.7	6.6	17.5	31.1	36.5
2000.....	4.3	*	*	*	*	*	0.3	1.9	6.4	17.4	31.8	38.0
1999.....	4.3	*	*	*	*	*	0.4	1.8	7.1	17.4	31.2	36.8

Non-Hispanic 4/, male

2007.....	4.6	*	*	*	*	*	0.5	2.0	6.6	18.3	38.9	50.8
2006.....	4.5	*	*	*	*	*	0.4	2.0	7.0	18.2	38.2	46.6
2005.....	4.6	*	*	*	*	*	0.4	2.0	7.2	20.0	38.7	46.9
2004.....	4.4	*	*	*	*	*	0.4	1.9	7.0	19.2	38.6	45.9
2003.....	4.6	*	*	*	*	*	0.5	2.2	7.5	20.2	38.2	51.2
2002.....	4.6	*	*	*	*	*	0.4	2.2	7.6	21.2	40.0	50.6
2001.....	4.5	*	*	*	*	*	0.4	2.2	7.9	20.8	38.6	48.4
2000.....	4.4	*	*	*	*	*	0.4	2.2	7.0	20.5	39.9	52.5
1999.....	4.5	*	*	*	*	*	0.4	2.1	8.5	20.4	38.9	52.2

Non-Hispanic 4/, female

2007.....	3.7	*	*	*	*	*	0.3	1.3	4.4	12.3	24.0	27.7
2006.....	3.8	*	*	*	*	*	0.3	1.4	4.7	12.4	23.5	28.6
2005.....	3.8	*	*	*	*	*	0.2	1.3	5.1	12.8	24.0	28.5
2004.....	3.9	*	*	*	*	*	0.3	1.3	5.0	12.6	25.9	28.3
2003.....	4.0	*	*	*	*	*	0.2	1.4	4.6	13.8	26.8	30.3
2002.....	4.0	*	*	*	*	*	0.3	1.5	5.2	14.5	25.5	30.8
2001.....	4.1	*	*	*	*	*	0.3	1.4	5.5	14.8	26.1	31.5
2000.....	4.2	*	*	*	*	*	0.3	1.6	5.7	14.8	26.5	32.1
1999.....	4.1	*	*	*	*	*	0.3	1.5	5.8	15.0	26.3	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic white, both sexes

2007 5/.....	4.2	*	*	*	*	*	0.3	1.3	4.8	13.9	28.7	34.2
2006 5/.....	4.2	*	*	*	*	*	0.3	1.4	5.1	13.9	28.5	33.3
2005 5/.....	4.3	*	*	*	*	*	0.3	1.3	5.4	14.8	29.0	33.5
2004 5/.....	4.2	*	*	*	*	*	0.3	1.4	5.3	14.2	29.6	32.6
2003 5/.....	4.4	*	*	*	*	*	0.3	1.4	5.4	15.6	29.8	35.3
2002.....	4.5	*	*	*	*	*	0.3	1.5	5.6	16.4	30.3	35.7
2001.....	4.4	*	*	*	*	*	0.3	1.5	5.7	16.3	29.7	35.1
2000.....	4.3	*	*	*	*	*	0.3	1.6	5.6	16.2	30.2	35.2
1999.....	4.3	*	*	*	*	*	0.3	1.5	6.1	16.2	29.6	35.0

Non-Hispanic white, male

2007 5/.....	4.8	*	*	*	*	*	0.4	1.7	5.9	17.4	37.5	50.6
2006 5/.....	4.6	*	*	*	*	*	0.4	1.7	6.1	17.0	37.2	45.9
2005 5/.....	4.7	*	*	*	*	*	0.4	1.6	6.4	18.7	37.6	46.5
2004 5/.....	4.5	*	*	*	*	*	0.3	1.6	6.3	17.9	37.2	44.8
2003 5/.....	4.7	*	*	*	*	*	0.4	1.8	6.8	19.3	37.0	50.5
2002.....	4.8	*	*	*	*	*	0.3	1.9	6.9	20.2	39.2	50.1
2001.....	4.6	*	*	*	*	*	0.3	1.9	6.9	19.5	37.4	47.3
2000.....	4.6	*	*	*	*	*	0.4	1.8	6.3	19.2	38.5	50.9
1999.....	4.6	*	*	*	*	*	0.3	1.9	7.5	19.2	37.4	50.6

Non-Hispanic white, female

2007 5/.....	3.7	*	*	*	*	*	0.2	1.0	3.7	10.8	22.5	26.4
2006 5/.....	3.8	*	*	*	*	*	0.2	1.1	4.1	11.2	22.5	27.5
2005 5/.....	3.9	*	*	*	*	*	0.2	1.0	4.5	11.4	23.1	27.5
2004 5/.....	3.9	*	*	*	*	*	0.2	1.1	4.3	11.1	24.4	27.1
2003 5/.....	4.0	*	*	*	*	*	0.2	1.1	4.0	12.4	24.9	28.7
2002.....	4.1	*	*	*	*	*	0.2	1.1	4.5	13.1	24.4	29.5
2001.....	4.1	*	*	*	*	*	0.2	1.1	4.7	13.6	24.6	30.0
2000.....	4.1	*	*	*	*	*	0.2	1.4	4.9	13.6	24.8	28.9
1999.....	4.1	*	*	*	*	*	0.2	1.2	4.7	13.7	24.6	28.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic black, both sexes

2007 5/.....	4.9	*	*	*	*	*	0.9	3.8	11.8	28.9	50.8	54.0
2006 5/.....	4.8	*	*	*	*	*	0.7	4.0	12.9	28.4	48.0	52.0
2005 5/.....	4.7	*	*	*	*	*	0.6	3.7	12.7	30.3	47.6	50.2
2004 5/.....	4.9	*	*	*	*	*	0.7	3.5	12.6	31.0	53.9	54.7
2003 5/.....	5.0	*	*	*	*	*	0.8	4.3	12.2	31.4	55.8	60.4
2002.....	4.8	*	*	*	*	*	0.7	4.3	13.4	31.8	49.2	55.7
2001.....	5.0	*	*	*	*	*	0.9	3.7	15.3	31.8	53.2	59.3
2000.....	5.2	*	*	*	*	*	0.8	4.5	14.1	32.5	55.6	72.8
1999.....	5.1	*	*	*	*	*	0.9	4.3	16.8	30.7	53.6	63.4

Non-Hispanic black, male

2007 5/.....	4.8	*	*	*	*	*	1.0	4.3	13.5	31.7	65.5	65.4
2006 5/.....	5.1	*	*	*	*	*	0.8	4.6	16.3	36.5	60.6	66.0
2005 5/.....	4.9	*	*	*	*	*	*	4.6	15.0	36.3	62.8	62.8
2004 5/.....	4.8	*	*	*	*	*	0.8	4.0	14.5	37.1	65.3	72.1
2003 5/.....	4.9	*	*	*	*	*	0.9	5.3	15.7	34.5	64.1	74.6
2002.....	4.8	*	*	*	*	*	0.8	4.8	15.9	37.6	60.1	71.1
2001.....	5.0	*	*	*	*	*	1.1	4.3	18.5	37.2	66.0	72.8
2000.....	5.0	*	*	*	*	*	*	5.0	15.4	39.1	68.5	81.0
1999.....	5.0	*	*	*	*	*	1.1	4.4	19.0	35.6	65.0	80.8

Non-Hispanic black, female

2007 5/.....	5.0	*	*	*	*	*	0.9	3.3	10.4	26.9	42.4	49.2
2006 5/.....	4.6	*	*	*	*	*	0.7	3.4	10.2	22.7	40.9	46.3
2005 5/.....	4.6	*	*	*	*	*	*	2.9	10.8	26.0	38.9	45.2
2004 5/.....	4.9	*	*	*	*	*	*	3.1	11.1	26.7	47.5	47.8
2003 5/.....	5.1	*	*	*	*	*	*	3.4	9.3	29.2	51.2	54.9
2002.....	4.9	*	*	*	*	*	*	3.8	11.3	27.7	43.1	49.7
2001.....	5.0	*	*	*	*	*	0.8	3.1	12.7	28.0	46.0	54.3
2000.....	5.4	*	*	*	*	*	0.9	4.1	13.1	27.9	48.3	69.8
1999.....	5.3	*	*	*	*	*	0.7	4.3	15.1	27.2	47.3	56.9











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes

2007.....	21.0	0.5	0.7	0.6	1.3	1.7	4.0	12.4	33.4	71.6	139.7	206.7
2006.....	21.1	0.5	0.7	0.6	1.2	1.8	4.1	12.7	33.5	75.0	140.1	210.1
2005.....	21.2	*	0.8	0.7	1.4	1.7	4.2	13.1	34.3	75.2	141.6	212.9
2004.....	21.1	0.6	0.8	0.7	1.3	1.7	4.3	12.7	34.4	76.8	143.0	213.1
2003.....	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
2002.....	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001.....	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000.....	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999.....	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4

All origins 3/, male

2007.....	22.1	*	0.7	0.6	1.7	2.1	4.2	14.2	40.1	86.7	174.0	260.9
2006.....	21.9	*	0.7	0.6	1.6	2.2	4.3	14.3	40.0	89.0	173.9	259.6
2005.....	22.1	*	0.8	0.7	1.7	2.2	4.6	14.9	41.0	90.1	173.1	263.5
2004.....	21.9	*	0.7	0.8	1.7	2.1	4.6	14.4	40.6	92.5	177.9	260.6
2003.....	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
2002.....	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001.....	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000.....	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999.....	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9

All origins 3/, female

2007.....	19.8	*	0.7	0.7	0.9	1.3	3.9	10.6	27.2	58.8	116.0	180.9
2006.....	20.2	*	0.6	0.6	0.9	1.4	3.9	11.3	27.5	63.2	117.0	186.9
2005.....	20.4	*	0.7	0.6	1.1	1.3	3.8	11.3	28.1	62.6	120.2	189.6
2004.....	20.3	*	0.9	0.6	0.9	1.4	3.9	11.0	28.6	63.7	119.5	191.8
2003.....	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
2002.....	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001.....	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000.....	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999.....	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Hispanic, both sexes

2007.....	7.5	*	*	0.6	1.5	1.6	3.1	9.4	22.7	48.8	95.5	136.2
2006.....	7.7	*	0.6	0.6	1.5	1.3	3.1	9.4	24.4	54.4	97.6	145.5
2005.....	7.6	*	0.6	0.6	1.3	1.7	3.1	8.0	22.7	55.8	108.9	139.5
2004.....	7.2	*	0.6	0.5	1.3	1.3	2.6	8.9	22.6	52.7	106.5	140.8
2003.....	7.7	*	0.6	0.7	1.3	1.5	3.3	9.8	23.7	58.7	109.4	168.9
2002.....	7.8	*	*	0.8	1.6	1.5	3.0	9.5	25.4	60.3	112.8	164.1
2001.....	7.7	*	*	0.6	1.3	1.2	3.4	9.8	26.0	60.7	110.9	193.8
2000.....	7.8	*	0.8	0.7	1.2	1.5	2.7	9.2	26.6	64.1	124.6	192.4
1999.....	8.1	*	*	0.7	1.3	1.6	3.5	10.6	28.8	64.8	122.3	202.1

Hispanic, male

2007.....	7.6	*	*	0.6	2.1	1.8	3.3	10.4	25.3	57.7	109.1	157.5
2006.....	7.6	*	*	0.6	1.9	1.7	2.7	9.2	27.0	67.1	113.8	163.5
2005.....	7.8	*	*	0.5	1.6	2.0	3.5	8.5	27.7	67.1	129.6	159.4
2004.....	7.1	*	*	0.6	1.8	1.6	2.9	9.3	25.3	59.4	131.3	147.1
2003.....	7.9	*	*	0.7	1.6	2.1	3.3	11.2	27.7	71.0	127.3	194.0
2002.....	7.8	*	*	0.8	1.9	1.7	3.1	10.4	29.6	69.2	138.7	204.1
2001.....	7.7	*	*	0.6	1.8	1.5	3.4	10.7	30.2	67.7	140.9	213.3
2000.....	7.7	*	*	0.7	1.4	1.9	2.9	9.5	29.2	73.6	151.9	231.8
1999.....	8.0	*	*	0.7	1.8	1.9	3.5	11.1	33.3	77.1	144.9	235.1

Hispanic, female

2007.....	7.5	*	*	0.6	0.8	1.3	2.9	8.3	20.3	41.5	86.0	124.7
2006.....	7.8	*	*	0.7	1.0	0.9	3.5	9.6	21.9	44.0	86.3	136.0
2005.....	7.4	*	*	0.6	1.0	1.4	2.6	7.6	18.1	46.7	94.4	129.0
2004.....	7.3	*	*	*	0.6	1.1	2.4	8.4	20.2	47.3	89.4	137.6
2003.....	7.6	*	*	0.7	1.1	0.8	3.2	8.4	20.1	48.8	97.2	156.3
2002.....	7.7	*	*	0.7	1.2	1.1	2.9	8.6	21.6	53.2	95.0	144.2
2001.....	7.7	*	*	0.6	0.7	0.8	3.5	8.9	22.4	55.3	90.7	184.3
2000.....	7.9	*	*	0.6	1.1	1.1	2.5	9.0	24.3	56.6	106.3	173.1
1999.....	8.2	*	*	0.7	0.8	1.2	3.6	10.1	24.7	55.1	107.2	186.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 4/, both sexes

2007.....	23.3	*	0.8	0.6	1.3	1.8	4.2	12.7	34.3	73.3	142.4	210.3
2006.....	23.4	*	0.7	0.6	1.2	1.9	4.3	13.1	34.2	76.5	142.6	213.1
2005.....	23.5	*	0.8	0.7	1.4	1.7	4.3	13.6	35.3	76.6	143.4	216.2
2004.....	23.3	0.6	0.8	0.7	1.3	1.8	4.5	13.1	35.2	78.5	144.8	215.9
2003.....	23.7	*	0.7	0.7	1.2	1.8	4.4	14.1	36.0	79.0	148.4	225.8
2002.....	24.0	*	0.9	0.7	1.2	1.9	4.8	14.1	37.3	81.9	148.0	232.1
2001.....	24.1	*	0.8	0.7	1.3	2.0	4.7	14.4	38.1	83.1	150.1	235.3
2000.....	24.4	0.7	0.8	0.6	1.3	1.8	4.8	14.5	39.1	86.5	153.3	241.0
1999.....	25.8	*	0.8	0.7	1.3	1.9	5.3	15.3	42.3	92.7	161.9	251.6

Non-Hispanic 4/, male

2007.....	24.8	*	0.7	0.6	1.6	2.2	4.4	14.7	41.3	88.8	178.1	266.7
2006.....	24.5	*	0.8	0.6	1.5	2.3	4.6	14.8	41.0	90.4	177.4	264.6
2005.....	24.5	*	0.8	0.8	1.7	2.2	4.7	15.6	42.1	91.7	175.5	268.7
2004.....	24.4	*	0.8	0.8	1.6	2.2	4.9	14.9	41.7	94.7	180.3	265.7
2003.....	24.6	*	0.8	0.7	1.5	2.1	4.9	16.1	42.4	94.7	183.4	278.2
2002.....	25.0	*	0.9	0.7	1.6	2.4	5.3	16.4	44.1	99.7	181.5	294.0
2001.....	24.9	*	1.0	0.8	1.7	2.4	5.1	16.9	44.7	100.0	182.7	297.9
2000.....	25.2	*	0.9	0.6	1.6	2.2	5.3	16.5	45.9	104.7	188.7	303.0
1999.....	26.8	*	1.0	0.8	1.6	2.5	6.1	17.8	50.1	112.0	197.6	321.9

Non-Hispanic 4/, female

2007.....	21.9	*	0.8	0.7	0.9	1.3	4.0	10.8	27.8	60.1	117.9	183.7
2006.....	22.2	*	0.6	0.5	0.8	1.5	3.9	11.4	28.0	64.7	118.9	189.1
2005.....	22.4	*	0.8	0.7	1.1	1.3	3.9	11.7	28.9	63.8	121.6	192.2
2004.....	22.3	*	0.8	0.6	1.0	1.5	4.1	11.3	29.3	64.8	121.0	193.7
2003.....	22.8	*	0.6	0.6	0.9	1.5	3.8	12.2	30.0	65.9	125.1	202.7
2002.....	22.9	*	0.8	0.7	0.9	1.4	4.3	11.8	31.0	67.1	125.8	205.4
2001.....	23.3	*	0.6	0.6	0.9	1.6	4.3	12.0	31.9	69.1	128.7	209.3
2000.....	23.7	*	0.8	0.6	1.0	1.4	4.4	12.5	32.8	71.5	130.2	215.9
1999.....	25.0	*	0.7	0.6	1.1	1.4	4.6	13.0	35.1	76.9	138.8	223.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic white, both sexes

2007 5/.....	25.5	*	0.8	0.6	1.3	1.7	4.1	12.1	33.6	73.6	145.2	216.2
2006 5/.....	25.6	*	0.6	0.6	1.2	1.9	4.2	12.5	33.5	77.1	145.3	217.1
2005 5/.....	25.6	*	0.8	0.7	1.4	1.7	4.3	12.9	34.5	76.9	145.8	220.9
2004 5/.....	25.4	*	0.8	0.7	1.3	1.9	4.4	12.3	34.4	78.7	146.6	220.6
2003 5/.....	25.9	*	0.6	0.7	1.3	1.7	4.2	13.4	35.4	79.5	150.3	229.9
2002.....	26.0	*	0.8	0.7	1.2	1.9	4.6	13.5	36.5	81.8	149.1	235.2
2001.....	26.0	*	0.9	0.7	1.3	2.0	4.5	13.6	36.8	82.5	150.8	238.9
2000.....	26.5	*	0.8	0.7	1.3	1.9	4.5	13.6	38.2	86.3	153.9	244.9
1999.....	27.9	*	0.9	0.7	1.3	2.0	5.0	14.6	40.9	92.7	162.3	253.6

Non-Hispanic white, male

2007 5/.....	27.2	*	0.7	0.6	1.6	2.1	4.2	14.0	40.4	89.1	182.2	276.0
2006 5/.....	27.0	*	0.7	0.7	1.5	2.4	4.6	14.1	40.3	91.5	180.8	271.8
2005 5/.....	26.8	*	0.8	0.7	1.7	2.1	4.7	14.8	41.0	91.3	178.8	276.1
2004 5/.....	26.5	*	0.9	0.8	1.6	2.4	4.7	13.9	40.3	94.4	183.2	272.4
2003 5/.....	26.8	*	0.7	0.7	1.6	2.1	4.8	15.2	41.5	94.8	186.1	284.6
2002.....	27.2	*	0.8	0.7	1.5	2.4	5.0	15.7	42.9	100.0	183.2	300.2
2001.....	26.8	*	1.1	0.7	1.7	2.3	5.0	15.9	42.8	99.3	183.7	302.9
2000.....	27.2	*	1.0	0.7	1.6	2.3	5.0	15.6	44.1	104.6	189.1	309.5
1999.....	28.8	*	1.1	0.8	1.5	2.6	5.8	16.6	48.3	111.5	198.9	326.9

Non-Hispanic white, female

2007 5/.....	23.9	*	0.8	0.7	0.9	1.3	4.0	10.3	27.1	60.2	119.3	187.9
2006 5/.....	24.3	*	0.5	0.6	0.8	1.4	3.8	11.0	27.0	64.6	120.7	191.5
2005 5/.....	24.5	*	0.7	0.7	1.1	1.2	3.8	11.1	28.4	64.3	123.1	195.6
2004 5/.....	24.4	*	0.8	0.5	1.0	1.4	4.0	10.8	28.8	65.2	121.7	197.4
2003 5/.....	25.0	*	0.6	0.6	0.9	1.3	3.7	11.7	29.6	66.4	126.2	205.8
2002.....	24.9	*	0.7	0.7	0.9	1.4	4.3	11.3	30.5	66.3	126.3	207.0
2001.....	25.3	*	0.7	0.6	1.0	1.6	4.0	11.4	31.3	68.3	129.1	212.3
2000.....	25.8	*	0.6	0.6	1.0	1.4	4.1	11.7	32.6	71.0	130.8	218.9
1999.....	27.0	*	0.8	0.6	1.1	1.3	4.2	12.6	33.9	77.0	138.3	224.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic black, both sexes

2007 5/.....	17.3	*	*	0.6	1.3	2.1	5.6	18.9	46.9	83.9	136.2	173.6
2006 5/.....	17.5	*	1.0	0.6	1.3	2.2	5.4	18.8	47.5	87.4	139.5	188.7
2005 5/.....	17.7	*	*	1.0	1.3	2.4	5.6	19.9	47.3	87.2	146.5	184.6
2004 5/.....	17.8	*	*	0.9	1.6	1.8	6.2	19.8	48.6	91.9	149.1	188.2
2003 5/.....	17.9	*	0.9	0.8	1.2	2.4	5.9	21.1	49.0	90.0	151.8	206.4
2002.....	18.7	*	1.1	0.8	1.3	2.4	6.5	20.5	51.8	97.9	159.6	223.2
2001.....	18.9	*	*	0.7	1.3	2.4	6.1	22.2	54.0	101.8	158.4	218.5
2000.....	18.9	*	0.9	0.6	1.5	1.7	7.7	22.6	52.8	101.2	163.6	211.1
1999.....	20.2	*	*	0.8	1.5	2.0	7.6	23.6	60.6	105.9	172.7	244.8

Non-Hispanic black, male

2007 5/.....	18.2	*	*	*	1.4	2.7	6.2	22.6	59.4	102.9	166.7	210.3
2006 5/.....	18.1	*	*	0.6	1.4	2.5	5.6	23.0	57.6	103.9	180.5	216.2
2005 5/.....	18.6	*	*	1.1	1.5	3.0	5.9	23.2	60.3	112.9	175.5	221.7
2004 5/.....	19.0	*	*	0.9	2.0	1.9	6.7	24.8	62.5	117.9	185.7	223.5
2003 5/.....	18.8	*	*	0.8	1.5	2.4	6.6	25.3	62.2	113.6	189.9	239.7
2002.....	19.4	*	*	0.9	1.8	2.7	7.6	24.5	64.7	119.0	197.8	265.8
2001.....	20.0	*	*	0.8	1.8	2.8	6.3	27.1	69.0	128.4	195.0	287.6
2000.....	20.0	*	*	0.7	2.0	1.9	8.7	26.9	69.1	123.8	212.0	253.6
1999.....	21.3	*	*	0.7	2.0	2.4	8.5	29.7	77.5	134.5	208.6	293.6

Non-Hispanic black, female

2007 5/.....	16.5	*	*	0.7	1.1	1.6	5.1	15.7	36.8	70.3	118.9	158.4
2006 5/.....	16.9	*	*	*	1.1	2.0	5.3	15.2	39.4	75.7	116.3	177.5
2005 5/.....	16.8	*	*	0.8	1.1	1.9	5.3	17.1	36.8	68.8	130.1	169.7
2004 5/.....	16.7	*	*	1.0	1.3	1.8	5.7	15.6	37.4	73.5	128.5	174.3
2003 5/.....	17.1	*	*	0.8	0.8	2.3	5.3	17.5	38.5	73.2	130.4	193.6
2002.....	18.0	*	*	0.8	0.8	2.1	5.5	17.1	41.4	83.0	137.9	206.9
2001.....	17.8	*	*	0.6	0.9	2.0	5.8	18.0	42.0	83.0	137.8	192.9
2000.....	17.8	*	*	*	1.0	1.5	6.8	18.9	39.6	85.2	136.3	195.4
1999.....	19.2	*	*	0.9	1.0	1.7	6.9	18.3	47.0	85.8	152.7	226.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes

2007.....	4.7	1.4	0.4	0.2	0.2	0.3	0.8	1.6	4.3	12.4	36.2	76.4
2006.....	4.7	1.6	0.4	0.2	0.2	0.4	0.7	1.8	4.2	12.5	36.4	78.0
2005.....	4.6	1.4	0.3	0.2	0.2	0.4	0.8	1.7	4.3	12.6	35.2	76.7
2004.....	4.6	1.7	0.3	0.2	0.3	0.4	0.7	1.7	4.5	12.7	35.8	77.7
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
2002.....	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000.....	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999.....	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9

All origins 3/, male

2007.....	4.9	1.1	0.3	0.2	0.2	0.4	0.8	1.8	5.3	15.2	48.4	95.7
2006.....	4.7	1.6	0.3	0.2	0.2	0.3	0.8	2.0	4.8	15.7	47.5	92.3
2005.....	4.7	1.0	0.4	0.2	0.2	0.4	0.8	1.9	5.2	15.7	45.8	92.7
2004.....	4.7	1.7	0.3	0.2	0.3	0.5	0.8	1.9	5.4	15.7	45.6	98.9
2003.....	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001.....	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000.....	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999.....	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2

All origins 3/, female

2007.....	4.5	1.7	0.4	0.2	0.2	0.2	0.7	1.4	3.4	10.0	27.8	67.2
2006.....	4.7	1.6	0.5	0.2	0.2	0.4	0.7	1.6	3.6	9.8	28.8	71.4
2005.....	4.6	1.8	0.3	0.2	0.2	0.4	0.8	1.5	3.6	10.0	28.0	69.4
2004.....	4.6	1.8	0.3	0.2	0.2	0.4	0.7	1.5	3.7	10.2	29.2	68.1
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
2002.....	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000.....	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999.....	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, both sexes

2007.....	1.5	1.9	*	*	*	0.2	0.5	1.2	3.1	7.4	24.7	46.5
2006.....	1.6	*	*	*	0.3	0.3	0.5	1.4	3.5	8.9	25.0	48.5
2005.....	1.3	*	*	*	*	0.3	0.4	1.3	3.2	8.1	20.8	34.2
2004.....	1.5	*	*	*	*	0.4	0.5	1.3	3.8	8.7	23.4	48.5
2003.....	1.5	*	*	*	0.3	0.4	0.4	1.1	3.1	8.7	28.9	48.2
2002.....	1.5	*	*	0.4	*	*	0.6	1.5	3.0	8.9	24.7	48.8
2001.....	1.5	*	*	0.3	*	0.3	0.7	1.5	3.6	8.8	23.1	57.5
2000.....	1.5	*	*	*	*	0.4	0.7	1.4	3.1	9.1	24.3	70.3
1999.....	1.5	*	*	*	*	0.4	0.6	1.3	3.7	10.5	23.1	51.9

Hispanic, male

2007.....	1.4	*	*	*	*	*	*	1.3	3.6	8.2	25.4	45.7
2006.....	1.6	*	*	*	*	*	*	1.8	4.1	11.2	30.4	53.0
2005.....	1.4	*	*	*	*	*	*	1.5	3.9	10.7	27.4	37.4
2004.....	1.5	*	*	*	*	*	*	1.6	4.1	9.7	27.9	57.4
2003.....	1.5	*	*	*	*	0.5	*	1.2	3.9	10.0	33.7	52.3
2002.....	1.4	*	*	*	*	*	*	1.4	3.7	10.3	31.6	56.9
2001.....	1.5	*	*	*	*	*	0.8	1.8	4.0	10.0	30.0	70.5
2000.....	1.3	*	*	*	*	*	*	*	3.1	9.7	26.6	86.7
1999.....	1.3	*	*	*	*	*	0.8	1.8	3.9	10.3	24.5	49.6

Hispanic, female

2007.....	1.7	*	*	*	*	*	*	1.0	2.6	6.7	24.2	46.9
2006.....	1.7	*	*	*	*	*	*	0.9	3.0	7.0	21.2	46.1
2005.....	1.3	*	*	*	*	*	*	1.0	2.5	6.0	16.3	32.6
2004.....	1.5	*	*	*	*	*	*	1.0	3.5	7.8	20.3	43.9
2003.....	1.6	*	*	*	*	*	*	1.0	2.3	7.6	25.5	46.1
2002.....	1.5	*	*	*	*	*	*	1.5	2.4	7.8	20.0	44.8
2001.....	1.5	*	*	*	*	*	*	1.2	3.3	7.9	18.4	51.2
2000.....	1.7	*	*	*	*	*	0.9	1.6	3.1	8.6	22.8	62.3
1999.....	1.6	*	*	*	*	*	*	*	3.5	10.7	22.2	53.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 4/, both sexes

2007.....	5.3	1.2	0.3	0.2	0.2	0.4	0.8	1.6	4.4	12.8	36.9	77.9
2006.....	5.2	1.5	0.4	0.2	0.2	0.4	0.8	1.9	4.2	12.8	37.1	79.5
2005.....	5.2	1.3	0.3	0.2	0.2	0.4	0.9	1.8	4.4	12.9	36.0	78.6
2004.....	5.1	1.7	0.4	0.2	0.3	0.4	0.8	1.7	4.6	12.9	36.4	78.9
2003.....	5.2	1.5	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.1	36.8	81.8
2002.....	5.1	1.6	0.4	0.2	0.2	0.5	0.8	1.8	4.6	13.1	36.6	79.8
2001.....	5.3	1.4	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.4	37.5	83.5
2000.....	5.3	1.7	0.4	0.2	0.3	0.4	0.8	1.9	5.2	14.1	38.1	85.2
1999.....	5.2	1.6	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.8	37.0	83.8

Non-Hispanic 4/, male

2007.....	5.5	*	0.3	0.2	0.2	0.5	0.9	1.8	5.4	15.8	49.8	98.5
2006.....	5.3	1.6	0.3	0.2	0.2	0.3	0.8	2.1	4.9	16.0	48.5	94.5
2005.....	5.2	*	0.4	0.2	0.3	0.4	0.9	2.0	5.2	16.1	46.8	95.4
2004.....	5.2	1.5	0.4	0.2	0.3	0.5	0.8	1.9	5.4	16.1	46.5	101.0
2003.....	5.2	1.6	0.4	0.2	0.2	0.5	0.9	1.9	5.2	16.3	46.7	103.9
2002.....	5.1	1.3	0.4	0.2	0.3	0.5	0.9	1.9	5.4	16.2	46.4	102.9
2001.....	5.3	1.5	0.4	0.2	0.3	0.4	0.9	2.1	5.6	17.8	48.9	109.0
2000.....	5.3	1.8	0.5	0.2	0.3	0.4	0.9	2.2	5.8	17.5	48.3	109.4
1999.....	5.3	1.6	0.5	0.2	0.3	0.5	0.9	2.0	5.9	18.8	46.9	111.4

Non-Hispanic 4/, female

2007.....	5.0	1.5	0.3	0.2	0.2	0.3	0.8	1.5	3.5	10.3	28.1	68.2
2006.....	5.2	1.4	0.5	0.2	0.2	0.4	0.7	1.7	3.6	10.1	29.2	72.5
2005.....	5.1	1.8	*	0.2	0.2	0.4	0.9	1.6	3.7	10.3	28.6	71.0
2004.....	5.0	1.9	0.4	0.2	0.3	0.4	0.7	1.5	3.7	10.3	29.7	69.0
2003.....	5.1	1.3	*	0.2	0.2	0.4	0.7	1.6	3.9	10.4	30.2	72.1
2002.....	5.1	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.8	10.5	30.0	69.8
2001.....	5.2	1.4	0.4	0.3	0.3	0.4	0.8	1.7	4.5	11.5	30.1	72.9
2000.....	5.3	1.7	*	0.3	0.2	0.4	0.8	1.6	4.6	11.4	31.4	75.4
1999.....	5.1	1.7	0.4	0.3	0.2	0.4	0.7	1.7	3.9	11.5	30.6	72.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic white, both sexes

2007 5/.....	5.9	1.2	0.4	0.2	0.2	0.3	0.8	1.6	4.4	13.0	38.2	81.2
2006 5/.....	5.9	1.5	0.5	0.2	0.2	0.3	0.7	1.8	4.2	13.2	38.2	81.6
2005 5/.....	5.8	1.2	0.3	0.2	0.2	0.4	0.9	1.7	4.4	13.2	37.2	81.7
2004 5/.....	5.7	1.8	0.4	0.2	0.3	0.4	0.7	1.6	4.4	13.2	37.4	81.6
2003 5/.....	5.8	1.4	0.3	0.2	0.2	0.5	0.8	1.7	4.5	13.4	38.1	84.4
2002.....	5.7	1.8	0.4	0.2	0.2	0.4	0.8	1.8	4.6	13.4	37.5	81.8
2001.....	5.9	1.3	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.7	38.6	84.8
2000.....	5.9	1.8	0.4	0.2	0.3	0.4	0.8	1.8	5.2	14.2	39.1	87.6
1999.....	5.7	1.5	0.4	0.3	0.2	0.4	0.7	1.8	4.8	15.0	37.8	86.3

Non-Hispanic white, male

2007 5/.....	6.2	*	*	0.2	0.2	0.4	0.9	1.8	5.3	15.8	51.6	102.5
2006 5/.....	6.0	*	*	0.2	0.2	0.3	0.8	2.1	4.9	16.5	50.1	96.0
2005 5/.....	5.9	*	0.4	0.2	0.3	0.5	0.9	1.9	5.2	16.6	48.7	100.0
2004 5/.....	5.9	*	0.4	0.2	0.3	0.5	0.8	1.8	5.3	16.5	47.7	105.6
2003 5/.....	5.9	1.7	*	0.2	0.3	0.5	0.9	1.9	5.1	16.8	48.4	107.9
2002.....	5.8	*	*	0.2	0.2	0.5	1.0	1.9	5.4	16.7	47.6	106.0
2001.....	6.0	*	*	0.2	0.2	0.5	0.9	2.1	5.7	18.3	50.1	111.4
2000.....	5.9	1.8	0.5	0.2	0.3	0.4	0.8	2.1	5.8	17.4	49.9	113.1
1999.....	5.9	*	*	0.2	0.2	0.4	0.9	2.0	5.9	19.2	48.1	116.8

Non-Hispanic white, female

2007 5/.....	5.7	*	*	0.2	0.2	0.3	0.8	1.4	3.5	10.5	28.8	71.1
2006 5/.....	5.8	*	0.6	*	0.2	0.4	0.6	1.6	3.5	10.3	29.9	74.9
2005 5/.....	5.7	*	*	0.2	0.2	0.4	0.8	1.5	3.6	10.3	29.3	73.4
2004 5/.....	5.6	2.0	*	0.2	0.2	0.3	0.6	1.4	3.6	10.4	30.4	70.9
2003 5/.....	5.8	*	*	*	0.2	0.5	0.6	1.5	4.0	10.4	31.1	74.0
2002.....	5.6	2.1	*	*	0.2	0.3	0.7	1.6	3.9	10.7	30.7	71.4
2001.....	5.8	*	*	0.3	0.3	0.4	0.7	1.7	4.5	11.6	30.9	73.7
2000.....	5.9	1.8	*	0.3	0.2	0.4	0.8	1.5	4.7	11.5	32.1	77.3
1999.....	5.6	*	*	0.3	0.2	0.4	0.6	1.5	3.7	11.5	31.1	74.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic black, both sexes

2007 5/.....	3.1	*	*	0.3	*	0.5	1.2	2.1	5.3	13.9	30.5	50.6
2006 5/.....	3.2	*	*	*	0.4	0.6	1.2	2.6	5.6	11.9	32.2	63.4
2005 5/.....	3.1	*	*	*	*	0.5	1.4	2.5	5.5	13.1	28.4	55.7
2004 5/.....	3.2	*	*	0.3	*	0.7	1.2	2.3	6.3	13.0	32.0	57.9
2003 5/.....	3.0	*	*	*	*	0.5	1.2	2.5	5.3	13.3	29.6	58.9
2002.....	3.1	*	*	*	*	0.6	1.2	2.5	5.6	12.8	31.8	64.4
2001.....	3.2	*	*	0.4	0.4	*	1.1	2.1	6.0	14.5	31.9	72.7
2000.....	3.2	*	*	*	*	0.6	1.1	2.7	5.6	15.1	31.3	64.2
1999.....	3.3	*	*	*	0.4	0.6	1.2	2.6	6.4	15.9	33.1	62.7

Non-Hispanic black, male

2007 5/.....	3.2	*	*	*	*	*	1.3	2.2	7.2	18.1	38.5	63.7
2006 5/.....	3.1	*	*	*	*	*	0.9	2.4	6.5	15.1	42.3	82.5
2005 5/.....	2.8	*	*	*	*	*	1.4	2.7	6.3	14.2	34.5	56.1
2004 5/.....	3.1	*	*	*	*	*	1.3	2.6	8.0	15.7	39.9	62.0
2003 5/.....	3.0	*	*	*	*	*	1.2	2.5	6.9	16.5	36.6	68.2
2002.....	2.9	*	*	*	*	*	1.0	2.2	7.3	14.3	39.3	81.0
2001.....	3.1	*	*	*	*	*	1.1	1.9	6.6	17.6	42.3	87.8
2000.....	3.1	*	*	*	*	*	1.2	2.8	7.0	19.5	36.4	76.3
1999.....	3.1	*	*	*	*	0.8	1.2	2.4	7.3	19.2	38.8	64.2

Non-Hispanic black, female

2007 5/.....	3.0	*	*	*	*	*	1.1	1.9	3.8	10.9	26.0	45.2
2006 5/.....	3.3	*	*	*	*	*	1.4	2.7	4.9	9.6	26.5	55.7
2005 5/.....	3.3	*	*	*	*	*	1.5	2.3	4.9	12.3	24.9	55.5
2004 5/.....	3.3	*	*	*	*	0.8	1.1	2.1	4.9	11.2	27.5	56.2
2003 5/.....	3.1	*	*	*	*	*	1.2	2.5	4.1	11.1	25.6	55.3
2002.....	3.4	*	*	*	*	0.8	1.4	2.7	4.3	11.7	27.5	58.1
2001.....	3.3	*	*	*	*	*	1.0	2.2	5.4	12.2	26.1	67.2
2000.....	3.3	*	*	*	*	0.8	1.0	2.6	4.5	12.1	28.5	59.7
1999.....	3.5	*	*	*	*	*	1.2	2.8	5.7	13.6	29.9	62.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Anemias (D50-D64)

##### All origins 3/, both sexes

2007.....	1.6	*	*	0.1	0.2	0.3	0.4	0.6	1.0	2.9	9.4	36.1
2006.....	1.3	*	0.2	0.1	0.2	0.3	0.4	0.5	1.0	2.1	7.5	30.0
2005.....	1.6	*	0.1	0.1	0.2	0.4	0.4	0.6	1.0	2.7	9.1	36.6
2004.....	1.6	*	0.2	0.1	0.3	0.4	0.5	0.6	1.1	2.6	8.6	38.1
2003.....	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002.....	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001.....	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000.....	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999.....	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8

##### All origins 3/, male

2007.....	1.3	*	*	0.1	0.2	0.3	0.5	0.6	1.0	3.0	10.1	34.3
2006.....	1.1	*	*	*	0.3	0.3	0.4	0.5	0.9	2.1	7.9	29.6
2005.....	1.2	*	*	*	0.3	0.4	0.4	0.5	1.0	2.6	9.5	33.5
2004.....	1.3	*	*	0.1	0.3	0.4	0.5	0.5	1.2	2.7	8.9	36.5
2003.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000.....	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999.....	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4

##### All origins 3/, female

2007.....	1.9	*	*	*	0.2	0.3	0.4	0.6	1.0	2.8	8.9	36.9
2006.....	1.6	*	*	*	0.2	0.4	0.5	0.6	1.0	2.2	7.1	30.2
2005.....	1.9	*	*	*	0.2	0.4	0.4	0.7	1.0	2.7	8.9	38.0
2004.....	1.8	*	*	0.1	0.3	0.4	0.4	0.6	1.0	2.5	8.4	38.8
2003.....	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002.....	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001.....	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000.....	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999.....	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Anemias (D50-D64)

##### Non-Hispanic 4/, both sexes

2007.....	1.8	*	*	0.1	0.2	0.4	0.5	0.6	1.0	3.0	9.5	37.0
2006.....	1.5	*	0.2	0.1	0.2	0.4	0.5	0.6	1.0	2.2	7.5	30.3
2005.....	1.7	*	*	0.1	0.3	0.5	0.5	0.6	1.0	2.7	9.3	37.2
2004.....	1.7	*	0.2	0.1	0.3	0.5	0.5	0.6	1.1	2.6	8.7	38.5
2003.....	1.8	0.6	*	0.1	0.3	0.5	0.4	0.6	1.0	2.7	8.9	41.2
2002.....	1.8	*	*	0.1	0.3	0.5	0.5	0.7	1.0	2.7	9.2	40.5
2001.....	1.8	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.5
2000.....	1.8	*	0.3	0.1	0.2	0.4	0.5	0.6	1.1	2.8	9.5	43.2
1999.....	1.8	*	0.2	0.1	0.4	0.4	0.5	0.7	1.0	2.6	9.6	44.0

##### Non-Hispanic 4/, male

2007.....	1.5	*	*	*	0.3	0.4	0.5	0.6	1.0	3.0	10.1	35.0
2006.....	1.2	*	*	*	0.3	0.4	0.5	0.5	1.0	2.1	8.1	30.2
2005.....	1.4	*	*	*	0.3	0.5	0.5	0.6	1.0	2.7	9.7	34.1
2004.....	1.4	*	*	*	0.4	0.5	0.5	0.6	1.2	2.8	9.0	37.1
2003.....	1.4	*	*	0.1	0.3	0.5	0.5	0.7	1.2	3.1	9.4	38.3
2002.....	1.4	*	*	0.1	0.4	0.6	0.4	0.6	1.0	2.9	9.0	37.6
2001.....	1.4	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	10.0	40.8
2000.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	2.8	9.6	41.0
1999.....	1.4	*	*	0.1	0.4	0.5	0.5	0.7	1.0	2.6	10.1	43.3

##### Non-Hispanic 4/, female

2007.....	2.1	*	*	*	0.2	0.4	0.5	0.6	1.0	2.9	9.0	38.0
2006.....	1.7	*	*	*	0.2	0.4	0.5	0.6	1.0	2.2	7.2	30.4
2005.....	2.1	*	*	*	0.3	0.4	0.5	0.7	1.1	2.6	9.0	38.6
2004.....	2.0	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.5	8.5	39.1
2003.....	2.1	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.6	42.4
2002.....	2.1	*	*	0.2	0.2	0.4	0.5	0.7	1.0	2.6	9.3	41.8
2001.....	2.2	*	*	0.2	0.3	0.4	0.6	0.6	1.0	2.6	9.7	43.1
2000.....	2.2	*	*	*	0.3	0.4	0.5	0.6	1.0	2.8	9.5	44.1
1999.....	2.1	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.3	44.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Anemias (D50-D64)

Non-Hispanic white, both sexes

2007 5/.....	1.8	*	*	*	*	0.1	0.1	0.3	0.7	2.7	9.2	37.4
2006 5/.....	1.4	*	*	*	*	*	0.1	0.3	0.7	2.0	7.4	30.0
2005 5/.....	1.7	*	*	*	*	*	0.1	0.3	0.8	2.3	8.9	37.2
2004 5/.....	1.6	*	*	*	0.1	0.1	0.2	0.3	0.7	2.5	8.6	38.1
2003 5/.....	1.6	*	*	*	*	0.1	0.1	0.3	0.6	2.4	8.6	41.0
2002.....	1.6	*	*	*	0.1	0.1	0.1	0.3	0.7	2.4	8.7	39.9
2001.....	1.7	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	42.1
2000.....	1.7	*	*	*	*	*	0.1	0.3	0.8	2.5	9.3	42.9
1999.....	1.6	*	*	*	0.1	*	0.1	0.3	0.6	2.4	9.4	43.2

Non-Hispanic white, male

2007 5/.....	1.4	*	*	*	*	*	*	0.3	0.7	2.8	9.9	35.2
2006 5/.....	1.1	*	*	*	*	*	*	0.2	0.7	2.0	7.9	30.6
2005 5/.....	1.2	*	*	*	*	*	*	0.3	0.8	2.3	9.2	34.3
2004 5/.....	1.3	*	*	*	*	*	0.2	0.3	0.8	2.7	8.6	37.3
2003 5/.....	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.2	38.1
2002.....	1.2	*	*	*	*	0.2	*	0.3	0.7	2.5	8.5	36.3
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.8	41.2
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	9.3	39.9
1999.....	1.2	*	*	*	*	*	*	0.3	0.5	2.4	9.9	43.4

Non-Hispanic white, female

2007 5/.....	2.1	*	*	*	*	*	*	0.3	0.7	2.6	8.8	38.5
2006 5/.....	1.7	*	*	*	*	*	*	0.3	0.8	2.0	7.1	29.7
2005 5/.....	2.1	*	*	*	*	*	*	0.4	0.9	2.3	8.8	38.5
2004 5/.....	2.0	*	*	*	*	*	*	0.2	0.7	2.3	8.6	38.4
2003 5/.....	2.0	*	*	*	*	*	*	0.2	0.5	2.1	8.2	42.2
2002.....	2.1	*	*	*	*	*	*	0.3	0.7	2.4	8.8	41.4
2001.....	2.1	*	*	*	*	*	0.2	0.3	0.8	2.4	9.2	42.4
2000.....	2.1	*	*	*	*	*	*	0.3	0.7	2.4	9.4	44.1
1999.....	2.0	*	*	*	*	*	*	0.3	0.7	2.4	9.1	43.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Anemias (D50-D64)

##### Non-Hispanic black, both sexes

2007 5/.....	2.5	*	*	*	1.0	1.9	2.6	2.7	3.1	6.0	13.3	36.1
2006 5/.....	2.4	*	*	*	1.0	2.1	2.6	2.6	3.2	4.2	10.5	39.3
2005 5/.....	2.7	*	*	*	1.2	2.5	2.6	2.8	3.0	6.3	14.9	43.4
2004 5/.....	2.7	*	*	0.4	1.5	2.4	2.5	2.8	4.1	4.8	11.8	48.9
2003 5/.....	2.8	*	*	0.4	1.3	2.2	2.4	3.1	4.3	5.6	14.5	50.3
2002.....	2.9	*	*	0.4	1.3	2.5	2.5	3.1	3.7	6.1	16.2	52.9
2001.....	2.8	*	*	0.4	1.2	2.3	2.8	3.1	3.0	5.5	16.1	49.3
2000.....	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.5	5.5	13.4	51.8
1999.....	2.9	*	0.9	0.5	1.5	2.7	2.6	3.2	4.3	5.0	13.4	53.0

##### Non-Hispanic black, male

2007 5/.....	2.4	*	*	*	1.1	2.1	3.0	2.9	3.2	5.9	14.6	34.0
2006 5/.....	2.1	*	*	*	1.1	2.1	2.5	2.4	3.7	4.1	12.0	30.2
2005 5/.....	2.5	*	*	*	1.4	2.7	3.0	2.6	2.6	6.6	17.4	38.1
2004 5/.....	2.6	*	*	*	1.6	2.8	2.9	2.5	4.9	5.0	14.0	43.7
2003 5/.....	2.8	*	*	*	1.3	2.3	2.6	3.4	5.6	8.0	15.3	50.1
2002.....	2.7	*	*	*	1.7	2.9	2.5	2.6	4.0	7.5	17.0	58.0
2001.....	2.6	*	*	*	1.4	3.0	2.5	3.6	3.2	6.8	15.3	42.7
2000.....	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5	4.6	13.9	56.4
1999.....	2.7	*	*	*	1.8	2.9	2.8	3.4	5.1	5.6	12.8	44.0

##### Non-Hispanic black, female

2007 5/.....	2.6	*	*	*	0.9	1.6	2.2	2.6	3.0	6.1	12.5	36.9
2006 5/.....	2.6	*	*	*	0.9	2.0	2.7	2.8	2.8	4.3	9.6	43.0
2005 5/.....	2.9	*	*	*	1.0	2.3	2.3	2.9	3.4	6.2	13.5	45.6
2004 5/.....	2.8	*	*	*	1.3	2.0	2.2	3.0	3.3	4.7	10.6	51.0
2003 5/.....	2.9	*	*	*	1.3	2.1	2.2	2.8	3.3	4.0	14.0	50.3
2002.....	3.0	*	*	*	1.0	2.0	2.6	3.5	3.4	5.1	15.8	51.0
2001.....	2.9	*	*	*	1.1	1.6	3.0	2.7	2.9	4.6	16.5	51.8
2000.....	3.1	*	*	*	1.1	2.5	2.8	2.9	3.5	6.1	13.2	50.2
1999.....	3.1	*	*	*	1.2	2.4	2.4	3.0	3.6	4.7	13.7	56.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

All origins 3/, both sexes

2007.....	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.6	78.1	162.7	276.2
2006.....	24.2	*	*	0.1	0.4	1.7	4.8	13.2	36.2	81.8	166.8	285.2
2005.....	25.3	*	*	0.1	0.5	1.5	4.7	13.4	37.2	86.8	177.2	312.1
2004.....	24.9	*	*	0.1	0.4	1.5	4.6	13.4	37.1	87.2	176.9	307.0
2003.....	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002.....	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001.....	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2

All origins 3/, male

2007.....	23.9	*	*	*	0.4	1.7	5.7	16.3	42.1	93.2	185.9	307.4
2006.....	24.4	*	*	*	0.4	1.8	5.9	16.5	43.8	96.2	194.1	321.0
2005.....	25.0	*	*	*	0.5	1.8	5.8	16.6	44.9	99.7	202.7	344.4
2004.....	24.4	*	*	*	0.4	1.7	5.7	16.5	44.0	100.0	202.5	333.9
2003.....	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002.....	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001.....	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000.....	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999.....	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7

All origins 3/, female

2007.....	23.5	*	*	*	0.3	1.3	3.5	10.1	27.5	65.3	146.7	261.4
2006.....	24.0	*	*	*	0.3	1.5	3.7	9.9	29.1	69.7	148.1	268.4
2005.....	25.7	*	*	0.1	0.4	1.2	3.5	10.3	30.1	75.9	159.9	297.2
2004.....	25.4	*	*	0.1	0.4	1.3	3.5	10.4	30.7	76.4	159.6	294.9
2003.....	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002.....	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001.....	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000.....	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999.....	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

Hispanic, both sexes

2007.....	14.1	*	*	*	*	0.7	3.3	13.3	42.0	107.3	232.5	337.5
2006.....	14.2	*	*	*	*	0.9	2.8	13.2	42.6	118.0	228.9	369.7
2005.....	15.6	*	*	*	*	0.9	3.4	14.8	49.1	126.4	267.0	404.7
2004.....	14.5	*	*	*	*	0.8	3.0	13.3	48.3	124.9	258.4	365.8
2003.....	15.5	*	*	*	*	0.9	3.3	15.5	53.7	131.8	272.1	428.1
2002.....	15.3	*	*	*	*	0.8	3.5	14.0	52.1	134.7	285.6	437.9
2001.....	15.3	*	*	*	*	0.7	3.1	14.9	53.2	141.2	279.2	489.6
2000.....	15.1	*	*	*	*	1.0	3.0	14.0	55.5	147.9	278.5	465.1
1999.....	15.3	*	*	*	*	0.7	3.2	13.9	56.6	149.7	285.2	493.9

Hispanic, male

2007.....	13.6	*	*	*	*	0.9	4.2	15.9	51.1	127.2	244.9	324.2
2006.....	13.7	*	*	*	*	0.8	3.3	16.2	49.9	135.6	257.7	381.2
2005.....	14.9	*	*	*	*	0.9	4.5	17.5	58.2	143.9	293.0	404.7
2004.....	13.4	*	*	*	*	0.9	3.4	16.0	56.9	134.2	279.3	325.0
2003.....	14.4	*	*	*	*	0.8	3.9	18.1	62.4	146.3	297.2	426.4
2002.....	13.9	*	*	*	*	0.9	4.1	16.7	58.9	143.7	306.9	425.0
2001.....	13.6	*	*	*	*	0.8	3.9	18.5	58.7	151.4	285.3	422.8
2000.....	13.8	*	*	*	*	1.4	3.6	15.5	64.7	160.8	299.4	449.4
1999.....	13.4	*	*	*	*	0.6	3.3	17.2	62.5	155.8	293.0	468.0

Hispanic, female

2007.....	14.6	*	*	*	*	0.5	2.4	10.6	33.7	91.1	223.8	344.8
2006.....	14.7	*	*	*	*	1.0	2.1	10.1	35.8	103.7	208.7	363.7
2005.....	16.3	*	*	*	*	0.8	2.1	12.0	40.7	112.2	248.9	404.7
2004.....	15.7	*	*	*	*	0.8	2.6	10.6	40.5	117.5	244.0	386.8
2003.....	16.6	*	*	*	*	0.9	2.7	13.0	45.8	120.3	254.9	428.9
2002.....	16.7	*	*	*	*	0.7	2.9	11.4	46.0	127.6	270.9	444.3
2001.....	17.1	*	*	*	*	0.7	2.2	11.4	48.3	133.2	275.0	522.2
2000.....	16.5	*	*	*	*	*	2.4	12.6	47.4	137.7	264.5	472.8
1999.....	17.3	*	*	*	*	0.8	3.1	10.7	51.4	144.9	280.0	506.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

Non-Hispanic 4/, both sexes

2007.....	25.3	*	*	*	0.4	1.7	4.8	13.1	33.8	75.7	157.9	272.7
2006.....	25.9	*	*	0.1	0.4	1.8	5.1	13.1	35.5	78.9	162.6	280.5
2005.....	26.9	*	*	0.1	0.5	1.7	4.9	13.2	36.1	83.7	171.6	307.2
2004.....	26.5	*	*	0.1	0.4	1.6	4.8	13.3	36.0	84.2	172.0	303.5
2003.....	27.0	*	*	0.1	0.4	1.8	4.8	13.6	37.1	87.7	175.8	312.0
2002.....	26.9	*	*	0.1	0.5	1.8	5.0	13.6	36.4	88.2	177.2	315.0
2001.....	26.4	*	*	0.1	0.4	1.7	4.5	13.4	36.5	87.7	176.4	314.8
2000.....	25.9	*	*	0.1	0.5	1.7	4.4	13.0	36.4	86.8	174.8	313.4
1999.....	25.7	*	*	0.1	0.4	1.6	4.4	12.8	36.8	88.1	173.1	310.0

Non-Hispanic 4/, male

2007.....	25.8	*	*	*	0.5	2.0	5.9	16.3	41.2	90.4	181.7	306.4
2006.....	26.3	*	*	*	0.5	2.1	6.3	16.5	43.1	93.0	189.9	317.0
2005.....	26.8	*	*	*	0.6	2.1	6.0	16.4	43.7	96.3	196.9	340.9
2004.....	26.2	*	*	*	0.4	1.9	6.1	16.5	42.7	97.3	197.7	333.5
2003.....	26.4	*	*	*	0.5	2.1	6.0	16.5	43.5	100.9	201.8	335.9
2002.....	25.8	*	*	0.1	0.5	2.2	6.2	16.3	43.1	98.2	196.6	344.0
2001.....	25.0	*	*	0.1	0.4	1.9	5.5	16.1	41.8	98.0	194.3	338.9
2000.....	24.2	*	*	*	0.5	2.0	5.3	15.2	40.7	95.0	194.6	339.7
1999.....	24.1	*	*	*	0.4	1.8	5.3	15.3	41.0	96.3	193.8	339.5

Non-Hispanic 4/, female

2007.....	24.9	*	*	*	0.4	1.4	3.7	10.0	26.9	63.1	141.5	256.8
2006.....	25.5	*	*	*	0.4	1.6	4.0	9.9	28.4	66.8	144.0	263.5
2005.....	27.1	*	*	0.1	0.5	1.3	3.8	10.1	29.1	73.0	154.4	291.9
2004.....	26.8	*	*	0.1	0.4	1.4	3.6	10.3	29.8	73.1	154.7	290.1
2003.....	27.6	*	*	*	0.3	1.5	3.6	10.9	31.2	76.7	158.5	301.5
2002.....	27.9	*	*	*	0.4	1.4	3.8	11.1	30.1	79.8	164.3	302.5
2001.....	27.8	*	*	*	0.3	1.4	3.6	10.9	31.6	79.1	164.6	304.7
2000.....	27.6	*	*	*	0.4	1.4	3.5	11.0	32.4	80.1	161.8	302.8
1999.....	27.3	*	*	*	0.4	1.4	3.6	10.4	32.8	81.4	159.6	298.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

Non-Hispanic white, both sexes

2007 5/.....	24.9	*	*	*	0.3	1.3	4.0	11.1	28.9	66.3	144.8	257.0
2006 5/.....	25.4	*	*	*	0.3	1.4	4.4	11.0	30.3	68.9	149.0	264.2
2005 5/.....	26.5	*	*	0.1	0.4	1.3	4.1	11.1	30.9	73.1	158.4	289.8
2004 5/.....	26.1	*	*	*	0.3	1.4	4.1	11.3	30.7	73.7	157.8	284.5
2003 5/.....	26.6	*	*	*	0.3	1.5	4.1	11.3	31.4	76.8	161.6	293.8
2002.....	26.4	*	*	*	0.4	1.5	4.3	11.3	30.6	76.8	163.3	296.4
2001.....	26.0	*	*	*	0.3	1.3	3.9	11.3	30.7	77.0	162.7	297.4
2000.....	25.4	*	*	*	0.4	1.4	3.9	10.7	30.9	76.1	159.9	295.5
1999.....	25.0	*	*	*	0.4	1.4	3.8	10.5	30.1	76.8	159.1	290.2

Non-Hispanic white, male

2007 5/.....	25.9	*	*	*	0.4	1.6	5.0	14.1	36.1	82.2	172.2	298.1
2006 5/.....	26.3	*	*	*	0.3	1.6	5.4	14.2	37.6	83.9	179.6	306.8
2005 5/.....	26.8	*	*	*	0.5	1.5	5.2	14.1	38.1	86.5	186.6	331.5
2004 5/.....	26.3	*	*	*	0.3	1.6	5.3	14.3	37.3	87.7	187.1	321.7
2003 5/.....	26.6	*	*	*	0.4	1.7	5.2	14.1	37.9	91.3	191.7	327.3
2002.....	26.0	*	*	*	0.4	1.9	5.5	13.8	37.5	88.9	187.4	338.4
2001.....	25.1	*	*	*	0.3	1.5	4.8	13.7	36.4	89.8	183.5	332.1
2000.....	24.3	*	*	*	0.5	1.7	4.8	13.0	35.4	86.6	184.5	330.4
1999.....	24.0	*	*	*	0.4	1.5	4.6	13.0	35.0	87.7	184.6	326.0

Non-Hispanic white, female

2007 5/.....	23.9	*	*	*	0.2	1.0	2.9	8.1	22.0	52.5	125.6	237.5
2006 5/.....	24.5	*	*	*	0.3	1.2	3.3	7.9	23.3	55.8	127.8	244.3
2005 5/.....	26.3	*	*	*	0.4	1.0	3.1	8.1	24.1	61.5	139.1	270.7
2004 5/.....	25.9	*	*	*	0.3	1.2	3.0	8.3	24.3	61.7	137.8	267.8
2003 5/.....	26.5	*	*	*	0.3	1.3	2.9	8.5	25.2	64.4	141.3	279.0
2002.....	26.8	*	*	*	0.4	1.2	3.2	8.8	24.1	66.5	147.0	278.3
2001.....	26.9	*	*	*	0.2	1.2	2.9	8.8	25.3	66.0	148.9	283.0
2000.....	26.5	*	*	*	0.3	1.1	3.1	8.6	26.6	67.3	143.8	281.4
1999.....	26.0	*	*	*	0.3	1.2	3.0	8.1	25.4	67.7	142.4	275.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

Non-Hispanic black, both sexes

2007 5/.....	32.4	*	*	*	0.9	3.9	10.3	27.6	75.4	159.1	302.2	451.8
2006 5/.....	33.6	*	*	*	0.8	4.4	10.6	28.0	79.3	170.0	316.4	475.9
2005 5/.....	34.4	*	*	*	1.2	4.1	9.8	28.5	81.3	180.6	325.0	513.4
2004 5/.....	34.4	*	*	*	1.0	3.4	9.8	28.8	81.9	182.6	336.4	536.5
2003 5/.....	35.0	*	*	*	0.7	4.0	9.9	30.8	87.5	188.9	340.0	532.2
2002.....	34.8	*	*	*	1.0	3.8	9.9	30.9	85.9	191.5	340.2	544.7
2001.....	34.1	*	*	*	1.0	3.8	9.1	30.1	86.5	190.8	337.9	533.8
2000.....	34.0	*	*	*	1.0	3.6	8.2	31.0	83.9	190.6	354.8	530.1
1999.....	34.1	*	*	*	0.8	3.1	9.1	31.0	92.0	194.0	336.1	530.9

Non-Hispanic black, male

2007 5/.....	29.9	*	*	*	0.6	4.4	12.2	33.2	87.8	175.7	317.1	404.0
2006 5/.....	31.7	*	*	*	1.0	5.3	13.4	34.3	92.6	191.3	334.2	468.2
2005 5/.....	31.8	*	*	*	1.2	5.0	12.0	34.1	96.2	197.4	346.5	471.0
2004 5/.....	31.3	*	*	*	1.0	4.0	12.2	34.0	93.4	201.9	344.3	508.1
2003 5/.....	30.9	*	*	*	0.9	4.9	11.9	34.6	97.8	201.2	342.8	457.1
2002.....	30.0	*	*	*	1.1	4.6	12.1	35.8	96.6	193.3	332.6	423.4
2001.....	29.4	*	*	*	0.9	5.0	10.5	35.0	93.8	189.5	345.1	452.7
2000.....	28.3	*	*	*	1.0	4.0	9.7	34.2	89.7	186.9	339.4	456.8
1999.....	28.6	*	*	*	*	3.4	10.7	35.3	95.4	189.7	317.7	506.3

Non-Hispanic black, female

2007 5/.....	34.6	*	*	*	1.1	3.4	8.6	22.8	65.4	147.2	293.7	471.6
2006 5/.....	35.3	*	*	*	0.7	3.7	8.1	22.6	68.6	154.7	306.2	479.1
2005 5/.....	36.7	*	*	*	1.1	3.3	7.8	23.7	69.4	168.6	312.9	530.4
2004 5/.....	37.2	*	*	*	1.0	2.9	7.8	24.3	72.6	168.7	332.0	547.7
2003 5/.....	38.7	*	*	*	*	3.2	8.1	27.6	79.2	180.2	338.5	561.3
2002.....	39.2	*	*	*	0.8	3.0	8.0	26.8	77.3	190.3	344.5	591.1
2001.....	38.4	*	*	*	1.0	2.8	7.8	25.9	80.6	191.7	333.9	563.8
2000.....	39.1	*	*	*	1.0	3.3	7.0	28.2	79.1	193.3	363.5	557.3
1999.....	39.0	*	*	*	0.9	2.7	7.7	27.2	89.2	197.0	346.5	540.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Nutritional deficiencies (E40-E64)

##### All origins 3/, both sexes

2007.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.7	6.2	24.4
2006.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.5	5.5	22.7
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.9	7.0	30.8
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.6	1.6	6.9	30.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
2002.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9

##### All origins 3/, male

2007.....	0.7	*	*	*	*	*	*	0.2	0.6	1.9	6.3	22.1
2006.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	5.6	18.7
2005.....	0.8	*	*	*	*	*	*	0.3	0.6	1.8	6.5	27.9
2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.8	27.4
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001.....	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4

##### All origins 3/, female

2007.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.6	6.2	25.5
2006.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.4	5.4	24.5
2005.....	1.4	*	*	*	*	*	0.1	0.3	0.4	1.9	7.4	32.1
2004.....	1.3	*	*	*	*	*	0.1	0.3	0.5	1.5	7.0	31.7
2003.....	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
2002.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Nutritional deficiencies (E40-E64)

##### Non-Hispanic 4/, both sexes

2007.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.8	6.3	24.8
2006.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.5	5.5	22.8
2005.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.9	7.0	31.1
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0	30.8
2003.....	1.3	*	*	*	*	*	0.1	0.3	0.6	1.9	7.8	35.7
2002.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.1	8.9	41.8
2001.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	8.8	43.0
2000.....	1.7	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.3
1999.....	1.7	*	*	*	*	*	0.1	0.3	0.7	2.4	10.3	53.9

##### Non-Hispanic 4/, male

2007.....	0.8	*	*	*	*	*	*	0.2	0.6	2.0	6.4	22.6
2006.....	0.7	*	*	*	*	*	*	0.2	0.6	1.6	5.6	18.9
2005.....	0.8	*	*	*	*	*	*	0.3	0.6	1.8	6.4	28.0
2004.....	0.8	*	*	*	*	*	*	0.3	0.6	1.7	7.0	28.0
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.0	7.9	30.7
2002.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.2	8.5	37.1
2001.....	1.0	*	*	*	*	*	*	0.3	0.7	2.2	8.5	34.8
2000.....	1.1	*	*	*	*	*	*	0.3	0.9	2.7	9.6	43.9
1999.....	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	10.4	48.1

##### Non-Hispanic 4/, female

2007.....	1.3	*	*	*	*	*	0.1	0.2	0.6	1.6	6.3	25.8
2006.....	1.2	*	*	*	*	*	0.1	0.3	0.7	1.4	5.4	24.6
2005.....	1.5	*	*	*	*	*	0.1	0.3	0.5	2.0	7.4	32.5
2004.....	1.4	*	*	*	*	*	0.1	0.3	0.5	1.6	7.0	32.1
2003.....	1.6	*	*	*	*	*	*	0.3	0.7	1.8	7.7	37.9
2002.....	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.1	43.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.6	2.0	8.9	46.4
2000.....	2.2	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.8
1999.....	2.2	*	*	*	*	*	0.2	0.3	0.6	2.1	10.3	56.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Nutritional deficiencies (E40-E64)

##### Non-Hispanic white, both sexes

2007 5/.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.6	6.0	23.7
2006 5/.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.4	5.3	22.2
2005 5/.....	1.3	*	*	*	*	*	0.1	0.3	0.5	1.8	6.5	30.3
2004 5/.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003 5/.....	1.4	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	34.4
2002.....	1.5	*	*	*	*	*	0.1	0.3	0.6	1.9	8.5	40.6
2001.....	1.5	*	*	*	*	*	0.1	0.2	0.6	2.0	8.2	42.2
2000.....	1.8	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	49.0
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.6	52.8

##### Non-Hispanic white, male

2007 5/.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.0	21.7
2006 5/.....	0.7	*	*	*	*	*	*	0.2	0.5	1.4	5.1	18.4
2005 5/.....	0.9	*	*	*	*	*	*	0.3	0.5	1.7	5.8	27.4
2004 5/.....	0.9	*	*	*	*	*	*	0.2	0.5	1.5	6.5	26.9
2003 5/.....	0.9	*	*	*	*	*	*	0.2	0.5	1.8	7.2	29.6
2002.....	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.0	36.5
2001.....	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.9	33.3
2000.....	1.2	*	*	*	*	*	*	0.2	0.7	2.3	9.0	43.0
1999.....	1.2	*	*	*	*	*	*	0.2	0.7	2.2	9.4	46.2

##### Non-Hispanic white, female

2007 5/.....	1.4	*	*	*	*	*	*	0.2	0.6	1.5	6.0	24.6
2006 5/.....	1.3	*	*	*	*	*	0.1	0.2	0.7	1.4	5.4	24.0
2005 5/.....	1.6	*	*	*	*	*	*	0.3	0.5	1.8	7.0	31.7
2004 5/.....	1.5	*	*	*	*	*	*	0.3	0.5	1.5	6.8	30.8
2003 5/.....	1.8	*	*	*	*	*	*	0.2	0.6	1.7	7.6	36.5
2002.....	2.0	*	*	*	*	*	*	0.3	0.6	1.8	8.8	42.3
2001.....	2.0	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.9
2000.....	2.3	*	*	*	*	*	0.2	0.3	0.6	2.3	10.6	51.5
1999.....	2.3	*	*	*	*	*	0.2	0.3	0.5	1.9	9.8	55.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional deficiencies (E40-E64)

Non-Hispanic black, both sexes

2007 5/.....	1.0	*	*	*	*	*	*	*	0.9	3.9	10.7	37.1
2006 5/.....	0.9	*	*	*	*	*	*	0.5	1.1	3.0	8.8	31.3
2005 5/.....	1.1	*	*	*	*	*	*	0.6	1.1	3.4	12.8	43.1
2004 5/.....	1.1	*	*	*	*	*	*	0.7	1.2	3.4	10.7	47.2
2003 5/.....	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.9
2002.....	1.4	*	*	*	*	*	*	0.4	1.8	4.3	14.7	60.2
2001.....	1.3	*	*	*	*	*	*	0.8	1.2	3.7	16.1	55.2
2000.....	1.5	*	*	*	*	*	*	0.6	1.7	4.8	16.3	67.4
1999.....	1.7	*	*	*	*	*	*	0.7	2.0	5.6	19.6	70.8

Non-Hispanic black, male

2007 5/.....	0.9	*	*	*	*	*	*	*	1.4	4.8	12.9	34.0
2006 5/.....	0.9	*	*	*	*	*	*	*	1.4	4.1	12.2	29.3
2005 5/.....	0.9	*	*	*	*	*	*	*	1.7	3.1	12.6	37.1
2004 5/.....	1.0	*	*	*	*	*	*	*	1.7	4.6	12.3	45.7
2003 5/.....	1.0	*	*	*	*	*	*	*	1.8	4.5	17.9	49.0
2002.....	1.1	*	*	*	*	*	*	*	2.9	4.5	16.4	45.9
2001.....	1.2	*	*	*	*	*	*	1.2	1.9	5.6	16.5	58.9
2000.....	1.3	*	*	*	*	*	*	*	2.8	6.6	17.9	64.6
1999.....	1.5	*	*	*	*	*	*	*	2.9	8.3	23.1	76.1

Non-Hispanic black, female

2007 5/.....	1.1	*	*	*	*	*	*	*	*	3.3	9.4	38.3
2006 5/.....	1.0	*	*	*	*	*	*	*	*	2.3	6.8	32.1
2005 5/.....	1.4	*	*	*	*	*	*	*	*	3.6	12.9	45.6
2004 5/.....	1.3	*	*	*	*	*	*	*	*	2.6	9.8	47.8
2003 5/.....	1.4	*	*	*	*	*	*	*	*	2.4	10.6	58.6
2002.....	1.6	*	*	*	*	*	*	*	*	4.1	13.8	65.6
2001.....	1.4	*	*	*	*	*	*	*	*	2.4	15.8	53.9
2000.....	1.8	*	*	*	*	*	*	*	*	3.6	15.5	68.5
1999.....	1.8	*	*	*	*	*	*	*	*	3.7	17.7	68.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Malnutrition (E40-E46)

##### All origins 3/, both sexes

2007.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.6	5.8	22.4
2006.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.2	20.9
2005.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.8	6.6	28.9
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.6	6.7	28.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000.....	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0

##### All origins 3/, male

2007.....	0.7	*	*	*	*	*	*	0.2	0.6	1.7	5.9	20.3
2006.....	0.6	*	*	*	*	*	*	0.2	0.5	1.4	5.2	17.5
2005.....	0.7	*	*	*	*	*	*	0.3	0.6	1.7	6.2	26.3
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.7	6.6	25.3
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000.....	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4

##### All origins 3/, female

2007.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.5	5.7	23.3
2006.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	22.4
2005.....	1.3	*	*	*	*	*	0.1	0.3	0.4	1.8	6.9	30.2
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003.....	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002.....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001.....	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000.....	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malnutrition (E40-E46)

Non-Hispanic 4/, both sexes

2007.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.7	5.9	22.7
2006.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	21.0
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.8	6.6	29.2
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.6	6.8	28.6
2003.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.8	7.5	33.7
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	8.3	38.7
2001.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.2	40.2
2000.....	1.5	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.5
1999.....	1.6	*	*	*	*	*	0.1	0.3	0.7	2.2	9.6	50.0

Non-Hispanic 4/, male

2007.....	0.8	*	*	*	*	*	*	0.2	0.6	1.8	6.0	20.7
2006.....	0.6	*	*	*	*	*	*	0.2	0.5	1.5	5.2	17.9
2005.....	0.8	*	*	*	*	*	*	0.3	0.6	1.7	6.1	26.5
2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.7	25.8
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	1.8	7.5	28.4
2002.....	1.0	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.3
2001.....	0.9	*	*	*	*	*	*	0.3	0.6	2.0	8.0	33.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.8	2.5	8.8	40.9
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	9.5	44.3

Non-Hispanic 4/, female

2007.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.6	5.9	23.6
2006.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	22.5
2005.....	1.4	*	*	*	*	*	0.1	0.3	0.4	1.9	6.9	30.5
2004.....	1.4	*	*	*	*	*	0.1	0.3	0.5	1.5	6.8	29.9
2003.....	1.5	*	*	*	*	*	*	0.2	0.6	1.7	7.5	36.0
2002.....	1.7	*	*	*	*	*	0.1	0.3	0.5	1.8	8.6	40.1
2001.....	1.7	*	*	*	*	*	*	0.3	0.6	1.9	8.3	43.2
2000.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.2	10.3	48.7
1999.....	2.0	*	*	*	*	*	0.2	0.3	0.5	2.0	9.7	52.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malnutrition (E40-E46)

Non-Hispanic white, both sexes

2007 5/.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.5	5.6	21.6
2006 5/.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.3	4.9	20.3
2005 5/.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.6	6.1	28.5
2004 5/.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.4	6.4	27.3
2003 5/.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.6	7.2	32.3
2002.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.4
2001.....	1.4	*	*	*	*	*	*	0.2	0.5	1.8	7.6	39.4
2000.....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.3
1999.....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.9	8.9	48.8

Non-Hispanic white, male

2007 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.5	5.6	19.7
2006 5/.....	0.7	*	*	*	*	*	*	0.2	0.4	1.3	4.7	17.2
2005 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.6	5.6	25.9
2004 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.4	6.2	24.6
2003 5/.....	0.9	*	*	*	*	*	*	0.2	0.4	1.6	6.9	27.3
2002.....	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.5	34.5
2001.....	0.9	*	*	*	*	*	*	0.2	0.5	1.7	7.4	31.7
2000.....	1.1	*	*	*	*	*	*	0.2	0.6	2.1	8.3	39.8
1999.....	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.6	42.0

Non-Hispanic white, female

2007 5/.....	1.3	*	*	*	*	*	*	0.2	0.6	1.5	5.6	22.5
2006 5/.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.3	5.1	21.8
2005 5/.....	1.5	*	*	*	*	*	*	0.3	0.4	1.7	6.5	29.7
2004 5/.....	1.5	*	*	*	*	*	*	0.3	0.5	1.4	6.6	28.5
2003 5/.....	1.7	*	*	*	*	*	*	0.2	0.6	1.6	7.4	34.5
2002.....	1.8	*	*	*	*	*	*	0.3	0.5	1.7	8.3	38.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.6	1.9	7.7	42.7
2000.....	2.2	*	*	*	*	*	0.1	0.2	0.6	2.0	10.0	47.4
1999.....	2.2	*	*	*	*	*	0.2	0.3	0.4	1.8	9.2	51.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malnutrition (E40-E46)

Non-Hispanic black, both sexes

2007 5/.....	1.0	*	*	*	*	*	*	*	0.9	3.9	10.2	35.5
2006 5/.....	0.9	*	*	*	*	*	*	0.5	1.0	3.0	8.2	30.0
2005 5/.....	1.1	*	*	*	*	*	*	0.6	1.1	3.3	12.3	41.0
2004 5/.....	1.1	*	*	*	*	*	*	0.6	1.1	3.3	10.5	46.3
2003 5/.....	1.2	*	*	*	*	*	*	0.7	1.3	3.1	13.0	54.1
2002.....	1.3	*	*	*	*	*	*	*	1.7	3.6	14.2	57.2
2001.....	1.2	*	*	*	*	*	*	0.6	1.1	3.4	15.0	52.1
2000.....	1.4	*	*	*	*	*	*	0.6	1.6	4.8	15.1	64.2
1999.....	1.6	*	*	*	*	*	*	0.6	1.8	5.5	18.1	66.0

Non-Hispanic black, male

2007 5/.....	0.9	*	*	*	*	*	*	*	*	4.8	12.7	34.0
2006 5/.....	0.8	*	*	*	*	*	*	*	*	4.1	11.1	29.3
2005 5/.....	0.8	*	*	*	*	*	*	*	1.7	3.1	12.1	34.3
2004 5/.....	0.9	*	*	*	*	*	*	*	1.6	4.4	12.0	43.7
2003 5/.....	1.0	*	*	*	*	*	*	*	1.8	4.3	17.3	46.9
2002.....	1.0	*	*	*	*	*	*	*	2.7	4.0	15.2	44.9
2001.....	1.1	*	*	*	*	*	*	1.0	*	5.0	15.9	56.6
2000.....	1.2	*	*	*	*	*	*	*	2.5	6.4	17.3	62.2
1999.....	1.4	*	*	*	*	*	*	*	2.6	8.0	20.6	74.9

Non-Hispanic black, female

2007 5/.....	1.1	*	*	*	*	*	*	*	*	3.2	8.8	36.2
2006 5/.....	0.9	*	*	*	*	*	*	*	*	2.3	6.5	30.3
2005 5/.....	1.3	*	*	*	*	*	*	*	*	3.5	12.4	43.7
2004 5/.....	1.2	*	*	*	*	*	*	*	*	2.5	9.7	47.4
2003 5/.....	1.4	*	*	*	*	*	*	*	*	2.3	10.6	57.0
2002.....	1.5	*	*	*	*	*	*	*	*	3.4	13.6	61.9
2001.....	1.3	*	*	*	*	*	*	*	*	2.3	14.5	50.5
2000.....	1.6	*	*	*	*	*	*	*	*	3.6	13.9	65.0
1999.....	1.7	*	*	*	*	*	*	*	*	3.7	16.6	62.6











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningitis (G00,G03)

All origins 3/, both sexes

2007.....	0.2	1.9	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.7	0.7
2006.....	0.2	1.5	0.1	*	0.1	0.1	0.2	0.2	0.2	0.4	0.7	0.9
2005.....	0.2	1.4	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.6	0.9
2004.....	0.2	1.6	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003.....	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002.....	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001.....	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000.....	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999.....	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5

All origins 3/, male

2007.....	0.2	1.6	*	*	*	*	0.2	0.3	0.3	0.5	0.8	*
2006.....	0.2	1.4	*	*	*	0.1	0.2	0.3	0.3	0.4	0.7	*
2005.....	0.2	1.4	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2004.....	0.2	1.5	*	*	*	0.1	0.2	0.4	0.3	0.4	0.7	*
2003.....	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002.....	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001.....	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000.....	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999.....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7

All origins 3/, female

2007.....	0.2	2.3	*	*	*	*	0.1	0.2	0.2	0.3	0.6	0.7
2006.....	0.2	1.6	*	*	*	*	0.2	0.2	0.2	0.4	0.6	0.8
2005.....	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.9
2004.....	0.2	1.7	*	*	*	*	0.1	0.2	0.3	0.6	0.6	0.7
2003.....	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002.....	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001.....	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000.....	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999.....	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Meningitis (G00,G03)

##### Non-Hispanic 4/, both sexes

2007.....	0.2	2.0	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.7	0.6
2006.....	0.2	1.6	*	*	*	0.1	0.2	0.3	0.2	0.4	0.7	0.9
2005.....	0.2	1.5	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.4	0.6	0.9
2004.....	0.2	1.8	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003.....	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002.....	0.3	2.1	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.6	1.2
2001.....	0.3	2.2	0.2	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.7	0.9
2000.....	0.3	2.7	0.3	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999.....	0.3	3.2	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.4

##### Non-Hispanic 4/, male

2007.....	0.2	1.7	*	*	*	*	0.2	0.3	0.3	0.5	0.8	*
2006.....	0.2	1.7	*	*	*	*	0.2	0.3	0.3	0.4	0.8	*
2005.....	0.3	1.5	*	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2004.....	0.3	1.8	*	*	*	*	0.2	0.4	0.3	0.4	0.7	*
2003.....	0.3	1.6	*	*	*	0.1	0.2	0.4	0.5	0.6	0.8	*
2002.....	0.3	2.1	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001.....	0.3	2.1	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000.....	0.3	2.7	*	*	*	*	0.2	0.3	0.5	0.8	1.2	*
1999.....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.1	*

##### Non-Hispanic 4/, female

2007.....	0.2	2.4	*	*	*	*	0.1	0.2	0.2	0.3	0.6	0.6
2006.....	0.2	1.6	*	*	*	*	0.2	0.2	0.2	0.4	0.6	0.8
2005.....	0.2	1.5	*	*	*	*	0.2	0.2	0.3	0.4	0.6	0.8
2004.....	0.2	1.9	*	*	*	*	*	0.2	0.3	0.6	0.6	0.7
2003.....	0.2	2.2	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0
2002.....	0.2	2.0	*	*	*	*	*	0.2	0.3	0.5	0.5	1.2
2001.....	0.2	2.3	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.7
2000.....	0.3	2.7	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.1
1999.....	0.3	3.3	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningitis (G00,G03)

Non-Hispanic white, both sexes

2007 5/.....	0.2	1.5	*	*	*	0.1	0.2	0.2	0.2	0.4	0.7	0.6
2006 5/.....	0.2	1.5	*	*	*	0.1	0.1	0.3	0.2	0.4	0.6	0.9
2005 5/.....	0.2	1.1	*	*	*	*	0.1	0.2	0.3	0.4	0.5	0.8
2004 5/.....	0.2	1.6	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.8
2003 5/.....	0.2	1.5	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0
2002.....	0.2	1.9	*	0.1	*	0.1	0.1	0.2	0.2	0.5	0.5	1.2
2001.....	0.2	1.6	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.7
2000.....	0.2	2.1	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.1
1999.....	0.3	2.2	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3

Non-Hispanic white, male

2007 5/.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.5	0.8	*
2006 5/.....	0.2	*	*	*	*	*	0.1	0.3	0.2	0.4	0.8	*
2005 5/.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.4	0.6	*
2004 5/.....	0.2	1.9	*	*	*	*	0.2	0.3	0.2	0.3	0.7	*
2003 5/.....	0.3	1.8	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002.....	0.2	1.9	*	*	*	*	0.1	0.2	0.3	0.7	0.7	*
2001.....	0.3	*	*	*	*	*	*	0.3	0.4	0.5	0.8	*
2000.....	0.3	2.3	*	*	*	*	0.1	0.3	0.4	0.7	1.1	*
1999.....	0.3	2.8	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*

Non-Hispanic white, female

2007 5/.....	0.2	1.9	*	*	*	*	*	0.2	0.2	0.3	0.6	*
2006 5/.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.4	0.5	0.8
2005 5/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.3	0.5	0.8
2004 5/.....	0.2	*	*	*	*	*	*	*	0.3	0.5	0.5	*
2003 5/.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.6	0.8
2002.....	0.2	1.8	*	*	*	*	*	0.2	0.2	0.4	0.4	1.3
2001.....	0.2	1.9	*	*	*	*	*	0.2	0.3	0.4	0.6	*
2000.....	0.2	1.8	*	*	*	*	*	0.2	0.2	0.3	0.8	1.0
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.2



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

All origins 3/, both sexes

2007.....	6.7	*	*	*	*	*	*	0.1	1.2	11.9	71.9	143.5
2006.....	6.5	*	*	*	*	*	*	0.2	1.3	12.2	69.8	144.8
2005.....	6.6	*	*	*	*	*	*	0.2	1.4	13.0	71.2	143.7
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	12.0	67.5	135.8
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4

All origins 3/, male

2007.....	7.8	*	*	*	*	*	*	0.2	1.6	17.0	107.2	228.0
2006.....	7.7	*	*	*	*	*	*	0.2	1.7	17.6	104.7	231.7
2005.....	7.7	*	*	*	*	*	*	0.2	1.9	18.7	107.2	228.6
2004.....	7.2	*	*	*	*	*	*	0.2	1.6	17.3	101.7	220.0
2003.....	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002.....	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001.....	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000.....	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999.....	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6

All origins 3/, female

2007.....	5.6	*	*	*	*	*	*	0.1	0.9	7.7	47.6	103.3
2006.....	5.4	*	*	*	*	*	*	0.1	0.8	7.7	45.9	104.1
2005.....	5.5	*	*	*	*	*	*	0.1	1.0	8.1	46.8	104.7
2004.....	5.1	*	*	*	*	*	*	*	0.8	7.5	44.4	97.9
2003.....	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002.....	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001.....	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000.....	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999.....	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Hispanic, both sexes

2007.....	1.7	*	*	*	*	*	*	*	0.9	8.2	51.3	76.5
2006.....	1.6	*	*	*	*	*	*	*	*	8.9	41.5	93.9
2005.....	1.7	*	*	*	*	*	*	*	0.9	9.0	48.5	95.0
2004.....	1.4	*	*	*	*	*	*	*	1.0	6.8	44.1	85.1
2003.....	1.3	*	*	*	*	*	*	*	*	7.9	40.3	77.7
2002.....	1.3	*	*	*	*	*	*	*	1.1	8.6	37.2	76.5
2001.....	1.3	*	*	*	*	*	*	*	*	8.6	41.5	86.3
2000.....	1.2	*	*	*	*	*	*	*	*	7.8	37.7	86.3
1999.....	1.1	*	*	*	*	*	*	*	*	8.1	38.3	77.2

Hispanic, male

2007.....	1.9	*	*	*	*	*	*	*	*	10.9	74.1	106.7
2006.....	1.8	*	*	*	*	*	*	*	*	11.7	63.9	151.1
2005.....	1.9	*	*	*	*	*	*	*	*	13.5	71.2	138.2
2004.....	1.6	*	*	*	*	*	*	*	*	9.5	66.5	126.1
2003.....	1.5	*	*	*	*	*	*	*	*	12.4	58.3	129.3
2002.....	1.4	*	*	*	*	*	*	*	*	13.6	50.1	113.8
2001.....	1.4	*	*	*	*	*	*	*	*	11.2	63.0	127.6
2000.....	1.3	*	*	*	*	*	*	*	*	9.7	55.1	143.1
1999.....	1.3	*	*	*	*	*	*	*	*	11.4	58.9	105.7

Hispanic, female

2007.....	1.5	*	*	*	*	*	*	*	*	6.1	35.2	60.1
2006.....	1.3	*	*	*	*	*	*	*	*	6.6	25.9	63.5
2005.....	1.5	*	*	*	*	*	*	*	*	5.3	32.8	72.3
2004.....	1.2	*	*	*	*	*	*	*	*	4.7	28.6	64.1
2003.....	1.1	*	*	*	*	*	*	*	*	4.3	27.9	51.6
2002.....	1.1	*	*	*	*	*	*	*	*	4.7	28.4	58.0
2001.....	1.2	*	*	*	*	*	*	*	*	6.6	26.9	66.1
2000.....	1.1	*	*	*	*	*	*	*	*	6.3	26.1	58.4
1999.....	1.0	*	*	*	*	*	*	*	*	5.5	24.6	63.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Non-Hispanic 4/, both sexes

2007.....	7.5	*	*	*	*	*	*	0.1	1.2	12.2	73.2	147.0
2006.....	7.4	*	*	*	*	*	*	0.2	1.3	12.5	71.5	147.2
2005.....	7.4	*	*	*	*	*	*	0.2	1.5	13.2	72.5	145.9
2004.....	6.9	*	*	*	*	*	*	0.2	1.2	12.4	68.6	137.9
2003.....	7.0	*	*	*	*	*	*	0.2	1.3	13.0	69.1	140.6
2002.....	6.6	*	*	*	*	*	*	0.1	1.2	12.4	65.1	137.4
2001.....	6.5	*	*	*	*	*	*	0.1	1.2	11.9	65.5	135.6
2000.....	6.2	*	*	*	*	*	*	0.1	1.1	11.7	62.8	133.3
1999.....	5.8	*	*	*	*	*	*	0.1	1.1	11.1	58.9	125.8

Non-Hispanic 4/, male

2007.....	8.9	*	*	*	*	*	*	0.2	1.6	17.4	109.3	235.0
2006.....	8.7	*	*	*	*	*	*	0.2	1.8	18.0	107.2	235.9
2005.....	8.7	*	*	*	*	*	*	0.2	2.0	19.1	109.3	233.1
2004.....	8.1	*	*	*	*	*	*	0.3	1.6	17.8	103.4	224.4
2003.....	8.0	*	*	*	*	*	*	0.2	1.7	18.5	102.7	226.5
2002.....	7.6	*	*	*	*	*	*	0.2	1.5	18.2	96.7	224.4
2001.....	7.5	*	*	*	*	*	*	0.2	1.8	17.1	98.3	227.8
2000.....	7.1	*	*	*	*	*	*	0.2	1.5	16.5	94.2	225.2
1999.....	6.7	*	*	*	*	*	*	0.2	1.5	15.9	90.6	209.4

Non-Hispanic 4/, female

2007.....	6.2	*	*	*	*	*	*	0.1	0.9	7.8	48.4	105.4
2006.....	6.1	*	*	*	*	*	*	0.1	0.8	7.8	47.1	106.0
2005.....	6.1	*	*	*	*	*	*	*	1.0	8.3	47.5	106.1
2004.....	5.7	*	*	*	*	*	*	*	0.8	7.8	45.3	99.2
2003.....	5.9	*	*	*	*	*	*	0.2	0.9	8.4	46.6	102.8
2002.....	5.6	*	*	*	*	*	*	*	0.9	7.6	44.2	99.8
2001.....	5.4	*	*	*	*	*	*	*	0.7	7.6	44.0	97.3
2000.....	5.3	*	*	*	*	*	*	*	0.8	7.7	42.3	96.1
1999.....	4.9	*	*	*	*	*	*	*	0.7	7.1	38.3	92.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Non-Hispanic white, both sexes

2007 5/.....	9.0	*	*	*	*	*	*	0.2	1.3	13.1	78.3	155.7
2006 5/.....	8.9	*	*	*	*	*	*	0.1	1.4	13.3	76.8	157.0
2005 5/.....	8.9	*	*	*	*	*	*	0.2	1.6	14.2	77.4	155.5
2004 5/.....	8.2	*	*	*	*	*	*	0.2	1.3	13.5	73.3	145.5
2003 5/.....	8.3	*	*	*	*	*	*	0.2	1.3	14.0	73.7	148.8
2002.....	7.9	*	*	*	*	*	*	0.2	1.3	13.3	69.6	145.8
2001.....	7.7	*	*	*	*	*	*	0.1	1.3	12.8	69.9	143.5
2000.....	7.3	*	*	*	*	*	*	0.1	1.2	12.6	66.7	141.2
1999.....	6.8	*	*	*	*	*	*	0.1	1.1	11.9	62.4	132.8

Non-Hispanic white, male

2007 5/.....	10.6	*	*	*	*	*	*	0.2	1.7	18.4	116.4	249.0
2006 5/.....	10.4	*	*	*	*	*	*	0.2	1.9	19.1	114.0	252.4
2005 5/.....	10.4	*	*	*	*	*	*	0.2	2.1	20.3	116.1	247.9
2004 5/.....	9.7	*	*	*	*	*	*	0.3	1.7	19.2	110.0	236.3
2003 5/.....	9.5	*	*	*	*	*	*	0.2	1.7	19.8	109.0	237.9
2002.....	9.1	*	*	*	*	*	*	0.2	1.6	19.3	102.9	237.8
2001.....	9.0	*	*	*	*	*	*	0.2	2.0	18.3	104.5	241.7
2000.....	8.4	*	*	*	*	*	*	0.2	1.6	17.6	99.2	238.6
1999.....	7.9	*	*	*	*	*	*	0.2	1.6	17.1	95.7	221.0

Non-Hispanic white, female

2007 5/.....	7.5	*	*	*	*	*	*	*	0.9	8.5	51.7	111.5
2006 5/.....	7.4	*	*	*	*	*	*	*	0.9	8.4	51.0	112.5
2005 5/.....	7.4	*	*	*	*	*	*	*	1.1	9.0	50.7	113.2
2004 5/.....	6.8	*	*	*	*	*	*	*	0.9	8.5	48.3	104.8
2003 5/.....	7.1	*	*	*	*	*	*	0.2	1.0	9.0	49.8	109.6
2002.....	6.7	*	*	*	*	*	*	*	1.0	8.1	47.3	106.0
2001.....	6.5	*	*	*	*	*	*	*	0.7	8.2	46.9	102.7
2000.....	6.3	*	*	*	*	*	*	*	0.9	8.3	45.3	101.9
1999.....	5.8	*	*	*	*	*	*	*	0.7	7.6	40.5	97.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Non-Hispanic black, both sexes

2007 5/.....	2.1	*	*	*	*	*	*	*	1.1	7.3	35.7	66.2
2006 5/.....	1.8	*	*	*	*	*	*	*	1.0	7.5	30.6	51.8
2005 5/.....	1.9	*	*	*	*	*	*	*	0.8	7.3	34.4	57.3
2004 5/.....	1.7	*	*	*	*	*	*	*	0.7	5.5	29.8	61.6
2003 5/.....	1.7	*	*	*	*	*	*	*	1.2	7.2	29.4	53.5
2002.....	1.5	*	*	*	*	*	*	*	*	6.7	25.7	51.7
2001.....	1.5	*	*	*	*	*	*	*	0.8	5.6	25.7	59.0
2000.....	1.5	*	*	*	*	*	*	*	*	5.8	27.3	52.5
1999.....	1.4	*	*	*	*	*	*	*	0.9	5.9	24.8	51.7

Non-Hispanic black, male

2007 5/.....	2.4	*	*	*	*	*	*	*	1.4	12.5	54.7	100.3
2006 5/.....	2.1	*	*	*	*	*	*	*	*	12.1	51.4	71.5
2005 5/.....	2.3	*	*	*	*	*	*	*	*	12.6	54.9	99.0
2004 5/.....	1.9	*	*	*	*	*	*	*	*	8.6	44.5	106.7
2003 5/.....	2.0	*	*	*	*	*	*	*	1.8	10.6	45.6	99.1
2002.....	1.8	*	*	*	*	*	*	*	*	11.1	40.4	87.5
2001.....	1.7	*	*	*	*	*	*	*	*	8.8	38.1	99.3
2000.....	1.8	*	*	*	*	*	*	*	*	9.0	47.5	88.1
1999.....	1.6	*	*	*	*	*	*	*	*	8.7	38.5	87.9

Non-Hispanic black, female

2007 5/.....	1.8	*	*	*	*	*	*	*	*	3.6	24.9	52.1
2006 5/.....	1.5	*	*	*	*	*	*	*	*	4.3	18.8	43.7
2005 5/.....	1.5	*	*	*	*	*	*	*	*	3.6	22.7	40.6
2004 5/.....	1.5	*	*	*	*	*	*	*	*	3.3	21.6	43.8
2003 5/.....	1.4	*	*	*	*	*	*	*	*	4.8	20.2	35.9
2002.....	1.3	*	*	*	*	*	*	*	*	3.6	17.4	38.0
2001.....	1.4	*	*	*	*	*	*	*	*	3.3	18.7	44.1
2000.....	1.2	*	*	*	*	*	*	*	*	3.5	16.0	39.2
1999.....	1.3	*	*	*	*	*	*	*	*	3.9	17.2	38.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Alzheimer's disease (G30)

##### All origins 3/, both sexes

2007.....	24.7	*	*	*	*	*	*	0.2	2.2	20.6	176.7	849.1
2006.....	24.2	*	*	*	*	*	*	0.2	2.1	20.2	175.6	848.3
2005.....	24.2	*	*	*	*	*	*	0.2	2.1	20.5	177.3	861.6
2004.....	22.5	*	*	*	*	*	*	0.2	1.9	19.7	168.7	818.8
2003.....	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002.....	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3

##### All origins 3/, male

2007.....	14.7	*	*	*	*	*	*	0.2	2.1	19.1	155.3	646.0
2006.....	14.3	*	*	*	*	*	*	0.2	2.0	19.7	156.6	638.6
2005.....	14.1	*	*	*	*	*	*	0.1	2.0	19.3	157.7	639.9
2004.....	13.1	*	*	*	*	*	*	0.2	1.8	18.5	149.0	619.6
2003.....	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002.....	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001.....	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000.....	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999.....	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7

##### All origins 3/, female

2007.....	34.5	*	*	*	*	*	*	0.2	2.3	21.9	191.4	945.7
2006.....	33.8	*	*	*	*	*	*	0.2	2.1	20.5	188.7	946.4
2005.....	33.9	*	*	*	*	*	*	0.2	2.3	21.4	190.5	963.4
2004.....	31.5	*	*	*	*	*	*	0.3	1.9	20.8	181.9	908.4
2003.....	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002.....	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001.....	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000.....	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999.....	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Hispanic, both sexes												
2007.....	5.4	*	*	*	*	*	*	*	1.7	12.7	108.1	490.6
2006.....	5.4	*	*	*	*	*	*	*	1.9	12.5	110.4	514.5
2005.....	5.2	*	*	*	*	*	*	*	1.3	13.0	109.8	507.8
2004.....	4.6	*	*	*	*	*	*	*	1.0	12.0	99.5	490.9
2003.....	4.6	*	*	*	*	*	*	*	1.3	14.6	103.7	493.4
2002.....	3.8	*	*	*	*	*	*	*	1.0	10.8	89.2	438.5
2001.....	3.4	*	*	*	*	*	*	*	*	9.9	83.9	399.5
2000.....	3.2	*	*	*	*	*	*	*	*	11.9	79.2	380.2
1999.....	2.9	*	*	*	*	*	*	*	*	9.7	75.4	351.5
Hispanic, male												
2007.....	3.4	*	*	*	*	*	*	*	*	13.2	95.2	390.3
2006.....	3.5	*	*	*	*	*	*	*	2.1	13.1	107.2	404.8
2005.....	3.5	*	*	*	*	*	*	*	*	13.9	112.6	412.2
2004.....	2.8	*	*	*	*	*	*	*	*	11.5	93.0	374.1
2003.....	3.1	*	*	*	*	*	*	*	*	13.8	101.2	426.4
2002.....	2.3	*	*	*	*	*	*	*	*	10.5	80.4	341.3
2001.....	2.1	*	*	*	*	*	*	*	*	8.8	83.2	306.6
2000.....	2.1	*	*	*	*	*	*	*	*	13.3	81.6	284.2
1999.....	1.8	*	*	*	*	*	*	*	*	11.2	66.7	278.2
Hispanic, female												
2007.....	7.6	*	*	*	*	*	*	*	2.1	12.2	117.2	545.0
2006.....	7.4	*	*	*	*	*	*	*	1.6	12.0	112.6	572.8
2005.....	7.0	*	*	*	*	*	*	*	1.6	12.3	107.9	557.9
2004.....	6.5	*	*	*	*	*	*	*	*	12.5	104.0	551.0
2003.....	6.2	*	*	*	*	*	*	*	*	15.3	105.3	527.3
2002.....	5.4	*	*	*	*	*	*	*	*	10.9	95.3	486.6
2001.....	4.7	*	*	*	*	*	*	*	*	10.7	84.3	444.9
2000.....	4.3	*	*	*	*	*	*	*	*	10.8	77.6	427.3
1999.....	4.0	*	*	*	*	*	*	*	*	8.4	81.2	386.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

Non-Hispanic 4/, both sexes

2007.....	28.2	*	*	*	*	*	*	0.2	2.3	21.2	181.1	867.5
2006.....	27.4	*	*	*	*	*	*	0.2	2.1	20.7	179.5	864.3
2005.....	27.3	*	*	*	*	*	*	0.2	2.2	21.0	181.0	877.5
2004.....	25.4	*	*	*	*	*	*	0.2	1.9	20.2	172.3	832.6
2003.....	24.5	*	*	*	*	*	*	0.2	2.0	21.3	167.3	814.0
2002.....	22.9	*	*	*	*	*	*	0.1	2.0	20.3	161.1	763.2
2001.....	21.2	*	*	*	*	*	*	0.2	2.2	19.2	150.2	720.5
2000.....	19.6	*	*	*	*	*	*	0.2	2.1	19.0	142.0	676.7
1999.....	17.7	*	*	*	*	*	*	0.2	2.0	17.8	131.5	608.9

Non-Hispanic 4/, male

2007.....	16.8	*	*	*	*	*	*	0.2	2.2	19.5	159.1	660.3
2006.....	16.3	*	*	*	*	*	*	0.3	2.0	20.2	159.5	651.1
2005.....	15.9	*	*	*	*	*	*	0.2	2.0	19.7	160.2	650.9
2004.....	14.9	*	*	*	*	*	*	0.2	1.9	19.0	152.0	631.2
2003.....	14.4	*	*	*	*	*	*	0.2	1.9	20.5	149.8	613.6
2002.....	13.6	*	*	*	*	*	*	0.2	1.9	19.1	145.0	581.4
2001.....	12.7	*	*	*	*	*	*	0.2	2.0	18.3	138.6	565.2
2000.....	11.7	*	*	*	*	*	*	0.2	1.8	19.0	129.6	531.0
1999.....	10.9	*	*	*	*	*	*	0.2	2.0	17.7	124.8	492.8

Non-Hispanic 4/, female

2007.....	39.0	*	*	*	*	*	*	0.2	2.4	22.6	196.1	965.3
2006.....	38.0	*	*	*	*	*	*	0.2	2.2	21.2	193.2	963.4
2005.....	38.2	*	*	*	*	*	*	0.2	2.3	22.1	195.2	980.9
2004.....	35.3	*	*	*	*	*	*	0.3	2.0	21.3	186.0	922.7
2003.....	34.1	*	*	*	*	*	*	0.2	2.2	22.0	179.0	902.1
2002.....	31.9	*	*	*	*	*	*	0.1	2.1	21.2	171.8	841.6
2001.....	29.3	*	*	*	*	*	*	0.2	2.4	20.0	157.8	785.1
2000.....	27.2	*	*	*	*	*	*	0.2	2.3	19.1	150.0	735.6
1999.....	24.2	*	*	*	*	*	*	0.2	1.9	18.0	135.9	655.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

Non-Hispanic white, both sexes

2007 5/.....	33.0	*	*	*	*	*	*	0.3	2.5	22.1	188.3	899.8
2006 5/.....	32.2	*	*	*	*	*	*	0.3	2.3	21.7	187.1	896.5
2005 5/.....	31.9	*	*	*	*	*	*	0.2	2.4	21.7	187.5	909.1
2004 5/.....	29.6	*	*	*	*	*	*	0.3	2.1	21.1	178.1	861.9
2003 5/.....	28.7	*	*	*	*	*	*	0.2	2.2	22.2	173.9	846.9
2002.....	26.9	*	*	*	*	*	*	0.2	2.1	21.3	168.2	793.9
2001.....	24.8	*	*	*	*	*	*	0.2	2.4	20.3	155.6	749.2
2000.....	22.9	*	*	*	*	*	*	0.2	2.2	19.9	147.4	705.1
1999.....	20.7	*	*	*	*	*	*	0.2	2.1	18.6	136.7	636.6

Non-Hispanic white, male

2007 5/.....	19.6	*	*	*	*	*	*	0.2	2.3	20.2	164.7	686.4
2006 5/.....	19.1	*	*	*	*	*	*	0.3	2.1	21.0	165.9	676.4
2005 5/.....	18.5	*	*	*	*	*	*	0.2	2.2	20.1	164.7	673.6
2004 5/.....	17.3	*	*	*	*	*	*	0.2	2.0	19.6	157.0	653.3
2003 5/.....	16.8	*	*	*	*	*	*	0.2	2.0	21.2	154.8	638.1
2002.....	15.8	*	*	*	*	*	*	0.2	2.0	19.9	150.5	605.4
2001.....	14.7	*	*	*	*	*	*	0.2	2.1	19.1	143.0	587.2
2000.....	13.6	*	*	*	*	*	*	0.2	2.0	19.8	134.6	551.7
1999.....	12.7	*	*	*	*	*	*	0.2	2.2	18.3	129.8	515.1

Non-Hispanic white, female

2007 5/.....	46.0	*	*	*	*	*	*	0.3	2.7	23.8	204.9	1,000.9
2006 5/.....	44.8	*	*	*	*	*	*	0.2	2.4	22.4	201.8	999.2
2005 5/.....	44.8	*	*	*	*	*	*	0.3	2.5	23.0	203.3	1,016.8
2004 5/.....	41.4	*	*	*	*	*	*	0.4	2.2	22.3	192.5	955.3
2003 5/.....	40.2	*	*	*	*	*	*	0.2	2.5	23.0	186.8	938.8
2002.....	37.6	*	*	*	*	*	*	0.2	2.3	22.5	180.1	875.3
2001.....	34.4	*	*	*	*	*	*	0.2	2.7	21.3	163.9	816.6
2000.....	31.9	*	*	*	*	*	*	0.2	2.5	20.0	155.8	767.0
1999.....	28.4	*	*	*	*	*	*	0.2	2.0	18.8	141.2	685.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

Non-Hispanic black, both sexes

2007 5/.....	12.4	*	*	*	*	*	*	*	1.2	20.0	161.5	691.1
2006 5/.....	11.7	*	*	*	*	*	*	*	1.7	19.1	149.3	674.5
2005 5/.....	12.3	*	*	*	*	*	*	*	1.5	22.7	160.1	698.8
2004 5/.....	11.4	*	*	*	*	*	*	*	1.3	19.4	155.4	666.9
2003 5/.....	10.1	*	*	*	*	*	*	*	1.1	19.6	140.3	591.1
2002.....	9.1	*	*	*	*	*	*	*	1.2	16.8	124.2	548.7
2001.....	8.7	*	*	*	*	*	*	*	1.1	15.5	126.9	513.2
2000.....	7.7	*	*	*	*	*	*	*	1.3	16.1	111.6	450.9
1999.....	6.7	*	*	*	*	*	*	*	1.2	15.5	99.1	374.7

Non-Hispanic black, male

2007 5/.....	7.1	*	*	*	*	*	*	*	*	20.2	147.7	510.5
2006 5/.....	6.5	*	*	*	*	*	*	*	1.7	19.3	134.9	478.3
2005 5/.....	7.2	*	*	*	*	*	*	*	*	22.8	152.8	527.2
2004 5/.....	6.5	*	*	*	*	*	*	*	*	19.4	138.6	507.0
2003 5/.....	6.2	*	*	*	*	*	*	*	*	20.5	138.8	465.6
2002.....	5.5	*	*	*	*	*	*	*	*	17.9	122.5	414.6
2001.....	5.3	*	*	*	*	*	*	*	*	16.3	123.6	420.4
2000.....	4.6	*	*	*	*	*	*	*	*	16.9	98.7	378.1
1999.....	4.2	*	*	*	*	*	*	*	*	16.8	90.7	320.9

Non-Hispanic black, female

2007 5/.....	17.2	*	*	*	*	*	*	*	1.2	19.9	169.4	766.0
2006 5/.....	16.4	*	*	*	*	*	*	*	1.6	18.9	157.5	754.5
2005 5/.....	16.9	*	*	*	*	*	*	*	1.7	22.7	164.3	767.9
2004 5/.....	15.8	*	*	*	*	*	*	*	1.4	19.4	164.9	730.0
2003 5/.....	13.7	*	*	*	*	*	*	*	*	18.9	141.1	639.7
2002.....	12.5	*	*	*	*	*	*	*	*	16.1	125.2	599.9
2001.....	11.8	*	*	*	*	*	*	*	*	15.0	128.7	547.5
2000.....	10.6	*	*	*	*	*	*	*	*	15.6	118.8	478.0
1999.....	9.0	*	*	*	*	*	*	*	*	14.5	103.8	394.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes

2007.....	267.3	13.4	1.4	0.8	3.2	9.7	34.4	105.5	247.0	597.5	1,758.7	5,679.0
2006.....	275.1	12.3	1.3	0.9	3.2	10.1	35.3	108.7	255.5	629.6	1,846.2	5,928.2
2005.....	288.8	12.2	1.3	0.9	3.3	10.1	36.1	110.5	263.2	667.2	1,963.8	6,398.5
2004.....	293.3	13.8	1.5	0.8	3.2	10.0	36.6	110.9	268.2	697.7	2,035.6	6,616.6
2003.....	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
2002.....	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001.....	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000.....	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999.....	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1

All origins 3/, male

2007.....	261.0	14.8	1.2	0.9	4.0	12.4	46.8	148.4	346.4	781.3	2,127.2	5,894.7
2006.....	267.7	13.4	1.4	1.0	4.0	13.4	47.5	153.3	355.1	821.3	2,235.3	6,151.8
2005.....	277.9	13.4	1.5	1.0	4.2	13.0	48.6	155.8	365.7	864.5	2,368.4	6,615.9
2004.....	281.2	14.7	1.5	0.8	4.0	12.8	49.0	156.7	371.5	903.9	2,462.3	6,818.1
2003.....	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
2002.....	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001.....	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000.....	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999.....	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2

All origins 3/, female

2007.....	273.3	11.9	1.6	0.8	2.4	6.9	22.1	63.9	154.4	441.4	1,504.9	5,576.4
2006.....	282.4	11.2	1.3	0.8	2.4	6.7	23.1	65.5	162.9	467.4	1,580.1	5,823.6
2005.....	299.3	11.0	1.1	0.8	2.3	7.0	23.7	66.6	167.9	500.8	1,689.1	6,298.7
2004.....	304.9	12.9	1.5	0.8	2.3	7.1	24.2	66.6	172.2	524.5	1,748.4	6,526.0
2003.....	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
2002.....	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001.....	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000.....	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999.....	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Hispanic, both sexes

2007.....	85.0	11.9	1.2	0.7	2.6	6.3	19.7	66.5	183.8	465.1	1,358.8	3,845.6
2006.....	86.5	10.2	1.1	0.8	2.7	6.3	19.9	70.4	188.4	496.9	1,410.4	4,096.9
2005.....	91.4	10.5	1.1	0.8	2.8	5.9	20.9	71.1	197.8	545.3	1,564.7	4,469.4
2004.....	89.3	11.5	1.2	0.6	2.5	5.6	21.2	71.3	202.2	555.6	1,568.0	4,583.7
2003.....	93.4	11.0	1.3	0.6	2.5	6.4	20.5	71.9	212.2	594.9	1,683.5	5,176.6
2002.....	94.3	13.1	1.2	0.7	2.6	6.1	20.4	73.6	225.8	621.8	1,728.4	5,369.7
2001.....	96.2	12.0	1.5	0.8	2.4	6.4	19.8	75.4	216.4	644.9	1,828.4	6,036.2
2000.....	96.5	13.7	1.3	0.7	2.6	6.4	21.3	75.0	236.3	660.8	1,852.8	6,168.2
1999.....	99.2	15.2	1.4	0.7	2.5	6.5	21.4	80.5	242.7	690.0	1,935.5	6,335.9

Hispanic, male

2007.....	86.0	11.9	1.1	0.8	3.4	8.0	27.6	94.7	258.5	610.7	1,602.0	3,825.8
2006.....	87.1	11.1	1.3	1.0	3.6	8.5	27.4	97.7	258.6	645.2	1,697.6	4,103.7
2005.....	92.3	11.3	1.5	0.9	3.3	7.9	28.7	100.6	273.3	712.2	1,895.1	4,556.5
2004.....	90.6	11.7	1.6	0.5	3.5	7.8	29.2	101.1	282.3	730.6	1,928.2	4,600.7
2003.....	92.6	12.2	1.5	0.7	3.3	7.8	27.9	102.8	295.6	779.5	2,007.9	5,079.7
2002.....	94.2	12.2	*	0.6	3.3	8.1	27.4	102.9	316.5	822.6	2,067.7	5,494.9
2001.....	95.4	12.7	1.4	0.8	3.0	8.3	26.8	107.2	297.2	852.4	2,220.5	6,155.6
2000.....	96.0	13.7	1.4	0.9	3.1	8.1	27.9	107.7	330.6	870.1	2,272.6	6,304.3
1999.....	99.2	12.7	1.4	0.9	3.0	8.6	29.7	116.3	333.2	925.3	2,356.2	6,595.2

Hispanic, female

2007.....	84.0	11.9	1.4	0.6	1.8	4.2	10.7	37.3	114.5	346.3	1,188.1	3,856.3
2006.....	85.9	9.3	*	0.6	1.8	3.5	11.6	42.4	123.8	376.4	1,209.4	4,093.2
2005.....	90.4	9.7	*	0.7	2.1	3.5	12.2	41.1	128.8	410.4	1,334.6	4,423.7
2004.....	87.9	11.3	*	0.7	1.4	2.9	12.2	41.4	129.5	415.2	1,319.3	4,575.0
2003.....	94.3	9.7	*	*	1.6	4.6	12.3	41.0	137.0	447.8	1,460.7	5,225.8
2002.....	94.4	14.0	1.3	0.8	1.7	3.7	12.8	44.4	144.4	462.0	1,496.0	5,307.7
2001.....	97.1	11.2	1.6	0.8	1.8	4.2	12.3	44.1	144.6	481.2	1,563.4	5,977.9
2000.....	96.9	13.8	*	*	2.0	4.5	14.3	42.9	152.6	496.2	1,570.9	6,101.4
1999.....	99.1	17.9	1.4	*	2.0	4.0	12.7	45.7	162.6	505.6	1,655.5	6,210.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 4/, both sexes

2007.....	299.2	13.8	1.5	0.9	3.3	10.6	37.1	109.9	251.9	606.7	1,782.7	5,771.9
2006.....	307.3	12.9	1.4	0.9	3.3	11.1	38.0	112.8	260.5	638.1	1,870.0	6,013.2
2005.....	321.4	12.8	1.4	0.9	3.4	11.0	38.6	114.5	267.9	674.9	1,983.7	6,483.0
2004.....	325.9	14.4	1.5	0.9	3.3	11.0	38.9	114.7	272.6	705.9	2,057.5	6,698.1
2003.....	343.8	14.5	1.5	0.9	3.7	11.1	40.7	117.2	289.5	756.6	2,190.9	7,206.9
2002.....	352.2	16.5	1.5	0.9	3.3	10.8	40.2	117.8	299.1	796.7	2,276.3	7,451.4
2001.....	356.7	15.5	1.9	0.9	3.3	10.9	39.3	116.5	306.3	818.6	2,338.1	7,715.2
2000.....	365.6	17.3	1.6	0.9	3.5	10.1	38.6	118.8	323.8	857.8	2,412.2	8,076.4
1999.....	372.7	17.4	1.5	0.9	3.6	10.2	39.5	118.8	331.8	895.4	2,489.2	8,239.8

Non-Hispanic 4/, male

2007.....	293.3	15.6	1.3	0.9	4.1	13.6	50.5	154.6	353.1	792.4	2,158.9	6,010.3
2006.....	300.1	14.1	1.4	1.0	4.1	14.8	51.2	159.4	362.1	832.0	2,264.9	6,255.9
2005.....	310.3	14.0	1.5	1.0	4.4	14.3	52.1	161.6	372.1	873.3	2,391.7	6,716.9
2004.....	313.4	15.5	1.4	0.9	4.1	14.1	52.3	162.2	377.3	913.1	2,486.9	6,917.7
2003.....	329.5	15.9	1.4	1.0	4.4	14.1	54.6	166.3	399.0	986.2	2,662.3	7,419.6
2002.....	336.4	17.6	1.7	1.0	4.2	13.8	54.2	167.5	412.3	1,037.1	2,755.7	7,713.8
2001.....	338.1	15.8	2.0	0.9	4.1	13.7	53.0	164.7	420.6	1,066.8	2,842.6	7,990.6
2000.....	347.7	18.3	1.7	1.0	4.1	12.6	52.0	169.4	447.3	1,124.0	2,946.0	8,513.1
1999.....	354.2	19.1	1.7	1.0	4.2	12.7	53.6	169.3	457.9	1,177.2	3,041.8	8,657.6

Non-Hispanic 4/, female

2007.....	304.8	11.8	1.6	0.8	2.5	7.6	24.1	66.9	157.6	448.4	1,523.9	5,659.3
2006.....	314.1	11.7	1.4	0.8	2.5	7.4	25.0	68.0	165.9	473.6	1,600.4	5,900.4
2005.....	332.0	11.4	1.3	0.9	2.4	7.8	25.4	69.2	170.9	506.9	1,707.0	6,376.3
2004.....	337.9	13.2	1.6	0.9	2.5	8.0	25.9	69.0	175.3	531.4	1,768.8	6,600.0
2003.....	357.5	13.1	1.6	0.8	3.0	8.1	27.1	70.0	187.7	564.0	1,876.4	7,113.4
2002.....	367.2	15.3	1.3	0.8	2.3	7.9	26.5	70.0	194.1	595.9	1,958.5	7,338.2
2001.....	374.3	15.2	1.8	0.9	2.5	8.2	25.9	70.1	200.4	612.3	2,006.9	7,600.7
2000.....	382.6	16.3	1.5	0.8	2.9	7.6	25.4	70.0	209.6	638.0	2,064.5	7,899.8
1999.....	390.2	15.5	1.4	0.8	2.9	7.7	25.7	70.2	215.4	664.0	2,130.7	8,072.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic white, both sexes

2007 5/.....	324.4	10.4	1.2	0.8	2.7	8.4	32.0	96.8	228.3	572.1	1,767.5	5,875.7
2006 5/.....	333.1	9.5	1.2	0.8	2.9	8.9	32.8	98.7	236.4	602.6	1,851.6	6,119.6
2005 5/.....	348.4	9.8	1.1	0.8	2.8	8.8	33.3	99.5	242.6	640.3	1,963.3	6,596.0
2004 5/.....	353.1	12.0	1.2	0.7	2.8	8.8	33.9	99.7	247.1	670.5	2,036.7	6,790.0
2003 5/.....	372.4	10.9	1.2	0.8	3.0	9.0	34.8	101.4	262.4	717.8	2,169.7	7,304.7
2002.....	381.9	13.4	1.4	0.8	2.7	8.6	34.9	102.9	272.0	760.6	2,252.2	7,546.1
2001.....	385.8	12.1	1.5	0.8	2.7	8.9	33.3	101.3	278.5	779.4	2,313.0	7,811.4
2000.....	395.5	14.6	1.4	0.8	2.8	8.1	32.8	103.9	295.3	820.2	2,384.9	8,179.0
1999.....	402.7	14.2	1.3	0.8	3.0	8.3	33.3	104.1	302.5	856.8	2,462.9	8,342.9

Non-Hispanic white, male

2007 5/.....	315.2	11.7	0.9	0.8	3.2	11.0	44.2	139.8	324.1	753.2	2,146.8	6,162.7
2006 5/.....	322.3	10.2	1.1	0.9	3.7	12.1	45.6	143.1	332.0	790.3	2,252.6	6,398.5
2005 5/.....	333.6	10.7	1.3	0.8	3.6	11.6	46.2	144.4	341.7	834.9	2,380.1	6,880.5
2004 5/.....	336.9	13.0	1.1	0.8	3.5	11.6	47.1	145.3	346.5	876.3	2,476.1	7,045.5
2003 5/.....	354.1	11.3	0.9	0.9	3.6	11.7	48.5	148.4	368.3	945.5	2,648.3	7,561.6
2002.....	362.2	14.8	1.6	0.9	3.5	11.1	48.7	151.1	381.0	1,001.2	2,744.3	7,845.8
2001.....	363.0	13.0	1.6	0.8	3.3	11.4	46.7	148.1	389.7	1,027.1	2,827.4	8,116.3
2000.....	373.6	15.5	1.6	0.9	3.3	10.5	46.1	153.2	415.4	1,088.2	2,930.7	8,650.5
1999.....	379.8	16.1	1.5	0.8	3.4	10.6	46.9	152.7	424.4	1,139.8	3,030.2	8,776.1

Non-Hispanic white, female

2007 5/.....	333.4	9.0	1.4	0.7	2.2	5.8	19.8	54.4	137.0	414.5	1,501.9	5,739.7
2006 5/.....	343.6	8.8	1.3	0.6	2.0	5.6	20.0	55.0	145.3	439.9	1,573.4	5,989.5
2005 5/.....	362.8	8.9	0.9	0.7	1.9	5.9	20.3	55.3	148.4	472.1	1,676.5	6,465.9
2004 5/.....	368.7	10.9	1.2	0.7	2.0	6.1	20.6	54.8	152.6	493.5	1,737.3	6,675.5
2003 5/.....	390.0	10.4	1.4	0.6	2.4	6.2	21.2	55.1	161.9	522.8	1,846.4	7,191.6
2002.....	400.8	12.0	1.1	0.7	1.9	6.0	21.1	55.4	168.8	555.5	1,922.5	7,416.6
2001.....	407.6	11.0	1.5	0.8	2.0	6.4	19.9	55.3	173.3	569.4	1,971.8	7,684.6
2000.....	416.5	13.6	1.1	0.6	2.2	5.7	19.5	55.3	182.0	594.6	2,026.2	7,988.9
1999.....	424.7	12.1	1.1	0.7	2.5	6.0	19.8	56.2	187.7	620.0	2,091.4	8,170.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic black, both sexes

2007 5/.....	249.2	25.9	2.3	1.3	6.4	22.4	72.7	216.4	487.8	1,014.2	2,250.2	5,502.3
2006 5/.....	255.2	26.3	2.2	1.4	5.8	23.8	74.6	225.6	502.9	1,051.1	2,378.1	5,656.4
2005 5/.....	265.1	23.5	2.4	1.6	6.4	24.0	76.5	233.4	522.3	1,084.7	2,531.9	6,088.3
2004 5/.....	269.2	22.4	3.0	1.5	6.1	23.8	76.8	236.2	535.6	1,140.7	2,606.3	6,436.5
2003 5/.....	282.2	26.6	2.9	1.5	7.1	23.7	81.3	244.5	565.5	1,224.4	2,745.6	6,923.9
2002.....	285.1	28.3	2.1	1.4	6.0	24.3	78.2	242.6	578.0	1,243.1	2,851.9	7,136.9
2001.....	289.7	30.1	3.7	1.3	6.8	23.5	80.7	243.2	587.2	1,293.5	2,915.8	7,366.6
2000.....	294.2	28.8	2.5	1.5	7.0	22.3	79.1	246.4	609.6	1,333.1	2,995.1	7,576.5
1999.....	298.5	30.2	2.6	1.4	7.0	21.6	83.7	246.8	623.7	1,382.8	3,044.9	7,624.4

Non-Hispanic black, male

2007 5/.....	253.7	31.5	2.5	1.3	8.6	28.1	96.9	286.7	676.9	1,331.3	2,738.7	5,259.0
2006 5/.....	259.6	29.6	2.7	1.3	6.6	30.9	94.6	302.6	694.8	1,390.3	2,867.7	5,493.9
2005 5/.....	264.0	26.8	2.4	1.6	8.4	30.3	97.5	311.4	715.4	1,400.0	2,994.3	5,778.7
2004 5/.....	266.0	24.8	2.9	1.5	7.8	29.0	95.7	312.4	730.6	1,453.6	3,071.2	6,280.6
2003 5/.....	277.5	31.7	3.3	1.6	8.7	29.2	101.4	324.7	753.7	1,562.4	3,299.8	6,625.9
2002.....	277.2	29.1	2.0	1.5	7.3	30.1	97.5	319.3	775.5	1,576.9	3,318.5	6,982.7
2001.....	281.2	29.7	3.8	1.2	8.3	28.9	100.1	319.1	772.1	1,639.6	3,461.6	7,413.8
2000.....	284.3	30.0	2.3	1.4	7.8	26.5	97.6	324.0	807.5	1,658.1	3,527.8	7,755.5
1999.....	291.1	31.5	2.7	1.6	8.7	26.6	105.2	329.7	833.1	1,738.6	3,542.5	7,952.2

Non-Hispanic black, female

2007 5/.....	245.1	20.1	2.1	1.3	4.2	17.0	51.3	156.1	335.9	787.7	1,972.4	5,603.1
2006 5/.....	251.3	22.8	1.8	1.4	4.9	17.3	56.9	159.6	348.4	808.9	2,100.3	5,722.7
2005 5/.....	266.1	20.1	2.4	1.6	4.5	18.1	58.0	166.5	366.9	859.2	2,269.7	6,212.9
2004 5/.....	272.2	20.0	3.0	1.6	4.3	19.1	60.1	170.8	379.0	917.6	2,343.8	6,498.1
2003 5/.....	286.5	21.3	2.4	1.3	5.5	18.8	63.5	175.8	414.5	984.0	2,432.6	7,039.3
2002.....	292.3	27.4	2.2	1.4	4.7	18.9	61.2	176.8	419.5	1,006.4	2,588.0	7,195.9
2001.....	297.4	30.6	3.6	1.4	5.3	18.6	63.5	178.1	438.5	1,049.3	2,609.4	7,349.1
2000.....	303.1	27.5	2.9	1.5	6.3	18.6	62.8	179.8	450.3	1,104.6	2,695.3	7,510.0
1999.....	305.1	28.9	2.4	1.3	5.3	17.2	64.7	176.0	455.9	1,131.8	2,766.3	7,501.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes

2007.....	204.3	10.0	1.1	0.6	2.6	7.9	27.4	85.3	200.3	462.9	1,315.0	4,267.7
2006.....	211.0	8.4	1.0	0.6	2.5	8.2	28.3	88.0	207.3	490.3	1,383.1	4,480.8
2005.....	220.0	8.7	0.9	0.6	2.7	8.1	28.9	89.7	214.8	518.9	1,460.8	4,778.4
2004.....	222.2	10.3	1.2	0.6	2.5	7.9	29.3	90.2	218.8	541.6	1,506.3	4,895.9
2003.....	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
2002.....	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001.....	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000.....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999.....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0

All origins 3/, male

2007.....	208.4	10.9	1.0	0.6	3.2	10.5	38.6	124.6	288.8	624.9	1,656.5	4,621.8
2006.....	214.0	8.8	1.1	0.7	3.3	11.2	39.5	128.9	296.8	660.5	1,743.5	4,819.9
2005.....	221.1	9.4	1.0	0.6	3.6	10.8	40.7	131.5	306.9	692.3	1,829.4	5,143.4
2004.....	222.8	10.9	1.1	0.6	3.2	10.5	40.9	132.3	312.8	723.8	1,893.6	5,239.3
2003.....	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
2002.....	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001.....	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000.....	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999.....	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6

All origins 3/, female

2007.....	200.2	9.0	1.1	0.6	1.9	5.3	16.2	47.2	117.9	325.4	1,079.7	4,099.3
2006.....	208.0	7.9	0.9	0.6	1.8	5.1	17.0	48.5	124.1	346.3	1,136.7	4,322.1
2005.....	218.9	8.0	0.9	0.6	1.7	5.3	17.1	49.2	129.1	372.7	1,210.5	4,610.8
2004.....	221.6	9.7	1.2	0.6	1.7	5.2	17.7	49.6	131.5	388.6	1,245.6	4,741.5
2003.....	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
2002.....	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001.....	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000.....	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999.....	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>Hispanic, both sexes</b>												
2007.....	63.8	8.8	1.0	0.5	1.9	4.6	14.1	48.8	139.4	351.9	1,012.1	2,913.8
2006.....	65.3	6.5	0.8	0.6	1.8	4.6	14.0	51.7	142.2	377.2	1,064.9	3,140.6
2005.....	69.2	7.6	0.6	0.4	1.9	4.5	15.0	52.3	153.1	420.1	1,176.9	3,407.5
2004.....	67.2	8.9	1.1	0.4	1.8	4.0	15.2	52.4	155.7	424.2	1,177.6	3,446.8
2003.....	70.9	8.7	1.1	0.4	1.9	4.7	14.9	53.3	165.4	461.2	1,261.0	3,932.9
2002.....	71.9	10.2	0.8	0.4	1.9	4.4	14.7	55.3	176.8	484.3	1,297.6	4,121.9
2001.....	73.3	9.7	1.1	0.5	1.7	4.6	13.8	55.2	170.7	504.5	1,376.3	4,607.5
2000.....	73.1	11.2	1.0	0.6	1.9	4.3	14.8	55.9	183.3	506.5	1,395.9	4,701.8
1999.....	76.2	12.7	1.2	0.6	1.8	4.5	15.4	60.9	192.7	536.3	1,467.8	4,888.6
<b>Hispanic, male</b>												
2007.....	66.6	9.3	*	0.5	2.7	6.2	20.5	73.3	201.9	477.0	1,233.4	2,960.8
2006.....	67.7	7.1	*	0.7	2.5	6.5	20.1	75.6	202.3	505.6	1,308.4	3,257.9
2005.....	72.1	8.0	*	0.5	2.4	6.3	21.7	77.9	219.3	561.5	1,469.2	3,534.2
2004.....	70.2	8.9	1.4	*	2.7	5.9	22.1	77.6	224.6	572.2	1,489.0	3,496.8
2003.....	72.2	9.5	*	*	2.7	6.1	21.1	79.6	235.6	625.0	1,543.5	3,874.5
2002.....	74.0	9.6	*	*	2.6	6.1	20.6	80.5	256.0	657.7	1,599.5	4,301.8
2001.....	74.6	10.3	*	*	2.1	6.1	20.0	82.9	242.2	683.7	1,702.7	4,784.3
2000.....	74.7	11.2	1.4	0.7	2.4	5.7	20.7	84.3	264.8	684.8	1,733.2	4,897.5
1999.....	77.7	10.3	1.4	0.7	2.1	6.5	22.5	92.2	272.1	737.0	1,816.0	5,012.2
<b>Hispanic, female</b>												
2007.....	60.8	8.3	1.2	*	1.2	2.8	6.8	23.4	81.4	249.7	856.6	2,888.2
2006.....	62.6	5.9	*	*	0.9	2.3	7.1	27.3	86.9	273.0	894.5	3,078.3
2005.....	66.2	7.2	*	*	1.4	2.3	7.4	26.2	92.6	305.9	973.4	3,341.3
2004.....	64.1	9.0	*	*	0.8	1.7	7.5	27.0	93.1	305.5	962.7	3,421.2
2003.....	69.6	7.8	*	*	1.1	3.1	8.0	27.0	102.1	330.6	1,067.0	3,962.5
2002.....	69.7	10.8	*	*	1.1	2.4	8.2	30.2	105.7	346.4	1,090.8	4,032.8
2001.....	71.8	9.0	*	*	1.2	2.8	7.1	27.9	107.2	363.1	1,155.7	4,521.1
2000.....	71.5	11.2	*	*	1.3	2.7	8.5	28.2	111.2	366.3	1,169.4	4,605.8
1999.....	74.6	15.2	*	*	1.5	2.3	7.9	30.5	122.5	379.0	1,236.1	4,828.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic 4/, both sexes

2007.....	228.8	10.2	1.1	0.6	2.7	8.8	29.9	89.5	205.1	470.7	1,333.1	4,336.0
2006.....	235.8	8.9	1.0	0.6	2.7	9.0	30.7	92.0	212.2	497.6	1,400.4	4,542.5
2005.....	244.9	9.0	1.0	0.7	2.8	8.9	31.2	93.5	219.3	525.1	1,474.7	4,837.9
2004.....	247.0	10.7	1.2	0.6	2.6	8.8	31.5	93.9	223.1	548.4	1,521.3	4,953.4
2003.....	261.0	11.5	1.2	0.7	2.9	8.9	32.9	96.1	237.5	591.6	1,625.0	5,324.5
2002.....	267.1	12.9	1.1	0.7	2.6	8.7	32.7	97.1	245.5	622.2	1,690.6	5,484.9
2001.....	270.7	12.3	1.5	0.7	2.6	8.7	31.6	96.0	251.6	641.4	1,736.4	5,688.6
2000.....	277.3	13.3	1.2	0.7	2.8	8.0	30.8	97.1	265.7	672.7	1,791.1	5,953.6
1999.....	284.3	14.0	1.2	0.7	2.9	8.2	31.8	98.3	274.1	708.5	1,859.2	6,086.3

Non-Hispanic 4/, male

2007.....	234.6	11.3	1.0	0.7	3.3	11.7	42.1	130.7	295.5	634.6	1,682.1	4,714.6
2006.....	240.3	9.4	1.1	0.7	3.4	12.4	43.1	134.8	303.8	669.9	1,767.6	4,899.0
2005.....	247.1	9.8	1.0	0.7	3.8	12.0	44.0	137.2	313.2	699.9	1,847.0	5,222.0
2004.....	248.5	11.5	1.1	0.6	3.3	11.7	44.0	137.7	318.7	731.9	1,911.8	5,317.0
2003.....	261.4	12.8	1.1	0.8	3.5	11.6	46.1	141.5	337.4	793.4	2,050.1	5,690.6
2002.....	267.0	13.7	1.2	0.7	3.4	11.6	46.4	143.6	348.6	834.6	2,128.7	5,874.4
2001.....	268.0	12.0	1.7	0.7	3.3	11.2	44.6	141.1	356.1	858.7	2,192.5	6,077.8
2000.....	275.2	13.7	1.4	0.8	3.3	10.3	43.8	144.5	377.5	907.0	2,263.4	6,474.6
1999.....	282.0	14.8	1.3	0.8	3.6	10.5	45.4	145.7	388.7	958.3	2,349.8	6,602.0

Non-Hispanic 4/, female

2007.....	223.3	9.1	1.1	0.6	2.0	5.8	17.8	49.9	120.8	331.1	1,093.0	4,157.3
2006.....	231.4	8.4	1.0	0.6	1.9	5.7	18.6	50.8	127.0	351.4	1,149.7	4,376.7
2005.....	242.8	8.3	1.0	0.7	1.8	5.9	18.7	51.5	131.9	377.2	1,222.3	4,662.6
2004.....	245.5	9.8	1.3	0.6	1.9	5.9	19.2	51.8	134.3	393.8	1,258.8	4,790.8
2003.....	260.5	10.2	1.3	0.6	2.3	6.2	20.1	52.4	144.6	422.4	1,341.5	5,163.6
2002.....	267.2	12.0	1.1	0.6	1.8	5.8	19.3	52.4	150.0	444.8	1,400.2	5,316.8
2001.....	273.3	12.7	1.4	0.7	1.9	6.1	18.9	52.6	154.8	460.9	1,437.0	5,526.8
2000.....	279.3	12.8	1.1	0.6	2.2	5.7	18.1	51.5	162.2	479.3	1,483.4	5,742.9
1999.....	286.4	13.3	1.1	0.6	2.3	5.9	18.5	52.6	168.2	503.4	1,540.8	5,879.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic white, both sexes

2007 5/.....	249.9	7.2	0.8	0.5	2.1	7.1	26.6	81.5	189.9	449.0	1,329.7	4,432.5
2006 5/.....	257.4	6.5	0.9	0.5	2.3	7.4	27.6	83.1	196.8	475.5	1,394.5	4,642.7
2005 5/.....	267.1	7.0	0.8	0.5	2.3	7.2	27.8	84.0	202.4	504.3	1,466.8	4,942.5
2004 5/.....	269.1	9.1	0.9	0.5	2.1	7.2	28.2	84.2	206.6	526.3	1,514.3	5,038.3
2003 5/.....	284.0	8.7	0.9	0.6	2.3	7.3	29.2	85.8	219.6	566.7	1,616.7	5,412.2
2002.....	290.8	10.7	1.0	0.6	2.1	6.9	29.3	87.5	227.2	599.7	1,679.2	5,568.4
2001.....	294.1	9.8	1.2	0.6	2.1	7.1	27.6	86.4	233.0	616.7	1,724.0	5,776.0
2000.....	301.3	11.4	1.1	0.5	2.1	6.4	27.2	87.7	246.4	648.1	1,777.4	6,047.9
1999.....	308.2	11.3	1.0	0.6	2.4	6.7	27.6	88.7	254.0	683.2	1,846.1	6,176.6

Non-Hispanic white, male

2007 5/.....	254.3	8.1	0.6	0.6	2.5	9.6	37.8	121.6	276.6	610.9	1,685.0	4,858.5
2006 5/.....	260.3	6.8	0.9	0.6	3.0	10.3	39.8	124.5	284.5	644.3	1,767.4	5,032.8
2005 5/.....	267.8	7.5	0.8	0.6	3.1	9.9	40.3	126.2	293.0	677.6	1,849.3	5,374.1
2004 5/.....	269.1	9.8	0.8	0.5	2.7	9.7	40.6	126.9	298.8	709.5	1,915.1	5,430.9
2003 5/.....	282.9	9.4	0.7	0.7	2.8	9.8	42.2	129.8	317.7	767.3	2,049.9	5,821.0
2002.....	289.2	11.6	1.1	0.7	2.8	9.4	42.9	133.1	327.6	813.5	2,129.9	5,994.1
2001.....	289.5	10.3	1.3	0.6	2.6	9.4	40.4	130.7	335.8	834.7	2,190.4	6,195.4
2000.....	297.5	12.0	1.2	0.7	2.6	8.6	40.1	134.3	356.3	885.1	2,261.9	6,606.6
1999.....	304.0	12.4	1.1	0.6	2.8	8.7	40.6	135.0	365.6	935.2	2,352.6	6,713.7

Non-Hispanic white, female

2007 5/.....	245.5	6.4	1.0	0.5	1.7	4.5	15.3	42.1	107.1	308.1	1,081.0	4,230.8
2006 5/.....	254.7	6.1	0.9	0.5	1.5	4.4	15.5	42.2	113.2	329.1	1,135.8	4,460.8
2005 5/.....	266.4	6.5	0.7	0.5	1.4	4.5	15.2	42.4	116.1	354.6	1,203.6	4,745.1
2004 5/.....	269.1	8.3	1.0	0.5	1.5	4.6	15.6	42.2	118.9	368.6	1,241.2	4,862.4
2003 5/.....	285.1	8.0	1.1	0.5	1.9	4.7	16.2	42.4	126.6	394.8	1,324.0	5,232.2
2002.....	292.3	9.6	0.8	0.5	1.4	4.4	15.8	42.6	132.0	417.4	1,377.2	5,384.5
2001.....	298.4	9.2	1.1	0.6	1.5	4.7	14.8	42.7	136.0	431.8	1,414.7	5,601.6
2000.....	304.9	10.8	0.9	0.4	1.7	4.2	14.3	41.9	142.9	448.5	1,458.9	5,822.7
1999.....	312.3	10.2	0.8	0.6	1.9	4.6	14.5	43.1	149.0	472.3	1,514.5	5,962.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic black, both sexes

2007 5/.....	184.7	20.5	1.7	1.0	5.4	18.2	54.8	162.0	372.5	755.8	1,631.2	3,989.1
2006 5/.....	189.3	17.9	1.8	1.0	4.8	19.1	55.8	170.1	382.6	786.1	1,729.7	4,103.9
2005 5/.....	196.3	16.5	1.8	1.2	5.3	19.0	58.0	175.8	403.4	806.0	1,837.3	4,373.6
2004 5/.....	198.7	15.6	2.5	1.2	5.0	18.5	58.3	179.9	410.1	852.4	1,866.7	4,610.3
2003 5/.....	209.4	20.4	2.3	1.2	5.7	18.8	61.4	185.8	434.7	925.6	1,983.5	4,977.8
2002.....	212.2	21.9	1.7	1.0	5.1	19.2	59.4	184.7	450.2	936.3	2,067.0	5,151.0
2001.....	215.5	22.9	3.1	1.1	5.6	18.7	61.3	183.6	454.5	977.9	2,123.4	5,307.7
2000.....	218.6	20.5	2.0	1.2	5.9	17.9	58.7	185.4	473.2	1,014.8	2,177.9	5,424.6
1999.....	223.8	24.8	2.1	1.2	6.0	17.5	63.6	189.1	488.5	1,058.0	2,223.2	5,514.7

Non-Hispanic black, male

2007 5/.....	193.7	23.5	2.2	1.1	7.2	23.9	76.0	222.4	527.7	1,008.4	2,029.6	3,931.8
2006 5/.....	198.8	18.8	2.1	1.1	5.6	25.4	72.9	235.0	537.9	1,062.7	2,164.2	4,129.6
2005 5/.....	201.5	19.0	*	1.1	7.2	24.6	76.8	242.9	559.3	1,059.7	2,229.0	4,277.2
2004 5/.....	203.2	17.0	2.5	1.1	6.5	23.5	75.1	245.1	570.4	1,111.3	2,262.0	4,695.5
2003 5/.....	212.6	24.5	2.5	1.3	7.4	23.1	79.7	253.9	589.2	1,210.9	2,449.4	4,897.8
2002.....	212.9	23.0	*	1.0	6.3	25.0	77.6	250.9	615.0	1,207.5	2,470.0	5,161.2
2001.....	215.7	20.7	3.4	1.0	7.1	23.6	78.9	247.6	610.5	1,259.3	2,586.9	5,495.4
2000.....	217.6	20.1	2.0	1.2	6.9	22.1	75.6	252.5	639.6	1,281.7	2,615.1	5,677.2
1999.....	224.9	24.6	2.3	1.4	7.7	22.1	83.8	259.8	668.4	1,355.1	2,631.6	5,877.1

Non-Hispanic black, female

2007 5/.....	176.4	17.3	*	0.9	3.5	13.0	36.1	110.2	247.7	575.6	1,404.6	4,012.8
2006 5/.....	180.6	17.0	*	0.9	3.9	13.2	40.6	114.4	257.7	588.6	1,483.2	4,093.4
2005 5/.....	191.6	14.0	2.1	1.3	3.4	13.9	41.3	118.2	278.0	624.6	1,615.3	4,412.4
2004 5/.....	194.6	14.2	2.4	1.4	3.5	14.0	43.5	124.0	281.4	667.9	1,643.6	4,576.6
2003 5/.....	206.5	16.1	2.0	1.1	4.0	14.8	45.3	127.4	310.7	722.8	1,720.5	5,008.9
2002.....	211.6	20.8	1.9	1.0	3.9	14.1	43.3	128.0	317.9	744.0	1,839.1	5,147.0
2001.....	215.4	25.2	2.9	1.1	4.0	14.2	45.7	128.7	328.9	779.4	1,863.2	5,238.2
2000.....	219.5	20.8	2.0	1.1	4.9	14.1	43.7	127.8	339.3	827.1	1,931.9	5,330.8
1999.....	222.8	25.0	1.9	1.0	4.4	13.3	45.8	128.7	344.3	848.5	1,994.5	5,379.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins 3/, both sexes

2007.....	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.7	8.1	17.9
2006.....	1.1	*	*	*	*	0.1	0.2	0.4	1.1	3.2	8.6	16.7
2005.....	1.1	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.5	19.0
2004.....	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.1
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002.....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001.....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0

All origins 3/, male

2007.....	0.7	*	*	*	*	*	0.1	0.4	0.8	2.0	6.3	14.4
2006.....	0.7	*	*	*	*	*	0.1	0.4	0.9	2.4	6.2	12.0
2005.....	0.7	*	*	*	*	*	0.2	0.4	1.0	2.5	6.2	14.0
2004.....	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	7.3	11.5
2003.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001.....	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000.....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3

All origins 3/, female

2007.....	1.4	*	*	*	*	*	0.2	0.4	1.2	3.4	9.4	19.6
2006.....	1.5	*	*	*	*	*	0.2	0.4	1.3	3.9	10.3	18.9
2005.....	1.5	*	*	*	*	*	0.2	0.5	1.4	4.0	10.1	21.3
2004.....	1.5	*	*	*	*	*	0.3	0.6	1.4	4.1	9.9	19.5
2003.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002.....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001.....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000.....	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999.....	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Hispanic, both sexes

2007.....	0.3	*	*	*	*	*	*	0.5	1.1	*	5.9	7.9
2006.....	0.4	*	*	*	*	*	*	*	1.1	2.5	6.6	10.2
2005.....	0.4	*	*	*	*	*	*	0.5	1.1	3.0	5.6	8.6
2004.....	0.4	*	*	*	*	*	*	*	1.3	3.5	6.3	*
2003.....	0.4	*	*	*	*	*	*	0.5	1.0	3.3	6.7	*
2002.....	0.5	*	*	*	*	*	*	0.7	1.8	4.6	7.0	*
2001.....	0.4	*	*	*	*	*	*	*	1.5	4.2	5.4	*
2000.....	0.4	*	*	*	*	*	*	*	1.8	4.4	5.3	*
1999.....	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*

Hispanic, male

2007.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*	*	*

Hispanic, female

2007.....	0.5	*	*	*	*	*	*	*	1.7	*	6.8	*
2006.....	0.5	*	*	*	*	*	*	*	*	2.6	8.9	12.0
2005.....	0.6	*	*	*	*	*	*	*	*	3.7	8.1	*
2004.....	0.5	*	*	*	*	*	*	*	2.1	3.7	6.4	*
2003.....	0.5	*	*	*	*	*	*	*	*	4.0	8.4	*
2002.....	0.8	*	*	*	*	*	*	*	2.1	5.9	10.7	*
2001.....	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999.....	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 4/, both sexes

2007.....	1.2	*	*	*	*	0.1	0.2	0.4	1.0	2.9	8.2	18.5
2006.....	1.2	*	*	*	*	0.1	0.2	0.4	1.1	3.3	8.7	17.0
2005.....	1.3	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.7	19.5
2004.....	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.3	9.0	17.7
2003.....	1.4	*	*	*	*	*	0.2	0.5	1.3	3.8	9.7	20.9
2002.....	1.4	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.7	18.1
2001.....	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.6	18.0
2000.....	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.7	17.1
1999.....	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.1	18.3

Non-Hispanic 4/, male

2007.....	0.8	*	*	*	*	*	0.1	0.3	0.9	2.1	6.4	14.8
2006.....	0.8	*	*	*	*	*	0.1	0.4	0.9	2.4	6.3	12.3
2005.....	0.8	*	*	*	*	*	0.2	0.4	1.0	2.5	6.4	14.6
2004.....	0.8	*	*	*	*	*	0.2	0.4	0.8	2.3	7.3	11.9
2003.....	0.9	*	*	*	*	*	0.2	0.4	1.1	2.9	7.6	16.6
2002.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	7.0	12.0
2001.....	0.8	*	*	*	*	*	0.3	0.5	1.1	2.8	6.7	13.1
2000.....	0.9	*	*	*	*	*	0.3	0.6	1.4	2.9	6.7	12.5
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.6	6.9	10.7

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	*	0.2	0.5	1.2	3.5	9.5	20.2
2006.....	1.6	*	*	*	*	*	0.2	0.4	1.3	4.0	10.3	19.2
2005.....	1.7	*	*	*	*	*	0.2	0.5	1.4	4.0	10.3	21.8
2004.....	1.7	*	*	*	*	*	0.2	0.6	1.4	4.1	10.0	20.2
2003.....	1.8	*	*	*	*	*	0.3	0.7	1.5	4.6	11.1	22.8
2002.....	1.8	*	*	*	*	*	0.2	0.7	2.0	4.5	11.6	20.8
2001.....	1.8	*	*	*	*	*	0.3	0.7	1.7	5.1	11.5	20.0
2000.....	1.9	*	*	*	*	*	0.4	0.8	2.0	5.8	11.6	19.0
1999.....	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic white, both sexes

2007 5/.....	1.3	*	*	*	*	*	0.1	0.3	0.9	2.9	8.8	19.7
2006 5/.....	1.4	*	*	*	*	*	0.1	0.3	1.0	3.3	9.2	18.4
2005 5/.....	1.4	*	*	*	*	*	0.2	0.4	1.1	3.4	9.2	21.0
2004 5/.....	1.4	*	*	*	*	*	0.1	0.5	1.0	3.4	9.5	18.9
2003 5/.....	1.5	*	*	*	*	*	0.2	0.4	1.2	3.9	10.4	22.3
2002.....	1.5	*	*	*	*	*	0.2	0.5	1.5	3.9	10.2	19.3
2001.....	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.2	19.0
2000.....	1.6	*	*	*	*	*	0.3	0.6	1.7	4.6	10.3	18.0
1999.....	1.6	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.6	19.2

Non-Hispanic white, male

2007 5/.....	0.9	*	*	*	*	*	*	0.3	0.8	2.2	6.6	16.0
2006 5/.....	0.8	*	*	*	*	*	*	0.3	0.8	2.4	6.7	13.2
2005 5/.....	0.9	*	*	*	*	*	0.2	0.3	0.9	2.5	6.7	15.4
2004 5/.....	0.9	*	*	*	*	*	*	0.4	0.7	2.4	7.8	12.8
2003 5/.....	1.0	*	*	*	*	*	0.2	0.3	0.9	3.0	8.3	17.7
2002.....	0.9	*	*	*	*	*	0.2	0.4	1.1	3.0	7.3	12.9
2001.....	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	7.0	13.7
2000.....	0.9	*	*	*	*	*	0.3	0.5	1.5	2.9	7.2	13.0
1999.....	0.9	*	*	*	*	*	0.2	0.5	1.1	3.8	7.2	11.6

Non-Hispanic white, female

2007 5/.....	1.8	*	*	*	*	*	0.2	0.4	1.0	3.6	10.3	21.4
2006 5/.....	1.9	*	*	*	*	*	*	0.4	1.2	4.1	10.9	20.8
2005 5/.....	1.9	*	*	*	*	*	0.2	0.4	1.2	4.2	10.9	23.6
2004 5/.....	1.9	*	*	*	*	*	0.2	0.5	1.3	4.2	10.6	21.6
2003 5/.....	2.1	*	*	*	*	*	0.2	0.5	1.4	4.6	11.8	24.4
2002.....	2.1	*	*	*	*	*	0.2	0.6	1.9	4.6	12.2	22.0
2001.....	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.2	21.2
2000.....	2.1	*	*	*	*	*	0.3	0.7	1.9	6.0	12.4	20.0
1999.....	2.2	*	*	*	*	*	0.2	0.7	2.2	5.7	12.9	22.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic black, both sexes

2007 5/.....	0.6	*	*	*	*	*	*	0.8	1.7	2.4	4.6	8.7
2006 5/.....	0.7	*	*	*	*	*	0.4	0.8	1.7	3.0	5.2	*
2005 5/.....	0.6	*	*	*	*	*	*	1.0	1.5	2.4	4.5	6.3
2004 5/.....	0.7	*	*	*	*	*	0.6	1.0	1.6	2.7	4.1	6.0
2003 5/.....	0.8	*	*	*	*	*	0.5	1.4	2.7	4.0	4.2	7.1
2002.....	0.7	*	*	*	*	*	0.4	1.0	2.4	2.9	4.4	*
2001.....	0.8	*	*	*	*	*	0.7	1.4	2.1	4.2	4.3	6.9
2000.....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.6	4.0	8.6
1999.....	0.8	*	*	*	*	*	0.5	1.2	2.2	3.4	4.4	8.4

Non-Hispanic black, male

2007 5/.....	0.5	*	*	*	*	*	*	*	1.6	*	*	*
2006 5/.....	0.5	*	*	*	*	*	*	*	1.7	*	*	*
2005 5/.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 5/.....	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2003 5/.....	0.6	*	*	*	*	*	*	0.9	2.8	3.0	*	*
2002.....	0.5	*	*	*	*	*	*	*	1.8	2.9	*	*
2001.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic black, female

2007 5/.....	0.8	*	*	*	*	*	*	1.0	1.7	3.1	4.2	9.8
2006 5/.....	0.8	*	*	*	*	*	*	0.9	1.8	3.3	6.2	*
2005 5/.....	0.7	*	*	*	*	*	*	1.1	1.6	2.9	4.6	*
2004 5/.....	0.8	*	*	*	*	*	*	1.1	1.7	3.4	5.2	*
2003 5/.....	1.0	*	*	*	*	*	*	1.7	2.6	4.6	5.2	*
2002.....	0.8	*	*	*	*	*	*	1.1	2.9	2.9	5.1	*
2001.....	1.2	*	*	*	*	*	0.9	1.7	2.6	5.4	5.4	*
2000.....	1.0	*	*	*	*	*	0.8	1.4	2.7	4.4	4.9	*
1999.....	1.0	*	*	*	*	*	*	1.5	2.7	4.1	4.4	11.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Hypertensive heart disease (I11)

##### All origins 3/, both sexes

2007.....	10.2	*	*	*	0.1	0.8	3.2	8.2	13.7	20.7	48.6	192.3
2006.....	9.9	*	*	*	0.1	0.7	3.1	8.3	13.1	20.4	47.8	194.0
2005.....	9.9	*	*	*	0.1	0.8	3.1	7.8	13.0	20.4	48.8	198.6
2004.....	9.7	*	*	*	0.1	0.8	3.0	7.8	13.4	21.7	49.3	192.8
2003.....	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002.....	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001.....	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000.....	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999.....	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9

##### All origins 3/, male

2007.....	9.6	*	*	*	0.2	1.1	4.4	11.5	19.1	25.5	50.3	146.3
2006.....	9.3	*	*	*	0.2	1.0	4.2	11.7	18.0	25.7	49.7	143.0
2005.....	9.1	*	*	*	0.1	1.1	4.4	10.9	18.1	25.2	48.7	147.5
2004.....	8.9	*	*	*	0.1	1.1	4.1	10.7	18.5	27.3	48.0	145.4
2003.....	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002.....	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001.....	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000.....	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999.....	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6

##### All origins 3/, female

2007.....	10.8	*	*	*	*	0.5	1.9	5.1	8.7	16.7	47.3	214.1
2006.....	10.6	*	*	*	*	0.4	2.0	5.0	8.5	15.9	46.5	217.9
2005.....	10.7	*	*	*	*	0.5	1.9	4.7	8.3	16.4	48.9	222.0
2004.....	10.5	*	*	*	*	0.4	1.9	5.1	8.6	16.9	50.2	214.1
2003.....	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002.....	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001.....	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000.....	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999.....	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Hispanic, both sexes												
2007.....	3.8	*	*	*	*	0.5	1.8	5.2	10.7	18.4	43.3	143.7
2006.....	3.6	*	*	*	*	0.4	1.8	4.9	9.5	18.4	43.1	157.7
2005.....	3.5	*	*	*	*	0.4	1.7	4.9	10.8	16.5	42.8	145.0
2004.....	3.6	*	*	*	*	0.3	1.7	4.8	11.0	22.2	47.0	150.3
2003.....	4.0	*	*	*	*	0.4	1.8	5.7	11.8	22.9	54.2	179.3
2002.....	3.7	*	*	*	*	0.4	1.7	4.9	12.2	22.4	51.4	165.7
2001.....	3.3	*	*	*	*	0.3	1.7	4.9	10.0	21.0	46.9	151.9
2000.....	2.8	*	*	*	*	0.3	1.2	3.8	8.2	18.4	46.0	143.3
1999.....	2.7	*	*	*	*	*	1.2	3.3	8.9	16.8	44.8	152.2
Hispanic, male												
2007.....	3.9	*	*	*	*	0.7	2.7	7.2	15.1	23.2	47.2	109.8
2006.....	3.7	*	*	*	*	0.6	2.7	7.1	13.1	23.8	44.6	125.2
2005.....	3.5	*	*	*	*	0.6	2.5	7.0	14.3	19.5	45.8	118.3
2004.....	3.8	*	*	*	*	*	2.7	6.1	16.0	27.8	55.4	131.7
2003.....	4.3	*	*	*	*	0.5	2.9	8.3	17.0	30.8	63.7	144.7
2002.....	3.9	*	*	*	*	0.6	2.6	7.4	17.0	28.5	61.1	137.2
2001.....	3.4	*	*	*	*	*	2.7	7.0	13.8	26.4	48.3	144.7
2000.....	2.7	*	*	*	*	*	1.6	5.4	10.7	22.4	50.6	112.9
1999.....	2.6	*	*	*	*	*	1.8	4.7	12.5	20.9	42.8	107.8
Hispanic, female												
2007.....	3.6	*	*	*	*	*	0.7	3.0	6.5	14.5	40.6	162.2
2006.....	3.6	*	*	*	*	*	0.8	2.6	6.3	14.0	42.0	174.9
2005.....	3.4	*	*	*	*	*	0.9	2.8	7.6	14.1	40.7	159.0
2004.....	3.4	*	*	*	*	*	*	3.4	6.4	17.7	41.2	159.9
2003.....	3.7	*	*	*	*	*	*	3.1	7.2	16.6	47.7	196.9
2002.....	3.4	*	*	*	*	*	*	2.5	7.9	17.5	44.7	179.9
2001.....	3.1	*	*	*	*	*	*	2.7	6.5	16.8	45.9	155.5
2000.....	2.9	*	*	*	*	*	0.8	2.1	6.1	15.3	42.9	158.3
1999.....	2.9	*	*	*	*	*	*	2.0	5.8	13.6	46.1	173.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Hypertensive heart disease (I11)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	11.3	*	*	*	0.1	0.9	3.4	8.5	13.9	20.8	48.8	194.6
2006.....	11.0	*	*	*	0.1	0.8	3.3	8.6	13.3	20.5	47.9	195.6
2005.....	10.9	*	*	*	0.1	0.9	3.3	8.0	13.2	20.6	49.0	200.7
2004.....	10.7	*	*	*	0.1	0.9	3.2	8.1	13.5	21.5	49.3	194.3
2003.....	10.6	*	*	*	0.1	0.9	3.1	8.1	13.1	21.9	50.7	195.0
2002.....	10.0	*	*	*	0.1	0.8	2.8	7.4	12.4	21.0	49.3	187.1
2001.....	9.4	*	*	*	*	0.7	2.5	6.3	10.9	19.6	48.6	190.3
2000.....	9.2	*	*	*	0.1	0.6	2.3	5.7	11.2	19.5	48.1	194.3
1999.....	8.8	*	*	*	0.1	0.6	2.2	5.4	10.7	19.1	47.6	185.3
<b>Non-Hispanic 4/, male</b>												
2007.....	10.6	*	*	*	0.2	1.3	4.7	11.9	19.3	25.5	50.4	148.2
2006.....	10.3	*	*	*	0.2	1.1	4.5	12.2	18.3	25.7	49.8	143.8
2005.....	10.0	*	*	*	0.1	1.2	4.7	11.3	18.3	25.4	48.7	148.7
2004.....	9.8	*	*	*	0.2	1.3	4.3	11.1	18.5	27.1	47.4	145.6
2003.....	9.5	*	*	*	0.2	1.1	4.1	10.9	17.1	26.7	50.1	141.7
2002.....	8.9	*	*	*	0.1	1.2	4.0	10.0	16.8	24.6	48.5	133.7
2001.....	8.1	*	*	*	*	0.9	3.4	8.6	14.2	23.3	48.3	133.4
2000.....	7.8	*	*	*	0.1	0.7	3.0	7.9	14.6	23.5	46.0	139.6
1999.....	7.4	*	*	*	0.1	0.8	3.1	7.2	13.6	21.7	46.7	134.2
<b>Non-Hispanic 4/, female</b>												
2007.....	12.0	*	*	*	*	0.6	2.2	5.3	8.9	16.8	47.7	216.6
2006.....	11.7	*	*	*	*	0.4	2.2	5.2	8.7	16.1	46.7	219.6
2005.....	11.8	*	*	*	*	0.6	2.0	4.9	8.3	16.6	49.2	224.5
2004.....	11.6	*	*	*	*	0.5	2.1	5.2	8.8	16.7	50.6	216.0
2003.....	11.7	*	*	*	*	0.6	2.1	5.4	9.4	17.8	51.1	218.5
2002.....	11.1	*	*	*	*	0.5	1.7	4.8	8.4	18.0	49.9	210.2
2001.....	10.7	*	*	*	*	0.4	1.7	4.2	7.8	16.5	48.8	214.0
2000.....	10.6	*	*	*	*	0.4	1.6	3.6	8.0	16.2	49.5	216.5
1999.....	10.2	*	*	*	*	0.3	1.4	3.7	8.0	16.9	48.2	205.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Hypertensive heart disease (I11)

##### Non-Hispanic white, both sexes

2007 5/.....	10.6	*	*	*	*	0.5	2.3	6.3	10.4	16.4	42.6	186.6
2006 5/.....	10.3	*	*	*	0.1	0.5	2.4	6.3	10.0	15.5	41.9	188.0
2005 5/.....	10.1	*	*	*	*	0.6	2.3	5.6	10.0	16.1	42.3	190.6
2004 5/.....	9.8	*	*	*	*	0.5	2.2	5.6	9.8	17.0	42.7	184.6
2003 5/.....	9.7	*	*	*	*	0.5	2.0	5.3	9.5	16.7	44.3	184.9
2002.....	9.1	*	*	*	*	0.4	1.9	5.1	8.8	16.1	42.2	177.0
2001.....	8.6	*	*	*	*	0.4	1.6	4.1	7.6	14.8	42.2	180.4
2000.....	8.3	*	*	*	*	0.3	1.5	3.6	7.5	14.5	41.2	182.5
1999.....	7.9	*	*	*	*	0.3	1.3	3.4	7.3	14.1	40.6	175.2

##### Non-Hispanic white, male

2007 5/.....	9.4	*	*	*	*	0.7	3.3	9.2	14.8	20.2	43.3	140.0
2006 5/.....	9.0	*	*	*	0.2	0.7	3.5	9.4	14.0	19.3	42.4	134.7
2005 5/.....	8.8	*	*	*	*	0.9	3.4	8.3	14.2	19.9	42.0	137.6
2004 5/.....	8.6	*	*	*	*	0.8	3.2	8.3	13.6	21.6	40.9	135.9
2003 5/.....	8.2	*	*	*	*	0.8	3.0	7.8	13.0	20.4	43.5	130.8
2002.....	7.6	*	*	*	*	0.7	2.8	7.3	12.2	19.0	41.2	123.5
2001.....	6.9	*	*	*	*	0.6	2.3	5.9	10.5	18.1	41.5	122.0
2000.....	6.6	*	*	*	*	0.4	2.1	5.4	10.3	18.0	38.8	129.5
1999.....	6.2	*	*	*	*	0.6	2.0	4.9	9.6	16.2	39.4	123.8

##### Non-Hispanic white, female

2007 5/.....	11.7	*	*	*	*	0.3	1.3	3.5	6.3	13.2	42.2	208.6
2006 5/.....	11.4	*	*	*	*	0.2	1.3	3.2	6.2	12.3	41.6	212.9
2005 5/.....	11.3	*	*	*	*	0.3	1.2	2.9	5.9	12.8	42.5	214.9
2004 5/.....	11.0	*	*	*	*	*	1.2	3.0	6.2	13.0	43.9	206.5
2003 5/.....	11.1	*	*	*	*	0.3	1.1	2.8	6.3	13.6	44.9	208.7
2002.....	10.5	*	*	*	*	0.2	1.0	2.9	5.5	13.6	42.9	200.1
2001.....	10.2	*	*	*	*	0.2	0.8	2.3	4.9	11.9	42.6	204.7
2000.....	9.9	*	*	*	*	0.2	0.8	1.9	4.9	11.6	42.7	203.9
1999.....	9.6	*	*	*	*	*	0.6	2.0	5.0	12.4	41.4	195.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Hypertensive heart disease (I11)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	18.4	*	*	*	0.4	3.0	10.4	24.6	44.3	62.5	121.6	315.5
2006 5/.....	17.9	*	*	*	*	2.6	9.2	25.9	42.7	66.2	119.0	304.2
2005 5/.....	18.3	*	*	*	0.4	2.6	9.8	25.6	42.0	64.0	129.0	341.9
2004 5/.....	18.2	*	*	*	0.3	2.8	9.5	26.7	45.8	65.4	129.1	318.6
2003 5/.....	18.5	*	*	*	0.4	2.6	9.8	28.5	45.1	70.3	128.4	334.9
2002.....	17.7	*	*	*	*	2.9	8.8	24.8	45.1	68.3	134.2	322.1
2001.....	16.5	*	*	*	*	2.2	8.5	23.2	39.4	66.1	125.6	328.7
2000.....	16.6	*	*	*	*	2.2	7.5	21.6	43.1	67.6	132.7	344.1
1999.....	16.0	*	*	*	0.4	1.9	8.1	21.3	40.1	67.4	132.7	308.5
<b>Non-Hispanic black, male</b>												
2007 5/.....	19.9	*	*	*	*	4.1	14.2	33.0	62.5	82.2	143.6	268.8
2006 5/.....	19.5	*	*	*	*	3.7	11.3	34.3	59.6	90.2	147.1	278.5
2005 5/.....	19.1	*	*	*	*	3.0	13.2	34.7	59.1	82.6	138.8	304.5
2004 5/.....	19.0	*	*	*	*	3.7	12.1	34.0	65.4	87.5	133.8	271.3
2003 5/.....	18.7	*	*	*	*	3.2	11.5	35.6	57.5	93.4	139.1	291.9
2002.....	18.0	*	*	*	*	4.1	11.7	31.9	61.9	84.0	141.8	264.7
2001.....	16.3	*	*	*	*	2.7	10.5	30.4	50.1	79.8	137.4	288.7
2000.....	16.1	*	*	*	*	2.7	9.2	28.4	56.0	81.9	139.5	275.9
1999.....	15.6	*	*	*	*	2.2	10.3	26.8	51.1	80.6	143.2	266.2
<b>Non-Hispanic black, female</b>												
2007 5/.....	17.1	*	*	*	*	1.9	7.0	17.4	29.7	48.3	109.2	334.9
2006 5/.....	16.5	*	*	*	*	1.5	7.3	18.7	29.1	49.0	103.0	314.6
2005 5/.....	17.5	*	*	*	*	2.3	6.9	17.9	28.2	50.7	123.5	356.9
2004 5/.....	17.5	*	*	*	*	2.0	7.3	20.4	30.1	49.6	126.4	337.3
2003 5/.....	18.4	*	*	*	*	2.1	8.2	22.4	35.1	53.9	122.4	351.6
2002.....	17.4	*	*	*	*	1.8	6.2	18.7	31.7	57.1	129.8	344.1
2001.....	16.7	*	*	*	*	1.7	6.7	17.1	30.8	56.5	118.9	343.5
2000.....	17.1	*	*	*	*	1.8	5.9	15.8	32.7	57.5	128.9	369.4
1999.....	16.4	*	*	*	*	1.6	6.1	16.5	31.3	58.1	126.8	324.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

All origins 3/, both sexes

2007.....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.0	6.1	22.3
2006.....	1.0	*	*	*	*	0.1	0.2	0.4	0.8	1.9	6.2	22.6
2005.....	1.1	*	*	*	*	0.1	0.2	0.4	0.8	2.3	7.0	25.3
2004.....	1.0	*	*	*	*	0.1	0.2	0.5	0.9	2.1	6.6	24.8
2003.....	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000.....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999.....	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1

All origins 3/, male

2007.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.1	6.5	23.0
2006.....	0.9	*	*	*	*	0.1	0.3	0.5	1.1	2.0	6.9	23.8
2005.....	0.9	*	*	*	*	0.1	0.2	0.6	1.0	2.5	7.2	27.4
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.2	7.2	24.3
2003.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002.....	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001.....	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000.....	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999.....	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3

All origins 3/, female

2007.....	1.1	*	*	*	*	*	0.1	0.3	0.7	1.8	5.8	22.0
2006.....	1.1	*	*	*	*	*	0.1	0.4	0.6	1.8	5.7	22.1
2005.....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.1	6.8	24.3
2004.....	1.2	*	*	*	*	*	0.1	0.4	0.7	1.9	6.2	25.0
2003.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001.....	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999.....	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

Hispanic, both sexes

2007.....	0.4	*	*	*	*	*	*	0.4	1.0	2.5	5.2	20.4
2006.....	0.3	*	*	*	*	*	*	*	0.9	2.3	4.9	12.9
2005.....	0.5	*	*	*	*	*	*	*	1.2	3.3	7.1	22.2
2004.....	0.5	*	*	*	*	*	*	*	1.3	1.8	8.9	20.0
2003.....	0.4	*	*	*	*	*	*	*	*	2.7	5.9	25.4
2002.....	0.4	*	*	*	*	*	*	*	1.3	2.9	8.5	22.7
2001.....	0.4	*	*	*	*	*	*	*	1.1	3.0	7.4	26.3
2000.....	0.4	*	*	*	*	*	*	*	*	2.5	6.3	25.2
1999.....	0.4	*	*	*	*	*	*	*	*	3.1	8.1	23.2

Hispanic, male

2007.....	0.4	*	*	*	*	*	*	*	*	3.1	*	22.4
2006.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.4	*	*	*	*	*	*	*	*	*	8.4	*
2004.....	0.4	*	*	*	*	*	*	*	*	*	10.4	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	*	35.4
2002.....	0.4	*	*	*	*	*	*	*	*	*	8.2	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*

Hispanic, female

2007.....	0.4	*	*	*	*	*	*	*	*	*	5.9	19.3
2006.....	0.3	*	*	*	*	*	*	*	*	*	*	12.6
2005.....	0.5	*	*	*	*	*	*	*	*	3.3	6.3	22.8
2004.....	0.5	*	*	*	*	*	*	*	*	*	7.9	18.0
2003.....	0.4	*	*	*	*	*	*	*	*	3.1	*	20.3
2002.....	0.5	*	*	*	*	*	*	*	*	*	8.7	21.6
2001.....	0.5	*	*	*	*	*	*	*	*	*	7.3	24.2
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	26.7
1999.....	0.5	*	*	*	*	*	*	*	*	*	8.7	20.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

Non-Hispanic 4/, both sexes

2007.....	1.1	*	*	*	*	0.1	0.2	0.4	0.9	1.9	6.1	22.4
2006.....	1.1	*	*	*	*	0.1	0.2	0.4	0.8	1.9	6.2	23.0
2005.....	1.2	*	*	*	*	0.1	0.2	0.5	0.8	2.2	6.9	25.4
2004.....	1.1	*	*	*	*	*	0.2	0.5	0.9	2.1	6.5	24.9
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.1	25.6
2002.....	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.1	6.7	23.5
2001.....	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.7	23.1
2000.....	1.1	*	*	*	*	0.1	0.2	0.4	0.9	2.2	7.0	24.2
1999.....	1.3	*	*	*	*	0.1	0.3	0.4	1.1	2.5	8.9	29.3

Non-Hispanic 4/, male

2007.....	1.0	*	*	*	*	*	0.3	0.5	1.1	2.1	6.6	23.0
2006.....	1.0	*	*	*	*	0.1	0.3	0.5	1.0	2.0	6.9	24.3
2005.....	1.0	*	*	*	*	0.1	0.2	0.6	0.9	2.4	7.0	27.7
2004.....	1.0	*	*	*	*	*	0.3	0.6	1.1	2.2	7.0	24.4
2003.....	1.0	*	*	*	*	*	0.3	0.7	1.2	2.3	7.4	26.3
2002.....	0.9	*	*	*	*	*	0.2	0.6	1.2	2.3	6.9	22.9
2001.....	0.9	*	*	*	*	*	0.3	0.5	1.0	2.4	6.7	23.8
2000.....	0.9	*	*	*	*	*	0.2	0.5	0.9	2.4	7.0	27.9
1999.....	1.1	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.4

Non-Hispanic 4/, female

2007.....	1.2	*	*	*	*	*	0.1	0.3	0.7	1.8	5.8	22.2
2006.....	1.2	*	*	*	*	*	0.1	0.4	0.6	1.7	5.8	22.5
2005.....	1.3	*	*	*	*	*	0.2	0.3	0.6	2.0	6.8	24.4
2004.....	1.3	*	*	*	*	*	0.1	0.4	0.7	1.9	6.1	25.2
2003.....	1.3	*	*	*	*	*	0.1	0.4	0.7	2.1	6.8	25.3
2002.....	1.2	*	*	*	*	*	0.2	0.3	0.8	2.0	6.5	23.7
2001.....	1.2	*	*	*	*	*	0.2	0.2	0.9	2.0	6.7	22.8
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.8
1999.....	1.5	*	*	*	*	*	0.2	0.3	0.9	2.4	8.4	28.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

Non-Hispanic white, both sexes

2007 5/.....	0.9	*	*	*	*	*	*	0.1	0.4	1.2	5.0	20.9
2006 5/.....	0.9	*	*	*	*	*	*	0.2	0.4	1.1	5.1	21.3
2005 5/.....	1.0	*	*	*	*	*	*	0.2	0.3	1.3	5.5	24.0
2004 5/.....	0.9	*	*	*	*	*	*	0.1	0.4	1.3	5.3	23.0
2003 5/.....	1.0	*	*	*	*	*	*	0.1	0.4	1.3	5.9	23.2
2002.....	0.9	*	*	*	*	*	*	0.1	0.4	1.2	5.6	20.8
2001.....	0.9	*	*	*	*	*	*	0.1	0.5	1.4	5.6	20.7
2000.....	0.9	*	*	*	*	*	0.1	0.1	0.4	1.5	5.6	21.8
1999.....	1.1	*	*	*	*	*	*	0.2	0.5	1.4	7.3	26.1

Non-Hispanic white, male

2007 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.3	5.6	21.5
2006 5/.....	0.8	*	*	*	*	*	*	0.2	0.4	1.1	5.6	22.9
2005 5/.....	0.8	*	*	*	*	*	*	0.2	0.4	1.4	5.6	25.7
2004 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.5	5.8	22.8
2003 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.6	6.6	23.6
2002.....	0.7	*	*	*	*	*	*	*	0.5	1.4	5.9	20.2
2001.....	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.8	22.2
2000.....	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.1	24.3
1999.....	0.9	*	*	*	*	*	*	0.2	0.5	1.6	8.3	28.7

Non-Hispanic white, female

2007 5/.....	1.1	*	*	*	*	*	*	*	0.4	1.1	4.7	20.7
2006 5/.....	1.1	*	*	*	*	*	*	0.1	0.3	1.1	4.7	20.5
2005 5/.....	1.2	*	*	*	*	*	*	*	0.2	1.2	5.5	23.2
2004 5/.....	1.1	*	*	*	*	*	*	*	0.3	1.1	4.9	23.1
2003 5/.....	1.1	*	*	*	*	*	*	*	0.3	1.1	5.5	23.0
2002.....	1.1	*	*	*	*	*	*	*	0.3	1.1	5.4	21.0
2001.....	1.1	*	*	*	*	*	*	*	0.5	1.3	5.4	20.1
2000.....	1.1	*	*	*	*	*	*	*	0.3	1.4	5.2	20.8
1999.....	1.2	*	*	*	*	*	*	*	0.4	1.3	6.6	25.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Hypertensive heart and renal disease (I13)

##### Non-Hispanic black, both sexes

2007 5/.....	2.1	*	*	*	*	*	1.0	2.1	4.7	8.5	17.6	40.1
2006 5/.....	2.2	*	*	*	*	*	0.9	2.2	4.5	8.7	18.8	42.5
2005 5/.....	2.4	*	*	*	*	0.5	1.0	2.5	4.6	9.8	21.8	42.9
2004 5/.....	2.4	*	*	*	*	*	1.2	2.9	5.0	8.7	20.0	49.2
2003 5/.....	2.5	*	*	*	*	0.4	1.3	3.1	5.5	9.3	19.9	53.8
2002.....	2.5	*	*	*	*	*	1.0	2.8	6.2	10.0	19.8	57.2
2001.....	2.3	*	*	*	*	0.4	1.2	2.1	5.1	10.1	20.2	50.6
2000.....	2.3	*	*	*	*	*	0.8	2.2	4.7	10.0	22.6	50.9
1999.....	2.8	*	*	*	*	*	1.5	2.3	6.2	12.7	27.3	66.0

##### Non-Hispanic black, male

2007 5/.....	2.2	*	*	*	*	*	1.6	2.7	6.7	9.8	19.8	42.8
2006 5/.....	2.2	*	*	*	*	*	1.3	2.7	6.3	10.7	23.4	40.3
2005 5/.....	2.4	*	*	*	*	*	1.3	3.4	6.0	11.3	23.5	50.4
2004 5/.....	2.3	*	*	*	*	*	1.9	3.6	6.3	8.8	23.7	42.7
2003 5/.....	2.4	*	*	*	*	*	1.8	4.1	7.2	9.1	19.0	51.1
2002.....	2.3	*	*	*	*	*	1.0	3.9	7.7	10.2	19.3	62.4
2001.....	2.2	*	*	*	*	*	1.6	3.2	6.0	12.0	18.6	45.0
2000.....	2.1	*	*	*	*	*	1.0	2.9	5.4	11.1	20.1	64.6
1999.....	2.6	*	*	*	*	*	1.8	3.3	7.8	13.8	26.0	60.6

##### Non-Hispanic black, female

2007 5/.....	2.1	*	*	*	*	*	*	1.5	3.0	7.6	16.4	39.1
2006 5/.....	2.1	*	*	*	*	*	*	1.7	3.1	7.2	16.3	43.3
2005 5/.....	2.3	*	*	*	*	*	0.7	1.7	3.4	8.7	20.8	39.8
2004 5/.....	2.4	*	*	*	*	*	*	2.3	4.0	8.5	17.9	51.8
2003 5/.....	2.6	*	*	*	*	*	0.8	2.2	4.2	9.4	20.4	54.9
2002.....	2.6	*	*	*	*	*	0.9	1.8	5.0	9.9	20.1	55.2
2001.....	2.4	*	*	*	*	*	0.8	1.2	4.3	8.7	21.1	52.6
2000.....	2.4	*	*	*	*	*	*	1.6	4.2	9.2	24.0	45.8
1999.....	3.1	*	*	*	*	*	1.2	1.5	4.8	12.0	28.0	68.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Ischemic heart diseases (I20-I25)

All origins 3/, both sexes

2007.....	134.7	0.6	*	0.1	0.4	2.6	14.4	55.6	141.1	325.7	891.9	2,705.3
2006.....	142.1	0.5	*	*	0.3	2.6	15.3	57.7	147.9	352.5	959.8	2,904.7
2005.....	150.4	*	*	*	0.4	2.5	15.6	59.6	154.2	376.2	1,029.8	3,158.1
2004.....	153.7	*	*	*	0.3	2.7	16.2	59.9	158.2	395.9	1,072.6	3,287.4
2003.....	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002.....	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001.....	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000.....	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999.....	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1

All origins 3/, male

2007.....	145.3	*	*	*	0.5	3.9	21.9	85.8	212.2	458.8	1,180.1	3,089.2
2006.....	152.2	*	*	*	0.5	4.0	23.2	89.1	220.7	494.2	1,269.7	3,303.0
2005.....	159.0	*	*	*	0.6	3.8	23.6	92.0	229.4	521.7	1,351.5	3,576.8
2004.....	161.6	*	*	*	0.4	4.0	24.4	92.8	234.8	549.1	1,410.3	3,680.3
2003.....	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002.....	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001.....	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000.....	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999.....	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5

All origins 3/, female

2007.....	124.4	*	*	*	0.2	1.2	6.9	26.3	74.9	212.6	693.3	2,522.7
2006.....	132.3	*	*	*	0.2	1.1	7.4	27.4	80.1	232.6	747.9	2,718.3
2005.....	142.0	*	*	*	0.1	1.2	7.6	28.2	84.2	253.4	811.4	2,965.8
2004.....	146.1	*	*	*	0.1	1.4	8.0	28.0	87.1	267.3	845.3	3,110.6
2003.....	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002.....	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001.....	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000.....	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999.....	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Hispanic, both sexes												
2007.....	44.9	*	*	*	*	1.3	6.7	31.8	102.5	262.9	749.4	2,082.1
2006.....	47.2	*	*	*	0.3	1.3	7.2	34.3	106.8	286.6	813.3	2,296.7
2005.....	51.0	*	*	*	0.3	1.3	7.9	35.7	115.4	326.7	910.6	2,528.4
2004.....	49.6	*	*	*	*	1.2	7.7	35.3	117.4	326.5	906.9	2,613.4
2003.....	52.1	*	*	*	0.3	1.4	7.5	35.4	122.0	355.0	973.0	2,970.0
2002.....	54.0	*	*	*	*	1.2	7.6	38.0	135.9	382.0	1,010.8	3,182.9
2001.....	55.9	*	*	*	*	1.6	6.8	38.7	130.2	400.9	1,099.1	3,620.2
2000.....	55.9	*	*	*	*	1.1	7.6	39.5	143.6	405.3	1,111.3	3,697.2
1999.....	58.7	*	*	*	*	1.4	7.4	44.8	150.4	433.5	1,178.0	3,876.2
Hispanic, male												
2007.....	48.0	*	*	*	*	2.1	10.3	50.2	154.6	369.8	943.8	2,143.6
2006.....	49.9	*	*	*	*	1.9	10.9	52.7	156.5	394.1	1,022.8	2,453.9
2005.....	54.2	*	*	*	*	1.9	12.0	55.2	171.0	449.6	1,165.5	2,694.8
2004.....	52.7	*	*	*	*	1.9	11.9	55.2	175.1	455.7	1,175.3	2,695.4
2003.....	53.7	*	*	*	0.5	2.1	11.0	55.8	179.7	494.6	1,214.9	2,933.9
2002.....	56.5	*	*	*	*	1.9	11.4	58.0	204.0	530.4	1,271.7	3,374.9
2001.....	57.9	*	*	*	*	2.4	10.4	60.3	191.9	559.9	1,393.3	3,828.2
2000.....	58.2	*	*	*	*	1.6	12.1	62.7	214.8	561.5	1,420.0	3,865.6
1999.....	60.8	*	*	*	*	2.1	12.2	71.3	218.3	608.5	1,495.9	3,987.7
Hispanic, female												
2007.....	41.6	*	*	*	*	*	2.6	12.7	54.2	175.5	612.8	2,048.7
2006.....	44.4	*	*	*	*	0.6	2.9	15.5	60.9	199.2	666.6	2,213.2
2005.....	47.6	*	*	*	*	*	3.2	15.9	64.5	227.5	733.1	2,441.3
2004.....	46.2	*	*	*	*	*	3.1	15.4	65.0	222.8	721.6	2,571.3
2003.....	50.4	*	*	*	*	0.6	3.7	15.0	69.8	243.7	806.9	2,988.3
2002.....	51.4	*	*	*	*	*	3.5	18.1	74.6	263.9	832.2	3,087.8
2001.....	53.7	*	*	*	*	*	2.8	17.4	75.4	275.4	900.3	3,518.6
2000.....	53.6	*	*	*	*	*	2.8	16.8	80.6	282.5	904.0	3,614.6
1999.....	56.6	*	*	*	*	*	2.3	19.0	90.5	296.4	966.5	3,822.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic 4/, both sexes												
2007.....	150.4	*	*	*	0.4	2.9	15.8	58.3	144.1	329.9	899.8	2,736.0
2006.....	158.2	0.6	*	*	0.3	2.9	16.8	60.3	150.9	356.5	967.2	2,931.2
2005.....	166.7	*	*	*	0.4	2.8	16.9	62.0	156.9	379.0	1,034.7	3,183.8
2004.....	170.3	*	*	*	0.3	3.1	17.5	62.3	160.9	399.6	1,079.3	3,312.1
2003.....	182.5	0.6	*	0.1	0.4	3.0	18.4	64.7	172.8	433.9	1,167.1	3,629.3
2002.....	189.0	*	*	*	0.3	2.8	18.4	66.1	180.1	462.1	1,226.5	3,787.0
2001.....	193.6	0.7	*	*	0.4	3.1	18.0	66.1	187.3	480.9	1,274.9	3,959.4
2000.....	200.5	0.8	*	*	0.3	2.6	17.7	68.1	199.5	509.0	1,328.8	4,187.0
1999.....	207.1	0.8	*	0.1	0.4	2.7	18.3	69.4	206.3	540.3	1,388.1	4,319.6
Non-Hispanic 4/, male												
2007.....	163.3	*	*	*	0.5	4.4	24.1	90.0	216.6	464.4	1,193.7	3,141.5
2006.....	170.6	*	*	*	0.5	4.6	25.5	93.2	225.4	500.0	1,282.9	3,344.5
2005.....	177.2	*	*	*	0.6	4.3	25.7	95.9	233.5	525.5	1,359.6	3,618.5
2004.....	179.9	*	*	*	0.5	4.6	26.4	96.6	238.7	553.4	1,419.8	3,722.6
2003.....	191.5	*	*	*	0.5	4.5	27.9	101.1	255.9	603.6	1,536.0	4,045.5
2002.....	197.6	*	*	*	0.5	4.1	28.3	103.1	266.5	644.3	1,609.7	4,228.4
2001.....	200.4	*	*	*	0.6	4.5	27.5	102.8	276.1	667.7	1,670.1	4,408.5
2000.....	207.6	*	*	*	0.4	3.9	27.4	106.7	294.0	710.4	1,744.6	4,721.1
1999.....	214.1	*	*	*	0.5	4.0	28.3	108.8	303.4	754.2	1,820.2	4,854.5
Non-Hispanic 4/, female												
2007.....	138.1	*	*	*	0.2	1.4	7.6	27.9	76.5	215.3	697.6	2,544.6
2006.....	146.4	*	*	*	0.2	1.3	8.2	28.7	81.6	234.7	751.6	2,739.1
2005.....	156.7	*	*	*	0.1	1.4	8.3	29.4	85.7	254.9	814.5	2,985.4
2004.....	161.1	*	*	*	0.2	1.6	8.7	29.3	88.6	269.9	850.4	3,128.5
2003.....	173.9	*	*	*	0.3	1.5	9.0	29.6	95.6	291.5	920.9	3,446.4
2002.....	180.7	*	*	*	0.2	1.5	8.8	30.5	100.0	309.9	972.5	3,596.5
2001.....	187.2	*	*	*	0.2	1.6	8.7	30.8	105.0	325.6	1,015.5	3,772.7
2000.....	193.8	*	*	*	0.2	1.3	8.2	31.0	112.1	342.8	1,057.9	3,971.0
1999.....	200.6	*	*	*	0.3	1.5	8.5	31.5	116.6	364.6	1,107.8	4,105.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic white, both sexes												
2007 5/.....	166.0	*	*	*	0.3	2.6	15.5	56.1	137.8	319.6	901.3	2,794.7
2006 5/.....	174.5	*	*	*	0.3	2.8	16.5	57.4	144.0	345.7	966.4	2,992.4
2005 5/.....	183.9	*	*	*	0.3	2.6	16.4	58.8	149.0	369.5	1,033.7	3,253.9
2004 5/.....	187.7	*	*	*	0.2	2.8	17.2	59.1	153.5	388.1	1,079.6	3,369.7
2003 5/.....	200.8	*	*	*	0.4	2.8	17.9	61.1	164.5	421.6	1,164.6	3,691.6
2002.....	208.0	*	*	*	0.3	2.6	17.9	62.8	171.2	450.8	1,223.1	3,848.2
2001.....	212.6	*	*	*	0.3	2.9	17.3	62.6	177.8	467.8	1,271.3	4,022.8
2000.....	220.2	*	*	*	0.3	2.4	16.9	64.6	189.5	496.4	1,324.5	4,261.3
1999.....	227.2	*	*	*	0.3	2.5	17.3	65.8	196.4	527.7	1,386.5	4,390.2
Non-Hispanic white, male												
2007 5/.....	179.9	*	*	*	0.4	4.0	23.8	87.8	209.1	454.4	1,203.2	3,241.0
2006 5/.....	187.5	*	*	*	0.5	4.4	25.6	90.2	216.7	488.6	1,288.8	3,441.2
2005 5/.....	195.1	*	*	*	0.5	4.0	25.4	92.4	224.3	517.2	1,367.9	3,732.9
2004 5/.....	198.0	*	*	*	0.3	4.3	26.4	93.4	230.4	543.1	1,431.6	3,814.0
2003 5/.....	210.2	*	*	*	0.4	4.3	27.5	97.2	247.3	591.8	1,541.1	4,149.5
2002.....	217.2	*	*	*	0.4	4.0	28.0	100.0	256.8	636.0	1,618.3	4,324.5
2001.....	219.6	*	*	*	0.5	4.3	26.9	99.5	266.3	656.5	1,678.7	4,506.9
2000.....	227.6	*	*	*	0.4	3.7	26.8	103.5	283.5	701.7	1,752.3	4,833.4
1999.....	234.2	*	*	*	0.5	3.6	27.2	105.1	292.3	745.2	1,834.9	4,954.1
Non-Hispanic white, female												
2007 5/.....	152.6	*	*	*	0.2	1.3	7.2	24.9	69.8	202.3	690.0	2,583.4
2006 5/.....	161.9	*	*	*	*	1.1	7.3	25.1	74.6	221.9	742.7	2,783.1
2005 5/.....	173.1	*	*	*	*	1.3	7.4	25.7	77.3	241.8	803.8	3,034.8
2004 5/.....	177.7	*	*	*	*	1.3	7.9	25.4	80.4	254.7	839.7	3,170.6
2003 5/.....	191.8	*	*	*	0.3	1.4	8.2	25.5	85.9	275.9	910.4	3,490.0
2002.....	199.1	*	*	*	0.2	1.3	7.8	26.1	90.2	293.0	958.3	3,642.5
2001.....	205.8	*	*	*	0.2	1.4	7.6	26.4	94.2	307.7	1,001.1	3,821.4
2000.....	213.1	*	*	*	0.2	1.0	6.9	26.4	100.9	323.6	1,043.3	4,030.7
1999.....	220.5	*	*	*	*	1.4	7.4	27.2	106.1	345.6	1,092.9	4,165.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic black, both sexes												
2007 5/.....	110.5	*	*	*	0.8	4.6	21.0	86.6	227.1	482.2	1,046.5	2,494.5
2006 5/.....	116.5	*	*	*	0.5	4.3	23.1	92.8	238.7	516.6	1,143.3	2,640.8
2005 5/.....	121.7	*	*	*	0.6	4.1	24.0	96.3	255.2	530.1	1,226.6	2,830.2
2004 5/.....	124.8	*	*	*	0.7	4.9	24.0	97.9	259.1	576.3	1,255.5	3,038.5
2003 5/.....	133.2	*	*	*	0.7	4.4	25.5	101.8	277.3	622.3	1,375.4	3,328.7
2002.....	137.0	*	*	*	0.5	4.1	25.8	103.6	292.4	643.9	1,433.9	3,481.1
2001.....	141.2	*	*	*	0.9	4.8	26.4	104.6	302.1	681.4	1,487.0	3,628.7
2000.....	144.4	*	*	*	0.4	4.4	26.1	107.7	318.9	710.1	1,534.5	3,701.7
1999.....	147.9	*	*	*	0.8	4.1	28.2	110.8	324.7	741.6	1,559.0	3,817.9
Non-Hispanic black, male												
2007 5/.....	118.0	*	*	*	1.1	7.2	30.9	125.4	331.7	660.6	1,337.8	2,528.7
2006 5/.....	124.5	*	*	*	*	6.3	32.2	133.8	347.2	717.2	1,476.8	2,685.6
2005 5/.....	127.0	*	*	*	0.8	6.3	33.6	140.2	364.5	712.6	1,542.6	2,808.9
2004 5/.....	129.2	*	*	*	1.2	6.7	32.9	141.2	368.2	771.0	1,551.8	3,133.7
2003 5/.....	137.6	*	*	*	0.9	6.4	36.2	148.3	387.0	836.1	1,758.6	3,309.2
2002.....	139.5	*	*	*	*	5.3	36.2	146.9	413.2	850.8	1,762.3	3,572.8
2001.....	143.2	*	*	*	1.2	6.8	37.1	149.0	418.0	903.1	1,834.9	3,791.8
2000.....	145.7	*	*	*	*	6.1	36.3	152.2	444.2	916.5	1,887.6	3,911.2
1999.....	150.0	*	*	*	0.9	6.0	40.3	160.0	458.4	967.5	1,873.3	4,082.5
Non-Hispanic black, female												
2007 5/.....	103.6	*	*	*	*	2.3	12.3	53.3	143.1	354.8	880.8	2,480.4
2006 5/.....	109.2	*	*	*	*	2.5	15.2	57.7	151.4	373.4	954.0	2,622.6
2005 5/.....	116.8	*	*	*	*	2.1	15.5	58.7	167.2	399.7	1,047.6	2,838.7
2004 5/.....	120.8	*	*	*	*	3.2	16.0	60.7	171.5	437.6	1,088.3	3,000.9
2003 5/.....	129.1	*	*	*	*	2.6	16.0	62.0	189.2	470.4	1,159.1	3,336.2
2002.....	134.8	*	*	*	*	2.9	16.7	66.5	195.4	497.1	1,248.1	3,446.0
2001.....	139.4	*	*	*	*	2.9	16.9	66.5	208.9	525.0	1,291.6	3,568.3
2000.....	143.2	*	*	*	*	2.9	17.1	69.4	218.1	565.0	1,335.7	3,623.9
1999.....	146.0	*	*	*	0.7	2.4	17.4	68.7	217.5	582.3	1,383.0	3,719.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute myocardial infarction (I21-I22)

All origins 3/, both sexes

2007.....	44.1	*	*	*	0.1	1.0	5.6	21.6	54.5	121.1	288.9	756.7
2006.....	47.2	*	*	*	0.1	1.0	6.1	22.5	58.2	131.1	319.0	825.8
2005.....	50.9	*	*	*	0.2	1.0	6.2	23.7	61.1	143.1	348.2	923.1
2004.....	53.4	*	*	*	0.1	1.1	6.4	24.2	64.5	153.4	370.3	992.9
2003.....	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002.....	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001.....	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000.....	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999.....	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3

All origins 3/, male

2007.....	48.2	*	*	*	0.2	1.4	8.2	33.0	80.8	166.8	370.8	855.8
2006.....	51.6	*	*	*	0.2	1.5	9.2	34.2	85.6	181.9	411.8	936.3
2005.....	54.8	*	*	*	0.2	1.5	9.2	36.2	89.9	195.5	445.9	1,045.0
2004.....	57.4	*	*	*	0.2	1.6	9.6	37.3	95.3	209.5	477.0	1,120.5
2003.....	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002.....	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001.....	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000.....	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999.....	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1

All origins 3/, female

2007.....	40.0	*	*	*	0.1	0.5	2.9	10.5	30.1	82.3	232.5	709.5
2006.....	43.0	*	*	*	*	0.5	3.0	11.2	32.6	88.1	255.6	774.1
2005.....	47.2	*	*	*	*	0.5	3.2	11.6	34.3	98.9	281.8	867.1
2004.....	49.6	*	*	*	*	0.6	3.3	11.6	35.9	106.3	298.5	935.5
2003.....	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002.....	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001.....	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000.....	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999.....	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Hispanic, both sexes												
2007.....	14.9	*	*	*	*	0.4	2.3	11.4	35.7	89.8	252.0	636.8
2006.....	15.7	*	*	*	*	0.5	2.7	11.5	37.7	100.5	273.6	692.9
2005.....	17.6	*	*	*	*	0.5	2.7	12.5	43.2	119.4	318.7	787.2
2004.....	17.3	*	*	*	*	0.5	2.9	12.9	44.2	120.4	315.7	816.8
2003.....	18.5	*	*	*	*	0.5	3.0	13.3	44.5	137.7	341.6	952.0
2002.....	19.7	*	*	*	*	0.5	3.0	14.2	55.1	146.2	364.3	1,048.2
2001.....	20.0	*	*	*	*	0.7	2.5	14.8	49.5	149.8	394.9	1,185.5
2000.....	20.1	*	*	*	*	0.5	3.0	15.5	55.0	154.3	404.7	1,177.1
1999.....	21.1	*	*	*	*	0.5	2.9	17.8	56.1	164.0	433.7	1,227.8
Hispanic, male												
2007.....	16.0	*	*	*	*	0.5	3.4	18.2	51.6	123.3	324.4	652.5
2006.....	16.5	*	*	*	*	0.7	3.9	16.7	52.7	137.7	338.6	742.0
2005.....	18.8	*	*	*	*	0.6	3.8	18.9	62.2	160.9	406.2	875.4
2004.....	18.4	*	*	*	*	0.6	4.3	20.1	65.4	164.1	403.4	840.6
2003.....	18.9	*	*	*	*	0.7	4.1	20.6	64.1	187.5	409.2	963.6
2002.....	20.6	*	*	*	*	0.7	4.2	21.0	80.8	199.3	460.0	1,107.7
2001.....	20.9	*	*	*	*	1.0	3.9	22.7	69.4	206.9	488.7	1,331.3
2000.....	21.1	*	*	*	*	0.7	4.7	23.4	80.0	207.6	514.8	1,273.8
1999.....	22.0	*	*	*	*	0.8	4.6	26.7	79.0	225.8	558.9	1,276.8
Hispanic, female												
2007.....	13.8	*	*	*	*	*	1.1	4.4	21.1	62.4	201.1	628.3
2006.....	14.9	*	*	*	*	*	1.4	6.2	23.9	70.4	228.2	666.8
2005.....	16.4	*	*	*	*	*	1.6	6.0	25.8	85.8	257.7	741.0
2004.....	16.0	*	*	*	*	*	1.3	5.6	24.9	85.4	255.1	804.5
2003.....	18.0	*	*	*	*	*	1.7	6.0	26.8	98.1	295.2	946.1
2002.....	18.7	*	*	*	*	*	1.8	7.4	32.1	103.9	298.8	1,018.8
2001.....	19.1	*	*	*	*	*	1.1	7.0	31.8	104.7	331.5	1,114.2
2000.....	19.1	*	*	*	*	*	1.2	7.8	32.9	112.3	330.8	1,129.7
1999.....	20.1	*	*	*	*	*	1.1	9.1	35.9	115.5	350.4	1,204.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic 4/, both sexes												
2007.....	49.2	*	*	*	0.1	1.1	6.2	22.8	56.1	123.4	291.0	762.6
2006.....	52.6	*	*	*	0.1	1.2	6.7	23.7	59.8	133.1	321.3	831.0
2005.....	56.5	*	*	*	0.2	1.1	6.8	24.9	62.5	144.7	349.4	928.4
2004.....	59.2	*	*	*	0.1	1.3	7.0	25.4	66.0	155.4	372.7	999.2
2003.....	64.9	*	*	*	0.2	1.2	7.6	26.7	72.8	170.4	415.5	1,123.8
2002.....	68.7	*	*	*	0.2	1.1	7.6	27.7	76.3	188.2	447.3	1,194.0
2001.....	71.4	*	*	*	0.2	1.3	7.6	28.8	80.3	195.6	473.6	1,265.3
2000.....	75.3	*	*	*	0.1	1.0	7.4	30.0	88.6	211.9	506.3	1,340.8
1999.....	78.2	*	*	*	0.2	1.2	7.7	30.5	92.2	227.9	531.9	1,381.0
Non-Hispanic 4/, male												
2007.....	54.2	*	*	*	0.2	1.7	9.1	34.8	83.2	169.8	373.3	867.2
2006.....	57.9	*	*	*	0.2	1.8	10.2	36.2	88.2	184.6	415.9	945.2
2005.....	61.2	*	*	*	0.3	1.7	10.2	38.0	92.0	197.8	447.5	1,052.8
2004.....	64.0	*	*	*	0.2	1.8	10.5	39.1	97.5	212.1	480.2	1,132.5
2003.....	69.7	*	*	*	0.2	1.7	11.4	41.8	106.9	233.4	535.7	1,274.5
2002.....	73.5	*	*	*	0.2	1.6	11.5	42.6	111.3	259.2	573.4	1,374.2
2001.....	75.9	*	*	*	0.2	1.8	11.6	44.2	117.0	267.7	607.9	1,463.9
2000.....	80.3	*	*	*	0.2	1.5	11.4	46.5	128.7	290.9	652.6	1,576.2
1999.....	83.8	*	*	*	0.2	1.7	11.8	47.8	135.1	313.8	687.7	1,631.5
Non-Hispanic 4/, female												
2007.....	44.4	*	*	*	0.6	3.3	11.2	30.8	83.9	234.4	713.2	
2006.....	47.6	*	*	*	0.6	3.3	11.8	33.4	89.4	256.7	778.0	
2005.....	52.0	*	*	*	0.5	3.5	12.2	35.0	99.7	282.9	871.7	
2004.....	54.7	*	*	*	0.7	3.6	12.2	36.7	107.7	300.4	939.6	
2003.....	60.3	*	*	*	0.2	0.6	3.8	12.3	41.0	117.5	335.3	1,057.5
2002.....	64.1	*	*	*	0.6	3.8	13.3	44.0	128.9	363.8	1,116.3	
2001.....	67.1	*	*	*	0.8	3.8	13.9	46.3	135.7	385.4	1,182.7	
2000.....	70.6	*	*	*	0.1	0.6	3.5	14.1	51.6	146.8	411.0	1,245.6
1999.....	72.9	*	*	*	0.7	3.7	13.9	52.5	157.3	430.7	1,280.8	

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic white, both sexes												
2007 5/.....	54.2	*	*	*	0.1	1.0	6.2	22.6	55.3	121.6	291.2	772.5
2006 5/.....	58.0	*	*	*	0.1	1.1	6.8	23.4	58.8	131.4	321.4	841.4
2005 5/.....	62.3	*	*	*	0.1	1.0	6.8	24.5	61.3	143.7	349.1	941.9
2004 5/.....	65.3	*	*	*	0.1	1.2	7.0	24.8	64.9	152.9	374.2	1,010.1
2003 5/.....	71.5	*	*	*	0.2	1.1	7.5	25.9	71.4	168.3	415.1	1,137.4
2002.....	75.5	*	*	*	0.1	1.1	7.5	27.0	74.7	186.1	445.9	1,202.8
2001.....	78.4	*	*	*	0.1	1.2	7.5	28.1	78.7	193.0	473.0	1,276.8
2000.....	82.7	*	*	*	0.1	0.9	7.3	29.3	86.9	209.3	505.3	1,355.3
1999.....	86.0	*	*	*	0.1	1.1	7.5	30.0	90.7	226.1	532.4	1,394.2
Non-Hispanic white, male												
2007 5/.....	60.0	*	*	*	0.1	1.6	9.3	35.1	83.0	168.9	375.9	888.1
2006 5/.....	64.0	*	*	*	0.2	1.7	10.5	36.3	87.5	184.4	418.6	964.6
2005 5/.....	67.8	*	*	*	0.3	1.6	10.4	38.2	91.3	198.9	451.0	1,077.9
2004 5/.....	70.9	*	*	*	*	1.9	10.7	38.8	97.3	210.8	487.4	1,156.5
2003 5/.....	77.0	*	*	*	0.2	1.6	11.5	41.2	106.5	232.9	537.8	1,300.9
2002.....	81.2	*	*	*	0.2	1.6	11.5	42.5	110.8	260.2	575.8	1,393.0
2001.....	83.7	*	*	*	0.2	1.7	11.6	44.3	116.8	267.7	612.5	1,483.2
2000.....	88.5	*	*	*	0.2	1.4	11.4	46.7	128.0	291.0	656.7	1,600.7
1999.....	92.4	*	*	*	0.2	1.5	11.8	48.1	134.6	315.0	695.7	1,653.8
Non-Hispanic white, female												
2007 5/.....	48.7	*	*	*	*	0.5	3.1	10.3	28.8	80.4	231.8	717.7
2006 5/.....	52.2	*	*	*	*	0.5	3.1	10.7	31.6	85.4	254.0	783.9
2005 5/.....	57.0	*	*	*	*	0.4	3.2	11.1	32.7	95.9	279.1	879.7
2004 5/.....	59.9	*	*	*	*	0.6	3.3	10.9	34.2	103.1	297.1	944.5
2003 5/.....	66.2	*	*	*	*	0.6	3.5	10.8	38.0	113.0	332.2	1,065.3
2002.....	70.1	*	*	*	*	0.5	3.6	11.8	40.6	122.9	358.8	1,120.7
2001.....	73.4	*	*	*	*	0.8	3.4	12.2	42.7	129.6	380.6	1,190.9
2000.....	77.2	*	*	*	*	0.5	3.1	12.3	48.2	140.5	405.9	1,256.3
1999.....	79.8	*	*	*	*	0.7	3.3	12.3	49.3	151.8	425.5	1,290.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic black, both sexes												
2007 5/.....	36.6	*	*	*	*	1.8	7.6	29.6	77.6	166.4	342.4	763.0
2006 5/.....	38.7	*	*	*	*	1.7	8.5	32.4	82.5	175.7	374.2	811.7
2005 5/.....	41.1	*	*	*	*	1.8	9.1	33.6	86.9	182.8	413.4	901.2
2004 5/.....	43.4	*	*	*	0.3	1.8	9.1	36.2	93.0	208.2	424.3	991.3
2003 5/.....	47.3	*	*	*	*	1.7	10.4	38.6	100.6	223.1	486.5	1,103.6
2002.....	50.2	*	*	*	*	1.7	10.0	39.0	109.1	242.6	523.0	1,212.9
2001.....	51.6	*	*	*	0.4	1.9	10.2	40.2	111.6	256.0	543.7	1,247.0
2000.....	54.0	*	*	*	*	1.8	10.1	41.9	122.3	273.0	576.6	1,281.8
1999.....	55.1	*	*	*	*	1.7	10.5	42.0	123.2	282.4	589.9	1,328.6
Non-Hispanic black, male												
2007 5/.....	38.1	*	*	*	*	2.6	10.8	41.3	107.8	219.1	416.4	767.9
2006 5/.....	40.6	*	*	*	*	2.5	11.9	45.1	117.3	233.0	463.5	824.6
2005 5/.....	41.8	*	*	*	*	2.6	12.2	46.6	120.2	233.2	499.2	898.3
2004 5/.....	43.9	*	*	*	*	2.1	12.0	51.6	126.2	269.9	491.7	1,020.2
2003 5/.....	48.2	*	*	*	*	2.4	14.3	54.7	135.9	287.1	606.9	1,121.9
2002.....	50.1	*	*	*	*	2.2	13.7	53.4	144.5	305.4	631.3	1,296.3
2001.....	50.9	*	*	*	*	2.5	13.5	53.4	146.1	320.8	646.6	1,389.4
2000.....	53.7	*	*	*	*	2.2	13.6	56.3	163.1	341.2	693.3	1,426.6
1999.....	55.2	*	*	*	*	2.5	14.5	58.1	168.7	357.0	691.4	1,509.4
Non-Hispanic black, female												
2007 5/.....	35.3	*	*	*	*	1.0	4.8	19.6	53.3	128.7	300.3	761.0
2006 5/.....	37.0	*	*	*	*	1.0	5.5	21.6	54.5	134.8	323.4	806.4
2005 5/.....	40.5	*	*	*	*	1.0	6.3	22.5	60.2	146.8	364.8	902.3
2004 5/.....	43.0	*	*	*	*	1.5	6.5	23.0	66.2	164.3	386.3	979.8
2003 5/.....	46.5	*	*	*	*	1.0	6.9	24.7	72.2	177.6	418.5	1,096.5
2002.....	50.4	*	*	*	*	1.3	6.7	26.7	80.7	198.0	461.8	1,181.0
2001.....	52.3	*	*	*	*	1.3	7.4	28.8	83.8	210.3	485.9	1,194.3
2000.....	54.2	*	*	*	*	1.4	7.0	29.6	89.5	225.0	510.9	1,228.0
1999.....	54.9	*	*	*	*	1.0	7.0	28.2	86.8	229.7	533.1	1,261.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

All origins 3/, both sexes

2007.....	1.4	*	*	*	*	*	0.3	0.9	2.1	3.8	7.8	20.8
2006.....	1.3	*	*	*	*	0.1	0.2	0.8	1.9	3.6	8.0	21.5
2005.....	1.2	*	*	*	*	*	0.2	0.8	1.7	3.3	7.2	20.8
2004.....	0.8	*	*	*	*	*	0.2	0.5	1.2	2.2	5.2	14.8
2003.....	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002.....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001.....	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999.....	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9

All origins 3/, male

2007.....	1.5	*	*	*	*	*	0.4	1.2	3.1	5.4	9.4	18.7
2006.....	1.4	*	*	*	*	*	0.2	1.2	2.6	4.6	9.6	21.8
2005.....	1.2	*	*	*	*	*	0.3	1.2	2.5	4.3	8.5	20.2
2004.....	0.9	*	*	*	*	*	0.3	0.7	1.8	3.0	6.3	16.3
2003.....	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002.....	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001.....	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999.....	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3

All origins 3/, female

2007.....	1.3	*	*	*	*	*	0.2	0.5	1.1	2.5	6.8	21.8
2006.....	1.3	*	*	*	*	*	0.2	0.5	1.2	2.7	6.8	21.4
2005.....	1.2	*	*	*	*	*	0.1	0.5	0.9	2.4	6.2	21.1
2004.....	0.8	*	*	*	*	*	0.1	0.3	0.7	1.4	4.4	14.1
2003.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001.....	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000.....	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999.....	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

Non-Hispanic 4/, both sexes

2007.....	1.6	*	*	*	*	*	0.3	0.9	2.2	4.0	8.1	21.4
2006.....	1.5	*	*	*	*	*	0.2	0.9	2.0	3.7	8.2	22.1
2005.....	1.4	*	*	*	*	*	0.2	0.9	1.8	3.4	7.4	21.2
2004.....	1.0	*	*	*	*	*	0.2	0.6	1.3	2.3	5.4	15.3
2003.....	1.2	*	*	*	*	*	0.3	0.7	1.5	3.3	6.7	20.6
2002.....	1.3	*	*	*	*	*	0.2	0.9	1.8	3.6	7.3	21.8
2001.....	1.3	*	*	*	*	*	0.2	0.8	1.9	3.6	7.1	22.0
2000.....	1.3	*	*	*	*	*	0.2	0.8	2.1	3.6	7.5	22.2
1999.....	1.3	*	*	*	*	*	0.2	0.7	2.0	3.8	7.2	22.2

Non-Hispanic 4/, male

2007.....	1.7	*	*	*	*	*	0.4	1.4	3.2	5.6	9.7	19.2
2006.....	1.6	*	*	*	*	*	0.3	1.3	2.8	4.8	10.0	22.6
2005.....	1.4	*	*	*	*	*	0.3	1.3	2.7	4.4	8.7	20.9
2004.....	1.0	*	*	*	*	*	0.3	0.8	1.9	3.2	6.5	17.0
2003.....	1.3	*	*	*	*	*	0.4	1.1	2.2	4.5	8.4	20.2
2002.....	1.4	*	*	*	*	*	0.3	1.4	2.6	4.9	9.0	22.1
2001.....	1.4	*	*	*	*	*	0.3	1.3	2.7	4.8	8.7	20.9
2000.....	1.4	*	*	*	*	*	0.3	1.3	3.1	4.9	9.3	21.0
1999.....	1.4	*	*	*	*	*	0.3	1.1	2.9	5.2	9.0	22.9

Non-Hispanic 4/, female

2007.....	1.4	*	*	*	*	*	0.2	0.5	1.2	2.6	7.0	22.5
2006.....	1.4	*	*	*	*	*	0.2	0.5	1.3	2.8	7.0	21.9
2005.....	1.3	*	*	*	*	*	0.1	0.5	1.0	2.5	6.5	21.4
2004.....	0.9	*	*	*	*	*	0.1	0.3	0.8	1.5	4.6	14.6
2003.....	1.2	*	*	*	*	*	0.1	0.4	0.9	2.3	5.7	20.7
2002.....	1.3	*	*	*	*	*	0.1	0.5	1.1	2.5	6.1	21.6
2001.....	1.3	*	*	*	*	*	0.1	0.4	1.1	2.6	6.0	22.4
2000.....	1.2	*	*	*	*	*	0.1	0.3	1.2	2.5	6.3	22.7
1999.....	1.2	*	*	*	*	*	*	0.4	1.1	2.7	6.0	21.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

Non-Hispanic white, both sexes

2007 5/.....	1.7	*	*	*	*	*	0.3	0.9	2.1	3.8	8.0	21.6
2006 5/.....	1.6	*	*	*	*	*	0.2	0.8	1.9	3.5	8.1	22.3
2005 5/.....	1.4	*	*	*	*	*	0.2	0.7	1.6	3.2	7.1	20.9
2004 5/.....	1.0	*	*	*	*	*	0.2	0.5	1.2	2.1	5.1	15.6
2003 5/.....	1.3	*	*	*	*	*	0.2	0.7	1.4	3.1	6.6	20.9
2002.....	1.4	*	*	*	*	*	0.2	0.8	1.7	3.5	7.2	22.0
2001.....	1.4	*	*	*	*	*	0.2	0.8	1.7	3.4	7.0	22.0
2000.....	1.4	*	*	*	*	*	0.2	0.7	1.9	3.4	7.3	22.5
1999.....	1.4	*	*	*	*	*	0.2	0.6	1.8	3.6	7.0	22.7

Non-Hispanic white, male

2007 5/.....	1.7	*	*	*	*	*	0.4	1.3	3.0	5.3	9.5	19.4
2006 5/.....	1.6	*	*	*	*	*	0.2	1.2	2.6	4.6	9.6	22.7
2005 5/.....	1.5	*	*	*	*	*	0.3	1.1	2.4	4.1	8.4	20.3
2004 5/.....	1.0	*	*	*	*	*	0.3	0.7	1.7	2.8	6.3	16.9
2003 5/.....	1.3	*	*	*	*	*	0.4	1.0	2.0	4.4	7.9	20.4
2002.....	1.5	*	*	*	*	*	0.3	1.2	2.5	4.6	9.0	22.7
2001.....	1.4	*	*	*	*	*	0.3	1.2	2.5	4.5	8.7	20.9
2000.....	1.5	*	*	*	*	*	0.3	1.2	2.8	4.6	9.0	21.3
1999.....	1.4	*	*	*	*	*	0.2	0.9	2.6	4.9	8.7	23.6

Non-Hispanic white, female

2007 5/.....	1.6	*	*	*	*	*	*	0.5	1.2	2.4	7.0	22.7
2006 5/.....	1.5	*	*	*	*	*	0.2	0.4	1.2	2.6	7.0	22.2
2005 5/.....	1.4	*	*	*	*	*	*	0.3	0.8	2.4	6.3	21.2
2004 5/.....	1.0	*	*	*	*	*	*	0.3	0.7	1.5	4.3	15.0
2003 5/.....	1.3	*	*	*	*	*	*	0.4	0.8	2.1	5.7	21.1
2002.....	1.4	*	*	*	*	*	*	0.5	1.0	2.5	6.0	21.8
2001.....	1.4	*	*	*	*	*	*	0.3	1.0	2.5	5.9	22.4
2000.....	1.3	*	*	*	*	*	*	0.3	1.0	2.4	6.1	23.0
1999.....	1.3	*	*	*	*	*	*	0.3	1.1	2.6	5.8	22.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

Non-Hispanic black, both sexes

2007 5/.....	1.4	*	*	*	*	*	0.4	1.5	3.4	6.2	11.3	22.0
2006 5/.....	1.4	*	*	*	*	*	0.5	1.7	3.5	6.1	11.3	21.0
2005 5/.....	1.5	*	*	*	*	*	*	2.0	3.9	5.7	11.8	27.9
2004 5/.....	1.1	*	*	*	*	*	*	1.4	3.0	4.2	9.1	15.5
2003 5/.....	1.2	*	*	*	*	*	0.5	1.3	3.0	5.5	10.1	20.2
2002.....	1.2	*	*	*	*	*	*	1.7	3.0	6.1	9.4	21.5
2001.....	1.3	*	*	*	*	*	0.5	1.6	4.0	5.8	8.5	24.3
2000.....	1.3	*	*	*	*	*	0.4	1.5	4.3	5.6	10.8	21.9
1999.....	1.2	*	*	*	*	*	0.5	1.9	3.6	5.8	10.9	17.5

Non-Hispanic black, male

2007 5/.....	1.6	*	*	*	*	*	*	2.2	5.4	9.0	14.6	20.9
2006 5/.....	1.6	*	*	*	*	*	*	2.3	5.2	8.1	15.6	23.8
2005 5/.....	1.6	*	*	*	*	*	*	2.4	6.0	8.1	15.7	28.5
2004 5/.....	1.3	*	*	*	*	*	*	1.9	4.7	7.1	12.0	23.4
2003 5/.....	1.5	*	*	*	*	*	*	2.0	4.9	7.7	16.4	24.5
2002.....	1.4	*	*	*	*	*	*	2.8	4.4	9.4	11.7	21.9
2001.....	1.5	*	*	*	*	*	*	2.5	5.6	8.8	10.5	24.3
2000.....	1.5	*	*	*	*	*	*	2.3	6.3	8.7	14.2	*
1999.....	1.5	*	*	*	*	*	0.8	2.7	6.1	8.4	12.8	*

Non-Hispanic black, female

2007 5/.....	1.2	*	*	*	*	*	*	0.8	1.8	4.2	9.4	22.4
2006 5/.....	1.2	*	*	*	*	*	*	1.2	2.1	4.7	8.8	19.8
2005 5/.....	1.3	*	*	*	*	*	*	1.6	2.2	4.0	9.5	27.6
2004 5/.....	0.8	*	*	*	*	*	*	0.9	1.7	2.0	7.4	12.4
2003 5/.....	0.9	*	*	*	*	*	*	*	1.5	3.9	6.5	18.6
2002.....	1.1	*	*	*	*	*	*	0.8	2.0	3.7	8.1	21.3
2001.....	1.1	*	*	*	*	*	*	*	2.6	3.8	7.4	24.4
2000.....	1.1	*	*	*	*	*	*	*	2.7	3.5	8.9	22.2
1999.....	1.0	*	*	*	*	*	*	1.1	1.6	3.9	9.8	16.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes

2007.....	89.3	*	*	*	0.2	1.6	8.6	33.2	84.5	200.7	595.1	1,927.9
2006.....	93.5	*	*	*	0.2	1.5	9.0	34.4	87.8	217.9	632.8	2,057.4
2005.....	98.2	*	*	*	0.2	1.5	9.2	35.0	91.4	229.8	674.5	2,214.2
2004.....	99.5	*	*	*	0.2	1.6	9.6	35.1	92.5	240.3	697.1	2,279.7
2003.....	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002.....	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001.....	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000.....	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999.....	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0

All origins 3/, male

2007.....	95.6	*	*	*	0.3	2.4	13.4	51.5	128.3	286.6	799.9	2,214.7
2006.....	99.2	*	*	*	0.3	2.4	13.8	53.7	132.4	307.7	848.3	2,344.9
2005.....	102.9	*	*	*	0.3	2.3	14.1	54.7	136.9	321.9	897.1	2,511.6
2004.....	103.3	*	*	*	0.2	2.4	14.5	54.8	137.7	336.5	926.9	2,543.6
2003.....	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002.....	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001.....	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000.....	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999.....	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1

All origins 3/, female

2007.....	83.1	*	*	*	0.1	0.7	3.8	15.3	43.7	127.8	454.0	1,791.4
2006.....	88.0	*	*	*	*	0.6	4.2	15.7	46.3	141.9	485.5	1,922.9
2005.....	93.7	*	*	*	*	0.7	4.3	16.0	49.0	152.2	523.3	2,077.6
2004.....	95.7	*	*	*	*	0.7	4.6	16.2	50.5	159.6	542.4	2,161.0
2003.....	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002.....	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001.....	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000.....	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999.....	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

Hispanic, both sexes

2007.....	29.8	*	*	*	*	1.0	4.3	20.3	66.0	171.0	493.7	1,437.0
2006.....	31.3	*	*	*	*	0.8	4.4	22.6	68.5	184.6	536.0	1,595.2
2005.....	33.2	*	*	*	*	0.8	5.1	23.0	71.8	205.9	588.8	1,730.5
2004.....	32.2	*	*	*	*	0.8	4.8	22.4	72.9	205.5	589.6	1,795.7
2003.....	33.5	*	*	*	*	0.9	4.6	22.0	77.2	216.4	629.5	2,012.8
2002.....	34.2	*	*	*	*	0.7	4.5	23.6	80.1	234.6	644.0	2,129.7
2001.....	35.7	*	*	*	*	0.9	4.2	23.8	80.0	249.4	701.7	2,426.6
2000.....	35.6	*	*	*	*	0.6	4.6	23.9	88.0	249.7	703.2	2,508.8
1999.....	37.5	*	*	*	*	0.9	4.5	26.8	93.5	268.8	741.8	2,640.7

Hispanic, male

2007.....	31.8	*	*	*	*	1.5	6.8	31.9	101.7	243.7	614.6	1,481.9
2006.....	33.3	*	*	*	*	1.2	7.0	35.7	103.2	254.5	680.7	1,705.1
2005.....	35.2	*	*	*	*	1.3	8.2	36.0	108.3	287.2	754.3	1,811.9
2004.....	34.2	*	*	*	*	1.3	7.5	35.0	109.3	291.0	769.7	1,853.5
2003.....	34.6	*	*	*	*	1.4	6.9	35.1	115.3	306.1	804.2	1,965.7
2002.....	35.7	*	*	*	*	1.2	7.2	36.6	122.3	329.1	808.8	2,262.2
2001.....	36.8	*	*	*	*	1.4	6.5	37.5	121.4	350.2	901.5	2,479.8
2000.....	36.9	*	*	*	*	0.9	7.4	39.1	134.2	351.6	900.8	2,577.7
1999.....	38.7	*	*	*	*	1.3	7.6	44.3	138.5	381.8	932.8	2,706.7

Hispanic, female

2007.....	27.6	*	*	*	*	1.5	8.2	32.9	111.7	408.8	1,412.7	
2006.....	29.2	*	*	*	*	1.4	9.2	36.6	127.8	434.6	1,536.8	
2005.....	31.0	*	*	*	*	1.6	9.7	38.5	140.3	473.5	1,687.9	
2004.....	30.1	*	*	*	*	1.8	9.8	39.9	136.9	465.3	1,766.0	
2003.....	32.3	*	*	*	*	2.0	8.9	42.8	144.9	509.6	2,036.7	
2002.....	32.6	*	*	*	*	1.6	10.5	42.2	159.4	531.2	2,064.1	
2001.....	34.5	*	*	*	*	1.7	10.3	43.3	169.9	566.7	2,400.7	
2000.....	34.2	*	*	*	*	1.7	8.9	47.2	169.6	570.5	2,475.0	
1999.....	36.3	*	*	*	*	1.3	9.7	53.8	180.2	614.8	2,608.9	

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 4/, both sexes

2007.....	99.7	*	*	*	0.2	1.7	9.4	34.6	85.8	202.5	600.7	1,952.0
2006.....	104.1	*	*	*	0.2	1.7	9.8	35.7	89.1	219.7	637.6	2,078.0
2005.....	108.9	*	*	*	0.2	1.7	9.9	36.3	92.6	230.9	677.9	2,234.1
2004.....	110.1	*	*	*	0.2	1.8	10.3	36.3	93.6	241.8	701.2	2,297.5
2003.....	116.3	*	*	*	0.2	1.8	10.6	37.2	98.5	260.2	744.8	2,484.9
2002.....	118.9	*	*	*	0.2	1.6	10.6	37.5	102.0	270.3	771.9	2,571.2
2001.....	120.9	*	*	*	0.2	1.7	10.2	36.5	105.1	281.6	794.2	2,672.2
2000.....	123.9	*	*	*	0.2	1.5	10.1	37.3	108.8	293.5	815.0	2,824.0
1999.....	127.6	*	*	*	0.2	1.5	10.4	38.1	112.1	308.6	849.1	2,916.3

Non-Hispanic 4/, male

2007.....	107.4	*	*	*	0.4	2.6	14.6	53.8	130.2	289.0	810.7	2,255.1
2006.....	111.1	*	*	*	0.3	2.7	15.0	55.7	134.4	310.5	857.1	2,376.7
2005.....	114.6	*	*	*	0.3	2.6	15.2	56.6	138.8	323.3	903.3	2,544.8
2004.....	114.9	*	*	*	0.2	2.7	15.6	56.7	139.3	338.1	933.0	2,573.1
2003.....	120.4	*	*	*	0.3	2.8	16.2	58.3	146.7	365.6	991.9	2,750.7
2002.....	122.7	*	*	*	0.3	2.4	16.6	59.2	152.6	380.2	1,027.3	2,832.2
2001.....	123.1	*	*	*	0.3	2.6	15.7	57.3	156.4	395.1	1,053.5	2,923.7
2000.....	126.0	*	*	*	0.2	2.3	15.7	58.9	162.3	414.7	1,082.7	3,123.9
1999.....	128.9	*	*	*	0.3	2.3	16.2	59.9	165.4	435.2	1,123.5	3,200.1

Non-Hispanic 4/, female

2007.....	92.3	*	*	*	0.1	0.8	4.2	16.1	44.5	128.8	456.2	1,808.9
2006.....	97.4	*	*	*	*	0.6	4.7	16.4	47.0	142.6	487.8	1,939.2
2005.....	103.4	*	*	*	*	0.8	4.7	16.7	49.7	152.8	525.1	2,092.3
2004.....	105.6	*	*	*	*	0.8	5.0	16.7	51.1	160.7	545.5	2,174.3
2003.....	112.4	*	*	*	0.1	0.9	5.1	16.9	53.7	171.7	580.0	2,368.1
2002.....	115.4	*	*	*	*	0.8	4.8	16.7	55.0	178.4	602.6	2,458.6
2001.....	118.8	*	*	*	0.1	0.8	4.8	16.5	57.5	187.3	624.1	2,567.6
2000.....	122.0	*	*	*	*	0.7	4.6	16.5	59.3	193.5	640.6	2,702.7
1999.....	126.4	*	*	*	0.1	0.8	4.7	17.3	62.9	204.6	671.1	2,802.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic white, both sexes

2007 5/.....	110.1	*	*	*	0.2	1.6	9.0	32.6	80.4	194.3	602.2	2,000.6
2006 5/.....	114.9	*	*	*	0.2	1.6	9.4	33.2	83.2	210.8	636.9	2,128.7
2005 5/.....	120.2	*	*	*	0.2	1.6	9.5	33.6	86.1	222.7	677.4	2,291.1
2004 5/.....	121.4	*	*	*	0.1	1.6	10.0	33.9	87.4	233.1	700.2	2,343.9
2003 5/.....	128.0	*	*	*	0.2	1.7	10.2	34.6	91.7	250.2	743.0	2,533.4
2002.....	131.0	*	*	*	0.2	1.5	10.2	34.9	94.8	261.3	770.0	2,623.4
2001.....	132.7	*	*	*	0.2	1.6	9.6	33.8	97.4	271.4	791.2	2,724.1
2000.....	136.1	*	*	*	0.2	1.4	9.4	34.6	100.7	283.7	811.9	2,883.6
1999.....	139.9	*	*	*	0.2	1.4	9.6	35.2	103.9	297.9	847.2	2,973.2

Non-Hispanic white, male

2007 5/.....	118.2	*	*	*	0.3	2.4	14.1	51.4	123.1	280.2	817.7	2,333.5
2006 5/.....	121.9	*	*	*	0.3	2.6	14.8	52.8	126.6	299.6	860.6	2,453.9
2005 5/.....	125.9	*	*	*	0.3	2.4	14.8	53.1	130.7	314.2	908.5	2,634.7
2004 5/.....	126.1	*	*	*	0.2	2.4	15.4	53.9	131.5	329.5	937.9	2,640.6
2003 5/.....	131.9	*	*	*	0.2	2.6	15.7	55.1	138.9	354.5	995.5	2,828.2
2002.....	134.5	*	*	*	0.2	2.3	16.2	56.4	143.5	371.2	1,033.6	2,908.8
2001.....	134.4	*	*	*	0.3	2.5	15.0	54.1	147.0	384.3	1,057.5	3,002.8
2000.....	137.7	*	*	*	0.2	2.2	15.1	55.7	152.7	406.1	1,086.5	3,211.3
1999.....	140.4	*	*	*	0.2	2.1	15.2	56.1	155.1	425.3	1,130.5	3,276.7

Non-Hispanic white, female

2007 5/.....	102.4	*	*	*	*	0.7	3.9	14.1	39.8	119.5	451.2	1,843.0
2006 5/.....	108.1	*	*	*	*	0.5	4.1	14.0	41.9	133.8	481.7	1,977.0
2005 5/.....	114.7	*	*	*	*	0.8	4.2	14.4	43.7	143.6	518.4	2,133.9
2004 5/.....	116.9	*	*	*	*	0.7	4.5	14.2	45.6	150.1	538.3	2,211.1
2003 5/.....	124.3	*	*	*	*	0.8	4.6	14.4	47.0	160.9	572.4	2,403.6
2002.....	127.7	*	*	*	*	0.7	4.2	13.8	48.7	167.6	593.4	2,500.1
2001.....	131.1	*	*	*	*	0.7	4.1	13.9	50.5	175.6	614.6	2,608.1
2000.....	134.6	*	*	*	*	0.5	3.8	13.8	51.8	180.6	631.3	2,751.4
1999.....	139.3	*	*	*	*	0.7	4.0	14.6	55.7	191.3	661.7	2,852.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic black, both sexes

2007 5/.....	72.5	*	*	*	0.5	2.7	12.9	55.5	146.1	309.6	692.8	1,709.6
2006 5/.....	76.4	*	*	*	*	2.5	14.1	58.7	152.7	334.8	757.9	1,808.1
2005 5/.....	79.1	*	*	*	*	2.3	14.5	60.7	164.4	341.6	801.4	1,901.2
2004 5/.....	80.3	*	*	*	0.4	3.0	14.6	60.3	163.1	363.9	822.2	2,031.7
2003 5/.....	84.7	*	*	*	0.4	2.7	14.6	61.9	173.6	393.8	878.9	2,204.8
2002.....	85.5	*	*	*	*	2.2	15.5	62.9	180.3	395.2	901.5	2,246.7
2001.....	88.2	*	*	*	0.5	2.9	15.6	62.9	186.6	419.6	934.8	2,357.3
2000.....	89.1	*	*	*	*	2.5	15.6	64.3	192.3	431.5	947.1	2,398.0
1999.....	91.6	*	*	*	0.4	2.4	17.2	66.9	197.9	453.5	958.2	2,471.9

Non-Hispanic black, male

2007 5/.....	78.4	*	*	*	0.7	4.4	19.5	81.9	218.4	432.5	906.8	1,739.9
2006 5/.....	82.3	*	*	*	*	3.7	19.7	86.4	224.7	476.1	997.7	1,837.1
2005 5/.....	83.6	*	*	*	*	3.5	20.9	91.2	238.4	471.3	1,027.6	1,882.1
2004 5/.....	83.9	*	*	*	*	4.5	20.4	87.7	237.3	493.9	1,048.2	2,090.1
2003 5/.....	87.9	*	*	*	*	4.0	21.3	91.6	246.2	541.3	1,135.2	2,162.8
2002.....	88.0	*	*	*	*	3.0	22.2	90.8	264.4	536.0	1,119.3	2,254.6
2001.....	90.8	*	*	*	*	4.3	22.9	93.1	266.3	573.5	1,177.8	2,378.1
2000.....	90.4	*	*	*	*	3.7	22.1	93.6	274.7	566.6	1,180.1	2,463.5
1999.....	93.3	*	*	*	*	3.5	25.1	99.2	283.6	602.1	1,169.2	2,552.9

Non-Hispanic black, female

2007 5/.....	67.2	*	*	*	*	1.2	7.2	32.8	88.0	221.9	571.1	1,697.0
2006 5/.....	71.0	*	*	*	*	1.4	9.2	34.9	94.8	233.9	621.8	1,796.3
2005 5/.....	75.0	*	*	*	*	1.1	8.9	34.6	104.8	248.9	673.2	1,908.8
2004 5/.....	76.9	*	*	*	*	1.6	9.4	36.8	103.6	271.3	694.6	2,008.6
2003 5/.....	81.7	*	*	*	*	1.5	8.7	36.5	115.4	288.8	734.1	2,221.1
2002.....	83.3	*	*	*	*	1.6	9.6	39.0	112.8	295.4	778.3	2,243.7
2001.....	86.0	*	*	*	*	1.6	9.2	36.9	122.4	310.9	798.4	2,349.6
2000.....	88.0	*	*	*	*	1.5	10.0	39.1	125.9	336.5	815.9	2,373.7
1999.....	90.1	*	*	*	*	1.3	10.2	39.4	129.2	348.6	840.0	2,441.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins 3/, both sexes

2007.....	19.6	*	*	*	*	0.5	3.2	13.5	30.7	51.1	109.2	314.7
2006.....	20.4	*	*	*	0.0	0.5	3.3	13.4	31.3	55.1	114.2	345.5
2005.....	21.2	*	*	*	*	0.5	3.2	13.9	31.5	55.0	124.8	376.4
2004.....	21.7	*	*	*	*	0.5	3.3	13.6	32.0	58.7	129.0	400.4
2003.....	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002.....	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001.....	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000.....	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999.....	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4

All origins 3/, male

2007.....	22.5	*	*	*	*	0.9	5.1	21.4	48.1	73.7	142.2	326.1
2006.....	23.2	*	*	*	*	0.8	5.1	21.7	48.8	79.1	146.8	365.3
2005.....	23.6	*	*	*	*	0.7	5.1	22.2	49.0	78.7	158.8	390.7
2004.....	23.9	*	*	*	*	0.8	5.2	22.0	49.0	83.6	164.9	414.0
2003.....	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002.....	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001.....	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000.....	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999.....	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6

All origins 3/, female

2007.....	16.8	*	*	*	*	0.2	1.3	5.8	14.6	31.9	86.5	309.3
2006.....	17.7	*	*	*	*	0.2	1.5	5.5	15.1	34.8	92.0	336.1
2005.....	18.8	*	*	*	*	0.2	1.4	5.7	15.3	35.0	101.6	369.9
2004.....	19.5	*	*	*	*	0.2	1.4	5.5	16.1	37.7	104.8	394.3
2003.....	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002.....	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001.....	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000.....	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999.....	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hispanic, both sexes												
2007.....	7.1	*	*	*	*	0.3	1.7	8.5	26.7	46.8	88.4	210.6
2006.....	7.7	*	*	*	*	*	1.7	9.1	28.2	53.9	98.8	260.2
2005.....	8.2	*	*	*	*	*	1.7	10.9	29.1	55.2	113.8	274.7
2004.....	7.9	*	*	*	*	*	1.7	9.7	29.3	55.8	118.9	279.7
2003.....	8.3	*	*	*	*	0.3	1.6	9.8	29.4	57.3	124.6	357.1
2002.....	8.2	*	*	*	*	0.4	1.9	8.9	29.7	61.1	124.1	358.6
2001.....	8.5	*	*	*	*	0.4	1.5	10.1	29.7	65.9	132.5	417.0
2000.....	8.4	*	*	*	*	*	1.6	9.5	30.5	65.7	128.2	452.5
1999.....	9.0	*	*	*	*	*	1.5	10.4	33.0	72.1	147.5	469.4
Hispanic, male												
2007.....	8.8	*	*	*	*	0.5	2.7	13.2	42.8	74.5	109.4	226.7
2006.....	9.5	*	*	*	*	*	2.7	15.3	46.4	81.1	126.8	277.4
2005.....	10.2	*	*	*	*	*	2.8	17.9	47.8	82.8	156.0	308.8
2004.....	9.8	*	*	*	*	*	2.7	16.6	47.1	83.3	169.5	297.0
2003.....	10.2	*	*	*	*	0.5	2.6	17.1	48.7	88.6	169.5	397.1
2002.....	9.7	*	*	*	*	0.6	2.9	14.8	49.3	90.8	153.9	415.0
2001.....	10.1	*	*	*	*	0.6	2.2	17.1	47.9	100.2	181.1	440.0
2000.....	9.9	*	*	*	*	*	2.7	16.6	50.5	92.4	180.9	493.8
1999.....	10.6	*	*	*	*	*	2.5	17.8	50.0	115.8	194.0	485.3
Hispanic, female												
2007.....	5.3	*	*	*	*	*	*	3.6	11.7	24.3	73.7	201.9
2006.....	5.8	*	*	*	*	*	*	2.7	11.5	31.9	79.2	251.0
2005.....	6.0	*	*	*	*	*	*	3.7	11.9	32.8	84.4	256.8
2004.....	5.9	*	*	*	*	*	*	2.8	13.2	33.8	84.0	270.8
2003.....	6.2	*	*	*	*	*	*	2.5	12.0	32.4	93.7	336.7
2002.....	6.5	*	*	*	*	*	*	3.1	12.0	37.6	103.7	330.7
2001.....	6.8	*	*	*	*	*	*	3.2	13.5	38.7	99.7	405.8
2000.....	6.8	*	*	*	*	*	*	2.6	12.8	44.6	92.8	432.3
1999.....	7.4	*	*	*	*	*	*	3.2	18.0	37.9	116.6	461.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 4/, both sexes

2007.....	21.7	*	*	*	*	0.6	3.5	14.0	30.9	51.0	110.1	319.6
2006.....	22.5	*	*	*	*	0.6	3.6	13.9	31.4	54.8	114.7	349.3
2005.....	23.3	*	*	*	*	0.5	3.5	14.1	31.5	54.6	124.8	380.6
2004.....	23.8	*	*	*	*	0.5	3.6	14.0	32.0	58.3	128.9	404.7
2003.....	25.3	*	*	*	*	0.6	3.6	14.2	32.6	62.6	142.6	445.3
2002.....	25.9	*	*	*	*	0.5	3.6	13.7	32.9	64.3	146.4	477.8
2001.....	26.0	*	*	*	*	0.5	3.5	13.0	33.3	65.0	152.6	494.8
2000.....	26.9	*	*	*	*	0.4	3.2	13.4	33.8	67.9	158.3	535.9
1999.....	27.9	*	*	*	*	0.4	3.5	13.1	35.0	72.0	165.7	564.9

Non-Hispanic 4/, male

2007.....	24.9	*	*	*	*	0.9	5.6	22.3	48.2	73.0	143.5	330.8
2006.....	25.5	*	*	*	*	1.0	5.6	22.3	48.6	78.2	147.2	369.6
2005.....	25.8	*	*	*	*	0.8	5.5	22.6	48.7	77.7	158.1	394.0
2004.....	26.2	*	*	*	*	0.9	5.6	22.5	48.9	82.8	163.7	418.3
2003.....	27.5	*	*	*	*	1.0	5.6	23.1	49.3	88.2	180.4	463.3
2002.....	27.8	*	*	*	*	0.7	5.8	22.0	50.7	91.0	185.6	493.3
2001.....	27.5	*	*	*	*	0.7	5.4	21.2	51.1	91.0	193.2	503.0
2000.....	28.2	*	*	*	*	0.7	5.1	21.7	52.1	96.4	198.5	551.0
1999.....	28.9	*	*	*	*	0.6	5.6	21.0	53.3	101.1	205.6	580.2

Non-Hispanic 4/, female

2007.....	18.6	*	*	*	*	0.2	1.5	6.0	14.8	32.3	87.0	314.2
2006.....	19.6	*	*	*	*	0.2	1.6	5.8	15.4	34.8	92.5	339.9
2005.....	20.8	*	*	*	*	0.2	1.5	6.0	15.5	35.0	102.2	374.5
2004.....	21.5	*	*	*	*	0.2	1.6	5.8	16.3	37.7	105.5	398.6
2003.....	23.3	*	*	*	*	0.2	1.6	5.7	16.9	41.2	117.4	437.4
2002.....	24.0	*	*	*	*	0.2	1.4	5.7	16.5	42.0	120.5	471.1
2001.....	24.6	*	*	*	*	0.3	1.5	5.2	16.8	43.3	126.0	491.4
2000.....	25.6	*	*	*	*	0.2	1.3	5.4	16.9	44.4	132.2	529.8
1999.....	27.0	*	*	*	*	0.2	1.4	5.4	18.2	48.2	139.8	558.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic white, both sexes												
2007 5/.....	22.5	*	*	*	*	0.5	3.3	12.9	27.9	46.1	104.7	317.2
2006 5/.....	23.2	*	*	*	*	0.5	3.3	12.5	28.0	49.1	108.5	346.8
2005 5/.....	24.0	*	*	*	*	0.4	3.3	12.7	27.5	49.1	117.9	377.8
2004 5/.....	24.6	*	*	*	*	0.5	3.5	12.6	28.4	52.4	121.7	399.9
2003 5/.....	26.2	*	*	*	*	0.6	3.4	12.8	28.6	55.9	135.1	441.5
2002.....	26.8	*	*	*	*	0.4	3.3	12.4	28.9	57.2	138.2	473.5
2001.....	26.8	*	*	*	*	0.4	3.2	11.6	28.4	57.1	144.2	490.7
2000.....	27.6	*	*	*	*	0.4	2.9	11.9	28.9	60.2	149.0	530.2
1999.....	28.6	*	*	*	*	0.4	3.1	11.3	29.8	63.6	156.2	558.8
Non-Hispanic white, male												
2007 5/.....	25.5	*	*	*	*	0.8	5.4	20.8	43.8	66.3	136.4	330.0
2006 5/.....	25.9	*	*	*	*	0.9	5.3	20.4	43.5	69.8	139.0	366.7
2005 5/.....	26.3	*	*	*	*	0.7	5.3	20.7	43.0	69.8	148.8	396.1
2004 5/.....	26.7	*	*	*	*	0.7	5.5	20.8	43.4	75.2	154.3	412.6
2003 5/.....	28.0	*	*	*	*	0.9	5.3	21.0	43.9	78.5	170.6	459.5
2002.....	28.2	*	*	*	*	0.7	5.4	20.3	44.8	81.3	174.4	489.2
2001.....	27.8	*	*	*	*	0.6	5.0	19.1	44.3	80.4	182.9	499.4
2000.....	28.4	*	*	*	*	0.6	4.8	19.6	45.2	86.3	186.1	544.7
1999.....	29.0	*	*	*	*	0.6	5.2	18.4	46.0	90.0	193.5	572.8
Non-Hispanic white, female												
2007 5/.....	19.6	*	*	*	*	0.2	1.3	5.1	12.8	28.5	82.6	311.1
2006 5/.....	20.6	*	*	*	*	*	1.4	4.7	13.3	31.1	87.3	337.5
2005 5/.....	21.9	*	*	*	*	0.2	1.2	4.8	12.8	31.2	96.6	369.4
2004 5/.....	22.6	*	*	*	*	0.2	1.4	4.5	14.1	32.9	99.4	394.2
2003 5/.....	24.6	*	*	*	*	0.2	1.5	4.6	14.2	36.5	111.0	433.5
2002.....	25.3	*	*	*	*	0.2	1.1	4.6	13.8	36.6	113.9	466.8
2001.....	25.8	*	*	*	*	0.2	1.3	4.1	13.3	37.3	118.6	487.0
2000.....	26.9	*	*	*	*	*	1.1	4.3	13.4	38.2	124.6	524.4
1999.....	28.2	*	*	*	*	0.2	1.1	4.2	14.6	41.6	131.8	553.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic black, both sexes

2007 5/.....	23.1	*	*	*	*	1.0	5.0	24.3	60.5	103.4	182.2	402.7
2006 5/.....	24.4	*	*	*	*	1.1	5.8	25.5	63.7	111.6	201.2	433.7
2005 5/.....	25.3	*	*	*	*	0.8	5.4	26.1	68.6	110.2	218.2	475.2
2004 5/.....	25.7	*	*	*	*	0.9	4.9	26.1	67.3	120.8	229.1	509.2
2003 5/.....	26.7	*	*	*	*	0.9	5.1	26.1	70.1	129.6	244.1	546.7
2002.....	27.7	*	*	*	*	0.8	6.0	25.7	71.9	137.2	261.1	580.4
2001.....	28.5	*	*	*	*	0.7	5.8	25.7	78.1	144.8	266.7	595.9
2000.....	29.3	*	*	*	*	0.7	5.3	27.3	78.9	147.2	280.2	652.2
1999.....	30.7	*	*	*	*	0.8	6.5	28.1	82.0	158.7	291.0	670.3

Non-Hispanic black, male

2007 5/.....	27.9	*	*	*	*	1.8	7.8	37.1	95.3	154.4	257.2	415.3
2006 5/.....	29.5	*	*	*	*	1.5	8.4	38.9	100.8	170.9	275.8	471.0
2005 5/.....	29.7	*	*	*	*	1.3	7.9	39.8	106.1	168.1	296.8	452.0
2004 5/.....	29.7	*	*	*	*	1.6	7.1	38.5	105.6	173.1	308.9	540.6
2003 5/.....	31.2	*	*	*	*	1.6	7.8	41.1	106.0	191.8	334.2	588.1
2002.....	31.9	*	*	*	*	1.1	8.9	38.8	111.8	198.7	358.4	603.9
2001.....	32.3	*	*	*	*	1.0	8.8	39.9	118.1	209.4	352.3	609.8
2000.....	33.2	*	*	*	*	1.1	7.9	42.3	120.8	209.3	371.5	711.6
1999.....	34.5	*	*	*	*	1.1	9.8	43.6	123.7	228.2	378.4	714.3

Non-Hispanic black, female

2007 5/.....	18.7	*	*	*	*	2.6	13.3	32.5	67.0	139.5	397.5	
2006 5/.....	19.8	*	*	*	*	3.5	14.0	33.8	69.3	158.9	418.5	
2005 5/.....	21.3	*	*	*	*	3.2	14.3	38.4	68.9	173.7	484.5	
2004 5/.....	22.1	*	*	*	*	3.1	15.4	36.5	83.4	184.0	496.7	
2003 5/.....	22.7	*	*	*	*	2.7	13.3	41.2	85.3	193.3	530.7	
2002.....	24.0	*	*	*	*	3.4	14.4	39.8	93.6	206.1	571.5	
2001.....	25.0	*	*	*	*	3.2	13.6	45.9	99.3	218.7	590.7	
2000.....	25.8	*	*	*	*	2.9	14.4	45.1	103.4	228.9	630.2	
1999.....	27.1	*	*	*	*	3.6	14.7	48.5	109.7	242.1	653.8	

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes

2007.....	69.7	*	*	*	0.2	1.0	5.4	19.7	53.8	149.7	485.9	1,613.2
2006.....	73.1	*	*	*	0.1	1.0	5.7	21.0	56.5	162.8	518.6	1,712.0
2005.....	77.0	*	*	*	0.2	1.1	6.0	21.2	59.8	174.9	549.7	1,837.8
2004.....	77.8	*	*	*	0.1	1.1	6.2	21.5	60.5	181.7	568.1	1,879.3
2003.....	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002.....	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001.....	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000.....	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999.....	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6

All origins 3/, male

2007.....	73.2	*	*	*	0.3	1.6	8.2	30.2	80.3	213.0	657.7	1,888.6
2006.....	76.1	*	*	*	0.2	1.6	8.7	32.1	83.7	228.7	701.5	1,979.5
2005.....	79.3	*	*	*	0.2	1.6	9.0	32.5	88.0	243.3	738.3	2,120.9
2004.....	79.4	*	*	*	0.2	1.7	9.3	32.8	88.7	252.9	762.0	2,129.6
2003.....	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002.....	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001.....	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000.....	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999.....	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4

All origins 3/, female

2007.....	66.3	*	*	*	*	0.5	2.5	9.5	29.1	95.9	367.5	1,482.2
2006.....	70.3	*	*	*	*	0.4	2.7	10.2	31.2	107.1	393.5	1,586.8
2005.....	74.8	*	*	*	*	0.5	2.9	10.3	33.7	117.2	421.7	1,707.7
2004.....	76.2	*	*	*	*	0.5	3.2	10.6	34.3	121.9	437.6	1,766.7
2003.....	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002.....	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001.....	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000.....	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999.....	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Hispanic, both sexes

2007.....	22.7	*	*	*	*	0.6	2.6	11.7	39.3	124.2	405.3	1,226.4
2006.....	23.6	*	*	*	*	0.6	2.7	13.5	40.3	130.6	437.2	1,335.0
2005.....	25.0	*	*	*	*	0.6	3.4	12.1	42.8	150.8	475.0	1,455.8
2004.....	24.3	*	*	*	*	0.6	3.2	12.7	43.6	149.7	470.7	1,516.0
2003.....	25.2	*	*	*	*	0.6	3.0	12.2	47.8	159.1	504.9	1,655.7
2002.....	26.0	*	*	*	*	0.4	2.7	14.6	50.4	173.4	519.9	1,771.1
2001.....	27.2	*	*	*	*	0.6	2.7	13.7	50.4	183.6	569.1	2,009.6
2000.....	27.2	*	*	*	*	0.5	3.0	14.3	57.5	184.0	575.0	2,056.3
1999.....	28.5	*	*	*	*	0.6	3.0	16.4	60.5	196.7	594.3	2,171.4

Hispanic, male

2007.....	23.0	*	*	*	*	1.0	4.1	18.7	58.9	169.2	505.2	1,255.3
2006.....	23.8	*	*	*	*	0.9	4.3	20.4	56.8	173.4	553.9	1,427.7
2005.....	25.0	*	*	*	*	0.9	5.5	18.1	60.5	204.4	598.3	1,503.1
2004.....	24.3	*	*	*	*	0.9	4.8	18.3	62.2	207.8	600.2	1,556.5
2003.....	24.4	*	*	*	*	0.9	4.3	18.0	66.6	217.6	634.7	1,568.6
2002.....	25.9	*	*	*	*	0.6	4.3	21.8	73.0	238.3	654.9	1,847.2
2001.....	26.7	*	*	*	*	0.9	4.3	20.4	73.6	250.0	720.4	2,039.8
2000.....	27.0	*	*	*	*	0.7	4.7	22.6	83.6	259.1	719.8	2,084.0
1999.....	28.1	*	*	*	*	1.0	5.1	26.5	88.5	266.0	738.8	2,221.4

Hispanic, female

2007.....	22.3	*	*	*	*	0.9	4.6	21.1	87.4	335.1	1,210.8
2006.....	23.3	*	*	*	*	1.0	6.5	25.1	95.9	355.4	1,285.7
2005.....	25.0	*	*	*	*	1.0	6.0	26.6	107.4	389.1	1,431.1
2004.....	24.2	*	*	*	*	1.3	7.0	26.7	103.2	381.3	1,495.2
2003.....	26.1	*	*	*	*	1.6	6.4	30.7	112.5	415.8	1,700.0
2002.....	26.1	*	*	*	*	0.9	7.4	30.2	121.8	427.5	1,733.3
2001.....	27.7	*	*	*	*	1.0	7.1	29.8	131.2	466.9	1,994.8
2000.....	27.5	*	*	*	*	1.2	6.2	34.4	124.9	477.7	2,042.7
1999.....	28.9	*	*	*	*	0.8	6.5	35.8	142.3	498.2	2,147.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 4/, both sexes

2007.....	78.0	*	*	*	0.2	1.1	5.8	20.6	54.9	151.5	490.6	1,632.5
2006.....	81.6	*	*	*	0.1	1.1	6.2	21.8	57.7	164.9	522.9	1,728.7
2005.....	85.6	*	*	*	0.2	1.2	6.4	22.2	61.1	176.4	553.1	1,853.4
2004.....	86.3	*	*	*	0.2	1.2	6.7	22.4	61.6	183.5	572.4	1,892.8
2003.....	91.0	*	*	*	0.2	1.2	7.0	22.9	65.9	197.6	602.3	2,039.6
2002.....	93.1	*	*	*	0.1	1.1	7.1	23.8	69.0	206.0	625.4	2,093.4
2001.....	94.9	*	*	*	0.2	1.2	6.7	23.5	71.8	216.7	641.6	2,177.4
2000.....	97.0	*	*	*	0.1	1.1	6.9	23.9	75.0	225.6	656.7	2,288.1
1999.....	99.7	*	*	*	0.2	1.1	6.9	25.1	77.1	236.5	683.4	2,351.4

Non-Hispanic 4/, male

2007.....	82.5	*	*	*	0.3	1.7	9.0	31.5	82.0	216.0	667.1	1,924.3
2006.....	85.6	*	*	*	0.2	1.8	9.5	33.4	85.8	232.3	709.8	2,007.1
2005.....	88.8	*	*	*	0.3	1.8	9.7	34.0	90.1	245.6	745.1	2,150.8
2004.....	88.7	*	*	*	0.2	1.8	10.0	34.2	90.4	255.3	769.4	2,154.8
2003.....	92.9	*	*	*	0.2	1.8	10.6	35.2	97.4	277.5	811.5	2,287.4
2002.....	94.9	*	*	*	0.2	1.6	10.8	37.2	102.0	289.3	841.7	2,338.9
2001.....	95.6	*	*	*	0.2	2.0	10.3	36.1	105.3	304.1	860.3	2,420.7
2000.....	97.8	*	*	*	0.2	1.7	10.5	37.2	110.2	318.2	884.2	2,572.9
1999.....	100.0	*	*	*	0.2	1.6	10.6	38.9	112.1	334.1	917.9	2,619.9

Non-Hispanic 4/, female

2007.....	73.6	*	*	*	*	0.6	2.7	10.1	29.7	96.5	369.2	1,494.7
2006.....	77.9	*	*	*	*	0.4	3.0	10.6	31.6	107.7	395.3	1,599.3
2005.....	82.6	*	*	*	*	0.6	3.2	10.7	34.2	117.8	422.8	1,717.8
2004.....	84.1	*	*	*	*	0.6	3.4	11.0	34.9	123.0	439.9	1,775.7
2003.....	89.2	*	*	*	0.1	0.7	3.4	11.2	36.7	130.6	462.6	1,930.7
2002.....	91.4	*	*	*	*	0.6	3.4	11.0	38.5	136.4	482.1	1,987.5
2001.....	94.2	*	*	*	*	0.5	3.2	11.3	40.7	143.9	498.0	2,076.2
2000.....	96.3	*	*	*	*	0.5	3.3	11.1	42.4	149.1	508.5	2,172.9
1999.....	99.4	*	*	*	*	0.6	3.3	11.8	44.8	156.4	531.2	2,243.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic white, both sexes

2007 5/.....	87.6	*	*	*	0.2	1.1	5.7	19.7	52.5	148.2	497.4	1,683.4
2006 5/.....	91.6	*	*	*	0.1	1.1	6.1	20.7	55.2	161.7	528.4	1,781.9
2005 5/.....	96.2	*	*	*	0.1	1.2	6.2	20.9	58.6	173.6	559.5	1,913.3
2004 5/.....	96.8	*	*	*	0.1	1.1	6.5	21.3	59.1	180.6	578.5	1,944.1
2003 5/.....	101.8	*	*	*	0.1	1.1	6.7	21.8	63.1	194.3	607.9	2,091.9
2002.....	104.3	*	*	*	0.1	1.1	6.9	22.5	65.9	204.1	631.9	2,149.8
2001.....	106.0	*	*	*	0.1	1.1	6.4	22.2	69.0	214.3	647.0	2,233.4
2000.....	108.4	*	*	*	0.1	1.0	6.5	22.7	71.9	223.5	662.9	2,353.3
1999.....	111.3	*	*	*	0.1	1.0	6.5	23.9	74.1	234.2	690.9	2,414.4

Non-Hispanic white, male

2007 5/.....	92.7	*	*	*	0.2	1.6	8.7	30.6	79.2	213.9	681.4	2,003.4
2006 5/.....	95.9	*	*	*	0.2	1.7	9.5	32.3	83.2	229.7	721.6	2,087.2
2005 5/.....	99.6	*	*	*	0.2	1.7	9.4	32.5	87.7	244.4	759.7	2,238.6
2004 5/.....	99.3	*	*	*	0.2	1.7	9.9	33.1	88.1	254.3	783.6	2,228.0
2003 5/.....	103.9	*	*	*	0.2	1.7	10.4	34.1	94.9	276.0	824.8	2,368.6
2002.....	106.2	*	*	*	0.2	1.6	10.7	36.1	98.7	289.9	859.2	2,419.7
2001.....	106.6	*	*	*	0.2	1.8	10.0	34.9	102.7	303.9	874.6	2,503.4
2000.....	109.2	*	*	*	0.2	1.6	10.3	36.1	107.5	319.8	900.4	2,666.6
1999.....	111.4	*	*	*	0.2	1.5	10.1	37.7	109.1	335.3	936.9	2,703.9

Non-Hispanic white, female

2007 5/.....	82.7	*	*	*	*	0.5	2.6	9.0	27.0	91.0	368.6	1,531.9
2006 5/.....	87.5	*	*	*	*	0.4	2.7	9.3	28.6	102.8	394.5	1,639.5
2005 5/.....	92.9	*	*	*	*	0.6	2.9	9.5	30.9	112.4	421.8	1,764.6
2004 5/.....	94.3	*	*	*	*	0.5	3.1	9.7	31.5	117.3	438.8	1,816.9
2003 5/.....	99.8	*	*	*	*	0.6	3.1	9.7	32.8	124.3	461.4	1,970.1
2002.....	102.4	*	*	*	*	0.5	3.1	9.2	34.9	131.0	479.6	2,033.3
2001.....	105.3	*	*	*	*	0.5	2.9	9.8	37.2	138.2	496.0	2,121.0
2000.....	107.7	*	*	*	*	0.4	2.7	9.5	38.3	142.4	506.8	2,227.0
1999.....	111.1	*	*	*	*	0.5	2.8	10.4	41.1	149.7	529.9	2,299.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic black, both sexes

2007 5/.....	49.4	*	*	*	0.4	1.7	7.9	31.2	85.6	206.2	510.6	1,306.9
2006 5/.....	51.9	*	*	*	*	1.4	8.4	33.2	89.0	223.2	556.6	1,374.4
2005 5/.....	53.8	*	*	*	*	1.5	9.1	34.6	95.8	231.4	583.2	1,426.0
2004 5/.....	54.6	*	*	*	0.4	2.1	9.6	34.2	95.9	243.2	593.1	1,522.6
2003 5/.....	57.9	*	*	*	*	1.8	9.5	35.8	103.6	264.2	634.8	1,658.1
2002.....	57.8	*	*	*	*	1.5	9.6	37.2	108.4	258.0	640.3	1,666.3
2001.....	59.7	*	*	*	0.4	2.1	9.8	37.1	108.4	274.7	668.1	1,761.4
2000.....	59.8	*	*	*	*	1.8	10.4	37.0	113.4	284.4	666.8	1,745.8
1999.....	61.0	*	*	*	*	1.6	10.7	38.9	115.9	294.7	667.1	1,801.6

Non-Hispanic black, male

2007 5/.....	50.5	*	*	*	*	2.6	11.7	44.9	123.1	278.1	649.7	1,324.6
2006 5/.....	52.7	*	*	*	*	2.2	11.3	47.5	123.9	305.2	721.9	1,366.1
2005 5/.....	53.9	*	*	*	*	2.2	13.0	51.4	132.3	303.3	730.8	1,430.2
2004 5/.....	54.2	*	*	*	*	3.0	13.4	49.2	131.6	320.8	739.3	1,549.6
2003 5/.....	56.7	*	*	*	*	2.4	13.4	50.5	140.2	349.5	801.1	1,574.7
2002.....	56.1	*	*	*	*	1.9	13.3	52.0	152.5	337.3	760.8	1,650.8
2001.....	58.4	*	*	*	*	3.3	14.1	53.2	148.2	364.1	825.5	1,768.3
2000.....	57.3	*	*	*	*	2.6	14.1	51.3	153.9	357.2	808.7	1,751.9
1999.....	58.8	*	*	*	*	2.4	15.3	55.5	160.0	374.0	790.8	1,838.6

Non-Hispanic black, female

2007 5/.....	48.5	*	*	*	*	1.0	4.5	19.5	55.5	154.9	431.5	1,299.5
2006 5/.....	51.2	*	*	*	*	0.7	5.7	21.0	61.0	164.6	462.9	1,377.8
2005 5/.....	53.7	*	*	*	*	0.8	5.7	20.3	66.4	180.0	499.6	1,424.3
2004 5/.....	54.9	*	*	*	*	1.2	6.3	21.3	67.1	187.8	510.6	1,511.9
2003 5/.....	59.1	*	*	*	*	1.2	6.1	23.2	74.2	203.5	540.8	1,690.4
2002.....	59.3	*	*	*	*	1.1	6.2	24.5	73.0	201.8	572.2	1,672.2
2001.....	60.9	*	*	*	*	1.0	6.0	23.3	76.5	211.7	579.7	1,758.9
2000.....	62.1	*	*	*	*	1.1	7.1	24.7	80.7	233.1	587.0	1,743.5
1999.....	63.0	*	*	*	*	0.8	6.6	24.6	80.7	238.9	597.9	1,787.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

All origins 3/, both sexes

2007.....	57.3	9.3	1.0	0.5	2.1	4.4	9.5	20.7	43.5	111.8	360.4	1,329.8
2006.....	56.9	7.7	0.9	0.6	2.1	4.7	9.5	21.2	44.4	112.3	360.7	1,342.8
2005.....	57.6	8.3	0.9	0.6	2.2	4.6	9.8	21.5	45.7	116.8	366.7	1,377.4
2004.....	56.6	10.0	1.1	0.5	2.1	4.3	9.7	21.5	45.2	118.6	368.9	1,373.9
2003.....	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002.....	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001.....	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000.....	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999.....	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9

All origins 3/, male

2007.....	51.9	10.3	0.9	0.6	2.5	5.3	11.9	26.5	55.5	136.4	413.4	1,348.9
2006.....	51.0	8.0	1.0	0.6	2.5	5.9	11.6	27.2	56.2	136.2	411.1	1,338.1
2005.....	51.4	8.9	0.9	0.6	2.8	5.8	12.3	27.6	57.5	140.5	415.8	1,377.8
2004.....	50.6	10.5	1.1	0.5	2.6	5.3	12.0	27.7	57.7	142.8	420.8	1,377.7
2003.....	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002.....	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001.....	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000.....	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999.....	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9

All origins 3/, female

2007.....	62.5	8.3	1.1	0.5	1.6	3.4	7.1	15.1	32.4	91.0	323.9	1,320.8
2006.....	62.6	7.5	0.9	0.5	1.5	3.5	7.3	15.3	33.5	92.1	326.3	1,345.0
2005.....	63.5	7.7	0.8	0.6	1.5	3.4	7.3	15.5	34.6	96.7	333.3	1,377.3
2004.....	62.4	9.4	1.1	0.6	1.5	3.3	7.4	15.5	33.6	98.4	334.0	1,372.1
2003.....	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002.....	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001.....	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000.....	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999.....	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

Hispanic, both sexes

2007.....	14.3	8.1	1.0	0.4	1.7	2.6	5.4	11.0	24.1	67.2	208.3	659.7
2006.....	13.7	6.3	0.8	0.5	1.4	2.8	4.7	11.9	23.9	67.5	197.0	663.2
2005.....	13.9	7.5	0.6	0.4	1.5	2.7	5.1	10.9	24.7	70.6	210.8	703.3
2004.....	13.2	8.5	1.0	0.4	1.5	2.3	5.3	11.4	24.7	70.3	208.5	659.3
2003.....	14.1	8.3	1.0	0.3	1.4	2.8	5.3	11.3	29.7	77.3	221.2	749.9
2002.....	13.3	9.7	0.7	0.4	1.5	2.6	5.1	11.4	25.7	72.6	219.8	740.0
2001.....	13.3	9.2	1.0	0.5	1.3	2.6	4.9	10.9	27.9	75.4	217.5	798.5
2000.....	13.6	10.2	1.0	0.5	1.7	2.8	5.5	11.6	28.8	75.9	227.0	829.4
1999.....	13.8	12.2	1.2	0.5	1.6	2.8	6.3	11.6	30.6	78.7	229.6	827.9

Hispanic, male

2007.....	14.0	8.7	*	0.5	2.3	3.3	7.2	14.7	30.4	80.4	233.7	679.0
2006.....	13.5	6.9	*	0.6	1.9	3.9	6.1	15.1	30.6	82.9	231.4	658.6
2005.....	13.7	8.0	*	*	1.9	3.7	6.9	14.9	31.8	87.0	247.5	697.4
2004.....	13.0	8.4	1.3	*	2.2	3.3	7.1	15.5	31.6	83.6	241.8	641.6
2003.....	13.6	9.1	*	*	1.8	3.4	7.0	14.7	37.0	95.0	253.5	752.7
2002.....	12.9	9.1	*	*	2.1	3.4	6.4	14.3	31.7	92.5	256.9	758.0
2001.....	12.6	9.6	*	*	1.6	3.2	6.5	15.0	34.1	91.1	248.6	775.2
2000.....	13.2	9.9	1.3	0.7	2.1	3.5	6.7	15.1	37.2	94.1	250.8	890.8
1999.....	13.7	10.3	1.4	0.6	1.8	3.8	8.2	15.4	38.8	102.4	267.5	888.6

Hispanic, female

2007.....	14.7	7.5	1.1	*	1.0	1.9	3.3	7.1	18.3	56.4	190.5	649.2
2006.....	13.9	5.7	*	*	0.8	1.5	3.2	8.5	17.8	54.9	172.8	665.6
2005.....	14.1	7.0	*	*	1.2	1.5	3.1	6.8	18.3	57.3	185.1	706.4
2004.....	13.5	8.6	*	*	0.8	1.1	3.3	7.3	18.5	59.6	185.5	668.4
2003.....	14.5	7.6	*	*	0.9	2.2	3.3	8.0	23.2	63.2	199.1	748.4
2002.....	13.7	10.3	*	*	0.8	1.6	3.7	8.5	20.3	56.7	194.5	731.1
2001.....	13.9	8.7	*	*	0.9	1.9	3.2	6.9	22.5	63.0	196.4	809.8
2000.....	14.0	10.6	*	*	1.1	1.9	4.3	8.1	21.4	61.6	211.0	799.3
1999.....	13.9	14.1	*	*	1.3	1.6	4.2	8.0	23.4	60.1	204.3	798.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Other heart diseases (I26-I51)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	64.8	9.6	1.0	0.6	2.1	4.8	10.3	21.8	45.2	115.3	370.1	1,364.4
2006.....	64.3	8.2	1.0	0.6	2.2	5.2	10.3	22.2	46.1	115.5	370.4	1,375.6
2005.....	64.8	8.6	1.0	0.6	2.3	5.0	10.6	22.6	47.3	120.1	375.4	1,408.5
2004.....	63.6	10.3	1.1	0.6	2.1	4.7	10.4	22.5	46.7	122.0	377.3	1,404.5
2003.....	65.4	10.9	1.1	0.6	2.3	4.9	11.0	22.3	49.3	130.0	390.5	1,453.7
2002.....	65.7	12.2	1.1	0.6	2.2	4.9	11.0	22.7	50.4	133.2	398.4	1,469.2
2001.....	65.2	11.6	1.5	0.6	2.1	4.7	10.6	22.6	51.1	134.7	396.5	1,497.8
2000.....	65.1	12.4	1.1	0.6	2.3	4.7	10.4	22.3	52.4	137.5	397.5	1,530.9
1999.....	65.6	13.1	1.1	0.6	2.4	4.7	10.8	22.4	54.3	142.0	404.4	1,533.9
<b>Non-Hispanic 4/, male</b>												
2007.....	59.0	10.8	0.9	0.6	2.6	5.9	12.8	27.9	57.6	140.5	424.9	1,387.0
2006.....	57.8	8.3	1.0	0.6	2.7	6.5	12.6	28.6	58.3	139.8	421.7	1,374.1
2005.....	58.0	9.2	1.0	0.6	3.0	6.3	13.2	29.0	59.5	144.2	425.3	1,412.5
2004.....	57.0	11.0	1.0	0.6	2.7	5.8	12.8	28.9	59.6	146.8	430.2	1,412.5
2003.....	58.6	12.0	1.0	0.7	2.8	5.8	13.5	28.4	62.2	158.0	448.9	1,460.5
2002.....	58.7	12.8	1.1	0.7	2.8	6.2	13.7	29.5	62.9	160.4	456.6	1,477.3
2001.....	57.8	11.0	1.6	0.6	2.6	5.6	13.2	28.7	63.6	162.5	460.6	1,499.0
2000.....	58.0	12.7	1.2	0.7	2.7	5.5	12.9	28.9	66.4	167.8	458.9	1,573.6
1999.....	58.6	13.8	1.3	0.7	2.9	5.5	13.5	28.7	69.3	176.2	466.4	1,571.2
<b>Non-Hispanic 4/, female</b>												
2007.....	70.4	8.4	1.1	0.5	1.7	3.7	7.7	16.0	33.6	93.7	332.3	1,353.8
2006.....	70.4	8.0	0.9	0.5	1.7	3.9	8.0	16.1	34.8	94.9	335.4	1,376.4
2005.....	71.2	7.9	1.0	0.6	1.6	3.8	8.0	16.4	36.0	99.6	341.5	1,406.6
2004.....	69.9	9.6	1.2	0.6	1.6	3.7	8.0	16.4	34.8	101.1	341.7	1,400.9
2003.....	71.8	9.7	1.2	0.5	1.9	4.0	8.5	16.4	37.3	106.5	351.5	1,450.7
2002.....	72.3	11.6	1.0	0.5	1.5	3.7	8.4	16.1	38.7	110.4	359.7	1,465.7
2001.....	72.3	12.2	1.3	0.6	1.6	3.9	8.0	16.7	39.4	111.6	354.4	1,497.3
2000.....	71.8	12.0	1.0	0.5	1.9	3.8	7.9	15.9	39.4	112.5	357.5	1,513.7
1999.....	72.2	12.3	1.0	0.6	1.9	3.9	8.1	16.2	40.5	113.9	364.2	1,519.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Other heart diseases (I26-I51)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	71.0	6.6	0.8	0.5	1.8	3.8	8.5	18.6	40.4	108.9	372.0	1,410.6
2006 5/.....	70.4	5.8	0.8	0.5	1.9	4.1	8.6	18.9	41.4	109.8	372.0	1,422.6
2005 5/.....	70.6	6.7	0.7	0.5	1.9	3.9	8.8	19.1	42.0	114.0	376.1	1,453.0
2004 5/.....	69.3	8.7	0.8	0.5	1.8	3.8	8.6	19.0	41.9	116.5	377.3	1,442.1
2003 5/.....	71.0	8.1	0.8	0.5	1.9	3.8	9.1	18.8	44.1	123.1	391.4	1,490.1
2002.....	71.3	10.0	0.9	0.5	1.8	3.8	9.3	19.0	45.3	127.6	398.1	1,503.1
2001.....	70.6	9.1	1.1	0.5	1.7	3.7	8.5	19.0	45.8	128.7	394.8	1,533.2
2000.....	70.3	10.5	0.9	0.5	1.7	3.7	8.4	18.7	47.4	131.2	395.8	1,564.2
1999.....	70.4	10.5	0.9	0.5	2.0	3.7	8.7	18.7	48.3	135.2	401.1	1,565.9
<b>Non-Hispanic white, male</b>												
2007 5/.....	63.4	7.4	0.6	0.5	2.0	4.8	10.5	24.1	51.5	132.9	426.4	1,440.1
2006 5/.....	62.1	5.7	0.8	0.6	2.3	5.1	10.5	24.4	52.5	132.9	424.0	1,420.8
2005 5/.....	62.2	6.9	0.8	0.5	2.5	4.9	11.2	24.9	53.1	136.5	427.1	1,462.5
2004 5/.....	60.9	9.3	0.7	0.5	2.3	4.6	10.8	24.7	53.6	140.9	428.9	1,445.4
2003 5/.....	62.6	8.7	0.7	0.6	2.3	4.6	11.4	24.3	56.0	150.6	450.4	1,499.3
2002.....	62.7	10.7	1.1	0.6	2.3	4.7	11.8	25.2	57.1	154.1	457.1	1,512.9
2001.....	61.4	9.4	1.2	0.6	2.0	4.4	10.9	24.8	57.5	155.8	457.4	1,530.7
2000.....	61.6	10.9	1.1	0.6	2.0	4.4	10.7	24.6	60.7	161.0	457.6	1,606.4
1999.....	61.7	11.6	1.1	0.5	2.3	4.4	11.2	24.5	62.1	168.5	462.9	1,595.5
<b>Non-Hispanic white, female</b>												
2007 5/.....	78.3	5.8	1.0	0.5	1.5	2.9	6.6	13.3	29.7	88.0	333.9	1,396.7
2006 5/.....	78.4	5.8	0.8	0.5	1.4	3.0	6.7	13.5	30.9	89.9	335.9	1,423.5
2005 5/.....	78.8	6.4	0.6	0.4	1.3	2.8	6.4	13.3	31.4	94.6	340.9	1,448.6
2004 5/.....	77.3	8.2	0.9	0.4	1.3	2.9	6.4	13.3	30.7	95.6	342.1	1,440.6
2003 5/.....	79.1	7.5	1.0	0.4	1.6	3.0	6.8	13.5	32.7	99.6	351.5	1,486.1
2002.....	79.6	9.3	0.8	0.5	1.2	2.8	6.8	13.0	34.1	105.0	358.5	1,498.8
2001.....	79.4	8.8	1.0	0.5	1.3	3.1	6.1	13.4	34.8	105.7	353.3	1,534.2
2000.....	78.7	10.1	0.8	0.4	1.4	2.9	6.2	12.9	34.8	106.0	355.3	1,547.2
1999.....	78.8	9.4	0.8	0.5	1.7	3.0	6.2	13.1	35.3	107.3	360.6	1,554.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Other heart diseases (I26-I51)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	53.0	20.0	1.6	1.0	4.0	10.2	22.2	48.0	94.7	200.3	440.8	1,130.2
2006 5/.....	52.0	16.8	1.7	0.9	3.9	11.8	22.2	48.4	95.0	191.6	443.4	1,111.9
2005 5/.....	53.4	15.6	1.8	1.2	4.3	11.7	22.9	50.4	100.2	199.7	455.4	1,152.4
2004 5/.....	52.7	15.3	2.4	1.1	3.8	10.4	23.0	51.4	98.5	199.4	458.1	1,197.9
2003 5/.....	54.4	19.8	2.0	1.0	4.5	11.3	24.4	51.1	104.2	219.8	455.6	1,253.2
2002.....	54.3	20.9	1.6	0.9	4.2	11.8	23.4	52.6	104.1	211.2	474.7	1,285.5
2001.....	54.7	21.3	3.0	1.0	4.4	11.0	24.5	52.2	105.7	216.1	486.4	1,292.9
2000.....	54.6	19.6	2.0	1.1	4.9	10.8	23.7	52.8	104.5	223.6	484.3	1,319.4
1999.....	56.2	23.2	2.0	1.0	4.7	11.0	25.3	53.5	115.3	232.9	499.9	1,314.0
<b>Non-Hispanic black, male</b>												
2007 5/.....	53.1	23.2	2.1	1.1	5.4	12.1	29.2	60.6	125.2	254.2	523.1	1,085.5
2006 5/.....	52.1	17.9	2.0	1.0	4.5	14.8	28.0	63.5	123.2	242.0	513.3	1,122.4
2005 5/.....	52.6	18.4	*	1.1	5.8	14.6	28.4	63.9	128.3	251.6	520.0	1,107.6
2004 5/.....	52.3	16.4	2.5	1.0	4.8	12.6	27.7	65.3	128.9	242.3	550.5	1,244.7
2003 5/.....	53.4	23.6	2.2	1.2	5.8	13.0	29.6	65.1	134.7	269.3	530.4	1,239.1
2002.....	52.6	21.8	*	0.9	5.1	14.8	28.2	67.5	130.4	259.4	543.4	1,258.0
2001.....	53.5	18.8	3.3	0.9	5.5	13.4	29.3	63.9	134.9	261.9	593.5	1,362.9
2000.....	53.3	19.1	2.0	1.2	5.7	12.7	28.7	68.3	132.9	269.8	565.5	1,416.1
1999.....	56.2	22.9	2.3	1.2	6.0	13.4	30.9	68.7	149.5	291.0	584.7	1,466.6
<b>Non-Hispanic black, female</b>												
2007 5/.....	52.8	16.7	*	0.9	2.6	8.4	16.0	37.1	70.2	161.8	394.0	1,148.7
2006 5/.....	51.9	15.8	*	0.8	3.3	8.9	17.0	35.4	72.3	155.6	403.7	1,107.6
2005 5/.....	54.2	12.7	2.1	1.2	2.8	9.0	18.0	38.9	77.6	162.6	418.8	1,170.4
2004 5/.....	53.1	14.2	2.3	1.2	2.8	8.4	19.0	39.4	74.0	168.8	405.9	1,179.4
2003 5/.....	55.4	15.8	1.8	0.9	3.3	9.7	19.8	39.2	79.6	184.6	413.4	1,258.7
2002.....	55.9	19.9	1.9	0.8	3.2	9.0	19.2	39.9	83.0	177.0	435.9	1,296.0
2001.....	55.8	23.8	2.7	1.0	3.2	8.9	20.3	42.2	82.3	183.7	426.2	1,267.0
2000.....	55.8	20.1	1.9	1.0	4.2	9.0	19.2	39.6	81.6	191.1	438.5	1,283.4
1999.....	56.3	23.6	1.8	0.9	3.4	8.9	20.4	40.5	87.9	192.0	452.4	1,257.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

All origins 3/, both sexes

2007.....	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.2	2.4	2.4
2006.....	0.4	*	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	2.2
2005.....	0.4	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.5	3.0
2004.....	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.3	2.5	3.2
2003.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001.....	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000.....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9

All origins 3/, male

2007.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.2	3.4
2006.....	0.4	*	*	*	*	*	0.2	0.7	0.9	1.4	2.7	2.3
2005.....	0.5	*	*	*	*	0.1	0.2	0.5	0.9	1.5	3.3	5.1
2004.....	0.5	*	*	*	*	0.1	0.2	0.6	0.7	1.4	3.1	4.8
2003.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002.....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001.....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999.....	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7

All origins 3/, female

2007.....	0.3	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8	2.0
2006.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	2.2	2.1
2005.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.1	2.0	2.1
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.1	2.0	2.5
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

Non-Hispanic 4/, both sexes

2007.....	0.4	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.4	2.5
2006.....	0.4	*	*	*	*	0.1	0.2	0.5	0.8	1.2	2.5	2.1
2005.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.5	3.1
2004.....	0.4	*	*	*	*	0.1	0.2	0.5	0.5	1.3	2.5	3.2
2003.....	0.5	*	*	*	*	0.1	0.3	0.4	0.8	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.7
2001.....	0.5	*	*	*	*	0.1	0.3	0.5	0.6	1.4	2.5	2.6
2000.....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.5	2.9
1999.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9

Non-Hispanic 4/, male

2007.....	0.5	*	*	*	*	0.2	0.3	0.6	0.9	1.5	3.3	3.6
2006.....	0.5	*	*	*	*	*	0.2	0.7	1.0	1.4	2.8	2.3
2005.....	0.5	*	*	*	*	*	0.2	0.6	0.8	1.4	3.3	5.3
2004.....	0.5	*	*	*	*	*	0.2	0.6	0.7	1.4	3.1	4.8
2003.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.3	4.1
2002.....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.6
2001.....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.6	3.9
2000.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.6	4.2
1999.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.7

Non-Hispanic 4/, female

2007.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2006.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	2.2	2.1
2005.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.1	2.0	2.2
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.2	2.0	2.6
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	1.9	2.5
2002.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.3
2001.....	0.4	*	*	*	*	*	0.3	0.3	0.5	1.2	1.8	2.0
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	1.8	2.3
1999.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

Non-Hispanic white, both sexes

2007 5/.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.1	2.4	2.5
2006 5/.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.2	2.4	2.1
2005 5/.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	2.5	3.2
2004 5/.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	3.3
2003 5/.....	0.5	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.2
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001.....	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.5	2.7
2000.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.8
1999.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	3.0

Non-Hispanic white, male

2007 5/.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	3.6
2006 5/.....	0.5	*	*	*	*	*	0.2	0.5	0.8	1.4	2.7	2.5
2005 5/.....	0.5	*	*	*	*	*	0.2	0.4	0.6	1.4	3.4	5.6
2004 5/.....	0.5	*	*	*	*	*	0.2	0.4	0.6	1.4	3.1	4.9
2003 5/.....	0.5	*	*	*	*	*	0.3	0.4	0.8	1.3	3.4	4.5
2002.....	0.5	*	*	*	*	0.2	0.3	0.5	0.6	1.5	3.2	3.7
2001.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.4	3.6	4.0
2000.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.6	4.3
1999.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	5.1

Non-Hispanic white, female

2007 5/.....	0.4	*	*	*	*	*	*	0.3	0.4	1.0	1.8	2.0
2006 5/.....	0.4	*	*	*	*	*	0.2	0.3	0.4	1.0	2.2	1.9
2005 5/.....	0.4	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.1
2004 5/.....	0.4	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.5
2003 5/.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.7
2002.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001.....	0.4	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7	2.2
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute and subacute endocarditis (I33)

Non-Hispanic black, both sexes

2007 5/.....	0.6	*	*	*	*	*	0.4	1.1	1.5	2.7	2.4	*
2006 5/.....	0.6	*	*	*	*	*	0.4	1.1	2.0	1.7	3.5	*
2005 5/.....	0.6	*	*	*	*	*	0.5	1.1	2.0	1.8	3.7	*
2004 5/.....	0.5	*	*	*	*	*	*	1.4	1.2	2.1	2.8	*
2003 5/.....	0.6	*	*	*	*	*	0.6	1.3	2.0	2.2	3.0	*
2002.....	0.5	*	*	*	*	*	0.4	1.3	1.6	2.3	*	*
2001.....	0.7	*	*	*	*	*	0.7	1.3	1.7	2.6	4.0	*
2000.....	0.6	*	*	*	*	*	0.6	1.4	1.6	1.9	2.9	*
1999.....	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*

Non-Hispanic black, male

2007 5/.....	0.8	*	*	*	*	*	*	1.7	2.4	3.8	*	*
2006 5/.....	0.7	*	*	*	*	*	*	1.5	2.7	*	*	*
2005 5/.....	0.7	*	*	*	*	*	*	1.5	2.8	*	*	*
2004 5/.....	0.6	*	*	*	*	*	*	1.9	*	*	*	*
2003 5/.....	0.7	*	*	*	*	*	*	2.0	2.3	*	*	*
2002.....	0.6	*	*	*	*	*	*	1.6	1.9	*	*	*
2001.....	0.8	*	*	*	*	*	0.8	1.9	2.2	3.9	*	*
2000.....	0.7	*	*	*	*	*	0.8	1.8	*	*	*	*
1999.....	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*

Non-Hispanic black, female

2007 5/.....	0.4	*	*	*	*	*	*	*	*	2.0	*	*
2006 5/.....	0.5	*	*	*	*	*	*	*	*	1.4	*	3.2
2005 5/.....	0.5	*	*	*	*	*	*	0.8	1.3	*	4.0	*
2004 5/.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2003 5/.....	0.5	*	*	*	*	*	*	*	1.7	2.1	*	*
2002.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001.....	0.5	*	*	*	*	*	0.7	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	1.0	1.6	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	1.6	2.9	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All origins 3/, both sexes

2007.....	0.3	0.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.1	2.1
2006.....	0.3	*	0.1	0.1	0.1	0.1	0.2	0.2	0.4	0.6	1.2	1.8
2005.....	0.3	0.5	*	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2004.....	0.3	0.5	0.2	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003.....	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002.....	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001.....	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999.....	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0

All origins 3/, male

2007.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.1	2.3
2006.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.3	1.8
2005.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.4	0.8	1.5	1.9
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.3
2003.....	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000.....	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5

All origins 3/, female

2007.....	0.3	*	*	*	0.1	0.1	0.1	0.3	0.3	0.5	1.0	2.0
2006.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.0	1.8
2005.....	0.3	*	*	0.1	*	0.1	0.2	0.3	0.3	0.6	1.0	1.9
2004.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.2	1.6
2003.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.7	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.1	2.2
2006.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2005.....	0.3	*	*	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2004.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003.....	0.3	0.9	*	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.7
2002.....	0.3	0.7	0.2	*	0.1	0.1	0.2	0.3	0.4	0.8	1.3	2.1
2001.....	0.3	0.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999.....	0.3	0.8	0.2	*	0.1	0.1	0.3	0.3	0.6	0.8	1.3	2.1

Non-Hispanic 4/, male

2007.....	0.3	*	*	*	*	0.2	0.3	0.3	0.5	0.8	1.1	2.4
2006.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.4	1.8
2005.....	0.3	*	*	*	*	0.1	0.2	0.4	0.4	0.8	1.5	2.0
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	2.4
2003.....	0.3	*	*	*	0.2	0.2	0.3	0.4	0.4	1.0	1.3	1.6
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.9	1.2	2.3
2001.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.4	0.9	1.4	2.1
1999.....	0.3	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.4	2.6

Non-Hispanic 4/, female

2007.....	0.3	*	*	*	0.1	0.1	0.1	0.3	0.3	0.5	1.0	2.1
2006.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.1	1.9
2005.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.0	1.9
2004.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	1.2	1.6
2003.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	1.9
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic white, both sexes

2007 5/.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.1	2.2
2006 5/.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.6	1.3	1.9
2005 5/.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.3	0.7	1.1	1.9
2004 5/.....	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.2	1.8
2003 5/.....	0.3	*	*	*	0.1	*	0.2	0.2	0.4	0.9	1.2	1.7
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.1
2001.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999.....	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.3	2.0

Non-Hispanic white, male

2007 5/.....	0.3	*	*	*	*	0.2	0.3	0.3	0.4	0.7	1.1	2.4
2006 5/.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.8	1.5	1.9
2005 5/.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.8	1.4	2.0
2004 5/.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.2	2.3
2003 5/.....	0.3	*	*	*	0.2	*	0.2	0.3	0.4	1.0	1.4	1.8
2002.....	0.3	*	*	*	*	0.2	0.2	0.3	0.4	0.9	1.2	2.3
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.2	2.0
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.8	1.4	2.1
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.7	0.8	1.4	2.6

Non-Hispanic white, female

2007 5/.....	0.3	*	*	*	*	*	*	0.3	0.3	0.5	1.1	2.0
2006 5/.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.1	1.9
2005 5/.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	0.9	1.8
2004 5/.....	0.3	*	*	*	*	*	*	0.2	0.3	0.6	1.1	1.6
2003 5/.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.1
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.8



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All origins 3/, both sexes												
2007.....	18.8	0.5	*	*	0.1	0.2	0.7	2.4	8.4	29.7	122.4	554.4
2006.....	20.2	*	*	*	0.1	0.2	0.8	2.8	8.9	31.5	132.7	614.4
2005.....	19.9	0.7	*	*	0.1	0.2	0.7	2.6	8.6	31.6	132.8	618.6
2004.....	19.5	0.7	*	*	0.1	0.2	0.7	2.5	8.4	32.2	131.2	621.2
2003.....	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002.....	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001.....	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000.....	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999.....	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All origins 3/, male												
2007.....	15.4	*	*	*	0.1	0.3	1.0	3.0	10.4	36.0	137.1	552.8
2006.....	16.2	*	*	*	0.1	0.3	0.9	3.7	10.8	36.9	148.6	599.0
2005.....	15.8	*	*	*	0.1	0.3	0.9	3.1	10.3	36.0	147.8	603.1
2004.....	15.4	*	*	*	*	0.3	0.8	3.0	10.1	37.5	147.0	605.5
2003.....	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002.....	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001.....	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000.....	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999.....	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All origins 3/, female												
2007.....	22.0	*	*	*	*	0.2	0.5	1.9	6.6	24.3	112.2	555.1
2006.....	24.0	*	*	*	*	0.2	0.6	2.0	7.0	26.9	121.8	621.6
2005.....	23.9	*	*	*	*	0.2	0.5	2.0	7.0	27.9	122.5	625.6
2004.....	23.4	*	*	*	*	0.2	0.6	2.0	6.8	27.8	120.5	628.2
2003.....	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002.....	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001.....	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000.....	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999.....	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Hispanic, both sexes												
2007.....	4.2	*	*	*	*	*	0.4	1.3	5.0	19.6	73.2	280.0
2006.....	4.1	*	*	*	*	*	*	1.5	4.4	18.8	74.5	304.8
2005.....	4.0	*	*	*	*	*	*	1.1	4.5	18.5	75.6	316.6
2004.....	3.7	*	*	*	*	*	0.4	1.3	4.7	19.2	72.9	287.3
2003.....	4.0	*	*	*	*	*	*	1.3	5.7	22.5	77.1	330.1
2002.....	3.6	*	*	*	*	*	*	1.2	4.6	19.9	75.2	316.0
2001.....	3.7	*	*	*	*	*	0.4	1.3	5.6	18.8	72.9	349.5
2000.....	3.6	*	*	*	*	*	*	0.7	5.8	18.9	72.1	369.6
1999.....	3.7	*	*	*	*	*	*	1.0	5.3	19.5	73.5	377.4
Hispanic, male												
2007.....	3.6	*	*	*	*	*	*	1.7	6.3	24.3	75.9	288.7
2006.....	3.5	*	*	*	*	*	*	2.1	4.9	22.9	89.8	276.3
2005.....	3.3	*	*	*	*	*	*	1.4	4.2	21.5	90.2	295.1
2004.....	3.1	*	*	*	*	*	*	1.7	6.0	22.1	84.1	263.4
2003.....	3.5	*	*	*	*	*	*	1.3	7.8	26.0	92.8	312.5
2002.....	3.0	*	*	*	*	*	*	1.1	5.8	23.1	85.8	296.2
2001.....	3.0	*	*	*	*	*	*	1.8	6.0	21.8	75.6	333.3
2000.....	2.9	*	*	*	*	*	*	*	7.7	23.0	74.2	380.9
1999.....	3.0	*	*	*	*	*	*	*	6.1	24.6	82.9	358.0
Hispanic, female												
2007.....	4.8	*	*	*	*	*	*	*	3.9	15.8	71.3	275.3
2006.....	4.8	*	*	*	*	*	*	0.9	4.0	15.5	63.7	319.9
2005.....	4.8	*	*	*	*	*	*	*	4.7	16.0	65.4	327.8
2004.....	4.4	*	*	*	*	*	*	1.0	3.4	16.9	65.2	299.6
2003.....	4.6	*	*	*	*	*	*	1.2	3.7	19.7	66.4	339.1
2002.....	4.4	*	*	*	*	*	*	1.3	3.6	17.3	68.0	325.8
2001.....	4.5	*	*	*	*	*	*	*	5.2	16.4	71.0	357.4
2000.....	4.3	*	*	*	*	*	*	*	4.2	15.6	70.7	364.0
1999.....	4.4	*	*	*	*	*	*	*	4.6	15.5	67.3	386.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic 4/, both sexes												
2007.....	21.3	*	*	*	0.1	0.2	0.8	2.6	8.7	30.5	125.5	568.6
2006.....	22.9	*	*	*	0.1	0.2	0.8	3.0	9.2	32.4	136.1	629.3
2005.....	22.5	0.7	*	*	0.1	0.3	0.8	2.7	8.9	32.5	136.0	632.5
2004.....	22.0	0.8	*	*	0.1	0.3	0.8	2.6	8.7	33.1	134.2	635.3
2003.....	22.2	1.0	*	*	0.1	0.3	0.8	2.5	9.1	34.1	135.9	653.2
2002.....	22.0	*	*	*	0.1	0.3	0.7	2.6	9.1	34.6	135.3	656.8
2001.....	22.4	*	*	*	0.1	0.3	0.7	2.5	9.6	35.2	138.8	685.6
2000.....	22.1	0.8	*	*	0.1	0.2	0.6	2.3	9.3	35.7	138.5	694.8
1999.....	21.8	*	*	*	0.1	0.2	0.6	2.2	9.1	36.3	139.4	691.0
Non-Hispanic 4/, male												
2007.....	17.6	*	*	*	0.1	0.3	1.1	3.2	10.7	36.9	141.2	567.9
2006.....	18.5	*	*	*	0.1	0.3	1.0	3.9	11.3	37.9	152.1	616.3
2005.....	18.0	*	*	*	0.1	0.3	1.0	3.3	10.8	37.0	151.0	618.8
2004.....	17.5	*	*	*	*	0.3	0.9	3.1	10.4	38.6	150.3	621.3
2003.....	17.7	1.4	*	*	0.1	0.3	1.0	3.1	11.0	40.5	151.2	643.2
2002.....	17.3	*	*	*	*	0.4	0.9	3.3	10.9	40.2	151.8	634.7
2001.....	17.4	*	*	*	*	0.3	0.9	3.2	11.4	41.4	157.2	658.0
2000.....	17.2	*	*	*	0.1	0.2	0.8	2.9	11.0	42.1	156.9	683.3
1999.....	16.9	*	*	*	0.1	0.2	0.8	2.8	11.0	43.5	154.5	677.1
Non-Hispanic 4/, female												
2007.....	24.9	*	*	*	*	0.2	0.6	2.0	6.8	25.0	114.8	568.9
2006.....	27.1	*	*	*	*	0.2	0.7	2.2	7.3	27.8	125.2	635.3
2005.....	26.9	*	*	*	*	0.2	0.6	2.1	7.2	28.8	125.7	638.7
2004.....	26.3	*	*	*	*	0.2	0.6	2.1	7.1	28.6	123.3	641.6
2003.....	26.5	*	*	*	*	0.2	0.6	2.0	7.3	28.8	125.6	657.6
2002.....	26.5	*	*	*	*	0.2	0.5	1.9	7.5	29.9	124.3	666.4
2001.....	27.1	*	*	*	*	0.2	0.6	1.9	8.0	30.1	126.8	697.0
2000.....	26.7	*	*	*	*	0.2	0.5	1.7	7.7	30.5	126.5	699.5
1999.....	26.5	*	*	*	*	0.2	0.5	1.6	7.2	30.4	129.7	696.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Heart failure (I50)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	24.1	*	*	*	0.1	0.2	0.5	2.0	7.5	29.1	127.1	590.2
2006 5/.....	26.0	*	*	*	*	0.1	0.6	2.2	8.2	31.2	138.3	653.7
2005 5/.....	25.4	*	*	*	0.1	0.2	0.5	2.1	7.8	31.1	136.7	655.9
2004 5/.....	24.8	*	*	*	*	0.2	0.6	2.0	7.6	32.3	135.7	657.1
2003 5/.....	25.0	*	*	*	*	0.2	0.5	1.9	8.0	32.8	137.1	673.5
2002.....	24.7	*	*	*	*	0.2	0.5	2.0	8.2	33.3	135.9	677.3
2001.....	25.1	*	*	*	*	0.2	0.5	2.1	8.7	33.9	139.7	706.8
2000.....	24.7	*	*	*	*	0.1	0.5	1.9	8.4	34.3	139.2	714.3
1999.....	24.3	*	*	*	*	0.1	0.5	1.7	8.1	34.9	139.1	709.1
<b>Non-Hispanic white, male</b>												
2007 5/.....	19.7	*	*	*	*	0.2	0.7	2.5	9.2	35.5	142.8	592.5
2006 5/.....	20.7	*	*	*	*	0.2	0.7	2.8	10.2	36.4	154.9	641.8
2005 5/.....	20.0	*	*	*	*	0.2	0.7	2.5	9.4	35.2	152.3	646.8
2004 5/.....	19.5	*	*	*	*	0.2	0.7	2.4	9.0	37.8	151.7	643.0
2003 5/.....	19.7	*	*	*	*	0.3	0.7	2.4	9.7	39.3	153.2	665.3
2002.....	19.2	*	*	*	*	0.2	0.7	2.6	9.8	38.6	153.2	656.2
2001.....	19.3	*	*	*	*	0.2	0.7	2.7	10.5	40.0	158.1	679.1
2000.....	19.0	*	*	*	*	0.2	0.6	2.4	10.1	41.0	157.9	701.6
1999.....	18.6	*	*	*	*	*	0.6	2.2	10.1	42.2	154.7	690.8
<b>Non-Hispanic white, female</b>												
2007 5/.....	28.4	*	*	*	*	*	0.4	1.5	5.9	23.6	116.1	589.1
2006 5/.....	31.0	*	*	*	*	*	0.5	1.6	6.3	26.8	126.8	659.3
2005 5/.....	30.5	*	*	*	*	*	0.4	1.6	6.2	27.5	125.9	660.0
2004 5/.....	30.0	*	*	*	*	*	0.4	1.5	6.2	27.6	124.7	663.4
2003 5/.....	30.1	*	*	*	*	*	0.4	1.5	6.3	27.2	126.3	677.2
2002.....	30.0	*	*	*	*	0.2	0.4	1.4	6.7	28.8	124.2	686.4
2001.....	30.7	*	*	*	*	0.2	0.4	1.4	6.9	28.8	127.5	718.4
2000.....	30.2	*	*	*	*	*	0.3	1.3	6.8	28.7	126.9	719.4
1999.....	29.9	*	*	*	*	*	0.3	1.2	6.3	28.8	128.8	716.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Heart failure (I50)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	14.3	*	*	*	*	0.7	2.5	7.3	20.7	51.1	146.5	467.9
2006 5/.....	14.6	*	*	*	*	0.8	2.3	8.7	20.0	52.5	149.9	492.6
2005 5/.....	14.9	*	*	*	*	0.8	2.4	7.5	20.7	54.8	163.9	497.8
2004 5/.....	14.1	*	*	*	*	0.7	2.0	7.7	19.9	49.2	153.0	513.2
2003 5/.....	14.5	*	*	*	*	0.7	2.4	7.1	21.0	55.3	152.7	538.7
2002.....	14.2	*	*	*	*	0.9	1.9	7.4	19.0	54.1	158.3	534.5
2001.....	14.3	*	*	*	*	0.6	2.0	6.6	20.0	55.5	159.1	551.2
2000.....	14.1	*	*	*	*	0.5	1.7	5.9	19.0	56.2	156.3	564.8
1999.....	14.3	*	*	*	*	0.6	1.7	6.5	19.2	57.3	164.8	557.4
<b>Non-Hispanic black, male</b>												
2007 5/.....	12.8	*	*	*	*	0.8	3.4	9.2	27.5	62.5	168.1	442.4
2006 5/.....	12.9	*	*	*	*	1.0	3.1	11.5	24.1	64.6	168.5	473.7
2005 5/.....	12.7	*	*	*	*	1.1	3.5	9.4	26.0	65.0	182.8	431.0
2004 5/.....	12.2	*	*	*	*	1.0	2.3	9.5	25.6	56.6	178.5	510.1
2003 5/.....	12.3	*	*	*	*	0.8	2.8	8.5	26.8	64.3	165.9	521.0
2002.....	12.0	*	*	*	*	1.1	2.3	9.2	23.3	65.9	176.4	489.0
2001.....	12.0	*	*	*	*	0.9	2.3	7.6	22.6	66.3	185.4	547.5
2000.....	11.6	*	*	*	*	*	1.8	7.3	21.8	62.9	176.5	589.4
1999.....	12.2	*	*	*	*	*	1.9	8.6	23.0	66.4	180.4	609.7
<b>Non-Hispanic black, female</b>												
2007 5/.....	15.7	*	*	*	*	*	1.6	5.7	15.3	43.1	134.2	478.5
2006 5/.....	16.2	*	*	*	*	*	1.7	6.2	16.6	43.8	139.3	500.4
2005 5/.....	16.9	*	*	*	*	*	1.4	5.8	16.5	47.5	153.2	524.7
2004 5/.....	15.7	*	*	*	*	*	1.8	6.1	15.4	43.9	138.6	514.4
2003 5/.....	16.5	*	*	*	*	*	2.0	5.8	16.3	48.9	145.2	545.6
2002.....	16.3	*	*	*	*	*	1.5	5.7	15.5	45.8	148.1	551.8
2001.....	16.4	*	*	*	*	*	1.7	5.6	18.0	47.9	144.4	552.6
2000.....	16.3	*	*	*	*	*	1.6	4.7	16.8	51.5	145.0	555.6
1999.....	16.2	*	*	*	*	*	1.5	4.8	16.1	50.9	156.0	537.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes

2007.....	37.8	8.2	0.8	0.5	1.8	4.0	8.3	17.5	34.1	80.3	234.6	770.9
2006.....	36.0	6.9	0.7	0.5	1.9	4.3	8.3	17.6	34.4	79.0	224.5	724.4
2005.....	37.0	7.1	0.7	0.5	2.0	4.2	8.7	18.2	36.0	83.2	230.2	754.0
2004.....	36.5	8.7	0.9	0.5	1.9	3.9	8.7	18.3	35.8	84.5	234.1	747.7
2003.....	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
2002.....	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001.....	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000.....	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999.....	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1

All origins 3/, male

2007.....	35.7	9.0	0.7	0.5	2.3	4.8	10.4	22.6	43.8	98.2	271.9	790.5
2006.....	34.0	6.9	0.8	0.6	2.3	5.4	10.3	22.6	43.9	97.1	258.4	735.0
2005.....	34.9	7.7	0.8	0.5	2.6	5.2	10.9	23.6	46.0	102.3	263.3	767.7
2004.....	34.5	9.2	0.9	0.5	2.4	4.8	10.7	23.8	46.3	103.1	269.4	765.2
2003.....	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
2002.....	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001.....	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000.....	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999.....	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7

All origins 3/, female

2007.....	39.9	7.3	0.9	0.4	1.3	3.1	6.3	12.7	25.0	65.1	208.9	761.7
2006.....	38.0	6.8	0.7	0.4	1.4	3.1	6.4	12.8	25.6	63.7	201.3	719.5
2005.....	39.0	6.4	0.7	0.4	1.3	3.1	6.5	13.0	26.8	67.1	207.8	747.6
2004.....	38.5	8.2	0.8	0.5	1.3	2.9	6.6	13.0	26.0	68.9	210.3	739.9
2003.....	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
2002.....	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001.....	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000.....	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999.....	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Hispanic, both sexes

2007.....	9.8	7.4	0.9	0.3	1.5	2.4	4.6	9.1	18.4	46.7	132.2	378.7
2006.....	9.2	5.3	0.6	0.4	1.2	2.6	4.2	9.8	18.6	47.4	120.2	354.5
2005.....	9.5	6.4	*	0.3	1.4	2.5	4.5	9.2	19.3	50.3	132.5	384.2
2004.....	9.1	7.2	0.7	0.3	1.4	2.1	4.5	9.5	18.7	49.6	131.9	369.1
2003.....	9.7	6.8	0.9	0.3	1.2	2.6	4.8	9.5	23.2	53.0	141.3	418.2
2002.....	9.3	8.6	0.6	0.3	1.4	2.4	4.7	9.6	19.7	50.9	141.6	419.6
2001.....	9.3	8.0	0.8	0.4	1.2	2.2	4.3	9.2	21.6	55.3	141.6	444.6
2000.....	9.6	8.7	0.7	0.5	1.6	2.5	4.9	10.2	22.0	54.9	152.1	457.2
1999.....	9.8	11.5	0.9	0.4	1.3	2.5	5.5	10.1	24.6	57.2	151.9	447.6

Hispanic, male

2007.....	10.0	7.9	*	*	1.9	3.0	6.1	12.2	23.2	55.4	153.9	389.3
2006.....	9.6	5.8	*	0.5	1.7	3.6	5.5	12.2	24.6	58.1	139.8	378.9
2005.....	10.0	6.9	*	*	1.7	3.4	6.1	12.7	26.3	62.3	153.7	401.0
2004.....	9.4	6.9	*	*	2.0	3.1	6.2	12.8	23.5	59.9	153.8	374.1
2003.....	9.8	7.0	*	*	1.6	3.0	6.4	12.7	28.3	66.2	159.1	437.2
2002.....	9.6	8.0	*	*	2.0	3.1	6.0	12.2	24.2	67.1	168.2	455.1
2001.....	9.3	7.9	*	*	1.5	2.8	5.7	12.6	27.3	67.7	170.4	438.1
2000.....	9.9	8.1	*	0.7	2.1	3.3	6.0	13.7	28.6	68.8	172.6	505.9
1999.....	10.3	9.8	*	*	1.5	3.4	7.2	13.7	31.9	75.6	179.4	526.2

Hispanic, female

2007.....	9.7	6.7	1.1	*	1.0	1.7	2.9	6.0	14.0	39.6	117.0	372.9
2006.....	8.8	4.6	*	*	0.7	1.3	2.8	7.4	13.2	38.7	106.4	341.5
2005.....	9.0	5.9	*	*	1.0	1.4	2.7	5.6	12.9	40.5	117.7	375.4
2004.....	8.8	7.4	*	*	0.7	1.0	2.6	6.1	14.3	41.3	116.8	366.6
2003.....	9.6	6.6	*	*	0.8	2.0	3.1	6.3	18.6	42.4	129.0	408.6
2002.....	9.0	9.3	*	*	0.8	1.5	3.3	7.0	15.8	38.0	123.4	402.0
2001.....	9.2	8.0	*	*	0.9	1.6	2.8	5.8	16.5	45.5	122.1	447.7
2000.....	9.3	9.3	*	*	1.0	1.7	3.8	6.8	16.1	44.0	138.3	433.3
1999.....	9.2	13.3	*	*	1.1	1.5	3.7	6.6	18.1	42.9	133.6	409.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 4/, both sexes

2007.....	42.7	8.3	0.8	0.5	1.9	4.3	9.0	18.5	35.4	82.9	241.1	791.1
2006.....	40.6	7.3	0.8	0.5	2.0	4.7	9.0	18.5	35.7	81.3	230.6	742.4
2005.....	41.5	7.2	0.8	0.5	2.1	4.6	9.4	19.1	37.4	85.5	235.7	770.9
2004.....	40.9	9.1	0.9	0.5	2.0	4.3	9.3	19.2	37.1	87.0	239.4	764.1
2003.....	42.4	9.0	0.9	0.5	2.1	4.4	9.7	19.1	39.1	93.7	250.9	795.8
2002.....	42.9	10.9	0.8	0.5	2.0	4.4	9.9	19.4	40.2	96.5	259.4	807.6
2001.....	42.1	10.1	1.2	0.5	1.9	4.3	9.4	19.3	40.4	97.5	253.9	807.8
2000.....	42.3	11.0	0.9	0.5	2.0	4.3	9.3	19.3	42.0	99.8	255.4	831.4
1999.....	43.0	11.9	0.9	0.5	2.2	4.3	9.6	19.5	44.0	103.6	261.3	837.9

Non-Hispanic 4/, male

2007.....	40.5	9.3	0.7	0.5	2.3	5.3	11.2	23.8	45.6	101.4	279.4	813.2
2006.....	38.5	7.2	0.8	0.6	2.5	5.9	11.1	23.7	45.5	99.8	265.4	753.7
2005.....	39.2	7.9	0.8	0.5	2.8	5.7	11.7	24.7	47.4	105.0	269.4	786.5
2004.....	38.8	9.7	0.9	0.5	2.4	5.2	11.4	24.9	48.1	106.0	275.4	784.1
2003.....	40.0	9.9	0.8	0.6	2.5	5.2	12.0	24.3	49.8	115.1	293.1	811.6
2002.....	40.6	11.6	0.8	0.6	2.6	5.5	12.3	25.3	50.8	117.7	300.6	836.7
2001.....	39.6	9.3	1.4	0.5	2.4	5.1	11.8	24.5	50.9	118.8	298.5	835.3
2000.....	40.0	10.9	1.0	0.6	2.4	5.1	11.5	25.1	54.1	123.4	297.1	883.9
1999.....	40.9	12.7	1.0	0.6	2.6	5.1	12.1	25.0	56.7	130.5	307.4	886.8

Non-Hispanic 4/, female

2007.....	44.9	7.4	0.8	0.5	1.4	3.4	6.9	13.4	26.0	67.1	214.7	780.7
2006.....	42.7	7.4	0.8	0.4	1.5	3.6	7.0	13.4	26.6	65.5	206.9	737.2
2005.....	43.7	6.5	0.8	0.5	1.4	3.4	7.1	13.7	28.0	69.1	212.8	763.9
2004.....	43.0	8.3	0.9	0.5	1.4	3.4	7.2	13.7	27.0	70.9	215.2	755.1
2003.....	44.6	8.1	1.0	0.4	1.6	3.6	7.5	14.0	29.1	75.8	222.8	788.8
2002.....	45.1	10.1	0.8	0.4	1.4	3.3	7.6	13.8	30.3	78.7	232.2	795.0
2001.....	44.5	11.0	1.0	0.5	1.4	3.5	6.9	14.2	30.6	79.7	224.6	796.3
2000.....	44.5	11.0	0.8	0.4	1.7	3.4	7.1	13.6	30.9	80.4	228.3	810.2
1999.....	44.9	11.1	0.9	0.5	1.8	3.6	7.2	14.1	32.2	81.5	231.4	818.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic white, both sexes

2007 5/.....	46.1	5.9	0.6	0.4	1.5	3.5	7.6	16.1	31.9	78.0	241.3	815.7
2006 5/.....	43.7	5.4	0.6	0.4	1.7	3.7	7.6	16.1	32.2	76.8	230.0	765.0
2005 5/.....	44.5	5.9	0.7	0.4	1.7	3.6	8.0	16.4	33.4	81.0	235.8	792.0
2004 5/.....	43.7	7.9	0.7	0.4	1.7	3.4	7.8	16.5	33.5	82.4	238.0	779.9
2003 5/.....	45.2	7.1	0.7	0.5	1.7	3.5	8.2	16.3	35.1	88.3	250.6	811.6
2002.....	45.8	9.2	0.7	0.5	1.6	3.4	8.4	16.5	36.2	92.3	258.6	820.9
2001.....	44.7	8.1	0.9	0.4	1.5	3.4	7.6	16.3	36.2	92.9	251.4	821.8
2000.....	44.9	9.6	0.8	0.4	1.6	3.4	7.6	16.3	38.0	95.0	253.1	845.3
1999.....	45.4	9.8	0.8	0.5	1.9	3.4	7.8	16.4	39.0	98.3	258.4	851.7

Non-Hispanic white, male

2007 5/.....	42.9	6.5	*	0.4	1.8	4.2	9.3	21.0	41.2	95.3	279.2	841.6
2006 5/.....	40.5	5.2	0.6	0.5	2.2	4.7	9.4	20.7	40.9	94.4	264.8	774.7
2005 5/.....	41.4	6.2	0.7	0.5	2.3	4.5	10.2	21.6	42.7	99.1	270.0	808.1
2004 5/.....	40.6	8.4	0.7	0.4	2.0	4.2	9.7	21.7	43.6	101.0	272.8	795.1
2003 5/.....	42.0	7.5	0.6	0.6	2.0	4.2	10.2	21.0	45.1	108.9	292.5	827.7
2002.....	42.7	9.8	0.8	0.5	2.1	4.2	10.7	21.9	46.2	113.1	299.5	850.7
2001.....	41.3	8.1	1.0	0.5	1.9	4.0	9.9	21.3	45.8	113.8	294.4	845.6
2000.....	41.7	9.6	0.8	0.5	1.8	4.1	9.6	21.5	49.3	117.9	294.7	898.3
1999.....	42.4	10.8	0.9	0.5	2.1	4.1	10.1	21.5	50.6	124.1	303.5	897.0

Non-Hispanic white, female

2007 5/.....	49.3	5.3	0.7	0.4	1.2	2.7	5.9	11.2	23.1	62.9	214.9	803.5
2006 5/.....	46.7	5.6	0.6	0.4	1.2	2.8	5.8	11.5	23.9	61.6	205.8	760.4
2005 5/.....	47.6	5.5	0.6	0.3	1.2	2.6	5.8	11.2	24.5	65.4	212.2	784.6
2004 5/.....	46.7	7.3	0.7	0.4	1.2	2.6	5.8	11.3	23.8	66.3	214.4	773.1
2003 5/.....	48.3	6.7	0.8	0.3	1.4	2.7	6.1	11.6	25.6	70.6	222.3	804.5
2002.....	48.9	8.5	0.6	0.4	1.1	2.5	6.2	11.2	26.6	74.6	231.1	808.1
2001.....	47.9	8.1	0.7	0.4	1.1	2.7	5.3	11.5	27.2	75.2	222.8	811.9
2000.....	47.9	9.6	0.7	0.3	1.3	2.7	5.6	11.1	27.4	75.7	225.7	824.0
1999.....	48.2	8.7	0.6	0.5	1.6	2.8	5.5	11.4	28.1	76.7	228.9	833.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic black, both sexes

2007 5/.....	37.7	17.0	1.3	0.9	3.6	9.2	18.9	39.0	71.7	145.5	291.2	656.9
2006 5/.....	36.5	14.3	1.5	0.8	3.5	10.6	19.2	38.2	72.5	136.9	288.9	614.7
2005 5/.....	37.4	12.7	1.2	0.9	3.9	10.5	19.7	41.3	76.7	142.3	285.8	649.7
2004 5/.....	37.7	12.8	1.9	0.9	3.5	9.4	20.3	41.7	76.7	147.1	300.0	678.7
2003 5/.....	38.8	15.2	1.6	0.8	4.0	10.1	20.9	42.3	80.8	161.3	298.6	711.8
2002.....	39.1	17.8	1.2	0.7	3.9	10.5	20.9	43.4	82.7	153.6	312.3	747.1
2001.....	39.3	18.2	2.4	0.9	3.8	10.0	21.4	43.7	83.3	157.1	322.1	737.9
2000.....	39.5	16.1	1.5	0.9	4.5	9.8	21.0	45.1	83.1	164.4	323.9	748.9
1999.....	40.9	20.9	1.5	0.9	4.4	10.1	22.3	45.2	93.3	171.8	331.4	752.7

Non-Hispanic black, male

2007 5/.....	39.1	19.6	1.7	0.9	4.8	11.0	24.7	49.2	94.4	186.9	350.4	637.0
2006 5/.....	38.2	14.8	1.7	0.9	4.0	13.4	24.1	49.9	95.9	174.7	340.1	647.8
2005 5/.....	38.8	15.2	*	1.0	5.4	12.9	23.9	52.5	98.6	184.1	331.0	672.7
2004 5/.....	39.0	13.9	1.9	0.7	4.5	11.2	24.6	53.0	101.2	181.6	366.5	728.6
2003 5/.....	39.9	18.3	1.9	0.9	5.1	11.4	25.6	53.9	105.1	201.8	359.8	718.1
2002.....	39.6	19.7	*	0.8	4.7	13.3	25.0	55.8	104.0	189.7	363.1	763.6
2001.....	40.3	16.0	2.7	0.9	5.0	11.9	25.9	53.5	109.1	190.6	401.2	813.1
2000.....	40.5	15.0	1.7	1.0	5.1	11.7	25.4	58.6	108.9	202.9	384.1	821.9
1999.....	42.8	20.8	*	1.1	5.6	12.1	27.4	57.6	123.1	220.6	401.5	855.7

Non-Hispanic black, female

2007 5/.....	36.4	14.2	*	0.9	2.4	7.6	13.8	30.3	53.4	115.9	257.5	665.1
2006 5/.....	34.9	13.8	*	0.7	3.0	8.0	14.9	28.3	53.7	109.8	259.8	601.3
2005 5/.....	36.2	10.1	*	0.8	2.5	8.2	15.9	31.8	59.0	112.3	260.1	640.4
2004 5/.....	36.5	11.6	1.8	1.1	2.5	7.7	16.5	32.0	57.0	122.6	262.4	659.0
2003 5/.....	37.8	11.9	*	0.7	2.8	8.8	16.7	32.3	61.2	132.5	264.0	709.4
2002.....	38.7	15.8	*	*	3.0	8.0	17.2	32.8	65.5	127.9	283.5	740.8
2001.....	38.5	20.5	2.1	0.9	2.7	8.3	17.3	35.3	62.5	133.5	277.6	710.1
2000.....	38.6	17.3	*	0.8	3.8	8.1	17.0	33.4	62.4	137.3	290.0	721.7
1999.....	39.2	21.1	*	*	3.2	8.2	17.9	34.7	69.5	137.3	292.2	714.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All origins 3/, both sexes

2007.....	7.9	*	*	*	0.1	0.2	0.9	2.8	6.5	16.2	49.5	191.1
2006.....	8.0	*	*	*	0.0	0.3	0.9	3.0	6.9	16.8	51.0	189.4
2005.....	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.7	55.6	210.0
2004.....	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	17.1	52.6	198.5
2003.....	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
2002.....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001.....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000.....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999.....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1

All origins 3/, male

2007.....	6.3	*	*	*	*	0.2	1.2	3.5	8.1	18.0	50.2	157.9
2006.....	6.4	*	*	*	*	0.3	1.1	3.8	8.4	18.5	51.2	159.2
2005.....	6.5	*	*	*	*	0.2	1.1	3.5	7.9	19.2	55.3	169.3
2004.....	6.1	*	*	*	*	0.3	1.0	3.5	7.4	18.2	53.3	166.0
2003.....	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
2002.....	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001.....	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000.....	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999.....	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4

All origins 3/, female

2007.....	9.5	*	*	*	*	0.2	0.6	2.1	5.0	14.6	49.0	207.0
2006.....	9.5	*	*	*	*	0.2	0.7	2.2	5.5	15.3	50.8	203.5
2005.....	10.3	*	*	*	*	0.2	0.7	2.0	5.1	16.5	55.8	228.7
2004.....	9.5	*	*	*	*	0.2	0.6	2.0	5.3	16.1	52.1	213.2
2003.....	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
2002.....	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001.....	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000.....	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999.....	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Hispanic, both sexes

2007.....	3.1	*	*	*	*	*	0.6	2.4	6.3	16.7	48.0	157.0
2006.....	2.8	*	*	*	*	*	0.5	2.3	7.1	16.9	40.1	147.1
2005.....	3.1	*	*	*	*	*	0.6	1.9	6.3	18.6	52.9	169.4
2004.....	2.7	*	*	*	*	*	0.4	2.2	5.4	16.1	44.3	172.2
2003.....	2.6	*	*	*	*	*	0.4	2.0	5.8	16.7	44.5	171.0
2002.....	2.4	*	*	*	*	*	0.5	1.1	6.2	15.5	47.2	148.6
2001.....	2.4	*	*	*	*	*	*	1.8	4.5	16.5	48.9	160.1
2000.....	2.3	*	*	*	*	*	0.5	1.8	6.4	15.0	45.8	148.0
1999.....	2.2	*	*	*	*	*	*	2.2	5.0	15.2	48.3	129.8

Hispanic, male

2007.....	2.7	*	*	*	*	*	0.7	2.8	7.8	20.2	48.4	151.4
2006.....	2.4	*	*	*	*	*	0.6	2.8	8.4	19.0	38.9	131.9
2005.....	2.5	*	*	*	*	*	0.8	2.1	6.1	20.5	54.1	140.7
2004.....	2.3	*	*	*	*	*	*	2.9	6.4	18.4	48.6	147.1
2003.....	2.1	*	*	*	*	*	*	2.8	7.4	15.5	44.5	133.9
2002.....	1.9	*	*	*	*	*	*	1.2	7.1	15.3	46.4	132.2
2001.....	1.9	*	*	*	*	*	*	2.3	3.9	16.8	51.0	129.5
2000.....	2.1	*	*	*	*	*	0.8	2.8	8.1	15.2	52.6	141.1
1999.....	1.9	*	*	*	*	*	*	2.4	5.2	18.5	51.6	131.6

Hispanic, female

2007.....	3.4	*	*	*	*	*	*	1.9	4.9	13.8	47.8	160.0
2006.....	3.3	*	*	*	*	*	*	1.9	6.0	15.3	40.9	155.2
2005.....	3.7	*	*	*	*	*	*	1.7	6.5	17.1	52.1	184.4
2004.....	3.1	*	*	*	*	*	*	1.5	4.6	14.3	41.2	185.1
2003.....	3.2	*	*	*	*	*	*	1.2	4.4	17.6	44.5	189.8
2002.....	3.0	*	*	*	*	*	*	*	5.5	15.7	47.8	156.7
2001.....	2.9	*	*	*	*	*	*	1.2	5.0	16.2	47.4	175.0
2000.....	2.5	*	*	*	*	*	*	*	4.9	14.9	41.3	151.3
1999.....	2.5	*	*	*	*	*	*	1.9	4.8	12.7	46.1	128.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic 4/, both sexes

2007.....	8.8	*	*	*	*	0.2	1.0	2.9	6.5	16.1	49.5	192.8
2006.....	8.8	*	*	*	*	0.3	1.0	3.1	6.8	16.7	51.6	191.2
2005.....	9.3	*	*	*	*	0.2	1.0	2.8	6.4	17.6	55.6	211.8
2004.....	8.7	*	*	*	*	0.3	0.9	2.8	6.4	17.1	52.9	199.4
2003.....	8.3	*	*	*	*	0.3	0.9	2.6	6.3	16.8	51.9	189.4
2002.....	7.7	*	*	*	*	0.2	0.8	2.4	5.7	15.9	48.1	181.1
2001.....	7.4	*	*	*	0.1	0.3	0.8	2.4	5.8	15.3	47.5	172.0
2000.....	7.0	*	*	*	*	0.2	0.8	2.4	5.8	15.1	45.3	162.9
1999.....	6.6	*	*	*	*	0.2	0.7	2.2	5.6	15.1	43.2	152.3

Non-Hispanic 4/, male

2007.....	7.0	*	*	*	*	0.3	1.2	3.6	8.1	17.8	50.2	158.2
2006.....	7.1	*	*	*	*	0.4	1.2	3.9	8.3	18.4	51.9	160.5
2005.....	7.2	*	*	*	*	0.3	1.2	3.7	8.0	19.0	55.3	170.7
2004.....	6.8	*	*	*	*	0.3	1.1	3.5	7.4	18.1	53.5	166.7
2003.....	6.3	*	*	*	*	0.3	1.0	3.3	7.3	18.5	49.1	154.4
2002.....	5.9	*	*	*	*	0.3	1.0	3.0	6.6	17.1	47.2	153.9
2001.....	5.7	*	*	*	*	0.3	1.0	3.0	6.6	16.5	45.3	151.1
2000.....	5.4	*	*	*	*	0.2	1.0	3.0	6.9	15.7	44.7	142.0
1999.....	5.1	*	*	*	*	0.2	0.9	2.7	6.5	16.4	42.4	131.9

Non-Hispanic 4/, female

2007.....	10.5	*	*	*	*	0.2	0.7	2.2	5.0	14.7	49.0	209.2
2006.....	10.5	*	*	*	*	0.2	0.7	2.2	5.5	15.3	51.4	205.5
2005.....	11.3	*	*	*	*	0.2	0.8	2.0	5.0	16.4	55.9	230.6
2004.....	10.5	*	*	*	*	0.3	0.6	2.0	5.4	16.2	52.6	214.0
2003.....	10.2	*	*	*	*	0.2	0.8	1.8	5.4	15.5	53.8	204.7
2002.....	9.4	*	*	*	*	0.2	0.6	1.9	4.8	15.0	48.7	192.8
2001.....	9.0	*	*	*	*	0.3	0.6	1.9	5.1	14.4	48.9	180.7
2000.....	8.5	*	*	*	*	0.2	0.6	1.8	4.9	14.6	45.8	171.3
1999.....	8.0	*	*	*	*	0.2	0.6	1.7	4.7	14.1	43.7	160.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic white, both sexes

2007 5/.....	8.6	*	*	*	*	0.1	0.6	1.8	4.6	12.8	44.0	184.1
2006 5/.....	8.5	*	*	*	*	0.1	0.5	1.9	4.9	13.1	45.8	180.8
2005 5/.....	9.0	*	*	*	*	*	0.5	1.7	4.5	13.4	49.6	201.5
2004 5/.....	8.3	*	*	*	*	0.1	0.5	1.7	4.2	13.1	46.7	188.3
2003 5/.....	8.0	*	*	*	*	0.1	0.5	1.6	4.2	12.6	46.2	181.0
2002.....	7.4	*	*	*	*	0.1	0.4	1.5	3.9	12.5	42.2	170.6
2001.....	7.0	*	*	*	*	0.1	0.4	1.4	3.9	11.7	41.8	162.8
2000.....	6.5	*	*	*	*	0.1	0.4	1.3	3.8	11.6	39.9	152.7
1999.....	6.1	*	*	*	*	0.1	0.3	1.2	3.6	11.8	37.3	141.7

Non-Hispanic white, male

2007 5/.....	6.6	*	*	*	*	*	0.9	2.3	5.9	14.1	44.7	149.9
2006 5/.....	6.5	*	*	*	*	0.2	0.6	2.5	6.1	14.2	45.9	152.0
2005 5/.....	6.6	*	*	*	*	*	0.7	2.3	5.7	14.3	48.6	160.8
2004 5/.....	6.3	*	*	*	*	0.2	0.7	2.2	5.2	14.2	46.9	159.6
2003 5/.....	5.8	*	*	*	*	*	0.7	2.1	4.9	14.4	43.6	145.4
2002.....	5.5	*	*	*	*	*	0.5	2.0	4.7	13.5	41.0	145.4
2001.....	5.1	*	*	*	*	0.2	0.5	1.7	4.5	12.8	39.6	142.1
2000.....	4.9	*	*	*	*	*	0.6	1.8	4.7	12.4	39.5	131.8
1999.....	4.6	*	*	*	*	0.2	0.5	1.6	4.5	13.0	36.5	119.8

Non-Hispanic white, female

2007 5/.....	10.5	*	*	*	*	*	0.3	1.2	3.2	11.7	43.5	200.4
2006 5/.....	10.3	*	*	*	*	*	0.3	1.2	3.6	12.1	45.7	194.2
2005 5/.....	11.2	*	*	*	*	*	0.4	1.1	3.3	12.6	50.3	220.1
2004 5/.....	10.2	*	*	*	*	*	0.3	1.1	3.3	12.3	46.6	201.2
2003 5/.....	10.0	*	*	*	*	*	0.4	1.1	3.5	11.2	47.9	196.6
2002.....	9.2	*	*	*	*	*	0.3	1.1	3.1	11.5	42.9	181.4
2001.....	8.8	*	*	*	*	*	0.3	1.0	3.3	10.8	43.2	171.5
2000.....	8.1	*	*	*	*	*	0.3	0.9	3.1	10.9	40.1	161.1
1999.....	7.6	*	*	*	*	*	0.2	0.9	2.8	10.8	37.8	150.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic black, both sexes

2007 5/.....	12.1	*	*	*	*	0.9	3.2	10.4	22.8	46.0	108.2	301.7
2006 5/.....	12.8	*	*	*	*	1.2	3.8	11.3	23.4	50.1	118.9	322.8
2005 5/.....	13.1	*	*	*	*	1.0	3.6	10.8	23.0	55.2	124.3	345.2
2004 5/.....	12.8	*	*	*	*	1.0	3.2	10.4	24.3	52.8	124.9	339.6
2003 5/.....	12.2	*	*	*	*	1.0	3.2	9.5	24.7	55.1	121.0	304.3
2002.....	11.5	*	*	*	*	0.9	3.4	9.2	21.4	48.3	120.0	312.4
2001.....	11.3	*	*	*	*	1.1	3.2	10.1	22.4	48.4	113.9	291.5
2000.....	11.0	*	*	*	*	0.7	3.2	9.9	23.3	49.5	108.9	285.9
1999.....	10.7	*	*	*	*	0.7	3.2	9.4	21.8	47.5	112.9	272.2

Non-Hispanic black, male

2007 5/.....	10.8	*	*	*	*	0.9	3.9	12.9	27.6	54.8	118.5	251.3
2006 5/.....	11.7	*	*	*	*	1.4	4.9	14.2	28.7	60.6	129.0	274.9
2005 5/.....	11.8	*	*	*	*	1.1	4.0	13.6	29.8	65.0	139.0	294.0
2004 5/.....	11.1	*	*	*	*	1.2	4.0	13.2	28.8	57.6	139.2	268.3
2003 5/.....	10.8	*	*	*	*	1.3	3.5	13.0	30.0	61.3	120.9	280.2
2002.....	9.9	*	*	*	*	1.0	4.3	11.3	24.0	52.6	129.5	276.8
2001.....	10.0	*	*	*	*	1.2	3.9	13.1	25.8	55.5	119.7	277.2
2000.....	9.5	*	*	*	*	*	3.8	12.1	28.4	51.5	110.5	268.9
1999.....	9.4	*	*	*	*	*	3.8	12.0	24.8	52.3	119.1	268.6

Non-Hispanic black, female

2007 5/.....	13.2	*	*	*	*	0.9	2.6	8.2	19.0	39.7	102.3	322.6
2006 5/.....	13.9	*	*	*	*	1.0	2.8	8.8	19.1	42.7	113.1	342.3
2005 5/.....	14.3	*	*	*	*	0.8	3.1	8.3	17.6	48.2	116.0	365.8
2004 5/.....	14.3	*	*	*	*	1.0	2.5	8.0	20.8	49.4	116.9	367.8
2003 5/.....	13.4	*	*	*	*	*	2.8	6.5	20.5	50.8	121.1	313.6
2002.....	13.0	*	*	*	*	0.7	2.6	7.4	19.4	45.3	114.6	326.1
2001.....	12.4	*	*	*	*	1.0	2.5	7.5	19.7	43.4	110.7	296.9
2000.....	12.4	*	*	*	*	*	2.6	8.0	19.3	48.1	108.0	292.2
1999.....	11.8	*	*	*	*	0.7	2.7	7.2	19.4	44.1	109.4	273.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Cerebrovascular diseases (I60-I69)

##### All origins 3/, both sexes

2007.....	45.1	3.1	0.3	0.2	0.5	1.2	4.9	14.6	32.1	93.0	322.3	1,015.5
2006.....	45.8	3.4	0.3	0.2	0.5	1.3	5.1	14.7	33.3	96.3	335.1	1,039.6
2005.....	48.4	3.1	0.4	0.2	0.5	1.4	5.2	15.0	33.0	101.1	359.0	1,141.8
2004.....	51.1	3.1	0.3	0.2	0.5	1.4	5.4	14.9	34.3	107.8	386.2	1,245.9
2003.....	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002.....	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8

##### All origins 3/, male

2007.....	36.4	3.5	0.2	0.2	0.5	1.2	5.3	16.2	38.0	105.2	333.2	895.7
2006.....	37.0	3.9	0.3	0.3	0.5	1.4	5.3	16.4	38.7	108.0	345.5	932.4
2005.....	38.8	3.5	0.5	0.3	0.4	1.5	5.2	16.5	38.5	113.6	372.9	1,023.3
2004.....	40.7	3.4	0.3	0.2	0.5	1.4	5.6	16.7	39.5	121.1	402.9	1,118.1
2003.....	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002.....	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001.....	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000.....	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999.....	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8

##### All origins 3/, female

2007.....	53.5	2.6	0.4	0.2	0.4	1.3	4.6	12.9	26.6	82.7	314.9	1,072.4
2006.....	54.4	2.9	0.4	0.2	0.5	1.2	4.8	13.0	28.2	86.5	328.0	1,089.8
2005.....	57.8	2.6	0.3	0.2	0.5	1.2	5.1	13.6	27.9	90.5	349.5	1,196.1
2004.....	61.2	2.8	*	0.2	0.5	1.4	5.1	13.1	29.5	96.6	374.9	1,303.4
2003.....	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002.....	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001.....	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000.....	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999.....	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Hispanic, both sexes												
2007.....	15.6	2.8	*	*	0.5	1.2	4.4	13.8	33.9	81.9	252.2	652.9
2006.....	15.8	3.4	*	*	0.7	1.2	4.6	14.4	34.2	87.3	262.1	687.0
2005.....	16.0	2.6	*	*	0.6	1.0	4.6	15.0	33.4	89.4	275.5	738.4
2004.....	16.4	2.2	*	*	0.5	1.2	4.9	14.6	34.9	97.0	293.1	813.0
2003.....	16.7	*	*	*	0.4	1.2	4.5	14.9	35.3	98.4	316.8	880.5
2002.....	16.6	2.4	*	0.3	0.4	1.3	4.4	15.3	35.9	102.9	319.9	898.0
2001.....	17.4	*	*	*	0.6	1.4	5.0	15.9	35.4	106.2	341.2	1,042.9
2000.....	17.5	*	*	*	0.5	1.7	5.2	15.2	39.8	113.2	344.1	1,082.9
1999.....	17.4	*	*	*	0.5	1.6	4.9	15.7	37.7	116.9	348.3	1,077.6
Hispanic, male												
2007.....	14.1	*	*	*	0.5	1.3	5.3	16.5	42.9	94.6	263.6	594.6
2006.....	14.3	*	*	*	0.8	1.4	5.5	17.0	41.1	100.1	292.8	581.9
2005.....	14.4	*	*	*	0.7	1.2	5.2	17.8	40.3	106.2	294.0	692.4
2004.....	15.0	*	*	*	0.5	1.5	5.9	17.5	42.9	114.4	323.3	778.9
2003.....	14.9	*	*	*	*	1.3	5.2	18.1	43.5	113.9	337.1	837.4
2002.....	15.0	*	*	*	*	1.5	5.1	18.6	45.0	124.6	338.5	856.7
2001.....	15.7	*	*	*	0.7	1.6	5.3	18.7	43.5	127.2	386.3	1,005.6
2000.....	15.8	*	*	*	*	1.9	5.5	18.1	48.8	136.1	392.9	1,029.9
1999.....	16.1	*	*	*	0.6	1.8	5.7	19.5	46.1	140.8	395.2	1,136.6
Hispanic, female												
2007.....	17.1	*	*	*	*	1.1	3.3	11.0	25.4	71.6	244.2	684.5
2006.....	17.5	*	*	*	0.6	0.9	3.7	11.8	27.8	76.9	240.6	742.9
2005.....	17.7	*	*	*	*	0.9	3.9	12.1	27.1	75.8	262.6	762.5
2004.....	17.9	*	*	*	*	0.9	3.9	11.8	27.7	83.0	272.2	830.4
2003.....	18.6	*	*	*	*	1.1	3.7	11.7	27.8	86.0	302.8	902.3
2002.....	18.4	*	*	*	*	1.0	3.7	12.0	27.6	85.6	307.2	918.5
2001.....	19.1	*	*	*	*	1.2	4.7	13.1	28.2	89.6	310.7	1,061.2
2000.....	19.4	*	*	*	*	1.4	4.9	12.4	31.9	95.2	311.3	1,108.9
1999.....	18.8	*	*	*	*	1.4	4.0	11.9	30.3	98.1	317.1	1,049.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	50.3	3.2	0.3	0.2	0.5	1.2	5.0	14.6	31.9	93.7	326.6	1,034.0
2006.....	50.9	3.5	0.4	0.2	0.4	1.3	5.2	14.6	33.1	96.8	339.1	1,056.5
2005.....	53.8	3.2	0.4	0.2	0.4	1.4	5.2	15.0	33.0	101.7	363.3	1,159.8
2004.....	56.7	3.3	0.3	0.2	0.5	1.5	5.4	14.8	34.1	108.4	390.8	1,263.8
2003.....	60.1	2.6	0.3	0.2	0.6	1.5	5.7	15.0	35.6	113.7	415.0	1,388.4
2002.....	62.4	3.0	0.3	0.2	0.4	1.5	5.5	15.0	37.2	121.3	435.5	1,465.0
2001.....	63.2	2.8	0.3	0.2	0.5	1.5	5.6	14.9	38.0	124.3	447.7	1,514.0
2000.....	65.4	3.6	0.3	0.2	0.5	1.5	5.8	15.9	40.9	129.2	465.0	1,604.5
1999.....	65.7	2.9	0.3	0.2	0.5	1.4	5.8	15.1	40.6	131.2	473.8	1,630.4
<b>Non-Hispanic 4/, male</b>												
2007.....	40.5	3.8	*	0.2	0.5	1.2	5.3	16.1	37.4	105.8	337.3	912.5
2006.....	41.1	4.1	0.3	0.3	0.5	1.4	5.3	16.2	38.4	108.3	348.2	950.4
2005.....	43.0	3.6	0.5	0.3	0.4	1.6	5.2	16.3	38.3	113.9	377.0	1,039.8
2004.....	45.0	3.7	0.3	0.2	0.5	1.4	5.6	16.5	39.0	121.4	406.7	1,133.7
2003.....	47.6	2.8	*	0.2	0.6	1.6	5.9	16.5	40.4	128.4	435.7	1,252.9
2002.....	48.9	3.5	0.4	0.2	0.5	1.4	5.3	16.4	42.3	135.4	450.1	1,336.0
2001.....	49.7	3.3	0.3	0.2	0.5	1.5	5.7	16.4	43.2	140.9	469.8	1,392.9
2000.....	51.5	4.2	*	0.2	0.5	1.5	5.8	17.3	46.8	145.1	493.7	1,499.9
1999.....	51.5	3.7	0.3	0.2	0.5	1.4	5.8	16.4	46.3	147.3	503.6	1,524.8
<b>Non-Hispanic 4/, female</b>												
2007.....	59.6	2.5	0.5	0.2	0.4	1.3	4.8	13.1	26.7	83.5	319.2	1,091.4
2006.....	60.3	2.8	0.4	0.2	0.4	1.2	5.0	13.1	28.2	87.0	332.8	1,105.8
2005.....	64.1	2.8	*	0.2	0.5	1.3	5.2	13.7	28.0	91.4	354.1	1,214.5
2004.....	67.8	3.0	*	0.2	0.5	1.5	5.2	13.2	29.5	97.4	380.0	1,321.9
2003.....	72.0	2.4	0.3	0.1	0.5	1.4	5.5	13.5	31.0	101.4	401.1	1,448.0
2002.....	75.3	2.5	*	0.2	0.3	1.5	5.8	13.7	32.4	109.5	425.8	1,520.7
2001.....	76.1	2.4	0.4	0.2	0.5	1.5	5.5	13.5	33.2	110.4	433.2	1,564.4
2000.....	78.7	2.8	0.4	0.2	0.5	1.5	5.8	14.6	35.5	116.1	446.4	1,646.8
1999.....	79.2	2.0	*	0.2	0.5	1.4	5.8	13.9	35.4	118.1	454.4	1,672.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	53.5	2.8	0.3	0.2	0.4	1.0	3.8	11.0	26.0	84.2	319.3	1,045.4
2006 5/.....	54.2	2.4	0.4	0.2	0.4	1.0	3.8	11.1	26.8	87.1	331.9	1,068.3
2005 5/.....	57.4	2.4	0.3	0.2	0.3	1.1	3.9	11.1	26.9	92.3	355.8	1,171.8
2004 5/.....	60.6	2.6	0.3	0.2	0.4	1.1	4.2	10.9	27.8	98.7	381.8	1,276.4
2003 5/.....	64.2	2.0	0.2	0.1	0.5	1.2	4.1	10.9	28.7	103.6	406.6	1,401.1
2002.....	67.0	2.2	0.3	0.2	0.4	1.2	4.1	11.1	30.8	110.8	427.5	1,479.2
2001.....	67.6	1.9	0.3	0.2	0.4	1.3	4.2	10.8	31.2	113.0	439.5	1,525.0
2000.....	70.0	2.6	0.3	0.2	0.5	1.2	4.3	12.0	34.2	119.3	455.7	1,616.8
1999.....	70.4	2.4	0.3	0.2	0.4	1.1	4.3	11.2	33.4	120.7	465.1	1,648.0
<b>Non-Hispanic white, male</b>												
2007 5/.....	41.9	3.1	*	0.2	0.4	0.9	4.0	12.3	30.0	93.8	326.3	922.4
2006 5/.....	42.6	2.6	*	0.3	0.4	1.1	3.9	12.1	30.3	96.5	340.5	960.2
2005 5/.....	44.8	2.8	0.4	0.2	0.3	1.2	4.0	12.1	30.7	102.4	368.2	1,050.5
2004 5/.....	47.0	2.9	*	0.2	0.5	1.1	4.4	12.1	31.1	110.0	396.9	1,145.3
2003 5/.....	49.7	1.9	*	0.2	0.5	1.3	4.2	12.1	32.1	116.9	426.0	1,264.2
2002.....	51.3	2.7	*	0.2	0.5	1.1	4.0	12.1	34.5	123.2	441.1	1,345.9
2001.....	52.0	2.2	*	0.2	0.4	1.3	4.3	11.9	35.1	128.0	460.5	1,399.0
2000.....	53.9	3.3	*	0.2	0.5	1.2	4.3	13.0	38.7	133.1	482.3	1,505.9
1999.....	53.9	3.1	*	0.2	0.4	1.1	4.4	12.0	38.0	135.0	492.8	1,535.1
<b>Non-Hispanic white, female</b>												
2007 5/.....	64.7	2.4	*	0.2	0.4	1.0	3.6	9.8	22.1	75.9	314.4	1,103.7
2006 5/.....	65.5	2.2	0.4	*	0.4	1.0	3.7	10.1	23.5	79.0	325.9	1,118.7
2005 5/.....	69.6	2.0	*	*	0.4	1.1	3.9	10.2	23.3	83.6	347.2	1,227.3
2004 5/.....	73.7	2.2	*	0.2	0.4	1.2	4.0	9.8	24.7	89.0	371.6	1,335.1
2003 5/.....	78.2	2.1	*	*	0.4	1.1	4.0	9.7	25.5	92.1	393.6	1,461.3
2002.....	82.1	*	*	0.2	0.3	1.2	4.3	10.1	27.2	100.2	418.4	1,536.7
2001.....	82.6	*	*	*	0.3	1.3	4.1	9.8	27.4	100.3	425.6	1,577.4
2000.....	85.5	1.9	*	0.2	0.4	1.2	4.3	10.9	29.9	107.6	438.3	1,661.6
1999.....	86.1	*	*	*	0.4	1.1	4.3	10.5	29.1	108.8	447.0	1,693.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	44.4	5.2	*	*	0.7	2.7	12.0	37.7	78.8	180.4	433.3	1,012.8
2006 5/.....	44.7	7.7	*	0.3	0.7	2.8	12.7	37.9	83.0	181.3	442.7	1,023.0
2005 5/.....	46.5	6.2	*	0.4	0.8	3.1	12.5	40.5	81.5	186.0	475.7	1,122.3
2004 5/.....	48.7	6.2	*	*	0.8	3.3	12.7	40.0	86.0	199.4	519.1	1,226.1
2003 5/.....	51.1	4.9	*	*	1.0	3.2	14.2	42.0	91.3	204.9	539.1	1,366.0
2002.....	51.7	5.6	*	0.4	0.6	3.2	13.3	41.6	89.7	218.7	558.8	1,398.0
2001.....	52.9	6.9	*	*	0.9	3.0	13.6	43.5	92.8	226.0	567.2	1,465.5
2000.....	54.3	8.0	*	*	0.7	3.0	14.9	44.3	95.2	226.3	593.8	1,545.8
1999.....	53.9	4.7	*	*	0.7	2.9	14.7	42.2	97.7	235.7	596.4	1,501.5
<b>Non-Hispanic black, male</b>												
2007 5/.....	41.1	7.4	*	*	0.9	2.7	13.0	42.1	102.2	227.3	501.4	883.0
2006 5/.....	40.8	10.2	*	*	0.6	2.9	13.3	44.6	108.6	222.7	479.4	894.3
2005 5/.....	41.8	6.8	*	*	0.8	3.5	12.8	45.9	106.6	228.5	510.4	998.2
2004 5/.....	43.1	7.1	*	*	1.0	3.1	13.3	46.0	110.2	240.1	558.7	1,077.1
2003 5/.....	44.9	*	*	*	0.8	3.5	14.9	48.1	114.1	241.8	597.6	1,200.7
2002.....	45.1	*	*	*	0.7	2.9	12.7	47.5	112.7	267.3	595.2	1,266.8
2001.....	46.2	8.3	*	*	0.7	3.1	13.5	49.9	113.9	273.5	620.5	1,359.4
2000.....	47.7	9.2	*	*	*	2.8	15.1	50.4	116.9	272.1	664.9	1,479.5
1999.....	47.5	*	*	*	*	2.9	15.0	49.6	119.5	279.1	656.7	1,455.9
<b>Non-Hispanic black, female</b>												
2007 5/.....	47.4	*	*	*	*	2.8	11.1	33.9	60.0	146.9	394.6	1,066.6
2006 5/.....	48.3	*	*	*	0.8	2.7	12.2	32.1	62.5	151.8	421.9	1,075.5
2005 5/.....	50.8	*	*	*	0.9	2.7	12.2	35.9	61.3	155.6	456.0	1,172.3
2004 5/.....	53.8	*	*	*	*	3.5	12.2	34.8	66.6	170.4	496.7	1,285.0
2003 5/.....	56.7	*	*	*	1.1	3.0	13.7	36.8	73.0	178.6	506.0	1,430.0
2002.....	57.7	*	*	*	*	3.5	13.8	36.6	71.3	184.2	538.2	1,448.1
2001.....	59.0	*	*	*	1.1	2.9	13.6	38.1	75.9	192.5	537.2	1,504.8
2000.....	60.3	*	*	*	0.8	3.2	14.8	39.2	77.8	194.1	553.8	1,570.4
1999.....	59.8	*	*	*	0.8	2.8	14.4	35.9	80.2	205.1	562.7	1,518.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

All origins 3/, both sexes

2007.....	2.7	*	*	*	*	*	0.1	0.3	1.1	4.3	17.6	83.3
2006.....	2.9	*	*	*	*	*	0.1	0.3	1.2	4.5	18.9	89.9
2005.....	4.0	*	*	*	*	*	0.1	0.5	1.4	5.4	25.9	132.9
2004.....	4.0	*	*	*	*	*	0.1	0.4	1.6	6.1	26.3	137.0
2003.....	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002.....	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001.....	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000.....	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999.....	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0

All origins 3/, male

2007.....	2.2	*	*	*	*	*	*	0.4	1.5	5.3	19.3	76.5
2006.....	2.3	*	*	*	*	*	0.1	0.5	1.6	5.3	20.5	85.3
2005.....	3.1	*	*	*	*	*	0.1	0.6	1.8	6.6	29.0	121.7
2004.....	3.1	*	*	*	*	*	0.1	0.5	2.0	7.3	29.7	128.2
2003.....	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002.....	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001.....	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000.....	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999.....	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0

All origins 3/, female

2007.....	3.3	*	*	*	*	*	*	0.2	0.6	3.4	16.5	86.5
2006.....	3.5	*	*	*	*	*	*	0.2	0.9	3.8	17.9	92.1
2005.....	4.9	*	*	*	*	*	*	0.3	1.1	4.4	23.9	138.1
2004.....	4.9	*	*	*	*	*	*	0.2	1.2	5.1	24.0	141.0
2003.....	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002.....	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001.....	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000.....	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Atherosclerosis (I70)</b>												
Hispanic, both sexes												
2007.....	0.7	*	*	*	*	*	*	*	*	2.9	15.0	50.4
2006.....	0.6	*	*	*	*	*	*	*	*	2.6	11.5	50.5
2005.....	0.9	*	*	*	*	*	*	*	0.9	3.9	17.7	79.6
2004.....	0.9	*	*	*	*	*	*	*	1.0	4.3	18.6	74.7
2003.....	0.9	*	*	*	*	*	*	*	1.1	4.6	18.9	86.0
2002.....	1.1	*	*	*	*	*	*	*	1.0	4.2	22.7	108.6
2001.....	1.1	*	*	*	*	*	*	*	*	4.0	23.1	130.7
2000.....	1.2	*	*	*	*	*	*	*	*	5.9	24.5	132.7
1999.....	1.2	*	*	*	*	*	*	*	1.5	5.2	27.1	129.1
Hispanic, male												
2007.....	0.6	*	*	*	*	*	*	*	*	3.7	17.2	42.7
2006.....	0.5	*	*	*	*	*	*	*	*	3.4	12.3	45.1
2005.....	0.8	*	*	*	*	*	*	*	*	5.2	20.4	83.4
2004.....	0.8	*	*	*	*	*	*	*	*	5.1	22.9	88.3
2003.....	0.9	*	*	*	*	*	*	*	*	6.1	24.2	95.4
2002.....	0.9	*	*	*	*	*	*	*	*	5.2	26.7	102.1
2001.....	0.9	*	*	*	*	*	*	*	*	4.8	27.3	127.6
2000.....	1.0	*	*	*	*	*	*	*	*	8.2	32.0	104.8
1999.....	1.1	*	*	*	*	*	*	*	*	6.8	31.3	151.0
Hispanic, female												
2007.....	0.9	*	*	*	*	*	*	*	*	*	13.4	54.6
2006.....	0.8	*	*	*	*	*	*	*	*	*	10.8	53.3
2005.....	1.1	*	*	*	*	*	*	*	*	2.7	15.8	77.6
2004.....	1.0	*	*	*	*	*	*	*	*	3.7	15.6	67.7
2003.....	1.0	*	*	*	*	*	*	*	*	3.4	15.3	81.3
2002.....	1.3	*	*	*	*	*	*	*	*	3.5	20.0	111.9
2001.....	1.4	*	*	*	*	*	*	*	*	3.5	20.2	132.2
2000.....	1.4	*	*	*	*	*	*	*	*	4.0	19.5	146.4
1999.....	1.4	*	*	*	*	*	*	*	*	4.0	24.3	118.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

Non-Hispanic 4/, both sexes

2007.....	3.1	*	*	*	*	*	0.1	0.3	1.1	4.4	17.8	84.9
2006.....	3.3	*	*	*	*	*	0.1	0.3	1.3	4.6	19.4	91.8
2005.....	4.5	*	*	*	*	*	0.1	0.5	1.5	5.5	26.4	135.3
2004.....	4.5	*	*	*	*	*	0.1	0.4	1.6	6.3	26.7	139.7
2003.....	5.0	*	*	*	*	*	0.1	0.5	1.7	6.7	30.2	157.1
2002.....	5.4	*	*	*	*	*	0.1	0.5	1.9	7.3	32.2	170.2
2001.....	5.5	*	*	*	*	*	0.1	0.5	1.9	7.5	33.4	180.5
2000.....	5.7	*	*	*	*	*	0.1	0.5	1.9	7.8	35.0	190.3
1999.....	5.9	*	*	*	*	*	0.1	0.5	2.1	8.1	36.1	203.9

Non-Hispanic 4/, male

2007.....	2.5	*	*	*	*	*	*	0.5	1.6	5.4	19.5	78.4
2006.....	2.6	*	*	*	*	*	*	0.5	1.7	5.5	20.9	87.5
2005.....	3.5	*	*	*	*	*	0.1	0.7	1.9	6.7	29.5	123.7
2004.....	3.5	*	*	*	*	*	0.1	0.5	2.0	7.5	30.0	130.1
2003.....	3.8	*	*	*	*	*	0.2	0.7	2.2	7.9	34.0	140.3
2002.....	4.1	*	*	*	*	*	0.1	0.6	2.5	9.0	35.8	158.7
2001.....	4.1	*	*	*	*	*	0.1	0.6	2.4	9.2	36.1	168.2
2000.....	4.3	*	*	*	*	*	0.2	0.7	2.6	9.7	39.9	178.7
1999.....	4.4	*	*	*	*	*	0.2	0.7	2.7	9.7	40.3	189.9

Non-Hispanic 4/, female

2007.....	3.7	*	*	*	*	*	*	0.2	0.7	3.5	16.7	88.0
2006.....	3.9	*	*	*	*	*	*	0.2	0.9	3.9	18.3	93.9
2005.....	5.5	*	*	*	*	*	*	0.3	1.1	4.6	24.3	140.6
2004.....	5.5	*	*	*	*	*	*	0.2	1.2	5.2	24.4	144.0
2003.....	6.2	*	*	*	*	*	*	0.3	1.2	5.8	27.7	164.5
2002.....	6.6	*	*	*	*	*	*	0.3	1.4	5.9	29.8	175.1
2001.....	6.8	*	*	*	*	*	*	0.3	1.5	6.0	31.6	185.5
2000.....	7.0	*	*	*	*	*	*	0.3	1.3	6.3	31.8	195.1
1999.....	7.4	*	*	*	*	*	*	0.3	1.5	6.9	33.4	209.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

Non-Hispanic white, both sexes

2007 5/.....	3.5	*	*	*	*	*	*	0.3	1.1	4.4	18.2	87.9
2006 5/.....	3.8	*	*	*	*	*	*	0.3	1.2	4.6	19.6	95.2
2005 5/.....	5.2	*	*	*	*	*	0.1	0.4	1.5	5.4	27.1	141.3
2004 5/.....	5.3	*	*	*	*	*	0.1	0.4	1.6	6.2	27.6	145.0
2003 5/.....	5.8	*	*	*	*	*	0.1	0.5	1.6	6.6	30.8	162.2
2002.....	6.1	*	*	*	*	*	0.1	0.4	1.8	7.3	32.5	177.0
2001.....	6.3	*	*	*	*	*	0.1	0.5	1.8	7.5	34.1	187.4
2000.....	6.5	*	*	*	*	*	0.1	0.4	1.8	7.8	35.5	196.8
1999.....	6.7	*	*	*	*	*	0.1	0.5	2.0	8.2	36.7	209.9

Non-Hispanic white, male

2007 5/.....	2.8	*	*	*	*	*	*	0.4	1.5	5.4	20.1	81.3
2006 5/.....	2.9	*	*	*	*	*	*	0.5	1.6	5.3	20.9	90.3
2005 5/.....	4.0	*	*	*	*	*	*	0.7	1.9	6.5	30.3	130.6
2004 5/.....	4.1	*	*	*	*	*	*	0.6	2.0	7.6	30.8	136.8
2003 5/.....	4.3	*	*	*	*	*	0.2	0.7	2.2	7.7	34.1	145.2
2002.....	4.6	*	*	*	*	*	0.1	0.5	2.3	9.0	35.9	165.7
2001.....	4.7	*	*	*	*	*	0.1	0.6	2.3	9.2	36.6	175.2
2000.....	4.9	*	*	*	*	*	0.2	0.6	2.6	9.8	40.6	183.6
1999.....	4.9	*	*	*	*	*	0.1	0.7	2.6	9.6	40.4	194.6

Non-Hispanic white, female

2007 5/.....	4.3	*	*	*	*	*	*	0.2	0.7	3.5	16.9	91.0
2006 5/.....	4.6	*	*	*	*	*	*	0.2	0.9	3.9	18.7	97.6
2005 5/.....	6.4	*	*	*	*	*	*	0.2	1.1	4.5	24.9	146.2
2004 5/.....	6.4	*	*	*	*	*	*	0.2	1.2	5.1	25.4	148.7
2003 5/.....	7.2	*	*	*	*	*	*	0.3	1.1	5.8	28.5	169.6
2002.....	7.6	*	*	*	*	*	*	0.3	1.3	5.9	30.2	181.9
2001.....	7.9	*	*	*	*	*	*	0.3	1.3	6.1	32.5	192.6
2000.....	8.0	*	*	*	*	*	*	0.2	1.1	6.2	32.1	202.1
1999.....	8.5	*	*	*	*	*	*	0.3	1.4	6.9	34.3	215.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Atherosclerosis (I70)

##### Non-Hispanic black, both sexes

2007 5/.....	1.7	*	*	*	*	*	*	0.6	1.7	5.5	17.6	70.8
2006 5/.....	1.9	*	*	*	*	*	*	0.6	2.1	6.6	20.9	72.7
2005 5/.....	2.2	*	*	*	*	*	*	0.8	2.1	7.6	24.0	99.7
2004 5/.....	2.2	*	*	*	*	*	*	0.5	2.2	7.5	24.3	103.6
2003 5/.....	2.7	*	*	*	*	*	*	0.8	2.7	9.1	30.8	122.9
2002.....	2.8	*	*	*	*	*	*	1.0	3.0	8.1	37.0	118.0
2001.....	2.7	*	*	*	*	*	*	0.7	3.4	8.9	30.7	126.1
2000.....	3.0	*	*	*	*	*	*	1.1	3.3	9.1	35.6	140.2
1999.....	3.1	*	*	*	*	*	*	0.7	3.3	9.3	35.0	155.8

##### Non-Hispanic black, male

2007 5/.....	1.4	*	*	*	*	*	*	0.9	2.9	7.3	17.3	64.6
2006 5/.....	1.7	*	*	*	*	*	*	*	3.1	8.9	26.1	76.0
2005 5/.....	1.7	*	*	*	*	*	*	0.9	2.8	9.8	26.6	77.1
2004 5/.....	1.7	*	*	*	*	*	*	*	3.0	8.6	30.2	78.2
2003 5/.....	2.3	*	*	*	*	*	*	1.2	3.7	12.0	40.4	105.5
2002.....	2.4	*	*	*	*	*	*	1.4	4.3	10.2	44.8	109.4
2001.....	2.2	*	*	*	*	*	*	1.0	3.9	11.0	37.2	113.2
2000.....	2.5	*	*	*	*	*	*	1.5	4.1	11.1	40.4	150.3
1999.....	2.6	*	*	*	*	*	*	*	4.5	12.0	44.4	166.4

##### Non-Hispanic black, female

2007 5/.....	1.9	*	*	*	*	*	*	*	*	4.3	17.8	73.4
2006 5/.....	2.0	*	*	*	*	*	*	*	1.4	4.9	18.0	71.4
2005 5/.....	2.7	*	*	*	*	*	*	*	1.5	6.0	22.6	108.8
2004 5/.....	2.7	*	*	*	*	*	*	*	1.7	6.7	20.9	113.6
2003 5/.....	3.0	*	*	*	*	*	*	*	1.9	7.1	25.4	129.6
2002.....	3.2	*	*	*	*	*	*	*	2.0	6.6	32.5	121.2
2001.....	3.2	*	*	*	*	*	*	*	3.0	7.4	27.1	130.9
2000.....	3.5	*	*	*	*	*	*	*	2.7	7.7	32.8	136.5
1999.....	3.5	*	*	*	*	*	*	*	2.3	7.4	29.8	151.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

All origins 3/, both sexes

2007.....	7.3	*	*	*	0.2	0.3	1.1	2.5	7.0	21.0	54.3	121.4
2006.....	7.5	*	*	*	0.2	0.4	1.0	2.7	6.8	21.7	58.0	128.5
2005.....	8.0	*	*	*	0.1	0.4	1.1	2.6	7.5	24.1	62.6	135.5
2004.....	8.1	*	*	*	0.1	0.4	1.0	2.7	7.2	25.1	64.3	139.3
2003.....	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
2002.....	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001.....	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000.....	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999.....	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2

All origins 3/, male

2007.....	7.7	*	*	*	0.2	0.4	1.6	3.5	10.0	27.8	68.0	142.8
2006.....	8.0	*	*	*	0.2	0.5	1.5	3.8	9.6	29.0	74.5	155.0
2005.....	8.5	*	*	*	0.2	0.5	1.5	3.6	10.5	32.8	81.8	158.2
2004.....	8.5	*	*	*	0.2	0.6	1.4	3.8	9.9	33.4	82.8	166.5
2003.....	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
2002.....	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001.....	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000.....	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999.....	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5

All origins 3/, female

2007.....	6.8	*	*	*	*	0.2	0.6	1.5	4.2	15.2	44.9	111.3
2006.....	7.0	*	*	*	0.1	0.2	0.6	1.6	4.2	15.5	46.7	116.1
2005.....	7.4	*	*	*	*	0.3	0.7	1.6	4.7	16.7	49.5	125.0
2004.....	7.6	*	*	*	*	0.3	0.7	1.6	4.7	18.1	51.8	127.0
2003.....	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
2002.....	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001.....	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000.....	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999.....	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Hispanic, both sexes

2007.....	1.9	*	*	*	*	0.3	0.7	1.3	3.8	11.7	31.6	71.5
2006.....	2.0	*	*	*	*	0.4	0.8	1.7	4.2	12.8	31.9	71.6
2005.....	2.1	*	*	*	*	*	0.8	1.8	4.1	13.3	41.7	74.4
2004.....	2.0	*	*	*	*	0.3	0.7	2.0	5.2	13.9	34.5	77.1
2003.....	2.2	*	*	*	*	0.3	0.7	1.6	4.6	14.1	42.3	106.2
2002.....	2.2	*	*	*	*	0.3	0.8	1.7	5.9	14.8	41.0	92.6
2001.....	2.0	*	*	*	*	*	0.7	2.1	4.8	13.7	39.0	95.0
2000.....	2.3	*	*	*	*	0.4	0.7	1.8	5.8	20.2	42.5	102.8
1999.....	2.1	*	*	*	*	*	0.7	1.3	5.7	16.4	44.0	110.8

Hispanic, male

2007.....	2.1	*	*	*	*	*	1.1	1.8	5.3	15.2	39.3	76.2
2006.....	2.3	*	*	*	*	0.5	1.1	2.1	6.0	17.2	45.2	86.8
2005.....	2.5	*	*	*	*	*	1.0	2.6	6.5	18.8	57.5	105.9
2004.....	2.3	*	*	*	*	*	0.9	3.0	7.3	20.5	44.4	89.7
2003.....	2.5	*	*	*	*	*	0.9	2.2	7.4	19.0	58.7	138.5
2002.....	2.3	*	*	*	*	*	1.0	2.2	7.1	19.9	56.6	102.1
2001.....	2.2	*	*	*	*	*	0.9	2.7	6.3	20.0	53.2	108.6
2000.....	2.5	*	*	*	*	*	0.8	2.1	8.0	25.7	62.0	131.0
1999.....	2.4	*	*	*	*	*	0.9	1.7	7.7	22.2	62.0	163.9

Hispanic, female

2007.....	1.8	*	*	*	*	*	*	0.9	2.4	8.9	26.1	69.0
2006.....	1.8	*	*	*	*	*	*	1.3	2.6	9.2	22.6	63.5
2005.....	1.7	*	*	*	*	*	*	0.9	1.9	8.9	30.7	58.0
2004.....	1.8	*	*	*	*	*	*	1.0	3.3	8.7	27.7	70.6
2003.....	1.9	*	*	*	*	*	*	*	2.1	10.1	31.1	89.8
2002.....	2.0	*	*	*	*	*	*	1.1	4.9	10.8	30.4	87.9
2001.....	1.8	*	*	*	*	*	*	1.6	3.4	8.8	29.3	88.4
2000.....	2.1	*	*	*	*	*	*	1.5	3.9	15.8	29.4	89.0
1999.....	1.9	*	*	*	*	*	*	*	4.0	11.9	31.9	85.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic 4/, both sexes

2007.....	8.2	*	*	*	0.2	0.3	1.2	2.6	7.3	21.7	55.7	124.0
2006.....	8.4	*	*	*	0.2	0.4	1.0	2.8	7.0	22.3	59.5	131.2
2005.....	8.9	*	*	*	0.1	0.4	1.1	2.6	7.7	24.8	63.7	138.2
2004.....	9.0	*	*	*	0.1	0.5	1.1	2.8	7.4	25.9	65.8	141.8
2003.....	9.5	*	*	*	0.2	0.4	1.1	3.1	8.4	27.7	68.7	147.5
2002.....	9.6	*	*	*	0.2	0.4	1.0	2.8	8.8	30.0	69.8	150.3
2001.....	9.9	*	*	*	0.2	0.5	1.2	2.6	8.9	30.1	73.2	160.1
2000.....	10.2	*	*	*	0.2	0.4	1.0	2.8	9.4	32.9	75.8	165.0
1999.....	10.2	*	*	*	0.1	0.4	1.1	2.8	9.5	32.3	76.9	166.8

Non-Hispanic 4/, male

2007.....	8.8	*	*	*	0.2	0.5	1.7	3.8	10.4	28.8	69.8	146.6
2006.....	9.1	*	*	*	0.2	0.5	1.5	4.0	9.9	29.8	76.2	158.5
2005.....	9.5	*	*	*	0.2	0.5	1.6	3.7	10.8	33.7	83.0	160.6
2004.....	9.6	*	*	*	0.2	0.6	1.5	3.9	10.1	34.3	84.9	170.1
2003.....	10.4	*	*	*	0.2	0.5	1.5	4.3	11.6	38.2	93.4	181.5
2002.....	10.5	*	*	*	0.2	0.5	1.3	3.9	12.3	41.1	93.9	190.7
2001.....	10.7	*	*	*	0.2	0.6	1.6	3.6	12.3	41.5	98.8	200.5
2000.....	11.3	*	*	*	0.2	0.5	1.3	3.9	13.5	46.5	104.4	217.9
1999.....	11.2	*	*	*	0.2	0.6	1.4	3.8	13.7	45.5	105.8	209.0

Non-Hispanic 4/, female

2007.....	7.7	*	*	*	*	0.2	0.7	1.5	4.4	15.7	46.0	113.4
2006.....	7.9	*	*	*	*	0.2	0.6	1.6	4.4	16.0	48.2	118.5
2005.....	8.3	*	*	*	*	0.3	0.7	1.6	4.9	17.3	50.5	128.0
2004.....	8.5	*	*	*	*	0.3	0.7	1.7	4.9	18.8	53.1	129.2
2003.....	8.6	*	*	*	*	0.3	0.7	1.9	5.5	19.0	52.3	132.6
2002.....	8.8	*	*	*	*	0.3	0.7	1.7	5.5	20.7	53.9	132.8
2001.....	9.1	*	*	*	*	0.3	0.8	1.7	5.8	20.6	56.3	143.3
2000.....	9.2	*	*	*	0.1	0.2	0.8	1.8	5.7	21.7	57.1	143.6
1999.....	9.3	*	*	*	*	0.2	0.8	1.8	5.7	21.5	58.2	149.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic white, both sexes

2007 5/.....	9.0	*	*	*	0.1	0.3	1.0	2.2	6.8	21.7	56.2	125.7
2006 5/.....	9.3	*	*	*	0.2	0.3	0.9	2.4	6.7	22.4	59.8	132.6
2005 5/.....	9.8	*	*	*	0.1	0.3	0.9	2.2	7.4	24.9	64.0	138.9
2004 5/.....	9.9	*	*	*	0.2	0.4	0.9	2.4	6.9	26.2	66.3	141.9
2003 5/.....	10.4	*	*	*	0.2	0.3	0.9	2.6	8.2	28.2	69.5	148.3
2002.....	10.6	*	*	*	0.1	0.3	0.9	2.3	8.4	30.4	70.9	150.9
2001.....	10.8	*	*	*	0.2	0.4	1.0	2.3	8.5	30.5	73.5	160.1
2000.....	11.2	*	*	*	0.1	0.3	0.8	2.4	9.0	33.5	76.3	164.8
1999.....	11.3	*	*	*	0.1	0.4	1.0	2.5	9.4	32.9	77.7	166.7

Non-Hispanic white, male

2007 5/.....	9.6	*	*	*	0.2	0.4	1.4	3.2	10.0	28.9	70.7	150.7
2006 5/.....	10.0	*	*	*	0.2	0.5	1.3	3.5	9.5	30.1	77.8	163.3
2005 5/.....	10.4	*	*	*	0.2	0.4	1.2	3.1	10.4	34.2	83.7	164.5
2004 5/.....	10.5	*	*	*	0.2	0.5	1.3	3.5	9.4	35.0	86.4	173.0
2003 5/.....	11.5	*	*	*	0.2	0.4	1.2	3.8	11.3	39.2	94.7	185.8
2002.....	11.6	*	*	*	0.2	0.4	1.2	3.3	11.8	42.0	96.3	194.6
2001.....	11.8	*	*	*	0.2	0.6	1.3	3.1	12.0	42.4	100.3	204.7
2000.....	12.6	*	*	*	0.2	0.5	1.0	3.5	13.2	47.9	106.3	222.7
1999.....	12.5	*	*	*	0.2	0.5	1.3	3.4	13.8	47.0	107.9	212.8

Non-Hispanic white, female

2007 5/.....	8.4	*	*	*	*	0.2	0.6	1.2	3.9	15.4	46.1	113.9
2006 5/.....	8.6	*	*	*	*	0.2	0.5	1.3	4.0	15.7	47.3	118.3
2005 5/.....	9.1	*	*	*	*	0.3	0.6	1.4	4.6	16.9	50.4	127.2
2004 5/.....	9.3	*	*	*	*	0.2	0.6	1.4	4.5	18.6	52.6	128.0
2003 5/.....	9.4	*	*	*	*	0.2	0.5	1.5	5.2	18.8	52.4	131.8
2002.....	9.6	*	*	*	*	0.3	0.6	1.4	5.2	20.5	53.9	132.1
2001.....	9.9	*	*	*	*	0.3	0.6	1.5	5.3	20.4	55.7	141.6
2000.....	9.9	*	*	*	*	0.2	0.6	1.4	5.2	21.3	56.6	141.5
1999.....	10.1	*	*	*	*	0.2	0.7	1.5	5.3	21.1	57.8	148.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic black, both sexes

2007 5/.....	6.4	*	*	*	*	0.5	2.6	5.7	12.0	26.4	59.8	127.8
2006 5/.....	6.5	*	*	*	*	0.7	2.1	5.8	11.7	27.0	65.8	134.0
2005 5/.....	6.9	*	*	*	*	0.9	2.4	5.5	12.3	29.9	70.5	147.5
2004 5/.....	6.8	*	*	*	*	0.9	2.3	5.3	12.9	28.6	71.2	156.9
2003 5/.....	6.8	*	*	*	*	0.7	2.4	6.4	12.1	29.6	71.1	152.9
2002.....	6.9	*	*	*	*	0.9	2.0	6.0	13.6	31.7	69.1	157.6
2001.....	7.3	*	*	*	*	0.8	2.5	5.2	14.1	32.3	80.5	175.7
2000.....	7.3	*	*	*	*	0.7	2.2	5.7	14.5	33.4	78.9	180.0
1999.....	7.0	*	*	*	*	0.6	2.0	5.4	12.5	32.1	77.3	180.1

Non-Hispanic black, male

2007 5/.....	6.7	*	*	*	*	*	3.8	8.4	16.4	33.6	71.8	128.3
2006 5/.....	6.5	*	*	*	*	1.0	3.2	8.0	16.6	35.4	69.0	119.1
2005 5/.....	7.1	*	*	*	*	1.1	3.7	8.0	16.9	37.0	89.1	132.3
2004 5/.....	6.9	*	*	*	*	1.2	3.1	7.5	18.3	36.1	81.0	161.6
2003 5/.....	6.9	*	*	*	*	1.2	3.1	8.4	16.7	36.4	91.5	141.7
2002.....	7.0	*	*	*	*	1.2	2.7	8.2	19.5	39.2	78.8	168.5
2001.....	7.2	*	*	*	*	1.0	3.5	7.4	17.9	40.3	97.2	168.6
2000.....	7.2	*	*	*	*	0.9	2.8	7.6	18.7	41.8	96.9	179.7
1999.....	6.7	*	*	*	*	0.9	2.4	7.4	16.0	40.0	90.7	184.2

Non-Hispanic black, female

2007 5/.....	6.1	*	*	*	*	*	1.5	3.5	8.6	21.2	53.0	127.7
2006 5/.....	6.5	*	*	*	*	*	1.2	3.9	7.7	20.9	64.1	140.1
2005 5/.....	6.7	*	*	*	*	*	1.2	3.4	8.5	24.9	59.9	153.6
2004 5/.....	6.8	*	*	*	*	*	1.6	3.5	8.6	23.3	65.7	155.0
2003 5/.....	6.8	*	*	*	*	*	1.7	4.6	8.4	24.7	59.6	157.2
2002.....	6.9	*	*	*	*	*	1.4	4.2	8.9	26.3	63.7	153.4
2001.....	7.4	*	*	*	*	*	1.6	3.3	11.0	26.6	71.1	178.4
2000.....	7.5	*	*	*	*	*	1.6	4.0	11.2	27.5	68.8	180.1
1999.....	7.2	*	*	*	*	*	1.7	3.6	9.7	26.6	69.9	178.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Aortic aneurysm and dissection (I71)

All origins 3/, both sexes

2007.....	4.3	*	*	*	0.1	0.2	0.8	1.7	4.5	13.5	33.4	59.5
2006.....	4.4	*	*	*	0.1	0.3	0.8	1.8	4.3	13.9	35.4	62.7
2005.....	4.7	*	*	*	0.1	0.3	0.8	1.8	4.8	15.4	38.1	64.9
2004.....	4.7	*	*	*	0.1	0.3	0.7	1.8	4.5	15.9	39.6	64.8
2003.....	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002.....	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001.....	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000.....	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999.....	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3

All origins 3/, male

2007.....	5.1	*	*	*	0.1	0.4	1.3	2.6	6.9	19.1	44.1	81.8
2006.....	5.2	*	*	*	0.1	0.4	1.2	2.8	6.6	19.5	48.4	87.6
2005.....	5.6	*	*	*	0.1	0.4	1.2	2.8	7.3	22.4	52.6	91.0
2004.....	5.6	*	*	*	0.2	0.5	1.1	2.7	6.9	23.1	54.8	95.0
2003.....	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002.....	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001.....	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3

All origins 3/, female

2007.....	3.6	*	*	*	*	0.1	0.4	0.8	2.3	8.8	26.0	48.9
2006.....	3.6	*	*	*	*	0.1	0.3	0.9	2.2	9.3	26.5	51.0
2005.....	3.8	*	*	*	*	0.2	0.3	0.8	2.5	9.5	28.2	52.9
2004.....	3.8	*	*	*	*	0.2	0.3	0.8	2.3	9.8	29.3	51.2
2003.....	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002.....	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001.....	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000.....	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999.....	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Aortic aneurysm and dissection (I71)

##### Hispanic, both sexes

2007.....	1.0	*	*	*	*	0.2	0.4	0.9	2.0	6.4	16.2	34.3
2006.....	1.1	*	*	*	*	*	0.6	1.0	2.4	7.1	16.3	31.7
2005.....	1.1	*	*	*	*	*	0.5	1.1	2.6	7.3	19.9	29.1
2004.....	1.1	*	*	*	*	*	0.4	1.3	3.3	7.9	18.0	30.4
2003.....	1.3	*	*	*	*	*	0.4	0.9	3.1	8.3	26.7	49.2
2002.....	1.1	*	*	*	*	*	0.4	1.0	3.8	7.5	21.2	31.0
2001.....	1.1	*	*	*	*	*	0.4	1.2	2.5	8.2	22.9	31.9
2000.....	1.3	*	*	*	*	*	0.4	1.3	3.4	11.2	27.1	41.8
1999.....	1.1	*	*	*	*	*	0.5	0.9	3.4	9.2	24.0	40.7

##### Hispanic, male

2007.....	1.3	*	*	*	*	*	0.7	1.3	3.0	8.9	22.7	45.7
2006.....	1.4	*	*	*	*	*	1.0	1.5	3.9	10.9	24.0	47.4
2005.....	1.5	*	*	*	*	*	0.8	1.9	4.8	10.7	30.7	43.6
2004.....	1.5	*	*	*	*	*	0.7	2.2	5.5	12.6	25.4	40.6
2003.....	1.6	*	*	*	*	*	*	1.4	5.1	12.5	38.3	80.0
2002.....	1.4	*	*	*	*	*	0.7	1.6	5.5	12.0	32.0	43.5
2001.....	1.5	*	*	*	*	*	0.7	1.8	3.8	14.2	34.9	53.3
2000.....	1.6	*	*	*	*	*	*	1.8	5.2	16.0	42.3	64.5
1999.....	1.5	*	*	*	*	*	*	1.5	5.3	14.7	33.9	79.8

##### Hispanic, female

2007.....	0.8	*	*	*	*	*	*	*	*	4.4	11.7	28.1
2006.....	0.7	*	*	*	*	*	*	*	*	4.1	10.8	23.4
2005.....	0.7	*	*	*	*	*	*	*	*	4.5	12.3	21.5
2004.....	0.8	*	*	*	*	*	*	*	*	4.1	12.8	25.2
2003.....	0.9	*	*	*	*	*	*	*	*	5.0	18.7	33.6
2002.....	0.8	*	*	*	*	*	*	*	2.3	4.0	13.8	24.9
2001.....	0.7	*	*	*	*	*	*	*	*	3.5	14.8	21.4
2000.....	1.0	*	*	*	*	*	*	*	*	7.5	16.8	30.7
1999.....	0.8	*	*	*	*	*	*	*	*	4.8	17.3	21.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Aortic aneurysm and dissection (I71)

##### Non-Hispanic 4/, both sexes

2007.....	4.9	*	*	*	0.1	0.2	0.9	1.8	4.8	14.1	34.5	60.8
2006.....	5.0	*	*	*	0.1	0.3	0.8	1.9	4.5	14.4	36.6	64.2
2005.....	5.3	*	*	*	0.1	0.3	0.8	1.9	5.0	16.0	39.1	66.5
2004.....	5.3	*	*	*	0.1	0.3	0.8	1.8	4.6	16.4	40.7	66.2
2003.....	5.7	*	*	*	0.1	0.3	0.8	2.1	5.5	18.5	43.2	70.7
2002.....	5.7	*	*	*	0.1	0.3	0.7	1.8	5.6	19.8	44.3	70.8
2001.....	6.0	*	*	*	0.1	0.3	0.8	1.7	5.7	19.8	46.7	79.0
2000.....	6.2	*	*	*	0.1	0.3	0.7	1.8	6.3	22.4	47.9	80.3
1999.....	6.3	*	*	*	0.1	0.3	0.8	1.8	6.2	22.4	49.5	81.7

##### Non-Hispanic 4/, male

2007.....	5.8	*	*	*	0.1	0.4	1.4	2.8	7.3	19.9	45.4	83.9
2006.....	5.9	*	*	*	0.2	0.4	1.3	3.0	6.8	20.1	49.9	89.8
2005.....	6.3	*	*	*	0.1	0.4	1.3	2.9	7.4	23.2	53.8	93.3
2004.....	6.3	*	*	*	0.2	0.5	1.2	2.7	7.0	23.8	56.4	97.5
2003.....	7.0	*	*	*	0.2	0.4	1.2	3.2	8.5	26.9	63.0	106.6
2002.....	7.2	*	*	*	0.2	0.4	1.1	2.9	8.7	29.6	64.1	109.1
2001.....	7.3	*	*	*	0.2	0.5	1.2	2.7	8.8	29.5	67.7	117.0
2000.....	7.8	*	*	*	*	0.4	1.0	2.7	9.8	33.8	71.6	129.6
1999.....	7.8	*	*	*	*	0.4	1.1	2.8	9.8	33.6	74.3	127.0

##### Non-Hispanic 4/, female

2007.....	4.0	*	*	*	*	0.1	0.4	0.9	2.4	9.1	26.9	49.9
2006.....	4.1	*	*	*	*	*	0.4	0.9	2.3	9.6	27.5	52.3
2005.....	4.3	*	*	*	*	0.2	0.3	0.9	2.7	9.9	29.1	54.2
2004.....	4.2	*	*	*	*	0.2	0.4	0.9	2.4	10.2	30.2	52.3
2003.....	4.4	*	*	*	*	0.2	0.3	1.0	2.7	11.4	30.0	55.0
2002.....	4.4	*	*	*	*	0.1	0.4	0.9	2.7	11.6	31.2	54.3
2001.....	4.7	*	*	*	*	0.2	0.3	0.8	2.9	11.8	32.9	63.2
2000.....	4.7	*	*	*	*	0.1	0.4	0.9	3.1	13.0	32.5	60.4
1999.....	4.8	*	*	*	*	*	0.4	0.9	2.9	13.2	33.4	63.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Aortic aneurysm and dissection (I71)

Non-Hispanic white, both sexes

2007 5/.....	5.4	*	*	*	0.1	0.2	0.7	1.5	4.5	14.5	35.6	63.1
2006 5/.....	5.5	*	*	*	0.1	0.2	0.7	1.6	4.3	14.7	37.7	66.0
2005 5/.....	5.8	*	*	*	0.1	0.3	0.6	1.5	4.9	16.5	40.2	67.9
2004 5/.....	5.9	*	*	*	0.1	0.3	0.6	1.6	4.3	17.2	41.9	68.4
2003 5/.....	6.4	*	*	*	0.1	0.2	0.6	1.8	5.4	19.3	44.8	73.2
2002.....	6.5	*	*	*	0.1	0.2	0.6	1.5	5.5	20.7	46.0	72.8
2001.....	6.7	*	*	*	0.1	0.3	0.6	1.5	5.6	20.7	47.9	81.3
2000.....	7.0	*	*	*	*	0.2	0.6	1.6	6.3	23.4	49.3	82.0
1999.....	7.1	*	*	*	0.1	0.2	0.7	1.6	6.3	23.4	51.1	83.7

Non-Hispanic white, male

2007 5/.....	6.4	*	*	*	*	0.3	1.1	2.3	7.1	20.6	47.1	88.2
2006 5/.....	6.6	*	*	*	0.2	0.4	1.1	2.6	6.7	20.6	52.0	92.8
2005 5/.....	7.0	*	*	*	*	0.4	1.0	2.4	7.4	24.2	55.6	96.7
2004 5/.....	7.0	*	*	*	0.2	0.4	1.0	2.4	6.6	24.8	58.3	101.4
2003 5/.....	7.9	*	*	*	0.2	0.3	1.0	2.8	8.4	28.3	65.5	111.2
2002.....	8.0	*	*	*	*	0.3	0.9	2.4	8.6	31.0	67.1	114.7
2001.....	8.2	*	*	*	0.2	0.5	1.0	2.3	8.7	30.9	70.0	122.9
2000.....	8.8	*	*	*	*	0.3	0.8	2.5	9.8	35.6	74.2	135.3
1999.....	8.9	*	*	*	*	0.4	1.0	2.4	10.1	35.4	77.5	131.8

Non-Hispanic white, female

2007 5/.....	4.5	*	*	*	*	*	0.4	0.7	2.1	9.1	27.6	51.2
2006 5/.....	4.5	*	*	*	*	*	0.3	0.7	2.1	9.6	27.8	53.4
2005 5/.....	4.7	*	*	*	*	0.2	0.3	0.7	2.5	9.9	29.7	54.8
2004 5/.....	4.7	*	*	*	*	*	0.3	0.7	2.2	10.6	30.7	53.6
2003 5/.....	4.9	*	*	*	*	*	0.2	0.8	2.5	11.6	30.9	56.5
2002.....	4.9	*	*	*	*	*	0.3	0.7	2.6	11.9	31.9	54.7
2001.....	5.2	*	*	*	*	*	0.3	0.6	2.7	12.0	33.3	64.0
2000.....	5.2	*	*	*	*	*	0.3	0.7	2.9	13.1	32.9	60.5
1999.....	5.3	*	*	*	*	*	0.4	0.7	2.7	13.3	33.9	64.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Aortic aneurysm and dissection (I71)

##### Non-Hispanic black, both sexes

2007 5/.....	3.3	*	*	*	*	*	2.2	4.0	7.0	13.2	26.6	40.9
2006 5/.....	3.3	*	*	*	*	0.5	1.5	4.1	6.7	14.6	28.3	47.8
2005 5/.....	3.4	*	*	*	*	0.5	1.8	4.1	7.0	14.8	31.1	51.3
2004 5/.....	3.3	*	*	*	*	0.6	1.7	3.6	7.6	13.3	32.4	44.6
2003 5/.....	3.3	*	*	*	*	0.6	1.7	4.4	7.3	14.5	30.4	47.3
2002.....	3.3	*	*	*	*	0.6	1.4	4.3	7.5	14.8	30.2	50.5
2001.....	3.5	*	*	*	*	0.4	1.6	3.6	7.7	14.7	38.0	55.9
2000.....	3.6	*	*	*	*	0.5	1.5	3.3	8.2	17.3	36.8	63.0
1999.....	3.4	*	*	*	*	0.4	1.5	3.9	6.5	16.8	34.0	61.1

##### Non-Hispanic black, male

2007 5/.....	3.8	*	*	*	*	*	3.4	6.3	9.9	17.1	31.9	42.8
2006 5/.....	3.7	*	*	*	*	0.9	2.4	5.9	9.8	19.3	30.3	52.2
2005 5/.....	4.0	*	*	*	*	0.8	3.2	6.2	10.1	18.4	39.2	52.3
2004 5/.....	3.9	*	*	*	*	1.0	2.6	5.5	11.8	18.5	37.4	57.9
2003 5/.....	3.9	*	*	*	*	1.1	2.5	6.5	11.0	17.9	39.8	52.2
2002.....	3.9	*	*	*	*	1.0	2.3	6.6	12.1	19.3	34.9	49.2
2001.....	3.9	*	*	*	*	*	2.6	5.4	11.3	19.9	47.4	49.7
2000.....	4.0	*	*	*	*	*	2.3	4.7	11.6	22.6	47.2	70.5
1999.....	3.7	*	*	*	*	*	2.0	6.0	9.3	22.1	38.8	77.3

##### Non-Hispanic black, female

2007 5/.....	2.7	*	*	*	*	*	1.1	2.1	4.6	10.5	23.6	40.1
2006 5/.....	2.9	*	*	*	*	*	0.8	2.4	4.3	11.2	27.1	46.0
2005 5/.....	3.0	*	*	*	*	*	*	2.3	4.4	12.2	26.5	50.9
2004 5/.....	2.7	*	*	*	*	*	0.9	2.0	4.3	9.5	29.6	39.4
2003 5/.....	2.8	*	*	*	*	*	1.0	2.5	4.3	12.1	25.1	45.4
2002.....	2.8	*	*	*	*	*	*	2.2	3.8	11.6	27.5	51.0
2001.....	3.1	*	*	*	*	*	0.7	2.0	4.8	11.1	32.7	58.2
2000.....	3.3	*	*	*	*	*	0.9	2.1	5.5	13.5	30.9	60.2
1999.....	3.1	*	*	*	*	*	1.0	2.0	4.3	13.1	31.3	55.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes

2007.....	3.0	*	*	*	0.1	0.1	0.3	0.8	2.5	7.5	20.9	62.0
2006.....	3.1	*	*	*	0.1	0.1	0.2	0.8	2.5	7.8	22.6	65.8
2005.....	3.3	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.5	70.6
2004.....	3.4	*	*	*	*	0.1	0.3	1.0	2.7	9.2	24.7	74.5
2003.....	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002.....	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001.....	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000.....	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999.....	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9

All origins 3/, male

2007.....	2.6	*	*	*	*	*	0.3	0.9	3.1	8.7	23.9	61.0
2006.....	2.8	*	*	*	*	*	0.2	1.0	3.0	9.5	26.1	67.3
2005.....	2.9	*	*	*	*	*	0.3	0.8	3.2	10.4	29.2	67.2
2004.....	2.9	*	*	*	*	0.1	0.3	1.1	3.0	10.3	28.0	71.6
2003.....	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002.....	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001.....	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000.....	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999.....	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2

All origins 3/, female

2007.....	3.3	*	*	*	*	0.1	0.2	0.6	1.9	6.4	18.9	62.4
2006.....	3.4	*	*	*	*	0.1	0.2	0.7	2.0	6.3	20.2	65.0
2005.....	3.7	*	*	*	*	0.1	0.4	0.7	2.2	7.2	21.3	72.2
2004.....	3.9	*	*	*	*	0.1	0.4	0.8	2.4	8.3	22.5	75.8
2003.....	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002.....	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001.....	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000.....	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999.....	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Hispanic, both sexes

2007.....	0.9	*	*	*	*	*	*	0.4	1.8	5.3	15.3	37.2
2006.....	0.9	*	*	*	*	*	*	0.7	1.8	5.7	15.6	39.9
2005.....	1.0	*	*	*	*	*	*	0.6	1.4	6.1	21.8	45.3
2004.....	0.9	*	*	*	*	*	*	0.7	1.8	6.1	16.5	46.6
2003.....	0.9	*	*	*	*	*	*	0.7	1.5	5.7	15.6	57.0
2002.....	1.1	*	*	*	*	*	0.3	0.7	2.1	7.3	19.8	61.5
2001.....	0.9	*	*	*	*	*	*	0.9	2.2	5.5	16.0	63.2
2000.....	1.0	*	*	*	*	*	*	*	2.4	8.9	15.4	61.0
1999.....	1.0	*	*	*	*	*	*	*	2.4	7.2	20.0	70.2

Hispanic, male

2007.....	0.8	*	*	*	*	*	*	*	2.2	6.4	16.6	30.5
2006.....	0.9	*	*	*	*	*	*	*	2.1	6.3	21.2	39.5
2005.....	1.0	*	*	*	*	*	*	*	1.8	8.1	26.7	62.3
2004.....	0.8	*	*	*	*	*	*	*	*	8.0	19.0	49.0
2003.....	0.9	*	*	*	*	*	*	*	2.2	6.5	20.3	58.5
2002.....	0.9	*	*	*	*	*	*	*	*	7.8	24.6	58.6
2001.....	0.8	*	*	*	*	*	*	*	2.5	5.8	18.3	55.2
2000.....	0.9	*	*	*	*	*	*	*	2.7	9.7	19.7	66.5
1999.....	0.9	*	*	*	*	*	*	*	*	7.5	28.2	84.1

Hispanic, female

2007.....	1.0	*	*	*	*	*	*	*	1.4	4.4	14.4	40.8
2006.....	1.0	*	*	*	*	*	*	*	*	5.1	11.7	40.1
2005.....	1.0	*	*	*	*	*	*	*	*	4.4	18.4	36.5
2004.....	1.0	*	*	*	*	*	*	*	1.9	4.5	14.8	45.4
2003.....	1.0	*	*	*	*	*	*	*	*	5.1	12.4	56.3
2002.....	1.3	*	*	*	*	*	*	*	2.6	6.8	16.6	63.0
2001.....	1.1	*	*	*	*	*	*	*	*	5.4	14.5	67.0
2000.....	1.1	*	*	*	*	*	*	*	*	8.3	12.5	58.4
1999.....	1.1	*	*	*	*	*	*	*	2.4	7.1	14.6	63.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 4/, both sexes

2007.....	3.3	*	*	*	0.1	0.1	0.3	0.8	2.5	7.6	21.3	63.2
2006.....	3.5	*	*	*	*	0.1	0.2	0.9	2.6	7.9	23.0	67.0
2005.....	3.7	*	*	*	*	0.1	0.3	0.8	2.8	8.8	24.6	71.7
2004.....	3.8	*	*	*	*	0.1	0.3	1.0	2.8	9.4	25.1	75.6
2003.....	3.8	*	*	*	0.1	0.1	0.3	1.0	2.9	9.3	25.5	76.7
2002.....	3.9	*	*	*	0.1	0.1	0.3	0.9	3.2	10.2	25.6	79.5
2001.....	3.9	*	*	*	*	0.1	0.4	0.9	3.2	10.3	26.5	81.1
2000.....	4.0	*	*	*	0.1	0.1	0.3	1.0	3.1	10.5	27.9	84.7
1999.....	3.9	*	*	*	*	0.1	0.3	1.0	3.3	9.9	27.5	85.1

Non-Hispanic 4/, male

2007.....	3.0	*	*	*	*	*	0.3	1.0	3.2	8.9	24.4	62.7
2006.....	3.1	*	*	*	*	*	0.2	1.0	3.1	9.7	26.3	68.7
2005.....	3.2	*	*	*	*	*	0.3	0.8	3.3	10.5	29.2	67.3
2004.....	3.2	*	*	*	*	*	0.3	1.1	3.1	10.5	28.5	72.6
2003.....	3.3	*	*	*	*	*	0.3	1.0	3.1	11.3	30.3	74.9
2002.....	3.4	*	*	*	*	*	0.3	1.0	3.5	11.5	29.8	81.7
2001.....	3.4	*	*	*	*	0.1	0.3	0.9	3.5	12.0	31.1	83.5
2000.....	3.5	*	*	*	*	0.1	0.2	1.2	3.6	12.7	32.8	88.2
1999.....	3.4	*	*	*	*	0.1	0.3	1.0	3.9	11.9	31.5	82.0

Non-Hispanic 4/, female

2007.....	3.7	*	*	*	*	*	0.3	0.7	2.0	6.6	19.1	63.5
2006.....	3.8	*	*	*	*	*	0.2	0.7	2.0	6.4	20.7	66.2
2005.....	4.1	*	*	*	*	0.1	0.4	0.7	2.3	7.4	21.4	73.7
2004.....	4.3	*	*	*	*	*	0.4	0.8	2.5	8.5	22.9	76.9
2003.....	4.2	*	*	*	*	*	0.4	0.9	2.8	7.6	22.3	77.6
2002.....	4.3	*	*	*	*	0.2	0.3	0.8	2.8	9.0	22.8	78.5
2001.....	4.4	*	*	*	*	0.1	0.4	0.9	2.9	8.8	23.5	80.1
2000.....	4.4	*	*	*	*	*	0.4	0.9	2.6	8.7	24.6	83.3
1999.....	4.5	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.8	86.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic white, both sexes

2007 5/.....	3.6	*	*	*	*	0.1	0.3	0.7	2.3	7.2	20.6	62.6
2006 5/.....	3.7	*	*	*	*	*	0.2	0.8	2.3	7.7	22.1	66.6
2005 5/.....	3.9	*	*	*	*	0.1	0.3	0.7	2.6	8.4	23.7	70.9
2004 5/.....	4.0	*	*	*	*	0.1	0.3	0.9	2.5	9.0	24.4	73.6
2003 5/.....	4.0	*	*	*	*	0.1	0.3	0.8	2.8	8.9	24.6	75.1
2002.....	4.1	*	*	*	*	0.1	0.3	0.8	2.9	9.7	24.9	78.2
2001.....	4.2	*	*	*	*	0.1	0.3	0.8	2.9	9.8	25.6	78.8
2000.....	4.3	*	*	*	*	0.1	0.3	0.9	2.8	10.1	27.1	82.8
1999.....	4.2	*	*	*	*	0.1	0.3	0.9	3.1	9.6	26.5	83.0

Non-Hispanic white, male

2007 5/.....	3.2	*	*	*	*	*	0.3	0.8	2.9	8.3	23.6	62.5
2006 5/.....	3.4	*	*	*	*	*	0.2	0.9	2.8	9.5	25.8	70.5
2005 5/.....	3.4	*	*	*	*	*	0.2	0.7	3.1	10.0	28.1	67.8
2004 5/.....	3.5	*	*	*	*	*	0.3	1.1	2.8	10.2	28.1	71.6
2003 5/.....	3.5	*	*	*	*	*	0.2	0.9	2.9	10.8	29.2	74.6
2002.....	3.6	*	*	*	*	*	0.2	0.9	3.2	11.0	29.3	79.9
2001.....	3.6	*	*	*	*	*	0.2	0.8	3.3	11.5	30.3	81.8
2000.....	3.8	*	*	*	*	*	0.2	1.0	3.3	12.3	32.1	87.4
1999.....	3.6	*	*	*	*	*	0.3	1.0	3.6	11.6	30.4	81.0

Non-Hispanic white, female

2007 5/.....	3.9	*	*	*	*	*	0.2	0.6	1.8	6.3	18.5	62.6
2006 5/.....	4.0	*	*	*	*	*	0.2	0.6	1.9	6.1	19.5	64.9
2005 5/.....	4.4	*	*	*	*	*	0.4	0.7	2.1	7.0	20.8	72.4
2004 5/.....	4.6	*	*	*	*	*	0.3	0.7	2.3	8.0	21.9	74.4
2003 5/.....	4.5	*	*	*	*	*	0.3	0.7	2.7	7.3	21.6	75.4
2002.....	4.7	*	*	*	*	*	0.3	0.7	2.6	8.6	21.9	77.4
2001.....	4.7	*	*	*	*	*	0.4	0.9	2.6	8.3	22.4	77.6
2000.....	4.7	*	*	*	*	*	0.3	0.7	2.2	8.2	23.8	81.0
1999.....	4.8	*	*	*	*	*	0.4	0.8	2.5	7.9	24.0	83.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic black, both sexes

2007 5/.....	3.1	*	*	*	*	*	0.4	1.7	5.1	13.2	33.2	86.9
2006 5/.....	3.2	*	*	*	*	*	0.6	1.7	5.0	12.4	37.5	86.3
2005 5/.....	3.4	*	*	*	*	*	0.6	1.4	5.3	15.1	39.3	96.1
2004 5/.....	3.5	*	*	*	*	*	0.6	1.7	5.3	15.3	38.8	112.2
2003 5/.....	3.5	*	*	*	*	*	0.7	2.0	4.8	15.0	40.7	105.6
2002.....	3.6	*	*	*	*	*	0.6	1.8	6.2	16.8	39.0	107.1
2001.....	3.8	*	*	*	*	*	0.9	1.7	6.4	17.5	42.5	119.9
2000.....	3.7	*	*	*	*	*	0.6	2.4	6.3	16.1	42.1	117.0
1999.....	3.6	*	*	*	*	*	0.6	1.5	5.9	15.3	43.3	119.0

Non-Hispanic black, male

2007 5/.....	2.9	*	*	*	*	*	*	2.1	6.5	16.5	39.9	85.5
2006 5/.....	2.8	*	*	*	*	*	*	2.1	6.8	16.0	38.7	66.9
2005 5/.....	3.1	*	*	*	*	*	*	1.9	6.8	18.6	49.9	79.9
2004 5/.....	3.0	*	*	*	*	*	*	2.0	6.5	17.5	43.6	103.6
2003 5/.....	3.0	*	*	*	*	*	*	1.9	5.7	18.5	51.7	89.5
2002.....	3.1	*	*	*	*	*	*	1.6	7.4	19.9	44.0	119.2
2001.....	3.3	*	*	*	*	*	0.9	2.0	6.6	20.5	49.8	119.0
2000.....	3.2	*	*	*	*	*	*	2.9	7.1	19.2	49.7	109.2
1999.....	3.0	*	*	*	*	*	*	1.4	6.7	17.9	51.9	107.0

Non-Hispanic black, female

2007 5/.....	3.4	*	*	*	*	*	*	1.3	4.0	10.7	29.4	87.5
2006 5/.....	3.6	*	*	*	*	*	*	1.4	3.5	9.8	36.9	94.2
2005 5/.....	3.7	*	*	*	*	*	*	1.1	4.1	12.7	33.4	102.6
2004 5/.....	4.1	*	*	*	*	*	0.8	1.5	4.4	13.7	36.1	115.7
2003 5/.....	4.0	*	*	*	*	*	0.8	2.1	4.1	12.5	34.5	111.8
2002.....	4.0	*	*	*	*	*	0.8	1.9	5.1	14.6	36.1	102.4
2001.....	4.4	*	*	*	*	*	0.9	1.4	6.3	15.5	38.4	120.2
2000.....	4.2	*	*	*	*	*	0.7	1.9	5.7	14.0	37.9	119.9
1999.....	4.2	*	*	*	*	*	0.8	1.6	5.4	13.5	38.5	123.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

All origins 3/, both sexes

2007.....	1.4	1.2	*	*	0.1	0.3	0.7	1.3	1.8	2.9	6.8	17.4
2006.....	1.3	0.8	*	*	0.1	0.3	0.6	1.2	1.7	2.9	7.0	18.4
2005.....	1.6	0.6	*	*	0.1	0.3	0.8	1.5	2.1	3.8	8.6	23.0
2004.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.1	3.6	9.0	23.3
2003.....	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001.....	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000.....	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999.....	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4

All origins 3/, male

2007.....	1.3	1.4	*	*	0.1	0.3	0.9	1.6	2.1	3.1	6.9	15.8
2006.....	1.2	*	*	*	*	0.3	0.7	1.5	2.1	2.9	7.0	16.2
2005.....	1.5	*	*	*	*	0.3	0.7	1.8	2.6	4.4	8.4	19.2
2004.....	1.4	*	*	*	*	0.4	0.7	1.6	2.5	3.9	9.7	19.6
2003.....	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002.....	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001.....	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000.....	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999.....	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8

All origins 3/, female

2007.....	1.4	1.1	*	*	*	0.3	0.6	1.1	1.5	2.6	6.8	18.2
2006.....	1.5	*	*	*	0.1	0.3	0.6	1.0	1.4	3.0	7.0	19.5
2005.....	1.8	*	*	*	0.1	0.4	0.8	1.1	1.7	3.3	8.8	24.8
2004.....	1.7	*	*	*	*	0.3	0.7	1.1	1.8	3.4	8.5	25.0
2003.....	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002.....	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001.....	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000.....	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999.....	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Hispanic, both sexes

2007.....	0.5	*	*	*	*	*	0.3	0.8	1.6	2.2	5.4	9.7
2006.....	0.5	*	*	*	*	*	*	0.9	1.2	2.3	4.7	11.3
2005.....	0.7	*	*	*	*	*	0.6	1.0	1.5	3.5	8.1	15.0
2004.....	0.5	*	*	*	*	*	0.3	1.2	1.6	2.0	4.8	17.6
2003.....	0.6	*	*	*	*	*	0.4	1.1	1.4	2.6	6.4	17.6
2002.....	0.6	*	*	*	*	*	0.4	1.0	1.4	3.1	6.3	18.3
2001.....	0.6	*	*	*	*	*	*	0.9	*	4.2	7.4	32.5
2000.....	0.6	*	*	*	*	*	0.6	1.1	1.3	3.3	8.7	24.6
1999.....	0.5	*	*	*	*	*	*	1.1	1.5	4.1	5.4	24.6

Hispanic, male

2007.....	0.5	*	*	*	*	*	*	1.0	1.9	*	*	*
2006.....	0.4	*	*	*	*	*	*	1.2	*	*	*	*
2005.....	0.6	*	*	*	*	*	*	1.2	2.0	3.9	*	*
2004.....	0.5	*	*	*	*	*	*	1.4	2.2	*	*	*
2003.....	0.6	*	*	*	*	*	*	1.4	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	1.3	*	4.0	8.9	*
2000.....	0.6	*	*	*	*	*	0.8	*	*	*	10.8	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Hispanic, female

2007.....	0.5	*	*	*	*	*	*	*	*	*	6.2	*
2006.....	0.5	*	*	*	*	*	*	*	*	*	2.6	4.9
2005.....	0.8	*	*	*	*	*	0.7	*	*	3.1	10.7	13.7
2004.....	0.6	*	*	*	*	*	*	*	*	*	4.9	23.8
2003.....	0.6	*	*	*	*	*	*	*	*	*	6.6	19.5
2002.....	0.7	*	*	*	*	*	*	*	*	3.2	6.2	23.2
2001.....	0.6	*	*	*	*	*	*	*	*	4.4	6.3	33.5
2000.....	0.7	*	*	*	*	*	*	*	*	*	7.3	23.7
1999.....	0.6	*	*	*	*	*	*	*	*	4.3	*	23.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic 4/, both sexes

2007.....	1.5	1.1	*	*	0.1	0.3	0.8	1.4	1.8	2.9	6.9	17.8
2006.....	1.5	0.8	*	*	0.1	0.3	0.7	1.3	1.7	3.0	7.1	18.8
2005.....	1.8	0.8	*	*	0.1	0.4	0.8	1.5	2.1	3.8	8.6	23.4
2004.....	1.8	0.8	*	*	0.1	0.4	0.8	1.4	2.2	3.7	9.2	23.5
2003.....	1.8	0.8	*	*	0.1	0.4	0.8	1.4	2.1	4.1	9.1	24.3
2002.....	1.8	0.6	*	*	0.1	0.4	0.7	1.4	2.3	3.9	10.0	24.7
2001.....	1.8	*	*	*	0.1	0.4	0.7	1.2	2.3	4.6	9.5	25.8
2000.....	1.8	0.7	*	*	0.1	0.4	0.7	1.4	2.2	4.4	9.4	25.6
1999.....	1.6	0.7	*	*	0.1	0.3	0.6	1.0	1.9	3.7	9.3	23.3

Non-Hispanic 4/, male

2007.....	1.4	1.3	*	*	*	0.4	1.0	1.6	2.1	3.1	7.0	16.3
2006.....	1.3	*	*	*	*	0.3	0.8	1.5	2.1	2.9	7.1	16.5
2005.....	1.6	*	*	*	*	0.3	0.8	1.9	2.6	4.5	8.7	19.3
2004.....	1.6	*	*	*	*	0.4	0.8	1.6	2.5	4.0	9.9	20.2
2003.....	1.5	*	*	*	*	0.3	0.8	1.6	2.4	4.2	8.6	19.3
2002.....	1.5	*	*	*	*	0.4	0.7	1.5	2.5	3.9	9.6	21.1
2001.....	1.5	*	*	*	*	0.3	0.7	1.4	2.6	4.9	9.8	21.6
2000.....	1.5	*	*	*	*	0.3	0.8	1.6	2.3	4.8	9.6	22.1
1999.....	1.4	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.7	20.5

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	0.3	0.7	1.1	1.5	2.7	6.8	18.5
2006.....	1.6	*	*	*	0.1	0.3	0.6	1.0	1.4	3.0	7.1	19.8
2005.....	1.9	*	*	*	*	0.4	0.8	1.1	1.7	3.3	8.6	25.3
2004.....	1.9	*	*	*	*	0.4	0.7	1.1	1.9	3.5	8.7	25.0
2003.....	2.1	*	*	*	0.1	0.4	0.9	1.2	1.7	4.1	9.5	26.5
2002.....	2.1	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.2	26.2
2001.....	2.0	*	*	*	0.1	0.4	0.6	1.0	2.1	4.3	9.2	27.6
2000.....	2.0	*	*	*	*	0.4	0.6	1.2	2.1	4.1	9.3	27.0
1999.....	1.8	*	*	*	*	0.3	0.5	0.9	1.8	3.3	9.0	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic white, both sexes

2007 5/.....	1.5	1.0	*	*	*	0.3	0.7	1.2	1.6	2.8	6.9	17.8
2006 5/.....	1.5	0.9	*	*	0.1	0.3	0.6	1.1	1.6	2.8	6.9	18.5
2005 5/.....	1.8	*	*	*	*	0.3	0.6	1.3	1.9	3.6	8.3	23.9
2004 5/.....	1.8	*	*	*	*	0.3	0.6	1.2	2.0	3.5	9.0	23.4
2003 5/.....	1.8	*	*	*	0.1	0.3	0.7	1.3	1.9	3.9	8.7	23.8
2002.....	1.8	*	*	*	*	0.3	0.6	1.2	2.1	3.6	9.6	24.5
2001.....	1.8	*	*	*	*	0.3	0.6	1.0	2.1	4.3	9.2	25.7
2000.....	1.8	*	*	*	*	0.3	0.6	1.2	1.9	4.0	9.2	25.5
1999.....	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.5	9.0	23.0

Non-Hispanic white, male

2007 5/.....	1.4	*	*	*	*	0.3	0.8	1.4	1.8	3.1	7.0	16.4
2006 5/.....	1.3	*	*	*	*	0.2	0.7	1.3	1.9	2.9	6.8	16.3
2005 5/.....	1.6	*	*	*	*	0.3	0.7	1.7	2.4	4.1	8.3	19.4
2004 5/.....	1.6	*	*	*	*	0.3	0.7	1.4	2.4	3.8	9.6	20.3
2003 5/.....	1.5	*	*	*	*	0.3	0.7	1.3	2.2	3.9	8.0	18.7
2002.....	1.5	*	*	*	*	0.3	0.6	1.3	2.2	3.7	9.1	20.5
2001.....	1.5	*	*	*	*	0.2	0.6	1.2	2.3	4.6	9.4	21.2
2000.....	1.5	*	*	*	*	0.2	0.7	1.5	2.0	4.5	9.4	21.8
1999.....	1.4	*	*	*	*	0.2	0.6	0.9	2.0	3.9	9.3	20.3

Non-Hispanic white, female

2007 5/.....	1.7	*	*	*	*	0.2	0.5	1.0	1.4	2.5	6.8	18.4
2006 5/.....	1.7	*	*	*	*	0.3	0.5	0.8	1.3	2.8	7.0	19.5
2005 5/.....	2.0	*	*	*	*	0.3	0.6	1.0	1.5	3.2	8.4	26.0
2004 5/.....	2.0	*	*	*	*	0.3	0.6	0.9	1.6	3.3	8.5	24.8
2003 5/.....	2.1	*	*	*	*	0.3	0.6	1.2	1.6	3.9	9.3	26.1
2002.....	2.2	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.9	26.3
2001.....	2.1	*	*	*	*	0.3	0.5	0.8	1.9	4.1	9.0	27.6
2000.....	2.0	*	*	*	*	0.3	0.5	1.0	1.9	3.6	9.0	27.0
1999.....	1.8	*	*	*	*	0.2	0.5	0.8	1.7	3.1	8.8	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic black, both sexes

2007 5/.....	1.8	*	*	*	*	0.7	1.8	2.8	3.8	4.6	9.2	23.0
2006 5/.....	1.9	*	*	*	*	0.8	1.7	2.9	3.5	4.9	11.1	26.0
2005 5/.....	2.1	*	*	*	*	0.8	1.8	3.0	4.6	6.5	14.0	24.3
2004 5/.....	2.1	*	*	*	*	1.0	2.0	3.0	4.2	6.8	13.9	29.1
2003 5/.....	2.2	*	*	*	*	0.9	2.0	2.7	4.4	7.4	15.5	33.6
2002.....	2.2	*	*	*	*	1.0	1.7	2.7	5.0	6.9	16.8	29.9
2001.....	2.0	*	*	*	*	0.8	1.6	2.3	4.9	7.9	14.3	29.7
2000.....	2.1	*	*	*	*	0.9	1.5	2.8	4.6	9.0	14.4	29.9
1999.....	1.8	*	*	*	*	0.5	1.5	2.4	3.0	6.8	14.0	28.5

Non-Hispanic black, male

2007 5/.....	1.9	*	*	*	*	0.8	2.1	3.4	5.2	5.0	9.9	20.1
2006 5/.....	1.8	*	*	*	*	0.8	1.5	3.5	4.2	5.1	12.0	22.9
2005 5/.....	2.1	*	*	*	*	*	1.7	3.8	5.7	8.9	16.0	23.8
2004 5/.....	2.0	*	*	*	*	1.1	2.0	3.5	4.5	7.3	15.4	25.4
2003 5/.....	2.1	*	*	*	*	0.9	1.4	3.5	5.4	8.4	17.9	33.0
2002.....	2.1	*	*	*	*	1.1	1.8	3.0	5.6	7.8	18.5	31.7
2001.....	2.0	*	*	*	*	*	1.8	2.4	6.5	9.2	15.9	28.9
2000.....	2.0	*	*	*	*	0.9	1.8	3.5	5.3	9.1	13.3	30.5
1999.....	1.7	*	*	*	*	*	1.9	2.9	2.7	8.4	17.2	*

Non-Hispanic black, female

2007 5/.....	1.7	*	*	*	*	*	1.6	2.3	2.7	4.3	8.8	24.2
2006 5/.....	1.9	*	*	*	*	0.8	1.8	2.3	2.9	4.8	10.6	27.3
2005 5/.....	2.1	*	*	*	*	0.9	1.9	2.4	3.6	4.8	12.9	24.5
2004 5/.....	2.3	*	*	*	*	0.8	1.9	2.6	3.9	6.5	13.0	30.5
2003 5/.....	2.3	*	*	*	*	1.0	2.6	1.9	3.6	6.8	14.2	33.8
2002.....	2.3	*	*	*	*	1.0	1.6	2.4	4.5	6.2	15.9	29.3
2001.....	2.1	*	*	*	*	0.9	1.4	2.1	3.7	7.0	13.5	29.9
2000.....	2.3	*	*	*	*	1.0	1.3	2.3	4.0	8.8	15.1	29.7
1999.....	1.8	*	*	*	*	*	1.1	2.0	3.2	5.6	12.3	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza and pneumonia (J09-J18)

All origins 3/, both sexes

2007.....	17.5	5.2	0.7	0.3	0.4	0.8	1.8	4.4	9.6	28.7	114.1	463.2
2006.....	18.8	6.4	0.8	0.2	0.4	0.8	1.9	4.6	10.0	32.0	127.8	502.5
2005.....	21.3	6.5	0.7	0.3	0.4	0.9	2.1	5.1	11.3	35.5	142.2	593.9
2004.....	20.3	6.7	0.7	0.2	0.4	0.8	2.0	4.6	10.8	34.6	139.3	582.6
2003.....	22.4	8.0	1.0	0.4	0.5	0.9	2.2	5.2	11.2	37.3	151.1	666.1
2002.....	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001.....	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000.....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999.....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8

All origins 3/, male

2007.....	16.2	6.2	0.7	0.3	0.4	0.9	2.0	5.2	11.9	34.3	139.5	542.7
2006.....	17.4	7.7	0.8	0.2	0.5	0.9	2.1	5.6	12.3	38.9	156.9	586.4
2005.....	19.2	7.0	0.7	0.3	0.5	1.0	2.5	6.2	13.6	42.8	174.3	673.5
2004.....	18.6	8.0	0.6	0.2	0.6	1.0	2.2	5.6	13.0	42.8	172.7	677.2
2003.....	20.1	9.4	1.1	0.3	0.6	1.1	2.5	6.4	13.2	46.5	186.8	757.8
2002.....	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001.....	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000.....	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999.....	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7

All origins 3/, female

2007.....	18.7	4.2	0.7	0.3	0.3	0.7	1.6	3.5	7.5	23.9	96.6	425.5
2006.....	20.2	5.0	0.8	0.2	0.4	0.8	1.7	3.7	7.8	26.3	107.8	463.3
2005.....	23.2	5.8	0.7	0.3	0.3	0.8	1.8	4.1	9.1	29.4	120.4	557.4
2004.....	22.0	5.4	0.9	0.2	0.3	0.6	1.8	3.6	8.8	27.7	116.8	540.0
2003.....	24.6	6.6	0.9	0.4	0.5	0.7	1.9	4.1	9.4	29.5	127.2	625.6
2002.....	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001.....	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000.....	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999.....	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J09-J18)												
Hispanic, both sexes												
2007.....	6.0	4.5	0.6	0.2	*	0.5	1.2	2.7	8.4	27.5	100.5	340.4
2006.....	6.7	6.2	0.7	*	0.3	0.6	1.2	3.7	9.8	30.8	114.2	390.9
2005.....	7.2	7.1	0.9	*	*	0.5	1.5	4.0	10.4	34.4	122.3	456.9
2004.....	7.0	5.8	0.7	*	0.5	0.6	1.3	3.8	9.6	33.2	125.2	482.3
2003.....	7.4	7.6	1.1	0.3	0.5	0.7	1.3	4.0	10.7	35.0	134.3	516.2
2002.....	7.3	5.8	*	*	*	0.6	1.3	3.4	10.5	33.8	142.6	566.0
2001.....	7.4	7.0	1.1	*	*	0.3	1.3	3.6	9.1	35.4	137.2	660.3
2000.....	7.4	7.8	*	0.3	0.3	0.7	1.6	3.8	11.2	37.6	136.3	643.6
1999.....	6.6	8.5	*	*	0.4	0.4	1.5	3.5	9.5	31.7	118.5	614.6
Hispanic, male												
2007.....	5.8	5.3	*	*	*	0.5	1.6	3.3	11.0	33.6	126.7	370.0
2006.....	6.3	7.3	*	*	*	0.7	1.4	4.5	12.6	36.7	131.5	440.9
2005.....	6.8	7.8	*	*	*	0.6	1.7	5.1	12.9	42.3	149.7	496.9
2004.....	6.6	8.0	*	*	0.6	0.7	1.5	4.9	11.5	38.9	152.0	528.2
2003.....	6.9	9.1	1.4	*	0.5	0.9	1.3	5.2	11.6	43.4	165.7	577.2
2002.....	7.0	7.7	*	*	*	0.7	1.7	4.7	13.9	43.8	173.2	655.9
2001.....	6.9	7.4	*	*	*	*	1.7	5.0	11.4	40.7	173.5	765.6
2000.....	7.2	7.6	*	*	*	0.8	2.1	5.3	14.8	46.0	180.4	763.9
1999.....	6.3	10.3	*	*	*	*	2.1	4.9	10.4	41.1	148.1	720.3
Hispanic, female												
2007.....	6.3	*	*	*	*	*	0.8	2.2	5.9	22.5	82.2	324.3
2006.....	7.1	5.1	*	*	*	*	0.9	2.8	7.3	26.0	102.0	364.3
2005.....	7.7	6.4	*	*	*	*	1.1	2.9	8.1	28.1	103.3	436.0
2004.....	7.6	*	*	*	*	*	1.2	2.6	7.9	28.7	106.7	458.8
2003.....	7.9	6.1	*	*	*	*	1.2	2.7	9.9	28.4	112.7	485.2
2002.....	7.6	*	*	*	*	*	0.9	2.1	7.4	25.9	121.7	521.4
2001.....	7.9	6.5	*	*	*	*	1.0	2.1	7.1	31.2	112.7	608.8
2000.....	7.7	8.0	*	*	*	0.7	1.0	2.4	7.9	31.0	106.6	584.6
1999.....	7.0	6.5	*	*	*	*	0.8	2.2	8.6	24.3	98.9	563.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Influenza and pneumonia (J09-J18)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	19.5	5.4	0.7	0.3	0.4	0.9	1.9	4.5	9.7	28.7	114.8	469.2
2006.....	20.9	6.3	0.8	0.2	0.5	0.9	2.1	4.7	10.0	32.0	128.4	507.5
2005.....	23.6	6.2	0.6	0.3	0.5	1.0	2.2	5.2	11.3	35.5	143.1	599.9
2004.....	22.4	6.9	0.8	0.2	0.4	0.8	2.1	4.6	10.9	34.5	139.7	586.0
2003.....	24.7	8.1	1.0	0.4	0.5	1.0	2.4	5.3	11.2	37.2	151.6	671.1
2002.....	25.1	6.7	0.7	0.2	0.4	0.9	2.3	4.9	11.3	37.6	157.1	700.2
2001.....	23.8	7.5	0.6	0.2	0.5	1.0	2.3	4.7	10.7	36.1	148.6	684.8
2000.....	25.4	7.5	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.0	160.7	745.7
1999.....	25.0	8.4	0.9	0.2	0.5	0.9	2.4	4.7	11.0	37.4	158.0	754.0
<b>Non-Hispanic 4/, male</b>												
2007.....	18.1	6.5	0.7	0.3	0.5	1.0	2.1	5.4	12.0	34.2	140.1	552.2
2006.....	19.4	7.8	0.8	0.2	0.5	0.9	2.2	5.7	12.3	38.9	158.2	593.8
2005.....	21.4	6.7	0.6	0.3	0.6	1.0	2.6	6.3	13.6	42.6	175.4	682.2
2004.....	20.6	8.0	0.6	0.2	0.5	1.0	2.3	5.6	13.0	42.7	173.5	683.2
2003.....	22.3	9.4	1.1	0.3	0.6	1.2	2.7	6.4	13.2	46.4	187.6	764.6
2002.....	22.5	7.8	0.8	0.2	0.5	1.0	2.5	5.8	13.5	46.3	195.1	810.7
2001.....	21.5	8.2	0.6	0.2	0.5	1.1	2.6	5.7	13.1	44.9	186.5	808.2
2000.....	22.7	8.5	0.7	0.2	0.6	1.1	2.9	5.7	14.3	48.2	199.7	894.9
1999.....	22.2	8.4	0.9	0.3	0.5	1.1	2.7	5.9	12.9	47.0	197.7	897.9
<b>Non-Hispanic 4/, female</b>												
2007.....	20.8	4.3	0.7	0.3	0.4	0.8	1.7	3.7	7.6	24.0	97.4	430.0
2006.....	22.3	4.8	0.8	0.2	0.4	0.9	1.9	3.8	7.8	26.2	108.0	467.4
2005.....	25.7	5.7	0.6	0.3	0.3	0.9	1.9	4.2	9.2	29.5	121.2	562.3
2004.....	24.2	5.9	0.9	0.2	0.3	0.6	1.9	3.7	8.9	27.5	117.0	542.5
2003.....	27.1	6.8	1.0	0.4	0.5	0.8	2.1	4.3	9.3	29.5	127.7	630.0
2002.....	27.5	5.4	0.7	0.3	0.4	0.8	2.0	4.0	9.2	30.3	131.8	652.6
2001.....	26.1	6.7	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.9	123.8	633.5
2000.....	27.9	6.5	0.6	0.2	0.5	0.8	2.0	3.8	9.6	31.4	135.3	685.4
1999.....	27.6	8.4	0.9	0.2	0.5	0.8	2.2	3.5	9.4	29.5	132.2	696.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Influenza and pneumonia (J09-J18)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	21.5	3.8	0.5	0.2	0.4	0.8	1.8	4.0	9.0	27.2	114.2	475.8
2006 5/.....	23.1	4.5	0.6	0.2	0.4	0.8	1.9	4.2	9.5	30.9	128.0	514.4
2005 5/.....	26.2	4.1	0.5	0.3	0.5	0.9	2.0	4.7	10.4	33.9	143.3	612.6
2004 5/.....	24.8	4.8	0.6	0.2	0.4	0.8	1.8	4.0	10.1	33.2	138.8	593.8
2003 5/.....	27.4	6.2	0.9	0.3	0.5	0.8	2.1	4.7	10.1	35.7	151.9	684.5
2002.....	27.9	4.9	0.5	0.2	0.4	0.8	2.0	4.4	10.3	36.7	156.9	712.7
2001.....	26.2	4.9	0.5	0.2	0.4	0.8	1.9	4.0	9.9	35.0	147.2	693.9
2000.....	27.9	5.0	0.5	0.2	0.5	0.8	2.1	4.1	10.7	36.8	160.2	759.8
1999.....	27.5	5.3	0.6	0.2	0.4	0.8	2.0	4.0	10.0	36.1	158.8	766.9
<b>Non-Hispanic white, male</b>												
2007 5/.....	19.6	4.8	0.4	0.2	0.5	0.9	1.9	4.6	10.9	31.8	137.7	557.8
2006 5/.....	21.2	5.8	0.6	0.2	0.5	0.8	2.1	4.9	11.5	37.1	156.9	598.8
2005 5/.....	23.3	5.1	0.5	0.2	0.6	1.0	2.3	5.5	12.2	40.1	174.6	693.4
2004 5/.....	22.4	5.5	0.5	0.2	0.5	0.9	2.0	4.9	12.0	40.3	170.7	691.1
2003 5/.....	24.1	7.4	0.9	0.3	0.6	1.1	2.3	5.6	11.4	43.6	185.5	774.1
2002.....	24.6	6.1	0.5	0.2	0.5	0.9	2.2	5.2	12.1	44.9	193.2	820.6
2001.....	23.1	5.8	0.6	0.2	0.5	0.9	2.1	4.8	11.7	42.7	182.5	813.9
2000.....	24.5	5.8	0.5	0.2	0.5	0.9	2.4	4.7	12.6	44.7	197.5	908.1
1999.....	23.9	5.5	0.5	0.3	0.4	0.9	2.1	4.8	11.5	44.6	197.3	909.3
<b>Non-Hispanic white, female</b>												
2007 5/.....	23.3	2.7	0.7	0.3	0.4	0.7	1.6	3.4	7.1	23.3	97.7	436.9
2006 5/.....	25.0	3.0	0.6	*	0.4	0.8	1.8	3.5	7.5	25.6	107.9	475.0
2005 5/.....	28.9	3.0	0.5	0.3	0.3	0.9	1.8	3.8	8.8	28.6	121.7	575.7
2004 5/.....	27.1	4.2	0.7	0.2	0.3	0.6	1.7	3.2	8.3	27.0	117.0	550.2
2003 5/.....	30.6	5.0	0.8	0.4	0.5	0.6	1.8	3.8	8.8	28.9	129.3	645.1
2002.....	31.1	3.6	0.5	0.3	0.3	0.8	1.8	3.7	8.6	29.7	132.5	666.1
2001.....	29.2	3.9	0.5	0.2	0.4	0.8	1.7	3.3	8.1	28.5	123.7	644.0
2000.....	31.3	4.2	0.6	0.2	0.4	0.8	1.8	3.4	9.0	30.1	135.7	700.0
1999.....	31.0	5.1	0.8	0.2	0.4	0.6	1.9	3.1	8.7	29.0	133.5	710.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Influenza and pneumonia (J09-J18)

##### Non-Hispanic black, both sexes

2007 5/.....	13.3	11.2	1.1	0.3	0.4	1.7	3.2	8.5	17.4	44.3	129.4	406.0
2006 5/.....	13.9	13.7	1.4	*	0.6	1.4	3.4	9.0	16.2	45.5	141.2	439.6
2005 5/.....	15.3	15.1	1.1	0.4	0.5	1.5	3.9	10.1	20.3	53.2	151.8	475.4
2004 5/.....	15.2	14.7	1.4	0.3	0.7	1.2	4.1	9.3	19.1	50.4	158.4	511.7
2003 5/.....	15.9	14.4	1.1	0.4	0.7	1.9	4.5	10.5	21.1	55.0	157.7	528.3
2002.....	16.1	13.3	1.7	*	0.6	1.6	4.5	8.9	21.6	50.9	167.7	579.8
2001.....	16.0	16.9	1.0	*	0.8	1.7	4.9	9.9	19.1	48.8	169.5	583.1
2000.....	16.9	17.5	1.2	*	0.8	1.8	4.6	10.3	22.3	60.8	172.2	604.5
1999.....	16.7	20.0	1.7	*	0.9	2.0	5.5	10.8	20.9	54.7	158.3	634.7

##### Non-Hispanic black, male

2007 5/.....	13.5	12.5	*	*	*	2.0	3.9	11.3	22.6	60.6	174.4	480.8
2006 5/.....	13.6	16.3	*	*	0.7	1.4	3.7	11.7	21.8	59.5	182.1	512.2
2005 5/.....	15.1	14.0	*	*	0.7	1.7	5.0	13.2	28.1	71.2	191.5	537.6
2004 5/.....	15.5	17.0	*	*	*	1.6	4.9	12.1	24.8	73.2	221.0	607.6
2003 5/.....	16.5	16.1	*	*	0.8	2.3	5.5	13.3	29.8	79.1	220.8	642.5
2002.....	15.9	14.2	1.8	*	0.7	1.8	5.2	11.5	29.6	66.5	231.2	695.8
2001.....	16.3	16.6	*	*	0.8	2.1	5.9	13.5	26.6	70.0	242.5	718.4
2000.....	17.3	18.8	1.7	*	0.8	2.4	5.8	14.3	30.3	84.1	235.4	763.2
1999.....	16.9	20.1	2.1	*	0.8	2.3	6.7	15.3	27.3	77.9	215.1	803.4

##### Non-Hispanic black, female

2007 5/.....	13.2	9.9	*	*	*	1.5	2.6	6.1	13.2	32.6	103.9	375.1
2006 5/.....	14.1	10.9	*	*	*	1.4	3.0	6.6	11.7	35.4	118.0	409.9
2005 5/.....	15.5	16.2	*	*	*	1.2	3.0	7.4	14.0	40.3	129.3	450.4
2004 5/.....	15.0	12.2	1.8	*	0.7	0.8	3.3	6.9	14.5	34.1	123.0	473.8
2003 5/.....	15.4	12.6	*	*	*	1.6	3.7	8.1	14.0	37.8	122.1	484.1
2002.....	16.2	12.3	*	*	*	1.4	3.9	6.6	15.2	39.8	131.8	535.5
2001.....	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	128.5	533.0
2000.....	16.5	16.2	*	*	0.8	1.2	3.5	6.9	15.9	44.4	136.7	545.6
1999.....	16.5	20.0	*	*	0.9	1.8	4.5	7.0	15.8	38.4	126.4	571.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza (J09-J11)

All origins 3/, both sexes

2007.....	0.1	*	*	0.1	*	*	*	*	0.1	0.2	0.6	2.5
2006.....	0.3	*	*	0.1	*	*	*	0.1	0.1	0.4	1.8	7.0
2005.....	0.6	*	*	0.0	*	*	*	0.1	0.2	0.7	3.7	19.3
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.2	11.4
2003.....	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000.....	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999.....	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2

All origins 3/, male

2007.....	0.1	*	*	*	*	*	*	*	0.1	*	0.7	3.0
2006.....	0.2	*	*	*	*	*	*	*	0.1	0.5	2.0	8.1
2005.....	0.5	*	*	*	*	*	*	0.1	0.3	0.7	3.6	18.2
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	11.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000.....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9

All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2006.....	0.3	*	*	*	*	*	*	*	0.1	0.4	1.7	6.5
2005.....	0.8	*	*	*	*	*	*	*	0.2	0.8	3.7	19.8
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.1	11.5
2003.....	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002.....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999.....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza (J09-J11)

Non-Hispanic 4/, both sexes

2007.....	0.1	*	*	0.1	*	*	*	*	0.1	0.2	0.6	2.6
2006.....	0.3	*	*	*	*	*	*	0.1	0.1	0.5	1.9	7.3
2005.....	0.7	*	*	*	*	*	*	0.1	0.2	0.8	3.8	19.8
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.3	11.8
2003.....	0.7	0.7	0.4	0.1	0.1	*	0.1	0.2	0.3	1.1	3.6	17.5
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	9.0
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.3
2000.....	0.7	*	*	*	*	*	0.0	0.1	0.4	1.1	4.1	20.5
1999.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.6

Non-Hispanic 4/, male

2007.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.1
2006.....	0.3	*	*	*	*	*	*	*	*	0.5	2.1	8.4
2005.....	0.5	*	*	*	*	*	*	0.1	0.3	0.7	3.8	18.7
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.5	11.6
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.5
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	9.2
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2000.....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.1	21.1
1999.....	0.5	*	*	*	*	*	*	*	0.3	0.8	4.3	20.7

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	*	*	*	*	*	*	0.6	2.3
2006.....	0.4	*	*	*	*	*	*	*	0.1	0.4	1.8	6.7
2005.....	0.9	*	*	*	*	*	*	*	0.2	0.8	3.8	20.3
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.2	11.8
2003.....	0.8	*	0.4	0.1	*	*	*	0.1	0.3	1.0	3.6	17.6
2002.....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000.....	0.9	*	*	*	*	*	*	0.1	0.4	1.0	4.1	20.3
1999.....	0.8	*	*	*	*	*	*	*	0.3	0.9	3.4	22.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza (J09-J11)

Non-Hispanic white, both sexes

2007 5/.....	0.2	*	*	0.1	*	*	*	*	0.1	0.2	0.7	2.8
2006 5/.....	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.0	7.8
2005 5/.....	0.8	*	*	*	*	*	*	0.1	0.2	0.8	4.1	21.5
2004 5/.....	0.5	*	*	*	*	*	*	*	0.2	0.6	2.5	12.7
2003 5/.....	0.8	*	0.3	0.1	*	*	*	0.2	0.3	1.1	3.9	18.8
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.6	9.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	1.2	4.3	21.7
1999.....	0.8	*	*	*	*	*	*	0.1	0.3	0.9	3.9	23.3

Non-Hispanic white, male

2007 5/.....	0.2	*	*	*	*	*	*	*	*	*	0.8	3.4
2006 5/.....	0.3	*	*	*	*	*	*	*	*	0.6	2.2	8.9
2005 5/.....	0.6	*	*	*	*	*	*	*	0.3	0.8	4.1	19.9
2004 5/.....	0.4	*	*	*	*	*	*	*	0.2	0.8	2.5	12.7
2003 5/.....	0.6	*	*	*	*	*	*	0.2	0.3	1.2	4.0	18.9
2002.....	0.3	*	*	*	*	*	*	*	*	0.4	1.7	10.0
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000.....	0.6	*	*	*	*	*	*	*	0.4	1.4	4.3	22.0
1999.....	0.6	*	*	*	*	*	*	*	0.3	0.9	4.6	22.5

Non-Hispanic white, female

2007 5/.....	0.2	*	*	*	*	*	*	*	*	*	0.6	2.6
2006 5/.....	0.4	*	*	*	*	*	*	*	0.2	0.4	2.0	7.3
2005 5/.....	1.0	*	*	*	*	*	*	*	0.2	0.9	4.2	22.2
2004 5/.....	0.6	*	*	*	*	*	*	*	0.2	0.5	2.4	12.8
2003 5/.....	1.0	*	*	*	*	*	*	0.1	0.3	1.1	3.9	18.7
2002.....	0.4	*	*	*	*	*	*	*	*	*	1.4	9.5
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000.....	1.0	*	*	*	*	*	*	0.1	0.5	1.0	4.3	21.6
1999.....	1.0	*	*	*	*	*	*	*	0.3	1.0	3.5	23.5



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All origins 3/, both sexes												
2007.....	17.3	4.9	0.5	0.2	0.4	0.8	1.8	4.3	9.5	28.5	113.5	460.7
2006.....	18.5	5.9	0.7	0.1	0.4	0.8	1.9	4.6	9.8	31.6	125.9	495.5
2005.....	20.6	6.0	0.6	0.2	0.4	0.9	2.1	5.0	11.1	34.8	138.5	574.7
2004.....	19.9	6.3	0.6	0.2	0.4	0.7	2.0	4.5	10.7	34.0	137.0	571.2
2003.....	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002.....	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001.....	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000.....	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999.....	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All origins 3/, male												
2007.....	16.1	5.8	0.5	0.2	0.4	0.9	2.0	5.2	11.8	34.0	138.8	539.7
2006.....	17.1	7.2	0.6	0.1	0.4	0.8	2.1	5.5	12.2	38.4	155.0	578.3
2005.....	18.8	6.6	0.6	0.2	0.5	0.9	2.4	6.1	13.3	42.1	170.7	655.3
2004.....	18.3	7.5	0.5	0.2	0.5	0.9	2.2	5.5	12.8	42.1	170.3	666.1
2003.....	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002.....	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001.....	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000.....	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999.....	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All origins 3/, female												
2007.....	18.6	4.0	0.6	0.2	0.3	0.7	1.6	3.5	7.4	23.7	96.0	423.2
2006.....	19.9	4.6	0.7	*	0.4	0.8	1.7	3.7	7.7	25.9	106.1	456.7
2005.....	22.5	5.3	0.6	0.2	0.2	0.8	1.8	4.0	8.9	28.7	116.7	537.6
2004.....	21.5	5.0	0.8	0.2	0.3	0.5	1.8	3.5	8.6	27.3	114.6	528.5
2003.....	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002.....	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001.....	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000.....	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999.....	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonia (J12-J18)

Hispanic, both sexes

2007.....	5.9	4.1	0.5	*	*	0.5	1.2	2.7	8.3	27.3	100.4	339.3
2006.....	6.6	5.7	0.5	*	0.3	0.6	1.2	3.6	9.7	30.7	113.8	388.9
2005.....	7.1	6.2	0.7	*	*	0.5	1.5	3.9	10.3	34.1	120.9	450.5
2004.....	7.0	5.2	0.6	*	0.4	0.6	1.3	3.7	9.6	32.9	124.9	479.5
2003.....	7.2	6.5	0.7	*	0.5	0.7	1.2	3.9	10.4	34.8	132.4	511.0
2002.....	7.2	5.5	*	*	*	0.6	1.3	3.4	10.5	33.6	142.1	563.2
2001.....	7.3	7.0	1.0	*	*	0.3	1.3	3.5	9.1	35.3	136.3	660.3
2000.....	7.3	7.5	*	*	0.3	0.7	1.6	3.8	11.0	37.2	134.3	633.0
1999.....	6.5	7.9	*	*	0.4	0.4	1.5	3.4	9.2	30.9	116.7	607.6

Hispanic, male

2007.....	5.7	4.5	*	*	*	0.5	1.6	3.1	10.9	33.4	126.7	370.0
2006.....	6.2	6.4	*	*	*	0.7	1.4	4.4	12.5	36.5	131.5	438.7
2005.....	6.7	6.7	*	*	*	0.6	1.7	4.9	12.8	42.0	149.0	488.2
2004.....	6.5	7.3	*	*	0.6	0.7	1.5	4.9	11.5	38.5	151.3	526.8
2003.....	6.7	7.5	*	*	*	0.8	1.3	5.2	11.4	42.8	163.0	571.1
2002.....	6.9	7.3	*	*	*	0.6	1.7	4.7	13.9	43.6	172.8	654.2
2001.....	6.8	7.4	*	*	*	*	1.7	5.0	11.4	40.7	172.6	765.6
2000.....	7.1	7.6	*	*	*	0.7	2.1	5.2	14.6	45.6	178.0	751.8
1999.....	6.1	10.3	*	*	*	*	2.0	4.7	9.9	39.5	147.0	718.2

Hispanic, female

2007.....	6.2	*	*	*	*	*	0.8	2.2	5.9	22.4	82.0	322.7
2006.....	7.0	4.8	*	*	*	*	0.9	2.8	7.1	26.0	101.4	362.5
2005.....	7.5	5.7	*	*	*	*	1.1	2.9	8.0	27.8	101.4	430.8
2004.....	7.5	*	*	*	*	*	1.2	2.5	7.9	28.4	106.7	455.2
2003.....	7.7	5.4	*	*	*	*	1.1	2.6	9.6	28.4	111.4	480.5
2002.....	7.5	*	*	*	*	*	0.9	2.1	7.4	25.7	121.1	518.1
2001.....	7.8	6.5	*	*	*	*	0.9	2.1	7.1	31.0	111.8	608.8
2000.....	7.6	7.4	*	*	*	*	1.0	2.3	7.9	30.7	105.0	574.7
1999.....	6.8	5.4	*	*	*	*	0.8	2.2	8.5	24.1	96.4	554.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Pneumonia (J12-J18)</b>												
<b>Non-Hispanic 4/, both sexes</b>												
2007.....	19.3	5.2	0.5	0.2	0.4	0.9	1.9	4.5	9.6	28.5	114.2	466.6
2006.....	20.6	5.9	0.7	0.1	0.4	0.9	2.0	4.7	9.9	31.6	126.5	500.2
2005.....	22.9	5.9	0.5	0.2	0.4	0.9	2.2	5.1	11.1	34.7	139.2	580.1
2004.....	22.0	6.6	0.6	0.2	0.4	0.8	2.1	4.6	10.7	33.9	137.4	574.2
2003.....	24.0	7.4	0.6	0.3	0.5	0.9	2.3	5.2	10.9	36.2	148.0	653.6
2002.....	24.8	6.5	0.7	0.2	0.4	0.9	2.3	4.9	11.1	37.3	155.6	691.2
2001.....	23.8	7.2	0.6	0.2	0.5	0.9	2.3	4.6	10.6	36.0	148.2	682.5
2000.....	24.7	7.3	0.6	0.2	0.5	0.9	2.4	4.6	11.4	37.9	156.6	725.2
1999.....	24.3	8.1	0.8	0.2	0.4	0.9	2.4	4.6	10.7	36.5	154.2	732.4
<b>Non-Hispanic 4/, male</b>												
2007.....	18.0	6.2	0.5	0.2	0.5	1.0	2.1	5.4	11.8	34.0	139.4	549.1
2006.....	19.1	7.4	0.7	0.1	0.5	0.9	2.2	5.6	12.2	38.4	156.1	585.5
2005.....	20.9	6.5	0.5	0.2	0.6	1.0	2.5	6.2	13.3	41.9	171.5	663.5
2004.....	20.3	7.5	0.5	0.2	0.5	1.0	2.3	5.6	12.8	42.0	171.0	671.6
2003.....	21.7	8.7	0.7	0.2	0.5	1.1	2.7	6.3	12.9	45.3	183.9	747.1
2002.....	22.3	7.6	0.8	0.2	0.5	1.0	2.5	5.8	13.4	45.9	193.5	801.5
2001.....	21.4	7.8	0.6	0.2	0.5	1.0	2.6	5.7	13.0	44.8	186.1	805.9
2000.....	22.2	8.2	0.7	0.2	0.5	1.0	2.8	5.6	13.9	46.8	195.7	873.8
1999.....	21.7	8.0	0.8	0.2	0.5	1.1	2.6	5.8	12.5	46.2	193.4	877.2
<b>Non-Hispanic 4/, female</b>												
2007.....	20.6	4.1	0.6	0.2	0.3	0.7	1.7	3.6	7.5	23.8	96.9	427.7
2006.....	21.9	4.4	0.7	*	0.4	0.9	1.9	3.8	7.7	25.8	106.2	460.6
2005.....	24.8	5.2	0.5	0.3	0.3	0.8	1.9	4.1	9.0	28.7	117.3	542.0
2004.....	23.7	5.6	0.8	0.2	0.3	0.6	1.8	3.6	8.7	27.1	114.8	530.7
2003.....	26.3	6.1	0.6	0.3	0.4	0.7	2.0	4.1	9.0	28.5	124.1	612.4
2002.....	27.2	5.3	0.6	0.2	0.4	0.8	2.0	4.0	9.1	30.1	130.6	643.7
2001.....	26.0	6.6	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.8	123.3	631.3
2000.....	27.1	6.4	0.5	0.1	0.4	0.8	2.0	3.7	9.2	30.4	131.2	665.2
1999.....	26.8	8.2	0.9	0.2	0.4	0.8	2.2	3.5	9.1	28.5	128.8	674.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Pneumonia (J12-J18)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	21.3	3.6	0.4	0.1	0.4	0.8	1.7	4.0	8.8	27.0	113.5	472.9
2006 5/.....	22.7	4.1	0.6	0.1	0.4	0.8	1.9	4.2	9.3	30.4	125.9	506.5
2005 5/.....	25.3	3.8	0.4	0.2	0.4	0.9	2.0	4.6	10.2	33.1	139.2	591.2
2004 5/.....	24.3	4.6	0.5	0.1	0.4	0.7	1.8	4.0	9.9	32.6	136.3	581.1
2003 5/.....	26.6	5.7	0.6	0.3	0.5	0.8	2.0	4.5	9.8	34.6	148.0	665.7
2002.....	27.6	4.7	0.5	0.2	0.4	0.8	2.0	4.4	10.2	36.4	155.3	703.1
2001.....	26.1	4.6	0.5	0.2	0.4	0.8	1.9	4.0	9.8	34.9	146.7	691.5
2000.....	27.1	4.8	0.5	0.2	0.4	0.8	2.1	3.9	10.3	35.6	155.9	738.1
1999.....	26.8	5.2	0.6	0.2	0.4	0.7	1.9	3.9	9.7	35.2	154.8	743.7
<b>Non-Hispanic white, male</b>												
2007 5/.....	19.4	4.6	*	*	0.5	0.9	1.9	4.6	10.8	31.6	137.0	554.4
2006 5/.....	20.9	5.4	0.6	*	0.5	0.8	2.0	4.9	11.4	36.5	154.7	589.9
2005 5/.....	22.7	4.7	*	0.2	0.6	0.9	2.2	5.4	11.9	39.3	170.5	673.6
2004 5/.....	22.0	5.3	*	*	0.5	0.9	2.0	4.8	11.8	39.5	168.2	678.4
2003 5/.....	23.5	6.8	0.6	0.2	0.5	1.0	2.3	5.4	11.1	42.5	181.5	755.2
2002.....	24.3	6.0	0.5	0.2	0.5	0.9	2.2	5.1	11.9	44.5	191.5	810.6
2001.....	23.0	5.3	0.6	0.2	0.5	0.9	2.1	4.8	11.6	42.6	182.0	811.5
2000.....	23.8	5.4	0.4	0.2	0.5	0.8	2.4	4.6	12.2	43.3	193.2	886.1
1999.....	23.4	5.4	*	0.2	0.4	0.9	2.0	4.7	11.1	43.8	192.7	886.8
<b>Non-Hispanic white, female</b>												
2007 5/.....	23.1	2.6	0.6	0.2	0.3	0.6	1.6	3.4	7.0	23.1	97.0	434.3
2006 5/.....	24.6	2.8	0.6	*	0.4	0.8	1.7	3.4	7.4	25.2	105.9	467.7
2005 5/.....	27.9	2.8	0.4	0.3	0.3	0.8	1.7	3.7	8.5	27.7	117.6	553.5
2004 5/.....	26.5	3.8	0.6	*	0.2	0.5	1.6	3.2	8.1	26.6	114.6	537.5
2003 5/.....	29.6	4.5	0.5	0.3	0.4	0.6	1.8	3.7	8.5	27.9	125.4	626.4
2002.....	30.7	3.4	0.5	0.2	0.3	0.8	1.8	3.7	8.5	29.5	131.1	656.6
2001.....	29.1	3.8	0.4	0.2	0.3	0.7	1.7	3.2	8.0	28.3	123.3	641.7
2000.....	30.3	4.2	0.5	*	0.3	0.8	1.7	3.3	8.5	29.1	131.4	678.4
1999.....	30.0	4.9	0.7	0.2	0.3	0.6	1.8	3.0	8.4	28.0	130.0	686.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic black, both sexes												
2007 5/.....	13.3	10.6	0.9	*	0.4	1.7	3.2	8.4	17.3	44.2	129.2	405.8
2006 5/.....	13.8	12.9	1.2	*	0.6	1.4	3.3	8.9	16.2	45.3	140.2	437.7
2005 5/.....	15.1	14.3	1.0	*	0.5	1.4	3.8	9.9	20.1	52.9	150.4	471.3
2004 5/.....	15.1	14.0	1.2	*	0.6	1.2	4.0	9.2	18.9	50.0	156.7	509.2
2003 5/.....	15.6	13.3	0.9	*	0.6	1.9	4.5	10.2	20.8	54.4	156.4	522.4
2002.....	16.0	13.1	1.5	*	0.6	1.6	4.5	8.9	21.4	50.5	167.3	577.4
2001.....	15.9	16.9	1.0	*	0.8	1.7	4.9	9.9	19.0	48.7	169.1	581.5
2000.....	16.6	17.2	1.2	*	0.8	1.8	4.5	10.1	22.1	59.9	170.8	595.0
1999.....	16.5	19.3	1.7	*	0.9	2.0	5.4	10.7	20.7	54.0	156.6	628.9
Non-Hispanic black, male												
2007 5/.....	13.5	11.6	*	*	*	2.0	3.9	11.3	22.4	60.5	174.1	480.8
2006 5/.....	13.5	15.7	*	*	0.7	1.4	3.7	11.6	21.8	59.3	180.5	509.4
2005 5/.....	15.0	14.0	*	*	0.7	1.7	4.8	13.0	27.7	71.0	190.6	530.0
2004 5/.....	15.4	15.8	*	*	*	1.6	4.9	12.0	24.7	72.7	218.2	606.6
2003 5/.....	16.2	15.2	*	*	0.7	2.3	5.4	13.0	29.4	78.5	219.9	639.3
2002.....	15.8	13.9	1.6	*	*	1.8	5.2	11.5	29.3	66.0	230.6	693.6
2001.....	16.3	16.6	*	*	0.8	2.0	5.9	13.5	26.4	69.7	242.5	716.1
2000.....	17.1	18.8	1.7	*	0.7	2.3	5.7	14.0	30.1	83.4	233.9	750.3
1999.....	16.8	19.1	2.0	*	0.8	2.3	6.6	15.1	27.2	77.0	213.3	799.9
Non-Hispanic black, female												
2007 5/.....	13.1	9.6	*	*	*	1.4	2.6	6.0	13.1	32.5	103.7	374.7
2006 5/.....	14.0	10.0	*	*	*	1.4	3.0	6.6	11.7	35.2	117.4	408.4
2005 5/.....	15.3	14.6	*	*	*	1.2	2.9	7.3	14.0	40.0	127.6	447.7
2004 5/.....	14.8	12.2	1.7	*	0.7	0.8	3.3	6.9	14.3	33.8	122.0	470.6
2003 5/.....	15.0	11.3	*	*	*	1.5	3.7	7.8	13.9	37.2	120.6	477.1
2002.....	16.1	12.3	*	*	*	1.4	3.9	6.6	15.1	39.5	131.5	533.0
2001.....	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	127.9	531.7
2000.....	16.3	15.5	*	*	0.8	1.2	3.5	6.8	15.6	43.3	135.3	537.3
1999.....	16.3	19.6	*	*	0.9	1.7	4.4	6.9	15.5	37.7	124.8	565.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other acute lower respiratory infections (J20-J22,U04)

All origins 3/, both sexes

2007.....	0.1	1.1	*	*	*	*	*	*	*	0.1	0.3	1.5
2006.....	0.1	1.3	*	*	*	*	*	*	*	0.1	0.5	2.0
2005.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.3
2004.....	0.1	1.5	*	*	*	*	*	*	*	0.1	0.4	2.6
2003.....	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002.....	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000.....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999.....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0

All origins 3/, male

2007.....	0.1	1.5	*	*	*	*	*	*	*	*	*	*
2006.....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	1.8
2005.....	0.1	1.4	*	*	*	*	*	*	*	*	0.8	3.1
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002.....	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0

All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	*	*	*	1.7
2006.....	0.1	1.1	*	*	*	*	*	*	*	*	0.5	2.1
2005.....	0.2	1.0	*	*	*	*	*	*	*	*	0.5	3.5
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.5	2.7
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999.....	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8



















Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified acute lower respiratory infections (J22,U04)

All origins 3/, both sexes

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*	0.4
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	0.9
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.2
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	0.3	1.3
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.2
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.4

All origins 3/, male

2007.....	*	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*

All origins 3/, female

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.1
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.1
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.8



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified acute lower respiratory infections (J22,U04)

Non-Hispanic 4/, both sexes

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*	0.4
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.0
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.5
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.2
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	0.3	1.4
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	0.2	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.2
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	0.2	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.4

Non-Hispanic 4/, male

2007.....	*	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.1
2005.....	0.1	*	*	*	*	*	*	*	*	*	*	*	1.5
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.4
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*	1.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.1
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	2.8





Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes

2007.....	42.4	1.0	0.3	0.3	0.4	0.6	1.8	9.5	39.1	148.1	368.9	596.1
2006.....	41.6	0.7	0.3	0.3	0.4	0.6	1.9	9.1	39.2	149.3	363.4	589.1
2005.....	44.2	0.8	0.3	0.3	0.4	0.6	2.0	9.4	42.0	160.5	385.6	637.2
2004.....	41.5	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.4	153.8	366.7	601.7
2003.....	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002.....	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0

All origins 3/, male

2007.....	41.2	1.2	0.4	0.4	0.4	0.7	1.8	9.8	42.7	165.2	441.2	758.7
2006.....	40.2	0.9	0.3	0.4	0.4	0.7	1.8	9.2	41.9	163.2	437.3	760.8
2005.....	42.8	*	0.4	0.3	0.4	0.6	1.9	9.7	45.0	180.7	463.7	819.9
2004.....	40.6	1.2	0.4	0.4	0.5	0.7	2.0	8.8	43.1	172.1	445.6	811.1
2003.....	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002.....	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001.....	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000.....	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999.....	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7

All origins 3/, female

2007.....	43.6	*	0.3	0.2	0.3	0.6	1.9	9.1	35.7	133.6	319.1	518.7
2006.....	43.0	*	*	0.2	0.3	0.6	2.0	8.9	36.6	137.5	312.8	508.7
2005.....	45.5	*	0.3	0.2	0.3	0.6	2.1	9.0	39.2	143.4	332.5	553.3
2004.....	42.5	*	*	0.2	0.3	0.5	2.1	8.1	38.0	138.4	313.6	507.4
2003.....	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002.....	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001.....	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000.....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999.....	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Hispanic, both sexes

2007.....	7.8	*	*	*	0.3	0.4	0.6	3.0	11.0	45.1	164.8	357.2
2006.....	7.5	*	*	*	*	0.3	0.7	2.6	12.0	45.5	153.5	374.4
2005.....	8.1	*	*	*	*	0.3	0.8	3.5	12.5	50.6	171.9	414.6
2004.....	7.5	*	*	*	*	0.4	1.1	3.1	12.2	51.7	158.7	391.5
2003.....	8.0	*	*	*	0.3	0.4	1.0	3.5	12.7	53.3	171.0	460.2
2002.....	7.9	*	*	*	*	0.4	1.2	3.5	13.9	53.8	176.0	457.3
2001.....	7.7	*	*	*	*	0.3	1.0	3.6	14.0	55.8	169.5	480.2
2000.....	7.6	*	*	*	*	0.3	1.1	3.3	14.3	53.1	181.9	481.1
1999.....	8.4	*	*	*	*	0.5	1.6	3.9	14.6	62.6	194.8	536.0

Hispanic, male

2007.....	8.1	*	*	*	*	0.5	0.6	3.5	13.4	60.2	217.1	461.5
2006.....	7.5	*	*	*	*	*	0.7	2.8	14.0	56.6	196.3	484.9
2005.....	8.3	*	*	*	*	*	1.0	3.6	15.8	64.5	234.2	526.8
2004.....	7.6	*	*	*	*	0.5	1.4	2.8	13.7	68.4	206.4	518.4
2003.....	8.2	*	*	*	*	*	1.0	3.2	16.6	68.1	231.2	646.5
2002.....	8.1	*	*	*	*	*	1.0	3.8	17.5	69.2	243.3	602.4
2001.....	7.8	*	*	*	*	*	0.7	3.2	16.1	75.5	224.0	676.1
2000.....	8.0	*	*	*	*	*	0.9	3.4	18.2	72.4	250.3	671.1
1999.....	8.9	*	*	*	*	*	1.4	3.6	17.6	81.9	272.7	836.8

Hispanic, female

2007.....	7.4	*	*	*	*	*	*	2.4	8.9	32.8	128.0	300.6
2006.....	7.5	*	*	*	*	*	0.7	2.5	10.1	36.5	123.5	315.7
2005.....	7.8	*	*	*	*	*	*	3.5	9.5	39.3	128.6	355.8
2004.....	7.4	*	*	*	*	*	0.7	3.4	10.9	38.3	125.7	326.3
2003.....	7.7	*	*	*	*	*	1.0	3.8	9.3	41.5	129.6	365.6
2002.....	7.6	*	*	*	*	*	1.4	3.1	10.6	41.5	129.8	385.5
2001.....	7.5	*	*	*	*	*	1.2	4.1	12.1	40.3	132.7	384.4
2000.....	7.2	*	*	*	*	*	1.3	3.3	10.8	38.0	136.0	387.8
1999.....	8.0	*	*	*	*	*	1.8	4.2	12.0	47.5	142.9	391.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 4/, both sexes

2007.....	48.5	1.0	0.4	0.3	0.4	0.7	2.1	10.2	41.5	156.1	381.7	608.3
2006.....	47.5	0.8	0.3	0.3	0.4	0.7	2.1	9.8	41.4	157.1	375.9	599.3
2005.....	50.2	0.8	0.4	0.3	0.4	0.7	2.2	10.0	44.4	168.5	397.6	647.1
2004.....	47.0	0.8	0.3	0.3	0.5	0.7	2.2	9.0	42.7	160.9	377.8	610.3
2003.....	49.0	0.8	0.3	0.3	0.5	0.8	2.3	9.1	45.6	170.6	393.3	641.2
2002.....	48.6	1.1	0.4	0.4	0.5	0.8	2.4	9.2	44.6	170.1	396.0	643.5
2001.....	48.4	1.1	0.3	0.3	0.5	0.8	2.3	8.9	46.3	174.8	388.6	649.4
2000.....	48.3	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.3	176.1	393.8	653.2
1999.....	49.4	1.0	0.4	0.4	0.6	0.9	2.0	8.9	49.8	183.5	405.1	648.0

Non-Hispanic 4/, male

2007.....	47.3	*	0.4	0.4	0.5	0.8	2.1	10.6	45.1	173.1	455.3	775.3
2006.....	46.1	*	0.3	0.4	0.5	0.8	2.0	9.9	44.1	171.0	451.8	775.2
2005.....	48.8	*	0.5	0.3	0.5	0.7	2.1	10.4	47.3	188.9	476.6	833.9
2004.....	46.2	*	0.4	0.4	0.6	0.8	2.1	9.5	45.3	178.9	458.4	824.6
2003.....	48.1	*	0.5	0.4	0.6	0.9	2.1	9.7	48.7	191.1	476.2	874.6
2002.....	48.4	1.2	0.6	0.4	0.6	0.9	2.4	9.6	47.3	191.9	491.2	906.0
2001.....	48.0	*	0.4	0.3	0.6	0.8	2.2	9.3	49.0	198.1	485.7	924.8
2000.....	48.7	*	0.4	0.4	0.6	0.9	2.1	9.4	49.7	201.9	497.6	978.0
1999.....	50.8	*	0.4	0.5	0.7	0.8	1.9	9.0	54.5	217.4	521.9	1,000.2

Non-Hispanic 4/, female

2007.....	49.6	*	0.4	0.2	0.3	0.6	2.1	9.9	38.0	141.7	331.1	529.5
2006.....	48.8	*	*	0.2	0.3	0.6	2.2	9.7	38.8	145.3	324.0	517.5
2005.....	51.5	*	0.3	0.2	0.3	0.7	2.3	9.6	41.6	151.3	344.0	561.8
2004.....	47.8	*	*	0.2	0.4	0.6	2.3	8.5	40.2	145.6	323.6	514.5
2003.....	49.9	*	*	0.2	0.4	0.7	2.5	8.6	42.7	153.3	338.0	538.6
2002.....	48.9	*	*	0.3	0.4	0.8	2.4	8.7	42.0	151.8	332.9	530.3
2001.....	48.7	*	*	0.2	0.4	0.8	2.4	8.5	43.8	155.4	324.9	534.8
2000.....	48.0	*	*	0.3	0.4	0.7	2.3	8.7	43.2	154.8	326.2	521.9
1999.....	48.0	*	*	0.2	0.5	0.9	2.1	8.7	45.4	155.7	329.3	507.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic white, both sexes

2007 5/.....	56.9	*	*	0.2	0.2	0.5	1.9	10.2	44.1	168.4	405.9	640.6
2006 5/.....	55.6	*	*	0.1	0.2	0.5	1.9	9.9	43.7	169.7	398.4	629.8
2005 5/.....	58.6	*	0.2	0.1	0.2	0.5	2.0	9.8	46.9	181.1	420.9	679.0
2004 5/.....	54.9	*	*	0.2	0.3	0.5	2.0	8.8	44.8	173.3	399.6	640.0
2003 5/.....	57.0	*	*	0.2	0.3	0.6	2.0	9.0	47.8	182.6	414.9	671.6
2002.....	56.4	0.9	0.2	0.2	0.4	0.6	2.0	9.0	46.7	181.6	416.3	673.1
2001.....	55.9	*	*	0.2	0.3	0.5	1.9	8.7	48.6	185.9	408.6	676.9
2000.....	55.7	*	*	0.2	0.4	0.6	1.8	8.6	48.2	186.5	413.2	680.8
1999.....	56.6	*	0.2	0.2	0.4	0.6	1.6	8.5	51.8	193.6	422.4	675.6

Non-Hispanic white, male

2007 5/.....	54.7	*	*	0.2	0.3	0.5	1.8	10.7	47.0	182.9	474.3	806.5
2006 5/.....	53.0	*	*	0.2	0.2	0.5	1.8	9.8	45.4	180.5	469.7	804.2
2005 5/.....	56.0	*	*	*	0.3	0.5	1.9	10.1	48.9	198.7	493.9	861.9
2004 5/.....	53.0	*	*	0.2	0.3	0.5	1.8	9.2	46.3	188.5	476.6	852.5
2003 5/.....	54.9	*	*	0.2	0.4	0.5	1.8	9.5	50.0	200.2	491.0	903.6
2002.....	55.1	*	*	0.2	0.4	0.6	2.0	9.3	48.3	200.4	506.7	935.4
2001.....	54.6	*	*	0.2	0.3	0.5	1.9	9.1	50.5	206.1	500.9	951.5
2000.....	55.1	*	*	0.2	0.4	0.5	1.7	8.9	50.8	208.8	513.6	1,008.6
1999.....	57.3	*	*	0.3	0.5	0.5	1.5	8.5	55.6	224.9	534.9	1,033.5

Non-Hispanic white, female

2007 5/.....	59.1	*	*	*	0.2	0.5	2.0	9.8	41.4	155.8	358.0	561.9
2006 5/.....	58.0	*	*	*	0.2	0.5	2.0	9.9	42.1	160.4	348.9	548.5
2005 5/.....	61.1	*	*	*	0.2	0.6	2.2	9.5	44.9	165.8	370.7	595.3
2004 5/.....	56.7	*	*	*	0.2	0.5	2.1	8.4	43.5	160.1	347.2	544.8
2003 5/.....	59.0	*	*	*	0.3	0.6	2.2	8.5	45.7	167.6	363.5	569.5
2002.....	57.7	*	*	*	0.4	0.6	2.1	8.6	45.1	165.5	355.7	559.8
2001.....	57.2	*	*	*	0.3	0.6	2.0	8.3	46.8	168.8	347.3	562.7
2000.....	56.2	*	*	0.2	0.3	0.6	1.9	8.3	45.8	167.6	347.2	548.7
1999.....	55.9	*	*	*	0.4	0.6	1.8	8.5	48.3	167.4	348.8	532.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic black, both sexes

2007 5/.....	20.5	*	1.2	1.0	1.0	1.7	3.6	13.1	35.7	105.5	230.8	342.9
2006 5/.....	20.3	*	*	1.1	1.2	2.0	4.1	12.3	38.1	102.9	229.9	337.4
2005 5/.....	21.8	*	1.2	0.8	1.2	1.8	4.0	14.2	40.3	116.4	244.8	369.2
2004 5/.....	19.9	*	0.8	1.0	1.3	1.9	4.0	13.0	39.1	103.9	225.9	335.9
2003 5/.....	20.9	*	1.0	1.0	1.3	2.0	4.4	12.9	41.3	115.3	238.5	354.9
2002.....	21.4	*	1.2	1.1	1.3	2.0	5.2	13.1	41.6	115.0	250.5	370.8
2001.....	21.1	*	0.9	0.8	1.3	2.5	4.9	12.7	41.3	120.4	236.3	379.3
2000.....	21.5	*	0.9	1.1	1.4	2.2	4.9	14.5	44.7	123.2	236.5	376.2
1999.....	22.6	*	*	1.0	1.4	2.7	4.7	14.1	46.7	129.9	265.3	377.0

Non-Hispanic black, male

2007 5/.....	22.9	*	*	1.3	1.4	2.1	3.9	12.8	45.6	141.7	352.9	522.7
2006 5/.....	22.7	*	*	1.3	1.4	2.4	4.0	13.6	48.8	139.7	345.1	522.3
2005 5/.....	24.7	*	1.6	1.0	1.5	2.0	4.0	15.5	49.9	160.1	381.2	596.6
2004 5/.....	22.8	*	*	1.4	1.6	2.6	4.2	14.4	51.4	142.5	342.3	569.0
2003 5/.....	24.2	*	*	1.3	1.6	2.9	4.1	13.7	51.5	158.0	386.7	608.4
2002.....	24.9	*	1.6	1.2	2.0	2.3	5.7	14.6	53.1	160.0	395.7	653.1
2001.....	24.5	*	*	1.0	1.7	2.8	4.7	13.5	50.4	170.1	385.6	682.6
2000.....	25.1	*	*	1.3	1.8	2.8	4.9	15.2	55.3	178.9	372.4	696.3
1999.....	27.0	*	*	1.4	1.9	2.7	4.7	15.6	60.4	186.1	438.4	707.2

Non-Hispanic black, female

2007 5/.....	18.4	*	*	*	0.7	1.3	3.3	13.3	27.8	79.7	161.4	268.4
2006 5/.....	18.1	*	*	0.8	0.9	1.6	4.2	11.1	29.5	76.6	164.6	262.0
2005 5/.....	19.1	*	*	*	0.9	1.6	3.9	13.1	32.5	85.1	167.5	277.7
2004 5/.....	17.2	*	*	0.6	1.0	1.3	3.9	11.9	29.3	76.3	160.2	243.8
2003 5/.....	17.9	*	*	*	1.1	1.2	4.7	12.2	33.2	85.0	154.8	256.7
2002.....	18.2	*	*	1.1	0.7	1.7	4.8	11.8	32.3	83.0	168.5	263.0
2001.....	18.0	*	*	0.7	1.0	2.2	5.0	11.9	34.0	85.3	152.5	266.9
2000.....	18.2	*	*	0.8	1.0	1.6	4.9	13.8	36.1	84.1	159.9	257.3
1999.....	18.6	*	*	0.6	0.9	2.7	4.7	12.8	35.8	90.3	168.4	253.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, chronic and unspecified (J40-J42)

All origins 3/, both sexes

2007.....	0.2	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.8
2006.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.0
2005.....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	7.4
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.5
2003.....	0.3	0.5	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002.....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001.....	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000.....	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999.....	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7

All origins 3/, male

2007.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.0
2006.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	5.9
2005.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.9	7.3
2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	2.2	7.0
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999.....	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9

All origins 3/, female

2007.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.2	4.8
2006.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	6.0
2005.....	0.3	*	*	*	*	*	*	*	0.2	0.7	1.4	7.4
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.3
2003.....	0.3	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic 4/, both sexes

2007.....	0.2	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.8
2006.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.0
2005.....	0.3	0.6	*	*	*	*	*	0.1	0.3	0.7	1.6	7.3
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.6
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002.....	0.4	0.6	*	*	*	*	*	0.1	0.2	0.8	2.2	8.0
2001.....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.1	9.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.7	11.8
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	10.7

Non-Hispanic 4/, male

2007.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	4.9
2006.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.1
2005.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	7.4
2004.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.6	7.0
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.2	6.9
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.3	9.3
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.6	12.8
1999.....	0.4	*	*	*	*	*	*	*	0.5	1.1	3.3	11.9

Non-Hispanic 4/, female

2007.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.3	4.8
2006.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	6.0
2005.....	0.4	*	*	*	*	*	*	*	0.3	0.7	1.3	7.2
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.4
2003.....	0.4	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	7.6
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.4	10.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic white, both sexes

2007 5/.....	0.3	*	*	*	*	*	*	*	0.1	0.2	0.5	1.5	4.9
2006 5/.....	0.3	*	*	*	*	*	*	*	0.1	0.2	0.5	1.6	6.3
2005 5/.....	0.4	*	*	*	*	*	*	*	0.1	0.2	0.7	1.6	7.8
2004 5/.....	0.3	*	*	*	*	*	*	*	0.1	0.2	0.6	1.6	6.7
2003 5/.....	0.4	*	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.5
2002.....	0.4	*	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.4
2001.....	0.4	*	*	*	*	*	*	*	0.1	0.3	0.8	2.2	9.7
2000.....	0.5	*	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.6
1999.....	0.5	*	*	*	*	*	*	*	0.1	0.5	1.0	2.8	11.2

Non-Hispanic white, male

2007 5/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	5.1
2006 5/.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	6.5
2005 5/.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	7.5
2004 5/.....	0.3	*	*	*	*	*	*	*	0.3	0.7	1.7	7.2
2003 5/.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	7.1
2002.....	0.4	*	*	*	*	*	*	*	0.2	1.0	2.6	9.3
2001.....	0.3	*	*	*	*	*	*	*	0.3	1.0	2.3	9.6
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.8	13.8
1999.....	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4

Non-Hispanic white, female

2007 5/.....	0.3	*	*	*	*	*	*	*	0.2	0.6	1.4	4.9
2006 5/.....	0.4	*	*	*	*	*	*	*	0.2	0.5	1.6	6.2
2005 5/.....	0.4	*	*	*	*	*	*	*	0.2	0.8	1.4	7.9
2004 5/.....	0.4	*	*	*	*	*	*	*	0.2	0.5	1.6	6.5
2003 5/.....	0.4	*	*	*	*	*	*	*	*	0.7	1.8	7.6
2002.....	0.5	*	*	*	*	*	*	*	0.2	0.7	2.1	8.0
2001.....	0.5	*	*	*	*	*	*	*	0.2	0.6	2.1	9.7
2000.....	0.6	*	*	*	*	*	*	*	0.3	0.8	2.1	12.2
1999.....	0.6	*	*	*	*	*	*	*	0.4	0.8	2.5	10.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Emphysema (J43)

##### All origins 3/, both sexes

2007.....	4.2	*	*	*	*	*	0.1	1.1	4.9	17.0	37.1	45.5
2006.....	4.2	*	*	*	*	*	0.2	1.0	4.8	17.3	36.8	45.3
2005.....	4.7	*	*	*	*	*	0.2	1.0	5.5	20.1	41.4	51.9
2004.....	4.6	*	*	*	*	*	0.2	1.0	5.4	20.5	41.0	49.9
2003.....	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002.....	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001.....	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000.....	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999.....	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1

##### All origins 3/, male

2007.....	4.4	*	*	*	*	*	0.2	1.4	5.7	20.1	46.2	61.6
2006.....	4.3	*	*	*	*	*	0.2	1.3	5.3	19.6	45.4	63.7
2005.....	5.0	*	*	*	*	*	0.2	1.3	6.7	23.9	52.4	71.5
2004.....	4.9	*	*	*	*	*	0.2	1.2	6.2	24.3	52.2	74.4
2003.....	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002.....	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001.....	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000.....	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999.....	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4

##### All origins 3/, female

2007.....	4.0	*	*	*	*	*	*	0.8	4.1	14.4	30.8	37.9
2006.....	4.1	*	*	*	*	*	0.2	0.9	4.4	15.4	30.9	36.7
2005.....	4.5	*	*	*	*	*	0.1	0.8	4.5	16.9	33.9	42.8
2004.....	4.4	*	*	*	*	*	0.1	0.8	4.7	17.4	33.5	38.9
2003.....	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002.....	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001.....	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000.....	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999.....	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Emphysema (J43)

Hispanic, both sexes

2007.....	0.6	*	*	*	*	*	*	*	1.1	5.1	13.7	21.5
2006.....	0.6	*	*	*	*	*	*	*	1.7	4.0	15.6	20.7
2005.....	0.7	*	*	*	*	*	*	*	1.3	5.4	16.6	24.8
2004.....	0.7	*	*	*	*	*	*	*	0.9	5.0	16.1	30.0
2003.....	0.8	*	*	*	*	*	*	*	1.4	6.0	20.0	34.7
2002.....	0.8	*	*	*	*	*	*	*	1.5	6.7	18.8	36.6
2001.....	0.8	*	*	*	*	*	*	*	1.4	6.6	20.4	39.4
2000.....	0.8	*	*	*	*	*	*	*	1.5	6.0	22.3	31.2
1999.....	0.9	*	*	*	*	*	*	*	1.3	7.1	23.5	44.2

Hispanic, male

2007.....	0.8	*	*	*	*	*	*	*	1.9	7.2	20.0	32.5
2006.....	0.8	*	*	*	*	*	*	*	2.2	6.2	21.8	31.6
2005.....	0.8	*	*	*	*	*	*	*	1.8	8.6	22.7	38.6
2004.....	0.7	*	*	*	*	*	*	*	*	6.5	20.4	47.6
2003.....	0.9	*	*	*	*	*	*	*	2.1	7.6	29.1	53.9
2002.....	0.8	*	*	*	*	*	*	*	*	9.2	26.7	53.5
2001.....	0.9	*	*	*	*	*	*	*	*	9.2	34.0	60.9
2000.....	0.9	*	*	*	*	*	*	*	*	9.1	32.5	58.4
1999.....	1.0	*	*	*	*	*	*	*	*	11.2	36.0	77.6

Hispanic, female

2007.....	0.5	*	*	*	*	*	*	*	*	3.4	9.3	15.4
2006.....	0.5	*	*	*	*	*	*	*	*	*	11.3	15.0
2005.....	0.6	*	*	*	*	*	*	*	*	2.9	12.3	17.6
2004.....	0.6	*	*	*	*	*	*	*	*	3.7	13.1	20.9
2003.....	0.7	*	*	*	*	*	*	*	*	4.7	13.7	25.0
2002.....	0.7	*	*	*	*	*	*	*	*	4.7	13.5	28.2
2001.....	0.6	*	*	*	*	*	*	*	*	4.6	11.2	28.9
2000.....	0.6	*	*	*	*	*	*	*	*	3.6	15.5	*
1999.....	0.7	*	*	*	*	*	*	*	*	4.0	15.3	28.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Emphysema (J43)

##### Non-Hispanic 4/, both sexes

2007.....	4.9	*	*	*	*	*	0.2	1.2	5.2	17.9	38.6	46.7
2006.....	4.8	*	*	*	*	*	0.2	1.1	5.1	18.3	38.1	46.5
2005.....	5.4	*	*	*	*	*	0.2	1.1	5.9	21.2	42.8	53.1
2004.....	5.3	*	*	*	*	*	0.2	1.1	5.8	21.7	42.4	50.7
2003.....	5.8	*	*	*	*	*	0.2	1.2	6.6	23.3	47.3	55.4
2002.....	6.1	*	*	*	*	*	0.2	1.2	6.8	25.1	49.3	60.1
2001.....	6.4	*	*	*	*	*	0.2	1.2	7.6	27.5	51.7	62.8
2000.....	6.7	*	*	*	*	*	0.2	1.3	7.9	29.1	53.8	64.3
1999.....	7.1	*	*	*	*	*	0.2	1.3	9.0	32.0	58.1	62.5

##### Non-Hispanic 4/, male

2007.....	5.1	*	*	*	*	*	0.2	1.5	6.0	21.1	47.9	63.2
2006.....	4.9	*	*	*	*	*	0.3	1.3	5.6	20.6	46.9	65.3
2005.....	5.7	*	*	*	*	*	0.3	1.4	7.0	25.0	54.1	73.1
2004.....	5.6	*	*	*	*	*	0.2	1.3	6.6	25.6	54.0	75.5
2003.....	6.0	*	*	*	*	*	0.2	1.5	7.3	26.8	59.5	80.4
2002.....	6.5	*	*	*	*	*	0.3	1.5	7.7	29.3	65.2	91.3
2001.....	6.8	*	*	*	*	*	0.3	1.5	8.4	32.1	66.6	102.7
2000.....	7.1	*	*	*	*	*	0.2	1.6	8.5	34.3	70.6	105.9
1999.....	7.8	*	*	*	*	*	0.2	1.5	10.0	39.2	77.8	106.1

##### Non-Hispanic 4/, female

2007.....	4.6	*	*	*	*	*	*	0.9	4.4	15.2	32.2	39.0
2006.....	4.7	*	*	*	*	*	0.2	0.9	4.6	16.4	32.0	37.8
2005.....	5.1	*	*	*	*	*	0.1	0.8	4.8	18.0	35.1	44.0
2004.....	5.0	*	*	*	*	*	0.2	0.9	5.1	18.4	34.6	39.6
2003.....	5.6	*	*	*	*	*	0.2	0.9	6.0	20.3	39.1	44.5
2002.....	5.7	*	*	*	*	*	0.1	1.0	6.0	21.7	38.8	46.6
2001.....	6.1	*	*	*	*	*	0.1	1.0	7.0	23.6	41.9	46.3
2000.....	6.3	*	*	*	*	*	0.2	1.1	7.3	24.8	42.9	47.5
1999.....	6.5	*	*	*	*	*	0.2	1.1	8.1	26.1	45.3	45.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Empysemata (J43)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	5.8	*	*	*	*	*	0.2	1.3	5.6	19.5	41.1	49.5
2006 5/.....	5.7	*	*	*	*	*	0.2	1.2	5.5	20.0	40.5	49.2
2005 5/.....	6.3	*	*	*	*	*	0.2	1.1	6.3	23.0	45.2	55.9
2004 5/.....	6.2	*	*	*	*	*	0.2	1.1	6.2	23.7	45.2	53.4
2003 5/.....	6.8	*	*	*	*	*	0.2	1.2	7.1	25.2	49.9	57.9
2002.....	7.1	*	*	*	*	*	0.2	1.3	7.3	27.0	52.3	63.1
2001.....	7.5	*	*	*	*	*	0.2	1.3	8.2	29.6	54.9	65.9
2000.....	7.8	*	*	*	*	*	0.2	1.3	8.4	31.2	57.0	66.9
1999.....	8.3	*	*	*	*	*	0.2	1.3	9.7	34.5	61.2	65.2
<b>Non-Hispanic white, male</b>												
2007 5/.....	5.9	*	*	*	*	*	0.2	1.6	6.3	22.5	49.7	65.6
2006 5/.....	5.7	*	*	*	*	*	0.3	1.4	5.7	22.0	48.7	68.4
2005 5/.....	6.5	*	*	*	*	*	0.2	1.4	7.3	26.4	55.8	75.5
2004 5/.....	6.5	*	*	*	*	*	0.2	1.2	6.8	27.2	56.5	78.2
2003 5/.....	6.9	*	*	*	*	*	0.2	1.5	7.6	28.5	61.3	81.6
2002.....	7.4	*	*	*	*	*	0.3	1.5	8.0	30.6	67.7	94.3
2001.....	7.8	*	*	*	*	*	0.3	1.5	8.7	33.8	69.6	106.6
2000.....	8.1	*	*	*	*	*	0.2	1.4	8.9	35.8	73.9	107.3
1999.....	8.9	*	*	*	*	*	0.2	1.4	10.4	41.1	80.6	110.0
<b>Non-Hispanic white, female</b>												
2007 5/.....	5.6	*	*	*	*	*	*	0.9	4.9	17.0	35.0	41.9
2006 5/.....	5.7	*	*	*	*	*	0.2	1.1	5.2	18.2	34.8	40.3
2005 5/.....	6.1	*	*	*	*	*	*	0.9	5.4	20.0	37.8	47.0
2004 5/.....	6.0	*	*	*	*	*	0.2	0.9	5.6	20.6	37.6	42.3
2003 5/.....	6.7	*	*	*	*	*	0.2	1.0	6.7	22.5	42.3	47.4
2002.....	6.8	*	*	*	*	*	*	1.1	6.7	24.0	41.9	49.7
2001.....	7.2	*	*	*	*	*	0.1	1.0	7.7	26.1	45.2	48.9
2000.....	7.4	*	*	*	*	*	0.2	1.1	8.0	27.3	45.8	50.6
1999.....	7.7	*	*	*	*	*	0.2	1.2	9.0	29.0	48.5	47.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Emphysema (J43)

Non-Hispanic black, both sexes

2007 5/.....	1.9	*	*	*	*	*	*	1.3	4.0	10.7	23.1	24.3
2006 5/.....	1.9	*	*	*	*	*	0.4	1.0	4.3	10.3	22.6	24.2
2005 5/.....	2.2	*	*	*	*	*	*	1.4	4.6	12.9	26.7	27.3
2004 5/.....	1.9	*	*	*	*	*	*	1.3	4.7	11.0	22.5	27.9
2003 5/.....	2.2	*	*	*	*	*	*	1.2	5.3	13.2	27.7	31.2
2002.....	2.3	*	*	*	*	*	0.5	1.2	4.6	16.2	28.2	32.1
2001.....	2.2	*	*	*	*	*	*	1.4	5.6	15.6	23.6	35.9
2000.....	2.5	*	*	*	*	*	*	2.0	6.1	17.1	27.0	37.5
1999.....	2.7	*	*	*	*	*	0.4	1.9	6.3	17.3	31.8	37.8

Non-Hispanic black, male

2007 5/.....	2.5	*	*	*	*	*	*	1.8	6.1	16.5	38.0	47.1
2006 5/.....	2.4	*	*	*	*	*	*	1.6	6.8	14.3	36.7	40.3
2005 5/.....	2.8	*	*	*	*	*	*	2.1	7.5	19.4	42.9	50.4
2004 5/.....	2.6	*	*	*	*	*	*	2.0	7.1	17.4	37.1	57.9
2003 5/.....	2.8	*	*	*	*	*	*	1.7	7.5	18.6	46.7	71.4
2002.....	3.1	*	*	*	*	*	*	1.8	6.7	25.3	49.5	65.6
2001.....	2.8	*	*	*	*	*	*	1.6	7.9	23.0	40.2	69.3
2000.....	3.3	*	*	*	*	*	*	3.0	8.2	27.4	41.3	89.2
1999.....	3.6	*	*	*	*	*	*	3.0	9.1	29.3	55.7	73.7

Non-Hispanic black, female

2007 5/.....	1.4	*	*	*	*	*	*	0.9	2.3	6.6	14.7	14.8
2006 5/.....	1.4	*	*	*	*	*	*	*	2.3	7.5	14.7	17.6
2005 5/.....	1.6	*	*	*	*	*	*	0.8	2.3	8.2	17.5	18.0
2004 5/.....	1.3	*	*	*	*	*	*	*	2.7	6.4	14.3	16.1
2003 5/.....	1.6	*	*	*	*	*	*	*	3.5	9.4	16.9	15.7
2002.....	1.6	*	*	*	*	*	*	*	2.8	9.8	16.1	19.2
2001.....	1.7	*	*	*	*	*	*	1.1	3.7	10.4	14.3	23.5
2000.....	1.8	*	*	*	*	*	*	1.3	4.5	9.8	18.9	18.3
1999.....	1.8	*	*	*	*	*	*	1.0	4.0	8.9	18.4	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Asthma (J45-J46)

##### All origins 3/, both sexes

2007.....	1.1	*	0.2	0.3	0.3	0.5	0.7	1.2	1.4	2.1	4.4	12.0
2006.....	1.2	*	0.2	0.2	0.3	0.5	0.9	1.3	1.6	2.3	4.8	12.3
2005.....	1.3	*	0.2	0.2	0.3	0.5	0.8	1.4	1.7	2.5	5.4	14.5
2004.....	1.3	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.1
2003.....	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002.....	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001.....	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000.....	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999.....	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3

##### All origins 3/, male

2007.....	0.9	*	0.2	0.3	0.4	0.6	0.6	1.0	1.2	1.5	3.2	7.1
2006.....	0.9	*	*	0.3	0.4	0.5	0.7	1.0	1.1	1.7	3.5	9.1
2005.....	0.9	*	0.3	0.3	0.4	0.5	0.7	1.0	1.2	1.8	4.1	8.7
2004.....	1.0	*	*	0.3	0.5	0.6	0.7	1.0	1.3	2.1	3.5	11.1
2003.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002.....	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001.....	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000.....	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999.....	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7

##### All origins 3/, female

2007.....	1.4	*	0.2	0.2	0.2	0.4	0.9	1.5	1.6	2.6	5.1	14.2
2006.....	1.5	*	*	0.1	0.3	0.4	1.0	1.6	2.0	2.9	5.7	13.9
2005.....	1.7	*	*	0.2	0.2	0.5	1.0	1.8	2.2	3.2	6.3	17.2
2004.....	1.6	*	*	0.2	0.3	0.4	1.0	1.6	2.2	3.2	6.3	15.4
2003.....	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002.....	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001.....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000.....	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999.....	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Asthma (J45-J46)</b>												
<b>Hispanic, both sexes</b>												
2007.....	0.5	*	*	*	*	0.3	0.3	0.7	1.3	1.7	4.6	9.7
2006.....	0.6	*	*	*	*	0.3	0.5	0.6	1.4	2.4	4.4	16.4
2005.....	0.6	*	*	*	*	*	0.5	0.8	1.6	2.9	4.8	13.7
2004.....	0.7	*	*	*	*	0.3	0.5	1.1	1.9	3.3	5.3	11.4
2003.....	0.7	*	*	*	*	0.3	0.5	1.6	2.2	3.4	6.1	14.5
2002.....	0.7	*	*	*	*	0.3	0.7	1.2	1.9	3.8	7.0	12.2
2001.....	0.7	*	*	*	*	*	0.6	1.4	2.4	3.8	7.9	13.8
2000.....	0.8	*	*	*	*	*	0.6	1.3	2.8	5.6	7.9	16.6
1999.....	0.9	*	*	*	*	0.3	0.9	1.6	2.4	5.8	10.4	17.5
<b>Hispanic, male</b>												
2007.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.6	*	*	*	*	*	*	1.4	2.2	*	*	*
2002.....	0.6	*	*	*	*	*	*	1.2	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	1.3	*	5.3	*	*
1999.....	0.7	*	*	*	*	*	0.9	1.4	*	*	*	*
<b>Hispanic, female</b>												
2007.....	0.6	*	*	*	*	*	*	*	*	*	5.5	11.0
2006.....	0.8	*	*	*	*	*	*	*	1.7	3.3	5.3	19.8
2005.....	0.7	*	*	*	*	*	*	*	2.1	3.4	5.6	13.0
2004.....	0.9	*	*	*	*	*	*	1.5	2.8	3.8	6.9	*
2003.....	0.9	*	*	*	*	*	*	1.8	2.1	4.1	6.3	17.2
2002.....	0.9	*	*	*	*	*	1.0	1.2	2.2	5.3	7.6	*
2001.....	1.0	*	*	*	*	*	*	2.0	2.9	3.9	9.4	18.6
2000.....	1.0	*	*	*	*	*	*	1.3	3.2	5.8	9.2	*
1999.....	1.2	*	*	*	*	*	1.0	1.9	3.1	7.1	13.2	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Asthma (J45-J46)

##### Non-Hispanic 4/, both sexes

2007.....	1.3	*	0.3	0.3	0.3	0.5	0.8	1.3	1.4	2.2	4.4	12.1
2006.....	1.3	*	0.2	0.3	0.3	0.5	0.9	1.4	1.6	2.3	4.8	12.1
2005.....	1.4	*	0.3	0.3	0.4	0.6	0.9	1.5	1.7	2.5	5.5	14.5
2004.....	1.4	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.2
2003.....	1.5	*	0.2	0.3	0.4	0.6	1.0	1.5	2.0	2.9	5.8	14.1
2002.....	1.6	*	0.3	0.3	0.5	0.6	1.1	1.5	2.0	3.1	6.3	14.7
2001.....	1.6	*	0.2	0.3	0.4	0.7	1.1	1.5	2.2	3.4	6.3	16.1
2000.....	1.7	*	0.2	0.3	0.5	0.7	1.1	1.6	2.2	3.9	6.8	16.6
1999.....	1.8	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.1	7.4	17.3

##### Non-Hispanic 4/, male

2007.....	0.9	*	*	0.4	0.4	0.6	0.7	1.0	1.2	1.5	3.2	7.1
2006.....	1.0	*	*	0.4	0.4	0.6	0.8	1.1	1.1	1.7	3.5	9.0
2005.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.1	1.2	1.7	4.1	8.4
2004.....	1.0	*	*	0.3	0.5	0.6	0.7	1.1	1.4	2.1	3.5	11.1
2003.....	1.1	*	0.4	0.4	0.5	0.7	0.8	1.1	1.5	2.0	4.5	10.7
2002.....	1.2	*	0.4	0.3	0.6	0.7	0.9	1.2	1.6	2.4	4.9	11.1
2001.....	1.1	*	*	0.3	0.4	0.6	0.9	1.2	1.5	2.6	4.4	12.2
2000.....	1.2	*	*	0.4	0.6	0.8	0.9	1.3	1.6	2.9	4.9	12.7
1999.....	1.3	*	*	0.4	0.6	0.7	0.8	1.1	1.8	3.1	5.6	12.6

##### Non-Hispanic 4/, female

2007.....	1.6	*	*	0.2	0.2	0.5	1.0	1.6	1.6	2.7	5.1	14.4
2006.....	1.6	*	*	0.2	0.3	0.5	1.1	1.7	2.0	2.8	5.7	13.6
2005.....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.2	3.2	6.3	17.3
2004.....	1.7	*	*	0.2	0.3	0.5	1.1	1.7	2.2	3.2	6.3	15.5
2003.....	1.9	*	*	0.2	0.4	0.5	1.2	1.9	2.4	3.6	6.7	15.6
2002.....	2.0	*	*	0.3	0.3	0.6	1.2	1.9	2.4	3.8	7.2	16.3
2001.....	2.1	*	*	0.2	0.4	0.7	1.2	1.9	2.8	4.1	7.5	17.7
2000.....	2.1	*	*	0.3	0.4	0.6	1.2	2.0	2.6	4.7	8.0	18.2
1999.....	2.2	*	*	0.2	0.4	0.7	1.2	2.2	3.0	4.9	8.6	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic white, both sexes												
2007 5/.....	1.1	*	*	0.1	0.2	0.4	0.6	0.9	1.0	1.8	4.0	11.9
2006 5/.....	1.1	*	*	0.1	0.2	0.3	0.7	1.0	1.1	1.9	4.5	11.8
2005 5/.....	1.2	*	*	0.1	0.2	0.4	0.7	1.1	1.3	1.9	4.9	14.5
2004 5/.....	1.2	*	*	0.1	0.2	0.3	0.7	0.9	1.3	2.3	4.6	13.8
2003 5/.....	1.3	*	*	0.1	0.2	0.4	0.7	1.0	1.5	2.4	5.3	14.1
2002.....	1.4	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.6	6.1	14.7
2001.....	1.4	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.9	5.8	15.7
2000.....	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.5	3.2	6.4	15.8
1999.....	1.5	*	*	0.2	0.4	0.4	0.7	1.2	1.9	3.4	7.1	16.5
Non-Hispanic white, male												
2007 5/.....	0.7	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.1	2.7	6.6
2006 5/.....	0.7	*	*	*	0.2	0.3	0.5	0.7	0.7	1.3	3.2	9.0
2005 5/.....	0.8	*	*	*	0.2	0.4	0.5	0.8	0.9	1.3	3.5	8.2
2004 5/.....	0.8	*	*	*	0.3	0.3	0.5	0.7	0.9	1.7	2.8	10.4
2003 5/.....	0.9	*	*	*	0.3	0.4	0.5	0.8	1.1	1.7	3.8	10.6
2002.....	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.1	1.8	4.5	10.8
2001.....	0.9	*	*	0.2	0.2	0.3	0.6	0.8	1.0	2.1	3.9	11.4
2000.....	1.0	*	*	0.2	0.3	0.5	0.6	0.9	1.0	2.3	4.6	11.1
1999.....	1.0	*	*	0.2	0.4	0.4	0.5	0.7	1.4	2.4	5.2	10.6
Non-Hispanic white, female												
2007 5/.....	1.4	*	*	*	*	0.4	0.7	1.1	1.2	2.4	4.9	14.4
2006 5/.....	1.5	*	*	*	*	0.3	0.8	1.3	1.6	2.4	5.4	13.2
2005 5/.....	1.7	*	*	*	0.2	0.5	0.8	1.3	1.7	2.5	5.9	17.4
2004 5/.....	1.6	*	*	*	0.2	0.4	0.8	1.1	1.7	2.9	5.9	15.3
2003 5/.....	1.7	*	*	*	0.2	0.5	0.9	1.3	2.0	2.9	6.3	15.7
2002.....	1.8	*	*	*	0.3	0.5	0.9	1.3	1.8	3.2	7.1	16.4
2001.....	1.9	*	*	*	0.2	0.4	0.8	1.4	2.1	3.6	7.1	17.5
2000.....	1.9	*	*	0.2	0.3	0.4	0.9	1.3	2.0	4.0	7.6	17.7
1999.....	2.1	*	*	*	0.3	0.5	0.9	1.6	2.3	4.2	8.3	18.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic black, both sexes												
2007 5/.....	2.3	*	1.0	0.9	0.9	1.4	2.2	4.0	4.4	5.1	6.5	11.8
2006 5/.....	2.5	*	*	1.0	1.1	1.7	2.5	4.0	5.2	5.5	6.1	12.7
2005 5/.....	2.7	*	1.1	0.8	1.1	1.6	2.4	4.3	5.5	7.2	9.9	12.0
2004 5/.....	2.7	*	*	0.9	1.2	1.7	2.5	4.6	5.8	5.8	9.4	15.3
2003 5/.....	2.8	*	0.8	0.9	1.2	1.7	2.8	4.9	5.2	7.2	9.3	11.9
2002.....	3.0	*	0.9	1.1	1.3	1.6	3.5	4.9	6.6	8.0	7.9	11.8
2001.....	3.1	*	*	0.8	1.2	2.2	3.3	4.7	6.9	7.7	10.7	16.5
2000.....	3.3	*	*	1.1	1.3	2.0	3.2	5.5	6.9	9.4	10.4	22.3
1999.....	3.3	*	*	1.0	1.3	2.4	3.0	5.4	7.0	9.8	10.6	20.7
Non-Hispanic black, male												
2007 5/.....	2.2	*	*	1.2	1.3	1.9	2.2	3.0	4.5	4.8	6.9	*
2006 5/.....	2.3	*	*	1.2	1.4	2.0	2.4	3.5	4.7	4.8	*	*
2005 5/.....	2.3	*	1.6	1.0	1.4	1.7	2.2	3.0	4.1	5.3	10.7	*
2004 5/.....	2.6	*	*	1.2	1.5	2.4	2.3	4.0	5.3	5.7	8.8	*
2003 5/.....	2.5	*	*	1.3	1.4	2.5	2.4	4.0	4.4	5.5	8.9	*
2002.....	2.8	*	*	1.1	1.9	2.0	3.7	4.0	5.9	7.5	6.7	*
2001.....	2.7	*	*	0.9	1.4	2.5	3.0	4.1	4.7	6.8	8.7	*
2000.....	2.8	*	*	1.3	1.6	2.5	3.0	3.7	6.1	7.9	6.5	24.7
1999.....	2.9	*	*	1.4	1.7	2.7	2.8	4.0	5.5	9.2	8.4	*
Non-Hispanic black, female												
2007 5/.....	2.5	*	*	*	*	1.0	2.3	4.8	4.4	5.4	6.3	14.1
2006 5/.....	2.7	*	*	0.7	0.9	1.3	2.6	4.4	5.5	6.1	7.1	15.3
2005 5/.....	3.1	*	*	*	0.8	1.5	2.6	5.4	6.6	8.6	9.5	13.8
2004 5/.....	2.8	*	*	*	1.0	1.1	2.7	5.1	6.2	5.9	9.7	16.9
2003 5/.....	3.0	*	*	*	1.1	1.0	3.2	5.7	5.9	8.4	9.5	13.6
2002.....	3.1	*	*	1.0	*	1.3	3.4	5.7	7.2	8.4	8.6	12.5
2001.....	3.4	*	*	0.7	0.9	2.0	3.6	5.1	8.6	8.4	11.8	17.1
2000.....	3.7	*	*	0.8	1.0	1.5	3.4	7.0	7.6	10.4	12.7	21.4
1999.....	3.6	*	*	0.6	0.9	2.2	3.2	6.5	8.1	10.2	11.7	21.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

All origins 3/, both sexes

2007.....	36.8	*	*	*	*	0.1	0.9	7.1	32.6	128.5	326.0	533.8
2006.....	36.0	*	*	*	*	0.1	0.9	6.6	32.6	129.1	320.2	525.4
2005.....	37.8	*	*	*	*	0.1	1.0	6.9	34.5	137.2	337.2	563.5
2004.....	35.3	*	*	*	*	0.1	0.9	6.0	33.0	130.0	319.0	531.1
2003.....	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
2002.....	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001.....	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000.....	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999.....	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8

All origins 3/, male

2007.....	35.7	*	*	*	*	0.1	1.0	7.4	35.6	143.1	390.1	684.9
2006.....	34.8	*	*	*	*	0.1	0.8	6.9	35.3	141.5	386.8	682.2
2005.....	36.6	*	*	*	*	0.1	1.0	7.3	37.0	154.4	405.2	732.4
2004.....	34.5	*	*	*	*	0.1	1.0	6.5	35.2	145.0	388.4	718.6
2003.....	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
2002.....	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001.....	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000.....	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999.....	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7

All origins 3/, female

2007.....	37.9	*	*	*	*	0.1	0.9	6.8	29.9	116.1	281.8	461.9
2006.....	37.1	*	*	*	*	0.1	0.9	6.4	30.1	118.7	274.8	452.1
2005.....	39.0	*	*	*	*	*	1.0	6.4	32.2	122.6	291.0	485.9
2004.....	36.2	*	*	*	*	*	0.9	5.5	30.8	117.3	272.2	446.8
2003.....	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
2002.....	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001.....	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000.....	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999.....	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

Hispanic, both sexes

2007.....	6.5	*	*	*	*	*	*	2.0	8.6	37.9	145.3	321.1
2006.....	6.2	*	*	*	*	*	*	1.6	8.9	39.0	132.1	331.8
2005.....	6.7	*	*	*	*	*	*	2.5	9.7	42.0	148.9	367.9
2004.....	6.1	*	*	*	*	*	0.5	1.7	9.3	43.2	135.4	345.4
2003.....	6.3	*	*	*	*	*	0.4	1.7	9.1	43.7	143.5	403.7
2002.....	6.3	*	*	*	*	*	0.4	2.1	10.4	43.1	147.8	397.4
2001.....	6.0	*	*	*	*	*	*	1.9	10.0	45.2	139.6	418.9
2000.....	5.9	*	*	*	*	*	0.4	1.7	9.9	40.7	149.1	422.7
1999.....	6.5	*	*	*	*	*	0.4	1.9	10.8	49.3	158.1	462.3

Hispanic, male

2007.....	6.7	*	*	*	*	*	*	2.5	10.2	50.9	191.7	416.7
2006.....	6.2	*	*	*	*	*	*	1.8	10.7	49.0	170.1	439.8
2005.....	6.9	*	*	*	*	*	*	2.6	13.0	53.5	205.8	468.2
2004.....	6.3	*	*	*	*	*	0.7	1.7	11.2	59.2	181.0	455.3
2003.....	6.6	*	*	*	*	*	*	1.5	12.2	57.8	194.4	574.2
2002.....	6.6	*	*	*	*	*	*	2.2	14.0	57.9	208.9	535.4
2001.....	6.3	*	*	*	*	*	*	2.1	12.2	62.5	181.5	605.7
2000.....	6.2	*	*	*	*	*	*	1.6	13.6	56.6	209.0	584.5
1999.....	7.0	*	*	*	*	*	*	1.6	13.9	65.9	228.9	735.4

Hispanic, female

2007.....	6.3	*	*	*	*	*	*	1.5	7.1	27.3	112.8	269.2
2006.....	6.1	*	*	*	*	*	*	1.4	7.3	30.8	105.6	274.4
2005.....	6.5	*	*	*	*	*	*	2.3	6.6	32.7	109.3	315.4
2004.....	5.9	*	*	*	*	*	*	1.7	7.6	30.4	104.0	288.8
2003.....	6.0	*	*	*	*	*	*	1.9	6.3	32.5	108.5	317.2
2002.....	5.9	*	*	*	*	*	*	1.9	7.1	31.3	106.0	329.1
2001.....	5.8	*	*	*	*	*	*	1.8	8.1	31.5	111.2	327.7
2000.....	5.6	*	*	*	*	*	*	1.7	6.6	28.2	109.0	343.3
1999.....	5.9	*	*	*	*	*	*	2.1	8.1	36.3	111.0	330.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 4/, both sexes

2007.....	42.1	*	*	*	*	0.1	1.1	7.7	34.7	135.6	337.4	544.7
2006.....	41.1	*	*	*	*	0.1	1.0	7.2	34.6	136.0	331.4	534.6
2005.....	43.0	*	*	*	*	0.1	1.1	7.3	36.5	144.2	347.8	572.1
2004.....	40.0	*	*	*	*	0.1	1.0	6.4	34.8	136.0	328.7	538.8
2003.....	41.4	*	*	*	*	0.1	1.1	6.3	36.7	143.8	338.3	564.3
2002.....	40.6	*	*	*	*	0.1	1.0	6.3	35.6	141.0	338.1	560.8
2001.....	40.0	*	*	*	0.1	0.1	1.0	6.0	36.2	143.2	328.6	561.3
2000.....	39.5	*	*	*	*	0.1	0.9	6.0	36.0	142.3	330.5	560.5
1999.....	40.0	*	*	*	*	0.1	0.8	5.8	37.9	146.4	336.8	557.6

Non-Hispanic 4/, male

2007.....	41.1	*	*	*	*	*	1.1	8.0	37.7	150.1	402.6	700.0
2006.....	40.0	*	*	*	*	0.1	0.9	7.5	37.3	148.2	399.8	694.8
2005.....	41.8	*	*	*	*	*	1.1	7.8	38.9	161.6	416.5	745.0
2004.....	39.3	*	*	*	*	0.2	1.0	7.0	37.1	150.6	399.4	730.9
2003.....	40.7	*	*	*	*	0.1	1.0	6.9	39.6	161.7	410.1	776.7
2002.....	40.4	*	*	*	*	0.1	1.1	6.9	37.8	159.2	418.5	794.9
2001.....	39.8	*	*	*	*	0.1	1.0	6.6	38.9	162.6	412.4	800.6
2000.....	40.0	*	*	*	*	*	0.9	6.5	39.3	163.8	418.4	846.6
1999.....	41.4	*	*	*	*	*	0.8	6.3	42.2	173.9	435.1	869.6

Non-Hispanic 4/, female

2007.....	43.1	*	*	*	*	0.1	1.0	7.4	31.9	123.2	292.5	471.4
2006.....	42.1	*	*	*	*	0.1	1.0	7.0	32.0	125.6	284.8	460.1
2005.....	44.1	*	*	*	*	*	1.1	6.9	34.3	129.4	301.2	493.2
2004.....	40.8	*	*	*	*	*	1.0	5.9	32.7	123.7	281.2	453.0
2003.....	42.0	*	*	*	*	*	1.1	5.8	34.1	128.9	290.3	471.0
2002.....	40.8	*	*	*	*	0.1	1.0	5.8	33.4	125.8	284.9	459.8
2001.....	40.1	*	*	*	*	*	1.0	5.5	33.8	127.1	273.6	461.7
2000.....	39.1	*	*	*	*	*	0.9	5.5	32.9	124.5	273.2	444.8
1999.....	38.7	*	*	*	*	0.2	0.7	5.2	33.9	123.9	272.9	432.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic white, both sexes

2007 5/.....	49.8	*	*	*	*	0.1	1.1	8.0	37.3	146.6	359.3	574.2
2006 5/.....	48.5	*	*	*	*	0.1	1.0	7.6	36.9	147.3	351.8	562.5
2005 5/.....	50.6	*	*	*	*	0.1	1.1	7.6	39.0	155.5	369.2	600.8
2004 5/.....	47.1	*	*	*	*	0.1	1.1	6.7	37.1	146.7	348.1	566.1
2003 5/.....	48.6	*	*	*	*	0.1	1.1	6.6	39.0	154.4	357.6	592.2
2002.....	47.5	*	*	*	*	0.1	1.0	6.5	37.7	151.2	355.7	586.9
2001.....	46.6	*	*	*	*	0.1	1.0	6.2	38.6	152.6	345.7	585.7
2000.....	46.0	*	*	*	*	*	0.8	6.1	37.9	151.2	347.1	585.5
1999.....	46.3	*	*	*	*	0.1	0.7	5.9	39.9	154.8	351.4	582.8

Non-Hispanic white, male

2007 5/.....	47.8	*	*	*	*	*	1.1	8.4	39.7	158.9	420.2	729.2
2006 5/.....	46.4	*	*	*	*	*	0.9	7.8	38.8	156.8	416.3	720.3
2005 5/.....	48.4	*	*	*	*	*	1.1	7.8	40.5	170.4	432.7	770.7
2004 5/.....	45.5	*	*	*	*	0.2	1.1	7.2	38.4	158.9	415.6	756.7
2003 5/.....	46.9	*	*	*	*	*	1.0	7.1	41.1	169.5	423.7	804.2
2002.....	46.4	*	*	*	*	*	1.1	6.9	39.0	167.0	431.9	821.0
2001.....	45.6	*	*	*	*	*	1.0	6.7	40.5	169.3	425.2	823.9
2000.....	45.7	*	*	*	*	*	0.9	6.4	40.5	169.8	431.4	876.5
1999.....	47.1	*	*	*	*	*	0.8	6.2	43.4	180.3	445.9	900.5

Non-Hispanic white, female

2007 5/.....	51.7	*	*	*	*	*	1.1	7.7	35.0	135.9	316.7	500.8
2006 5/.....	50.5	*	*	*	*	*	1.0	7.5	35.1	139.2	307.0	488.9
2005 5/.....	52.8	*	*	*	*	*	1.2	7.3	37.6	142.5	325.5	523.1
2004 5/.....	48.8	*	*	*	*	*	1.1	6.3	35.9	136.2	302.2	480.7
2003 5/.....	50.2	*	*	*	*	*	1.1	6.1	37.0	141.5	313.1	498.8
2002.....	48.6	*	*	*	*	*	1.0	6.2	36.4	137.7	304.6	485.7
2001.....	47.6	*	*	*	*	*	1.0	5.7	36.8	138.5	292.9	486.6
2000.....	46.3	*	*	*	*	*	0.8	5.8	35.5	135.5	291.6	468.2
1999.....	45.5	*	*	*	*	*	0.7	5.5	36.6	133.4	289.5	456.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic black, both sexes

2007 5/.....	16.1	*	*	*	*	*	1.2	7.7	27.2	89.1	200.3	303.3
2006 5/.....	15.8	*	*	*	*	*	1.2	7.2	28.4	86.5	199.5	296.8
2005 5/.....	16.8	*	*	*	*	*	1.2	8.5	29.8	95.7	207.2	328.2
2004 5/.....	15.0	*	*	*	*	*	1.1	6.9	28.4	86.8	193.2	288.1
2003 5/.....	15.7	*	*	*	*	*	1.3	6.5	30.5	94.6	200.3	306.1
2002.....	15.9	*	*	*	*	*	1.2	6.8	30.2	90.3	213.1	322.7
2001.....	15.6	*	*	*	*	*	1.3	6.5	28.7	96.1	200.8	323.4
2000.....	15.4	*	*	*	*	*	1.3	6.8	31.4	95.8	196.8	312.6
1999.....	16.4	*	*	*	*	*	1.2	6.6	33.2	101.9	220.1	314.0

Non-Hispanic black, male

2007 5/.....	18.1	*	*	*	*	*	1.5	8.0	35.1	119.7	307.0	466.0
2006 5/.....	17.9	*	*	*	*	*	1.1	8.3	36.9	120.1	301.7	473.7
2005 5/.....	19.5	*	*	*	*	*	1.2	10.3	38.0	134.5	325.7	534.8
2004 5/.....	17.4	*	*	*	*	*	1.3	8.0	38.8	119.0	295.8	495.9
2003 5/.....	18.6	*	*	*	*	*	1.4	7.7	39.3	133.6	329.0	526.3
2002.....	18.7	*	*	*	*	*	1.3	8.5	40.3	126.6	337.9	572.1
2001.....	18.8	*	*	*	*	*	1.4	7.5	37.6	139.1	334.9	592.5
2000.....	18.7	*	*	*	*	*	1.3	8.5	40.7	142.7	322.4	578.9
1999.....	20.3	*	*	*	*	*	1.5	8.3	45.4	146.5	369.3	610.9

Non-Hispanic black, female

2007 5/.....	14.4	*	*	*	*	*	0.9	7.5	20.9	67.2	139.7	235.8
2006 5/.....	13.8	*	*	*	*	*	1.3	6.2	21.6	62.5	141.5	224.6
2005 5/.....	14.4	*	*	*	*	*	1.2	6.8	23.2	68.0	140.0	245.1
2004 5/.....	12.9	*	*	*	*	*	1.0	5.9	20.1	63.8	135.2	206.0
2003 5/.....	13.0	*	*	*	*	*	1.2	5.5	23.5	66.8	127.6	220.8
2002.....	13.3	*	*	*	*	*	1.1	5.3	22.1	64.6	142.4	227.4
2001.....	12.7	*	*	*	*	*	1.3	5.7	21.5	65.8	125.5	223.7
2000.....	12.4	*	*	*	*	*	1.3	5.3	24.0	62.8	126.1	213.7
1999.....	12.9	*	*	*	*	*	1.0	5.0	23.3	70.4	136.6	203.0











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

All origins 3/, both sexes

2007.....	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	8.9	39.8	153.3
2006.....	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	9.3	40.5	155.8
2005.....	5.8	*	*	*	0.1	0.2	0.4	1.1	2.7	9.3	42.6	163.9
2004.....	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002.....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000.....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999.....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6

All origins 3/, male

2007.....	6.0	*	*	*	0.1	0.2	0.5	1.3	3.5	11.7	55.1	222.0
2006.....	6.0	*	*	*	*	0.3	0.5	1.3	3.5	12.6	57.2	224.8
2005.....	6.0	*	*	*	0.1	0.2	0.4	1.3	3.5	12.5	57.9	229.3
2004.....	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002.....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999.....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4

All origins 3/, female

2007.....	5.3	*	*	*	*	0.1	0.3	0.7	2.0	6.5	29.3	120.7
2006.....	5.2	*	*	*	*	0.1	0.3	0.8	1.9	6.4	29.1	123.5
2005.....	5.7	*	*	*	0.1	0.1	0.4	0.9	2.1	6.7	32.2	133.9
2004.....	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	132.4
2003.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002.....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001.....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000.....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999.....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Hispanic, both sexes

2007.....	1.2	*	*	*	*	*	*	*	2.0	5.2	19.3	75.1
2006.....	1.3	*	*	*	*	*	*	0.5	1.3	5.7	25.9	77.5
2005.....	1.4	*	*	*	*	*	*	0.7	1.4	5.3	24.3	109.9
2004.....	1.3	*	*	*	*	*	*	*	1.6	5.1	26.3	101.3
2003.....	1.3	*	*	*	*	*	*	*	1.7	6.5	22.8	113.0
2002.....	1.4	*	*	*	*	*	*	*	1.8	5.3	29.0	131.4
2001.....	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000.....	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999.....	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5

Hispanic, male

2007.....	1.2	*	*	*	*	*	*	*	2.1	7.0	26.3	100.6
2006.....	1.3	*	*	*	*	*	*	*	2.1	7.3	31.9	86.8
2005.....	1.4	*	*	*	*	*	*	*	2.2	6.1	32.1	154.4
2004.....	1.3	*	*	*	*	*	*	*	2.4	7.4	31.8	121.9
2003.....	1.3	*	*	*	*	*	*	*	2.0	8.7	32.2	123.1
2002.....	1.4	*	*	*	*	*	*	*	2.6	6.1	43.1	167.3
2001.....	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000.....	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999.....	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6

Hispanic, female

2007.....	1.2	*	*	*	*	*	*	*	1.9	3.8	14.4	61.2
2006.....	1.3	*	*	*	*	*	*	*	*	4.3	21.7	72.5
2005.....	1.3	*	*	*	*	*	*	*	*	4.7	18.8	86.7
2004.....	1.3	*	*	*	*	*	*	*	*	3.3	22.5	90.8
2003.....	1.4	*	*	*	*	*	*	*	*	4.9	16.3	107.8
2002.....	1.4	*	*	*	*	*	*	*	*	4.6	19.4	113.6
2001.....	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2
2000.....	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999.....	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic 4/, both sexes

2007.....	6.4	*	*	*	0.1	0.2	0.4	1.1	2.8	9.2	41.1	157.4
2006.....	6.4	*	*	*	0.1	0.2	0.4	1.1	2.8	9.5	41.4	159.6
2005.....	6.6	*	*	*	0.1	0.2	0.4	1.2	2.8	9.6	43.6	166.3
2004.....	6.4	*	*	*	0.1	0.2	0.4	1.0	2.6	9.9	43.7	167.4
2003.....	6.7	*	*	*	0.1	0.2	0.4	1.0	2.9	9.7	46.0	178.5
2002.....	6.8	*	*	*	0.1	0.2	0.5	0.9	2.6	10.1	47.1	187.8
2001.....	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	190.0
2000.....	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999.....	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.7	176.8

Non-Hispanic 4/, male

2007.....	6.9	*	*	*	0.2	0.3	0.5	1.4	3.6	12.1	56.9	228.8
2006.....	6.9	*	*	*	*	0.3	0.5	1.3	3.6	13.0	58.8	232.3
2005.....	6.8	*	*	*	0.1	0.2	0.5	1.4	3.6	12.9	59.3	232.8
2004.....	6.7	*	*	*	*	0.2	0.5	1.3	3.3	13.2	60.1	242.7
2003.....	6.9	*	*	*	0.1	0.2	0.4	1.2	3.6	13.0	63.1	251.5
2002.....	7.0	*	*	*	0.1	0.2	0.6	1.2	3.1	13.4	64.7	272.1
2001.....	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.2	271.6
2000.....	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.7
1999.....	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6

Non-Hispanic 4/, female

2007.....	5.9	*	*	*	*	0.2	0.3	0.8	2.0	6.7	30.3	123.6
2006.....	5.9	*	*	*	*	0.1	0.3	0.8	2.1	6.6	29.5	125.8
2005.....	6.3	*	*	*	0.1	0.1	0.4	1.0	2.2	6.8	32.9	136.0
2004.....	6.1	*	*	*	*	0.2	0.3	0.7	2.0	7.1	32.8	133.8
2003.....	6.5	*	*	*	*	0.2	0.4	0.8	2.1	7.0	34.6	146.5
2002.....	6.6	*	*	*	*	0.2	0.3	0.6	2.1	7.3	35.4	151.4
2001.....	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000.....	6.3	*	*	*	*	0.1	0.3	0.8	2.0	7.1	33.6	152.3
1999.....	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.4	142.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic white, both sexes

2007 5/.....	7.3	*	*	*	0.1	0.2	0.4	1.1	2.6	9.2	41.8	162.0
2006 5/.....	7.3	*	*	*	*	0.2	0.4	1.1	2.8	9.2	42.0	164.1
2005 5/.....	7.4	*	*	*	0.1	0.2	0.5	1.1	2.7	9.4	43.5	169.0
2004 5/.....	7.2	*	*	*	0.1	0.2	0.4	1.0	2.4	9.7	44.1	170.4
2003 5/.....	7.5	*	*	*	0.1	0.2	0.4	1.0	2.6	9.5	45.7	181.0
2002.....	7.7	*	*	*	0.1	0.2	0.4	0.9	2.5	9.7	47.3	191.3
2001.....	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.2
2000.....	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.5
1999.....	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3

Non-Hispanic white, male

2007 5/.....	7.8	*	*	*	0.2	0.2	0.5	1.4	3.4	12.0	57.2	236.9
2006 5/.....	7.8	*	*	*	*	0.3	0.5	1.3	3.6	12.5	59.2	239.9
2005 5/.....	7.6	*	*	*	*	0.2	0.5	1.3	3.4	12.4	58.7	237.8
2004 5/.....	7.5	*	*	*	*	0.2	0.5	1.3	2.9	12.7	60.2	247.4
2003 5/.....	7.6	*	*	*	*	0.2	0.4	1.2	3.3	12.6	62.4	256.5
2002.....	7.9	*	*	*	*	0.3	0.6	1.1	3.0	12.9	64.9	278.4
2001.....	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9
2000.....	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.1
1999.....	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4

Non-Hispanic white, female

2007 5/.....	6.8	*	*	*	*	0.2	0.3	0.7	1.9	6.8	30.9	126.5
2006 5/.....	6.7	*	*	*	*	*	0.3	0.8	2.1	6.4	30.1	128.7
2005 5/.....	7.2	*	*	*	*	0.2	0.4	1.0	2.1	6.8	33.1	137.5
2004 5/.....	6.9	*	*	*	*	*	0.3	0.7	2.0	7.1	33.1	135.9
2003 5/.....	7.3	*	*	*	*	0.2	0.3	0.8	2.0	6.8	34.4	147.8
2002.....	7.4	*	*	*	*	0.2	0.3	0.6	2.0	7.1	35.6	153.7
2001.....	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.5
2000.....	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999.....	6.5	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.3	144.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic black, both sexes

2007 5/.....	3.7	*	*	*	*	*	0.4	1.4	4.7	11.5	43.5	127.6
2006 5/.....	3.8	*	*	*	*	0.4	0.7	1.4	3.8	14.1	42.9	130.9
2005 5/.....	4.3	*	*	*	*	*	0.5	1.8	4.6	13.5	53.2	155.4
2004 5/.....	4.0	*	*	*	*	*	0.6	1.4	5.0	14.2	47.6	147.4
2003 5/.....	4.4	*	*	*	*	*	0.7	1.3	5.3	13.9	55.5	165.7
2002.....	4.2	*	*	*	*	*	0.7	1.4	3.8	15.2	52.8	167.3
2001.....	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000.....	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999.....	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.6	168.5

Non-Hispanic black, male

2007 5/.....	4.0	*	*	*	*	*	*	1.7	6.7	17.3	66.6	163.2
2006 5/.....	4.2	*	*	*	*	*	0.8	1.7	5.3	21.2	68.4	177.8
2005 5/.....	4.6	*	*	*	*	*	*	2.5	6.3	20.0	81.0	205.5
2004 5/.....	4.3	*	*	*	*	*	0.8	1.8	7.3	21.5	68.7	207.3
2003 5/.....	4.5	*	*	*	*	*	*	1.7	7.2	19.8	79.6	215.2
2002.....	4.4	*	*	*	*	*	0.9	2.0	4.9	22.3	74.1	237.4
2001.....	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000.....	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999.....	4.7	*	*	*	*	*	0.9	1.8	6.6	25.3	81.6	229.4

Non-Hispanic black, female

2007 5/.....	3.4	*	*	*	*	*	*	1.1	3.2	7.3	30.3	112.8
2006 5/.....	3.4	*	*	*	*	*	*	1.2	2.5	9.0	28.4	111.7
2005 5/.....	4.0	*	*	*	*	*	*	1.1	3.2	8.9	37.5	135.2
2004 5/.....	3.8	*	*	*	*	*	*	1.0	3.1	8.9	35.7	123.7
2003 5/.....	4.3	*	*	*	*	*	0.8	1.0	3.7	9.7	41.9	146.5
2002.....	4.1	*	*	*	*	*	*	0.9	3.0	10.2	40.8	140.5
2001.....	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000.....	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999.....	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.4	145.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes

2007.....	9.5	7.5	0.6	0.2	0.4	0.5	1.1	3.1	9.3	28.7	73.2	138.9
2006.....	9.2	8.1	0.6	0.2	0.2	0.5	1.2	3.1	8.9	28.2	73.4	136.6
2005.....	9.1	7.4	0.6	0.2	0.3	0.5	1.1	3.2	8.8	28.8	72.5	136.5
2004.....	9.3	6.5	0.7	0.2	0.3	0.4	1.2	3.2	9.5	29.9	74.2	139.4
2003.....	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
2002.....	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001.....	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000.....	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999.....	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6

All origins 3/, male

2007.....	9.6	8.7	0.8	0.2	0.4	0.6	1.2	3.7	10.8	35.1	92.7	169.2
2006.....	9.3	9.2	0.7	0.2	0.3	0.5	1.2	3.3	10.3	35.0	93.1	163.7
2005.....	9.0	8.9	0.7	0.2	0.3	0.4	1.0	3.5	10.2	34.9	90.7	157.7
2004.....	9.3	6.6	0.7	0.2	0.3	0.4	1.2	3.5	10.8	36.9	93.9	168.8
2003.....	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
2002.....	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001.....	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000.....	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999.....	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3

All origins 3/, female

2007.....	9.3	6.2	0.4	0.2	0.3	0.5	1.1	2.6	7.9	23.3	59.8	124.5
2006.....	9.2	6.9	0.5	0.2	0.2	0.5	1.2	3.0	7.5	22.6	59.9	124.0
2005.....	9.2	5.8	0.5	0.1	0.2	0.5	1.1	2.9	7.6	23.7	60.2	126.7
2004.....	9.3	6.4	0.6	0.1	0.3	0.4	1.2	3.0	8.4	24.1	61.0	126.1
2003.....	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
2002.....	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001.....	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000.....	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999.....	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, both sexes

2007.....	3.8	5.4	0.5	*	*	0.4	0.7	2.5	8.3	26.9	63.9	105.1
2006.....	3.6	6.3	*	*	0.3	0.3	0.7	2.6	7.6	25.4	59.5	106.8
2005.....	3.7	5.8	*	*	0.3	0.4	0.7	2.6	8.8	26.9	67.7	104.0
2004.....	3.6	5.8	*	*	*	*	0.9	2.4	7.8	27.2	71.0	112.7
2003.....	3.4	6.6	*	*	*	0.4	0.9	2.5	8.9	25.5	58.2	118.7
2002.....	3.3	5.2	*	*	*	0.4	0.7	2.7	8.3	26.1	58.0	112.0
2001.....	3.4	5.0	*	*	*	*	0.7	3.1	8.9	25.3	65.5	134.4
2000.....	3.5	5.6	*	*	*	0.5	1.1	2.6	9.8	25.4	64.2	135.4
1999.....	3.4	5.0	*	*	*	0.4	0.9	3.2	9.6	26.0	64.0	134.0

Hispanic, male

2007.....	3.8	7.2	*	*	*	0.4	0.8	3.4	9.8	31.9	71.6	128.1
2006.....	3.4	7.1	*	*	*	*	0.9	2.7	10.1	28.9	67.3	112.8
2005.....	3.5	6.3	*	*	*	0.5	0.7	3.0	9.9	29.1	81.8	114.6
2004.....	3.4	5.6	*	*	*	*	0.8	2.9	9.1	31.1	82.3	112.1
2003.....	3.3	6.6	*	*	*	*	1.1	3.1	11.3	30.8	63.3	127.8
2002.....	3.2	4.9	*	*	*	*	0.8	2.6	9.8	34.6	64.0	108.8
2001.....	3.4	5.0	*	*	*	*	0.9	3.4	10.6	30.2	78.3	167.6
2000.....	3.3	5.8	*	*	*	*	1.3	3.0	11.9	29.3	81.1	133.0
1999.....	3.3	5.2	*	*	*	*	1.1	3.1	11.6	30.5	82.4	161.8

Hispanic, female

2007.....	3.8	*	*	*	*	*	0.7	1.5	6.9	22.9	58.4	92.7
2006.....	3.7	5.5	*	*	*	*	*	2.6	5.3	22.7	54.0	103.6
2005.....	3.9	5.3	*	*	*	*	0.8	2.2	7.8	25.2	57.9	98.4
2004.....	3.9	6.1	*	*	*	*	0.9	1.9	6.6	24.1	63.2	113.1
2003.....	3.6	6.6	*	*	*	*	*	1.8	6.7	21.2	54.8	114.1
2002.....	3.4	5.4	*	*	*	*	*	2.8	6.9	19.3	54.0	113.6
2001.....	3.5	5.0	*	*	*	*	*	2.8	7.5	21.4	56.8	118.2
2000.....	3.6	5.3	*	*	*	*	0.9	2.2	7.8	22.2	52.8	136.5
1999.....	3.5	*	*	*	*	*	*	3.3	7.8	22.4	51.7	120.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 4/, both sexes

2007.....	10.4	8.1	0.6	0.2	0.4	0.6	1.2	3.2	9.4	28.8	73.8	140.6
2006.....	10.2	8.6	0.7	0.2	0.2	0.6	1.3	3.2	9.0	28.4	74.1	138.1
2005.....	10.0	7.8	0.6	0.2	0.3	0.5	1.1	3.3	8.8	28.9	72.7	138.0
2004.....	10.2	6.7	0.7	0.2	0.3	0.5	1.3	3.3	9.6	30.1	74.3	140.5
2003.....	9.6	7.0	0.6	0.1	0.4	0.6	1.2	3.3	8.7	28.2	68.9	138.7
2002.....	9.5	7.3	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.8	68.4	134.1
2001.....	10.0	7.8	0.6	0.2	0.4	0.6	1.3	3.5	9.7	30.3	73.3	142.7
2000.....	9.8	7.5	0.6	0.2	0.4	0.7	1.4	3.6	9.9	29.8	72.1	141.6
1999.....	9.7	6.9	0.6	0.2	0.3	0.7	1.3	3.3	9.4	29.5	70.8	147.7

Non-Hispanic 4/, male

2007.....	10.7	9.0	0.8	0.2	0.4	0.6	1.2	3.7	10.9	35.4	94.0	171.6
2006.....	10.4	9.8	0.8	0.2	0.3	0.5	1.3	3.3	10.3	35.4	94.7	166.4
2005.....	10.0	9.6	0.8	0.2	0.3	0.4	1.1	3.6	10.2	35.3	91.0	159.9
2004.....	10.3	6.9	0.8	0.2	0.3	0.5	1.3	3.6	10.9	37.2	94.4	171.6
2003.....	9.5	8.3	0.6	0.2	0.4	0.6	1.1	3.5	10.1	34.2	86.3	166.5
2002.....	9.4	8.7	0.7	0.2	0.3	0.4	1.3	3.4	10.5	35.3	84.7	158.5
2001.....	9.8	10.0	0.6	0.2	0.4	0.6	1.4	3.9	10.9	36.7	89.5	169.1
2000.....	9.8	8.4	0.6	0.2	0.4	0.6	1.5	3.9	11.3	36.0	90.7	181.3
1999.....	9.6	7.9	0.8	0.2	0.4	0.7	1.4	3.6	10.5	36.4	89.9	182.7

Non-Hispanic 4/, female

2007.....	10.2	7.1	0.4	0.2	0.3	0.5	1.2	2.8	7.9	23.3	59.8	126.0
2006.....	10.0	7.3	0.5	0.2	0.2	0.6	1.3	3.0	7.7	22.5	60.1	124.9
2005.....	10.1	5.9	0.4	0.2	0.2	0.5	1.2	3.0	7.5	23.6	60.3	128.0
2004.....	10.1	6.5	0.6	0.1	0.3	0.5	1.3	3.1	8.5	24.1	60.8	126.6
2003.....	9.7	5.7	0.5	*	0.3	0.5	1.2	3.2	7.5	23.2	57.3	126.4
2002.....	9.6	5.9	0.4	0.2	0.3	0.5	1.2	3.1	8.0	23.3	57.5	123.6
2001.....	10.2	5.5	0.6	0.2	0.3	0.6	1.3	3.1	8.5	25.0	62.7	131.8
2000.....	9.9	6.6	0.5	0.2	0.3	0.7	1.3	3.3	8.6	24.7	60.1	125.5
1999.....	9.8	5.9	0.5	0.2	0.2	0.7	1.2	3.1	8.5	23.8	58.4	133.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic white, both sexes

2007 5/.....	11.7	5.9	0.5	0.1	0.3	0.5	1.1	2.9	9.0	29.3	76.7	145.7
2006 5/.....	11.4	5.4	0.6	0.2	0.2	0.5	1.2	2.8	8.6	29.2	77.0	142.9
2005 5/.....	11.1	6.0	0.5	0.2	0.2	0.4	1.0	2.9	8.5	29.4	74.8	141.8
2004 5/.....	11.4	4.5	0.6	0.1	0.3	0.4	1.1	2.9	9.4	30.9	76.9	145.4
2003 5/.....	10.7	4.9	0.4	0.1	0.3	0.5	0.9	3.0	8.5	29.1	71.3	143.7
2002.....	10.5	4.7	0.4	0.2	0.3	0.4	1.1	2.9	8.8	29.3	70.3	138.6
2001.....	11.0	4.8	0.5	0.2	0.3	0.5	1.1	3.0	9.3	30.9	75.6	145.9
2000.....	10.8	5.4	0.5	0.2	0.3	0.6	1.2	3.2	9.6	30.2	73.8	145.7
1999.....	10.6	4.9	0.6	0.2	0.3	0.6	1.1	3.0	9.1	29.7	72.3	150.7

Non-Hispanic white, male

2007 5/.....	11.9	6.8	0.6	*	0.4	0.5	1.1	3.3	10.5	35.9	97.7	178.3
2006 5/.....	11.6	5.6	0.7	0.2	0.2	0.5	1.2	3.1	10.1	36.3	98.2	171.5
2005 5/.....	11.0	7.3	0.7	0.2	0.2	0.4	1.0	3.2	9.7	36.1	92.9	162.2
2004 5/.....	11.4	4.6	0.6	*	0.3	0.4	1.0	3.2	10.6	38.1	97.3	177.3
2003 5/.....	10.6	5.5	0.4	0.2	0.3	0.6	1.0	3.2	9.7	35.2	88.8	172.0
2002.....	10.3	5.6	0.5	*	0.3	0.4	1.1	3.0	10.1	36.0	87.0	164.1
2001.....	10.8	6.2	0.4	0.2	0.4	0.4	1.3	3.4	10.5	37.7	92.4	173.4
2000.....	10.7	6.6	0.5	0.2	0.3	0.6	1.3	3.5	11.0	36.4	92.6	185.9
1999.....	10.5	5.6	0.6	0.2	0.4	0.7	1.2	3.2	10.0	37.0	92.0	186.1

Non-Hispanic white, female

2007 5/.....	11.4	4.9	*	*	0.3	0.5	1.1	2.5	7.7	23.5	62.0	130.3
2006 5/.....	11.2	5.2	*	*	*	0.6	1.1	2.6	7.2	23.1	62.3	129.6
2005 5/.....	11.2	4.8	*	0.2	0.2	0.4	1.0	2.6	7.4	23.6	62.4	132.5
2004 5/.....	11.3	4.4	0.5	*	0.3	0.4	1.1	2.7	8.3	24.7	63.0	131.0
2003 5/.....	10.8	4.4	*	*	0.3	0.4	0.9	2.7	7.3	23.9	59.4	131.2
2002.....	10.7	3.7	*	0.2	0.2	0.5	1.0	2.7	7.7	23.6	59.1	127.6
2001.....	11.3	3.4	0.5	0.2	0.3	0.6	0.9	2.6	8.2	25.1	64.4	134.5
2000.....	11.0	4.1	0.5	*	0.3	0.6	1.1	3.0	8.4	25.0	61.5	129.5
1999.....	10.7	4.0	0.5	0.2	0.2	0.5	0.9	2.7	8.2	23.6	59.4	136.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic black, both sexes

2007 5/.....	6.7	16.4	1.2	*	0.6	1.0	2.1	5.9	13.6	29.7	54.0	96.1
2006 5/.....	6.4	20.3	0.9	*	0.5	0.8	2.2	6.0	13.4	26.0	51.9	96.6
2005 5/.....	6.6	15.7	0.9	*	0.4	0.9	2.2	6.3	12.3	29.6	57.1	104.0
2004 5/.....	6.4	15.3	1.1	*	0.4	0.9	2.7	6.4	13.6	27.9	53.9	94.4
2003 5/.....	6.0	15.3	1.1	*	0.7	1.0	2.5	6.6	12.3	24.6	51.3	91.0
2002.....	6.2	17.3	1.2	0.4	0.7	0.9	2.4	6.2	13.8	27.1	52.3	90.7
2001.....	6.9	19.0	1.1	*	0.7	1.3	2.9	7.5	13.9	29.6	56.9	110.8
2000.....	6.7	17.7	0.8	*	0.7	1.1	2.7	6.9	13.7	30.6	58.6	99.8
1999.....	6.7	15.8	1.1	*	0.4	1.4	2.9	6.6	13.6	31.3	58.7	112.8

Non-Hispanic black, male

2007 5/.....	6.8	18.1	1.7	*	0.6	1.0	2.3	7.1	16.6	35.1	67.1	116.1
2006 5/.....	6.4	24.7	*	*	0.6	0.8	2.3	5.8	14.4	33.4	67.6	119.1
2005 5/.....	6.7	20.5	*	*	*	0.8	2.0	6.9	15.4	34.4	76.5	134.2
2004 5/.....	6.5	16.4	*	*	*	1.1	3.2	6.9	15.5	34.8	69.9	117.9
2003 5/.....	6.2	18.6	*	*	0.9	1.1	2.3	6.4	15.3	31.4	69.0	116.1
2002.....	6.3	20.3	*	*	*	0.9	2.8	6.5	16.4	32.5	67.1	102.8
2001.....	6.6	24.3	*	*	*	1.4	2.6	7.8	15.7	32.9	69.0	131.7
2000.....	6.8	17.1	*	*	0.8	1.2	3.2	8.0	15.6	36.2	74.0	135.0
1999.....	6.6	18.0	*	*	*	1.1	2.9	7.3	15.9	35.6	73.5	146.2

Non-Hispanic black, female

2007 5/.....	6.6	14.5	*	*	*	1.1	2.0	5.0	11.3	25.9	46.6	87.9
2006 5/.....	6.4	15.8	*	*	*	0.8	2.1	6.2	12.5	20.7	43.1	87.4
2005 5/.....	6.5	10.7	*	*	*	1.0	2.4	5.9	9.8	26.2	46.1	91.9
2004 5/.....	6.3	14.2	*	*	*	0.8	2.3	6.0	12.2	23.0	44.9	85.1
2003 5/.....	5.9	11.9	*	*	*	1.0	2.7	6.6	9.8	19.8	41.4	81.3
2002.....	6.2	14.2	*	*	0.7	0.9	2.0	5.9	11.7	23.3	43.9	86.1
2001.....	7.2	13.4	*	*	*	1.2	3.2	7.2	12.4	27.2	50.0	103.1
2000.....	6.6	18.3	*	*	*	1.0	2.3	6.0	12.1	26.6	49.8	86.8
1999.....	6.9	13.6	*	*	*	1.7	2.9	5.9	11.9	28.2	50.4	100.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

All origins 3/, both sexes

2007.....	1.0	*	*	*	*	0.0	0.2	0.7	1.2	2.5	6.4	17.3
2006.....	1.1	*	*	*	*	*	0.2	0.7	1.2	2.7	7.2	20.7
2005.....	1.2	*	*	*	*	0.1	0.2	0.7	1.2	2.9	8.0	21.4
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9	24.4
2003.....	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002.....	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001.....	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000.....	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999.....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6

All origins 3/, male

2007.....	1.0	*	*	*	*	*	0.2	0.9	1.5	3.0	7.2	18.0
2006.....	1.1	*	*	*	*	*	0.2	0.9	1.6	3.2	8.2	22.0
2005.....	1.2	*	*	*	*	*	0.3	1.0	1.7	3.6	9.7	22.4
2004.....	1.2	*	*	*	*	*	0.3	0.9	1.6	3.6	10.2	24.6
2003.....	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002.....	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001.....	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000.....	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999.....	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5

All origins 3/, female

2007.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.1	5.8	16.9
2006.....	1.1	*	*	*	*	*	0.1	0.4	0.8	2.3	6.5	20.1
2005.....	1.2	*	*	*	*	*	0.2	0.4	0.8	2.4	6.8	21.0
2004.....	1.3	*	*	*	*	*	0.1	0.4	1.0	2.8	8.0	24.3
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002.....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001.....	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000.....	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999.....	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

Non-Hispanic 4/, both sexes

2007.....	1.1	*	*	*	*	*	0.2	0.7	1.2	2.5	6.5	17.7
2006.....	1.2	*	*	*	*	*	0.2	0.7	1.2	2.8	7.3	20.7
2005.....	1.3	*	*	*	*	0.1	0.3	0.7	1.3	3.0	8.0	21.6
2004.....	1.4	*	*	*	*	*	0.2	0.7	1.3	3.2	8.9	24.6
2003.....	1.5	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.7
2002.....	1.6	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.2	28.6
2001.....	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.0	11.5	33.6
2000.....	1.8	*	*	*	*	*	0.3	0.8	1.8	4.1	11.5	35.7
1999.....	1.8	*	*	*	*	*	0.3	0.7	1.8	4.6	12.4	34.8

Non-Hispanic 4/, male

2007.....	1.1	*	*	*	*	*	0.2	1.0	1.5	3.0	7.3	18.4
2006.....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.2	8.3	22.2
2005.....	1.3	*	*	*	*	*	0.4	1.0	1.7	3.7	9.7	23.0
2004.....	1.3	*	*	*	*	*	0.3	0.9	1.6	3.7	10.2	24.8
2003.....	1.4	*	*	*	*	*	0.3	0.9	1.8	4.0	11.7	29.6
2002.....	1.5	*	*	*	*	*	0.4	0.9	1.9	4.5	12.3	31.7
2001.....	1.6	*	*	*	*	0.1	0.4	1.0	2.2	4.8	12.8	35.5
2000.....	1.8	*	*	*	*	*	0.4	1.2	2.4	5.4	13.9	41.0
1999.....	1.7	*	*	*	*	*	0.4	1.0	2.3	6.1	14.8	37.7

Non-Hispanic 4/, female

2007.....	1.2	*	*	*	*	*	0.1	0.5	0.9	2.1	6.0	17.3
2006.....	1.3	*	*	*	*	*	0.2	0.4	0.9	2.4	6.6	20.0
2005.....	1.3	*	*	*	*	*	0.2	0.4	0.9	2.4	6.9	21.0
2004.....	1.5	*	*	*	*	*	0.2	0.5	1.0	2.8	8.1	24.6
2003.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.7	25.4
2002.....	1.6	*	*	*	*	*	0.2	0.5	1.2	3.0	8.8	27.3
2001.....	1.8	*	*	*	*	*	0.2	0.5	1.2	3.3	10.6	32.8
2000.....	1.8	*	*	*	*	*	0.1	0.5	1.3	3.1	9.9	33.5
1999.....	1.9	*	*	*	*	*	0.2	0.5	1.3	3.3	10.8	33.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

Non-Hispanic white, both sexes

2007 5/.....	1.2	*	*	*	*	*	0.2	0.6	1.1	2.5	6.5	18.1
2006 5/.....	1.3	*	*	*	*	*	0.1	0.6	1.2	2.6	7.3	21.2
2005 5/.....	1.4	*	*	*	*	*	0.2	0.6	1.2	2.9	8.0	21.8
2004 5/.....	1.5	*	*	*	*	*	0.2	0.6	1.2	3.1	9.0	25.4
2003 5/.....	1.6	*	*	*	*	*	0.2	0.6	1.3	3.2	9.8	27.3
2002.....	1.7	*	*	*	*	*	0.2	0.6	1.4	3.6	10.4	29.4
2001.....	1.9	*	*	*	*	*	0.2	0.7	1.5	3.9	11.6	34.7
2000.....	1.9	*	*	*	*	*	0.2	0.7	1.8	4.0	11.6	36.3
1999.....	2.0	*	*	*	*	*	0.2	0.6	1.7	4.5	12.4	35.6

Non-Hispanic white, male

2007 5/.....	1.2	*	*	*	*	*	0.2	0.8	1.3	2.9	7.2	18.5
2006 5/.....	1.2	*	*	*	*	*	0.1	0.7	1.6	2.9	8.2	22.2
2005 5/.....	1.4	*	*	*	*	*	0.3	0.8	1.6	3.5	9.7	23.1
2004 5/.....	1.4	*	*	*	*	*	0.2	0.8	1.5	3.6	10.0	25.7
2003 5/.....	1.5	*	*	*	*	*	0.2	0.8	1.6	4.0	11.3	29.8
2002.....	1.6	*	*	*	*	*	0.3	0.8	1.7	4.4	12.4	32.6
2001.....	1.7	*	*	*	*	*	0.3	0.8	2.0	4.5	12.7	36.3
2000.....	1.9	*	*	*	*	*	0.3	1.1	2.3	5.2	13.7	41.0
1999.....	1.8	*	*	*	*	*	0.3	0.8	2.1	5.9	14.9	38.6

Non-Hispanic white, female

2007 5/.....	1.3	*	*	*	*	*	*	0.4	0.8	2.2	6.0	18.0
2006 5/.....	1.4	*	*	*	*	*	0.2	0.4	0.8	2.4	6.7	20.7
2005 5/.....	1.4	*	*	*	*	*	0.2	0.4	0.9	2.4	6.8	21.2
2004 5/.....	1.7	*	*	*	*	*	0.1	0.4	1.0	2.7	8.3	25.3
2003 5/.....	1.7	*	*	*	*	*	0.2	0.4	1.0	2.6	8.8	26.2
2002.....	1.8	*	*	*	*	*	0.2	0.5	1.1	3.0	9.0	27.9
2001.....	2.1	*	*	*	*	*	0.1	0.5	1.1	3.4	10.9	34.0
2000.....	2.0	*	*	*	*	*	0.1	0.4	1.3	3.1	10.2	34.4
1999.....	2.1	*	*	*	*	*	0.2	0.4	1.3	3.2	10.8	34.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

Non-Hispanic black, both sexes

2007 5/.....	0.8	*	*	*	*	*	*	1.3	2.1	2.9	5.7	11.5
2006 5/.....	0.9	*	*	*	*	*	0.4	1.4	2.1	3.8	7.3	14.9
2005 5/.....	1.0	*	*	*	*	*	0.5	1.3	1.7	4.0	7.3	17.5
2004 5/.....	1.0	*	*	*	*	*	0.5	1.3	2.1	4.3	7.8	16.1
2003 5/.....	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.1	17.3
2002.....	1.0	*	*	*	*	*	0.6	1.2	2.6	4.2	7.9	18.8
2001.....	1.1	*	*	*	*	*	0.6	1.4	2.8	4.5	9.0	20.9
2000.....	1.2	*	*	*	*	*	0.5	1.7	2.6	4.9	10.1	26.7
1999.....	1.3	*	*	*	*	*	0.8	1.5	2.8	5.7	11.0	25.5

Non-Hispanic black, male

2007 5/.....	0.9	*	*	*	*	*	*	2.0	3.0	4.3	7.7	*
2006 5/.....	1.2	*	*	*	*	*	*	2.3	2.9	5.8	9.5	*
2005 5/.....	1.2	*	*	*	*	*	0.8	2.2	2.7	5.9	8.4	20.0
2004 5/.....	1.1	*	*	*	*	*	*	1.8	3.0	4.7	11.1	*
2003 5/.....	1.3	*	*	*	*	*	*	2.3	3.4	5.2	16.4	*
2002.....	1.2	*	*	*	*	*	0.8	2.0	3.9	6.0	11.1	*
2001.....	1.4	*	*	*	*	*	1.0	2.3	4.3	7.0	11.7	24.3
2000.....	1.4	*	*	*	*	*	0.8	2.4	3.8	7.2	13.9	31.7
1999.....	1.5	*	*	*	*	*	1.2	2.4	4.5	8.3	12.2	28.5

Non-Hispanic black, female

2007 5/.....	0.7	*	*	*	*	*	*	*	1.4	*	4.5	11.2
2006 5/.....	0.8	*	*	*	*	*	*	*	1.5	2.3	6.2	14.2
2005 5/.....	0.8	*	*	*	*	*	*	*	*	2.7	6.7	16.5
2004 5/.....	0.9	*	*	*	*	*	*	0.9	1.3	4.1	6.0	16.9
2003 5/.....	0.8	*	*	*	*	*	*	*	*	*	8.1	16.1
2002.....	0.9	*	*	*	*	*	*	*	1.6	2.9	6.1	18.4
2001.....	0.9	*	*	*	*	*	*	*	1.5	2.7	7.4	19.7
2000.....	1.0	*	*	*	*	*	*	1.1	1.7	3.3	8.0	24.9
1999.....	1.1	*	*	*	*	*	*	*	*	3.9	10.3	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diseases of appendix (K35-K38)

All origins 3/, both sexes

2007.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5
2006.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.8	1.7
2005.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	2.1
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.7
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999.....	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5

All origins 3/, male

2007.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.2	2.1
2006.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.2	2.3
2005.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.1	2.8
2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9

All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	1.2
2006.....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	1.5
2005.....	0.1	*	*	*	*	*	*	*	*	0.1	0.2	1.8
2004.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.4
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diseases of appendix (K35-K38)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.5
2006.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	1.7
2005.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.2
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	2.1
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.2
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.7
1999.....	0.1	*	*	*	*	*	0.0	0.1	0.2	0.4	0.8	1.5

Non-Hispanic 4/, male

2007.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.2	2.2
2006.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.2	2.3
2005.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.1	3.0
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.2
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.3	2.3
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	3.0
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.0	3.0

Non-Hispanic 4/, female

2007.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	1.2
2006.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.4	1.5
2005.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	1.8
2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.5
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.8	1.4
2002.....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

Non-Hispanic white, both sexes

2007 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6
2006 5/.....	0.2	*	*	*	*	*	*	*	*	0.1	0.5	0.7	1.8
2005 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.3
2004 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8
2003 5/.....	0.2	*	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.8
1999.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6

Non-Hispanic white, male

2007 5/.....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	1.2	2.3
2006 5/.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	1.1	2.4
2005 5/.....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.1	3.1
2004 5/.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	1.4	2.3
2003 5/.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.6	1.4	2.1
2001.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	1.0	2.6
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	0.4	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1

Non-Hispanic white, female

2007 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.6	1.3
2006 5/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.4	1.5
2005 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	0.7	1.9
2004 5/.....	0.2	*	*	*	*	*	*	*	*	*	0.4	0.8	1.6
2003 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	0.8	1.3
2002.....	0.2	*	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.4	2.0
2000.....	0.2	*	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7	1.0



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Hernia (K40-K46)

##### All origins 3/, both sexes

2007.....	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.3	3.6	11.3
2006.....	0.6	0.8	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.1
2005.....	0.6	1.0	*	*	*	*	0.1	0.2	0.6	1.1	3.5	12.1
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.3	3.9	11.0
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002.....	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001.....	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000.....	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999.....	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7

##### All origins 3/, male

2007.....	0.5	1.0	*	*	*	*	*	0.3	0.6	1.2	3.6	12.0
2006.....	0.5	0.9	*	*	*	*	0.1	0.2	0.5	1.3	4.1	13.5
2005.....	0.4	1.3	*	*	*	*	*	0.2	0.6	1.0	3.3	12.7
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.4	3.8	11.3
2003.....	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001.....	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000.....	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999.....	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2

##### All origins 3/, female

2007.....	0.6	*	*	*	*	*	*	0.2	0.5	1.3	3.7	11.0
2006.....	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.8	11.4
2005.....	0.7	*	*	*	*	*	0.1	0.2	0.6	1.2	3.6	11.8
2004.....	0.6	1.0	*	*	*	*	*	0.2	0.5	1.2	4.0	10.8
2003.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999.....	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hernia (K40-K46)

Non-Hispanic 4/, both sexes

2007.....	0.6	0.9	*	*	*	*	0.1	0.2	0.5	1.3	3.7	11.4
2006.....	0.6	0.9	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.3
2005.....	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.1	3.5	12.1
2004.....	0.6	0.9	*	*	*	*	0.1	0.2	0.5	1.3	3.9	11.2
2003.....	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.3	3.6	12.1
2002.....	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.2	3.8	13.0
2001.....	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.5
2000.....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.1
1999.....	0.6	1.3	*	*	*	*	0.1	0.1	0.5	1.1	3.7	12.7

Non-Hispanic 4/, male

2007.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3	3.7	12.3
2006.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	4.1	13.9
2005.....	0.5	1.5	*	*	*	*	*	0.2	0.6	1.0	3.4	12.8
2004.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.9	11.6
2003.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	3.7	11.9
2001.....	0.5	1.2	*	*	*	*	*	0.2	0.5	1.2	3.5	12.1
2000.....	0.5	1.4	*	*	*	*	*	0.2	0.4	1.1	3.9	12.8
1999.....	0.5	1.5	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.3

Non-Hispanic 4/, female

2007.....	0.7	*	*	*	*	*	*	0.2	0.5	1.3	3.7	11.0
2006.....	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.9	11.6
2005.....	0.7	*	*	*	*	*	0.1	0.2	0.6	1.2	3.6	11.8
2004.....	0.7	*	*	*	*	*	*	0.2	0.4	1.2	4.0	11.0
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.3	12.0
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.4
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.5	12.7
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.2
1999.....	0.7	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hernia (K40-K46)

Non-Hispanic white, both sexes

2007 5/.....	0.7	0.9	*	*	*	*	0.1	0.2	0.5	1.2	3.7	12.2
2006 5/.....	0.7	0.9	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.8
2005 5/.....	0.7	1.1	*	*	*	*	0.1	0.2	0.6	1.1	3.6	12.5
2004 5/.....	0.7	1.0	*	*	*	*	0.1	0.2	0.4	1.3	4.1	11.4
2003 5/.....	0.7	1.1	*	*	*	*	*	0.2	0.5	1.3	3.6	12.4
2002.....	0.7	0.9	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.6
2001.....	0.6	0.9	*	*	*	*	*	0.2	0.5	1.2	3.5	12.8
2000.....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.3
1999.....	0.6	1.4	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0

Non-Hispanic white, male

2007 5/.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.5	12.8
2006 5/.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	4.0	14.5
2005 5/.....	0.5	*	*	*	*	*	*	0.2	0.6	1.0	3.4	12.9
2004 5/.....	0.6	*	*	*	*	*	*	0.2	0.5	1.4	4.0	11.7
2003 5/.....	0.6	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.6
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.8	12.2
2001.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.5	11.9
2000.....	0.5	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.0
1999.....	0.5	*	*	*	*	*	*	0.2	0.6	1.1	4.1	13.0

Non-Hispanic white, female

2007 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.2	3.9	11.8
2006 5/.....	0.8	*	*	*	*	*	*	0.2	0.6	1.2	3.9	12.0
2005 5/.....	0.8	*	*	*	*	*	*	0.2	0.6	1.2	3.8	12.3
2004 5/.....	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	11.3
2003 5/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.3	3.3	12.4
2002.....	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	14.2
2001.....	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.5	13.1
2000.....	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.4
1999.....	0.7	*	*	*	*	*	*	*	0.4	1.1	3.4	12.9



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes

2007.....	9.7	*	*	*	0.1	0.9	6.0	18.7	24.5	26.7	28.4	19.8
2006.....	9.2	*	*	*	0.1	0.8	5.8	17.8	22.8	26.0	29.0	19.4
2005.....	9.3	*	*	*	0.1	0.8	6.1	17.7	23.5	27.2	29.0	19.7
2004.....	9.2	*	*	*	*	0.8	6.3	18.0	22.6	27.7	28.8	19.7
2003.....	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
2002.....	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001.....	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000.....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999.....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2

All origins 3/, male

2007.....	12.9	*	*	*	*	1.3	8.0	26.8	36.3	36.2	36.7	25.8
2006.....	12.1	*	*	*	*	1.0	7.6	25.9	33.5	34.7	35.9	26.1
2005.....	12.3	*	*	*	*	1.0	8.1	26.0	34.8	35.8	36.4	27.6
2004.....	12.1	*	*	*	*	1.1	8.7	26.3	32.8	37.0	36.7	25.5
2003.....	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
2002.....	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001.....	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000.....	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999.....	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1

All origins 3/, female

2007.....	6.5	*	*	*	*	0.6	3.9	10.9	13.4	18.6	22.6	17.0
2006.....	6.4	*	*	*	*	0.5	4.1	10.0	13.0	18.6	24.2	16.2
2005.....	6.4	*	*	*	*	0.6	4.1	9.7	12.9	19.9	23.9	16.0
2004.....	6.3	*	*	*	*	0.4	4.0	10.0	13.1	19.9	23.4	17.1
2003.....	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
2002.....	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001.....	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000.....	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999.....	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Hispanic, both sexes

2007.....	8.6	*	*	*	*	1.3	6.8	24.6	35.4	48.9	52.4	35.8
2006.....	8.1	*	*	*	*	1.1	6.7	23.7	34.2	43.0	54.4	40.3
2005.....	8.3	*	*	*	*	1.0	7.3	23.5	36.9	48.8	54.0	34.7
2004.....	8.2	*	*	*	*	1.0	7.6	24.8	34.6	47.3	60.2	26.2
2003.....	8.5	*	*	*	*	1.4	7.9	24.3	38.4	51.1	62.0	31.1
2002.....	8.8	*	*	*	*	1.1	8.7	26.9	38.4	56.2	53.5	45.5
2001.....	8.9	*	*	*	*	1.2	9.7	28.0	39.0	54.4	60.4	36.9
2000.....	9.0	*	*	*	*	1.4	9.8	27.2	41.4	58.3	63.8	48.4
1999.....	8.8	*	*	*	*	1.3	10.5	26.6	39.0	56.3	59.4	51.2

Hispanic, male

2007.....	11.9	*	*	*	*	1.8	10.5	38.5	55.8	68.6	59.0	43.7
2006.....	11.0	*	*	*	*	1.4	9.9	37.4	54.1	56.9	61.0	51.9
2005.....	11.6	*	*	*	*	1.4	10.9	37.8	58.4	69.1	68.8	36.1
2004.....	11.1	*	*	*	*	1.5	11.6	40.8	52.4	59.4	70.8	29.4
2003.....	11.6	*	*	*	*	2.0	12.3	40.2	62.2	61.3	71.3	35.4
2002.....	12.2	*	*	*	*	1.6	13.5	44.2	60.0	77.8	64.4	45.2
2001.....	12.7	*	*	*	*	1.8	15.0	46.4	63.9	77.1	76.0	41.9
2000.....	12.7	*	*	*	*	2.3	15.0	45.6	66.1	83.4	80.1	60.5
1999.....	12.3	*	*	*	*	1.8	16.5	44.3	62.6	79.1	70.9	66.9

Hispanic, female

2007.....	5.1	*	*	*	*	0.6	2.5	10.2	16.5	32.9	47.8	31.4
2006.....	5.0	*	*	*	*	0.6	3.0	9.7	16.0	31.6	49.8	34.2
2005.....	4.8	*	*	*	*	*	3.2	8.9	17.1	32.4	43.7	33.9
2004.....	5.0	*	*	*	*	*	3.1	8.6	18.5	37.6	52.9	24.5
2003.....	5.2	*	*	*	*	0.6	3.2	8.3	17.0	42.9	55.6	28.9
2002.....	5.2	*	*	*	*	*	3.5	9.7	18.9	38.9	46.1	45.6
2001.....	5.0	*	*	*	*	*	4.0	9.9	17.0	36.5	49.9	34.4
2000.....	5.1	*	*	*	*	*	4.1	9.3	19.5	38.7	52.8	42.5
1999.....	5.0	*	*	*	*	*	4.1	9.3	18.1	38.4	51.7	43.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 4/, both sexes

2007.....	9.8	*	*	*	0.1	0.9	5.8	18.0	23.4	24.9	26.8	19.0
2006.....	9.4	*	*	*	0.1	0.7	5.7	17.1	21.8	24.6	27.3	18.3
2005.....	9.4	*	*	*	0.1	0.7	5.9	17.0	22.3	25.5	27.4	18.9
2004.....	9.3	*	*	*	*	0.7	6.1	17.2	21.5	26.2	27.0	19.4
2003.....	9.6	*	*	*	*	0.8	6.6	17.6	21.7	27.9	28.2	19.6
2002.....	9.5	*	*	*	0.1	0.9	6.7	17.0	21.5	27.5	30.2	20.4
2001.....	9.5	*	*	*	0.1	0.9	7.1	17.5	21.3	28.2	28.7	21.6
2000.....	9.4	*	*	*	0.1	1.0	7.1	16.7	22.3	27.9	29.5	22.1
1999.....	9.5	*	*	*	*	1.0	6.9	16.5	22.5	29.0	30.7	22.1

Non-Hispanic 4/, male

2007.....	13.0	*	*	*	*	1.1	7.4	25.2	34.5	33.6	35.2	24.7
2006.....	12.3	*	*	*	*	0.9	7.1	24.5	31.5	32.9	34.2	24.6
2005.....	12.3	*	*	*	*	0.8	7.6	24.4	32.7	33.2	34.4	27.1
2004.....	12.3	*	*	*	*	1.0	8.2	24.6	31.0	35.3	34.8	25.3
2003.....	12.6	*	*	*	*	1.1	8.6	26.0	31.3	37.5	35.0	27.1
2002.....	12.2	*	*	*	*	1.1	8.6	24.9	30.7	35.7	37.0	27.7
2001.....	12.3	*	*	*	*	1.2	9.1	25.6	30.6	37.6	34.9	31.4
2000.....	12.4	*	*	*	*	1.2	9.6	25.0	31.8	37.1	36.5	33.0
1999.....	12.5	*	*	*	*	1.2	9.5	25.2	32.4	39.3	38.2	31.6

Non-Hispanic 4/, female

2007.....	6.8	*	*	*	*	0.6	4.2	11.0	13.1	17.4	21.0	16.3
2006.....	6.6	*	*	*	*	0.5	4.2	10.0	12.7	17.6	22.6	15.3
2005.....	6.6	*	*	*	*	0.6	4.2	9.8	12.5	18.9	22.7	15.2
2004.....	6.5	*	*	*	*	0.5	4.1	10.1	12.7	18.5	21.8	16.7
2003.....	6.7	*	*	*	*	0.5	4.6	9.5	12.7	19.8	23.7	16.3
2002.....	6.9	*	*	*	*	0.7	4.9	9.4	13.0	20.6	25.8	17.3
2001.....	6.9	*	*	*	*	0.7	5.1	9.8	12.7	20.4	24.7	17.5
2000.....	6.7	*	*	*	*	0.7	4.7	8.7	13.6	20.3	24.9	17.7
1999.....	6.6	*	*	*	*	0.8	4.4	8.1	13.3	20.6	25.8	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic white, both sexes

2007 5/.....	10.7	*	*	*	*	0.9	6.1	18.5	23.5	25.7	28.0	19.7
2006 5/.....	10.3	*	*	*	*	0.7	5.9	17.6	22.1	25.7	28.5	19.0
2005 5/.....	10.2	*	*	*	*	0.6	6.2	17.1	22.3	26.5	28.2	19.5
2004 5/.....	10.1	*	*	*	*	0.7	6.2	17.4	21.3	27.1	28.3	19.9
2003 5/.....	10.3	*	*	*	*	0.8	6.8	17.5	21.5	28.9	29.7	20.3
2002.....	10.2	*	*	*	*	0.8	6.9	16.8	21.4	28.5	31.5	21.2
2001.....	10.1	*	*	*	*	0.9	7.0	17.0	21.0	28.9	29.9	22.5
2000.....	10.0	*	*	*	*	0.8	7.0	16.2	22.1	28.4	30.6	22.8
1999.....	9.9	*	*	*	*	0.9	6.6	15.8	21.8	29.5	31.7	22.6

Non-Hispanic white, male

2007 5/.....	14.2	*	*	*	*	1.1	7.8	26.0	34.5	34.3	36.5	26.1
2006 5/.....	13.4	*	*	*	*	0.9	7.4	25.1	31.9	33.9	35.4	25.6
2005 5/.....	13.3	*	*	*	*	0.8	7.9	24.6	32.5	34.2	34.9	28.1
2004 5/.....	13.2	*	*	*	*	1.0	8.3	24.7	30.6	36.1	36.3	26.4
2003 5/.....	13.6	*	*	*	*	1.0	9.0	25.9	30.8	38.7	36.5	27.9
2002.....	13.2	*	*	*	*	1.0	8.8	24.7	30.5	36.6	38.7	28.7
2001.....	13.0	*	*	*	*	1.1	8.9	24.8	29.7	38.1	36.3	32.9
2000.....	13.1	*	*	*	*	1.1	9.4	24.4	31.5	37.7	37.8	34.1
1999.....	13.1	*	*	*	*	1.1	9.1	24.2	31.5	39.7	39.2	32.7

Non-Hispanic white, female

2007 5/.....	7.4	*	*	*	*	0.6	4.5	11.1	13.1	18.3	22.1	16.7
2006 5/.....	7.2	*	*	*	*	0.5	4.3	10.2	12.9	18.5	23.6	15.9
2005 5/.....	7.2	*	*	*	*	0.5	4.4	9.7	12.6	19.9	23.6	15.6
2004 5/.....	7.1	*	*	*	*	0.5	4.2	10.1	12.5	19.4	22.8	17.0
2003 5/.....	7.2	*	*	*	*	0.5	4.6	9.3	12.6	20.5	25.2	17.0
2002.....	7.5	*	*	*	*	0.6	5.0	9.0	12.9	21.5	26.8	18.0
2001.....	7.4	*	*	*	*	0.7	5.1	9.3	12.8	21.1	25.7	18.2
2000.....	7.0	*	*	*	*	0.6	4.5	8.0	13.2	20.6	25.9	18.2
1999.....	6.9	*	*	*	*	0.7	4.1	7.5	12.7	21.0	26.7	18.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic black, both sexes

2007 5/.....	6.6	*	*	*	*	0.5	3.7	16.5	26.3	22.3	15.5	10.0
2006 5/.....	6.1	*	*	*	*	0.4	4.6	15.8	22.4	19.0	16.3	10.9
2005 5/.....	6.7	*	*	*	*	0.6	4.3	18.0	25.3	20.5	19.3	12.0
2004 5/.....	6.8	*	*	*	*	0.4	5.6	18.7	25.9	22.0	14.4	12.1
2003 5/.....	7.1	*	*	*	*	0.8	5.8	20.2	26.1	24.1	14.1	12.5
2002.....	7.0	*	*	*	*	0.9	5.6	21.2	25.3	21.8	18.3	12.1
2001.....	7.7	*	*	*	*	0.9	7.5	22.9	26.3	25.3	18.2	10.6
2000.....	7.7	*	*	*	*	1.1	8.0	22.6	26.1	26.5	18.1	14.0
1999.....	8.0	*	*	*	*	1.0	8.2	23.0	29.7	26.9	21.7	16.8

Non-Hispanic black, male

2007 5/.....	9.1	*	*	*	*	*	4.8	23.4	41.1	36.4	24.8	*
2006 5/.....	8.3	*	*	*	*	*	5.9	22.9	33.9	29.3	24.5	*
2005 5/.....	9.2	*	*	*	*	*	5.5	26.5	39.9	31.4	29.4	*
2004 5/.....	9.4	*	*	*	*	*	7.6	27.6	40.0	34.5	20.5	*
2003 5/.....	9.8	*	*	*	*	1.0	7.3	31.0	41.2	33.7	20.2	22.4
2002.....	9.3	*	*	*	*	1.1	6.9	30.7	38.2	32.4	22.0	*
2001.....	10.7	*	*	*	*	1.0	10.4	35.1	42.0	39.7	22.5	*
2000.....	10.3	*	*	*	*	1.1	10.7	33.4	38.5	38.2	25.0	*
1999.....	11.0	*	*	*	*	1.1	11.3	35.8	43.8	41.0	30.0	*

Non-Hispanic black, female

2007 5/.....	4.3	*	*	*	*	*	2.8	10.6	14.3	12.3	10.2	10.1
2006 5/.....	4.2	*	*	*	*	*	3.5	9.7	13.2	11.6	11.7	9.0
2005 5/.....	4.4	*	*	*	*	*	3.2	10.8	13.6	12.7	13.5	10.7
2004 5/.....	4.4	*	*	*	*	*	3.8	11.0	14.5	13.1	10.9	11.2
2003 5/.....	4.6	*	*	*	*	*	4.6	10.9	14.0	17.3	10.6	8.7
2002.....	5.0	*	*	*	*	0.8	4.5	13.2	15.0	14.2	16.2	10.5
2001.....	4.9	*	*	*	*	0.7	5.0	12.5	13.7	15.1	15.8	9.8
2000.....	5.4	*	*	*	*	1.1	5.7	13.4	16.2	18.3	14.2	11.8
1999.....	5.4	*	*	*	*	1.0	5.5	12.1	18.5	16.9	17.0	15.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

All origins 3/, both sexes

2007.....	4.8	*	*	*	*	0.7	4.2	11.7	13.2	10.1	5.7	2.1
2006.....	4.4	*	*	*	*	0.6	4.0	10.5	12.1	9.4	5.9	2.2
2005.....	4.4	*	*	*	*	0.5	4.1	10.6	12.3	9.9	5.7	1.8
2004.....	4.3	*	*	*	*	0.6	4.2	10.6	11.7	10.0	5.5	2.1
2003.....	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002.....	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001.....	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000.....	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999.....	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8

All origins 3/, male

2007.....	7.1	*	*	*	*	1.0	5.6	17.0	20.9	16.9	10.4	4.7
2006.....	6.4	*	*	*	*	0.9	5.1	15.5	18.9	15.7	10.2	4.7
2005.....	6.5	*	*	*	*	0.7	5.4	15.7	19.5	15.9	10.2	4.2
2004.....	6.3	*	*	*	*	0.8	5.8	15.5	18.2	16.5	10.1	4.6
2003.....	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002.....	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001.....	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000.....	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999.....	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1

All origins 3/, female

2007.....	2.5	*	*	*	*	0.5	2.9	6.5	6.0	4.3	2.5	0.9
2006.....	2.4	*	*	*	*	0.4	2.8	5.8	5.8	4.2	3.0	1.1
2005.....	2.3	*	*	*	*	0.4	2.8	5.6	5.5	4.9	2.7	0.7
2004.....	2.3	*	*	*	*	0.3	2.7	5.7	5.6	4.5	2.3	1.0
2003.....	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002.....	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001.....	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000.....	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999.....	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Hispanic, both sexes

2007.....	4.7	*	*	*	*	0.9	4.7	16.1	20.2	19.9	14.2	7.2
2006.....	4.3	*	*	*	*	0.9	4.6	15.1	19.3	16.9	14.5	*
2005.....	4.2	*	*	*	*	0.8	4.9	14.4	18.9	20.2	11.4	*
2004.....	4.1	*	*	*	*	0.7	5.3	15.2	17.2	16.1	14.6	*
2003.....	4.3	*	*	*	*	0.9	5.6	14.4	21.4	17.7	14.1	*
2002.....	4.3	*	*	*	*	0.7	5.5	16.1	19.9	20.8	10.5	*
2001.....	4.7	*	*	*	*	0.9	6.3	17.1	21.4	21.1	17.0	*
2000.....	4.8	*	*	*	*	1.0	6.9	16.8	23.4	22.9	15.4	*
1999.....	4.6	*	*	*	*	0.8	6.9	16.8	21.7	20.6	14.4	*

Hispanic, male

2007.....	7.5	*	*	*	*	1.4	7.5	26.1	35.9	37.5	26.0	*
2006.....	6.9	*	*	*	*	1.3	6.9	24.7	33.8	32.1	27.2	*
2005.....	6.8	*	*	*	*	1.2	7.6	23.9	33.4	38.1	23.7	*
2004.....	6.6	*	*	*	*	1.2	8.2	26.6	29.7	28.6	28.3	*
2003.....	6.9	*	*	*	*	1.3	8.8	25.5	38.4	29.0	26.5	*
2002.....	7.1	*	*	*	*	1.2	9.0	27.0	36.3	40.3	18.1	*
2001.....	7.7	*	*	*	*	1.5	10.0	29.3	38.9	41.9	30.4	*
2000.....	7.9	*	*	*	*	1.8	10.6	29.3	42.8	46.6	33.9	*
1999.....	7.4	*	*	*	*	1.2	10.8	29.1	40.0	39.5	27.1	*

Hispanic, female

2007.....	1.6	*	*	*	*	*	1.6	5.7	5.7	5.5	5.9	*
2006.....	1.6	*	*	*	*	*	2.1	5.3	6.0	4.6	5.5	*
2005.....	1.4	*	*	*	*	*	2.0	4.8	5.6	5.7	*	*
2004.....	1.4	*	*	*	*	*	2.0	3.6	5.9	6.1	5.2	*
2003.....	1.5	*	*	*	*	*	2.1	3.4	6.1	8.7	5.5	*
2002.....	1.4	*	*	*	*	*	1.8	5.1	5.1	5.3	*	*
2001.....	1.5	*	*	*	*	*	2.2	5.1	5.9	4.7	7.9	*
2000.....	1.4	*	*	*	*	*	2.9	4.5	6.2	4.3	*	*
1999.....	1.5	*	*	*	*	*	2.8	4.9	5.6	5.7	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Non-Hispanic 4/, both sexes

2007.....	4.8	*	*	*	*	0.7	4.2	11.1	12.5	9.3	5.2	1.9
2006.....	4.4	*	*	*	*	0.5	3.8	10.0	11.4	8.8	5.4	1.9
2005.....	4.4	*	*	*	*	0.5	3.9	10.1	11.6	9.0	5.4	1.7
2004.....	4.3	*	*	*	*	0.5	4.1	10.0	11.2	9.5	4.9	2.1
2003.....	4.2	*	*	*	*	0.5	4.2	10.1	10.8	9.1	5.2	1.7
2002.....	4.2	*	*	*	*	0.6	4.2	9.7	10.8	9.3	5.5	1.9
2001.....	4.2	*	*	*	*	0.6	4.6	10.1	10.7	9.3	5.2	2.2
2000.....	4.2	*	*	*	*	0.6	4.7	9.7	11.3	9.5	5.7	2.2
1999.....	4.2	*	*	*	*	0.6	4.6	9.8	11.3	9.9	6.1	2.5

Non-Hispanic 4/, male

2007.....	7.0	*	*	*	*	0.8	5.3	15.8	19.5	15.3	9.4	3.9
2006.....	6.3	*	*	*	*	0.7	4.8	14.3	17.5	14.3	9.1	3.9
2005.....	6.4	*	*	*	*	0.6	5.0	14.6	18.2	14.1	9.4	3.8
2004.....	6.3	*	*	*	*	0.7	5.3	14.2	17.2	15.6	9.1	4.5
2003.....	6.2	*	*	*	*	0.8	5.5	15.0	16.6	15.1	9.2	3.4
2002.....	6.0	*	*	*	*	0.7	5.3	14.4	16.7	14.8	9.2	4.3
2001.....	6.2	*	*	*	*	0.8	5.9	14.8	16.7	14.9	9.2	5.1
2000.....	6.2	*	*	*	*	0.8	6.3	14.7	17.1	15.4	10.5	5.0
1999.....	6.4	*	*	*	*	0.7	6.2	15.1	17.7	16.7	10.5	5.5

Non-Hispanic 4/, female

2007.....	2.7	*	*	*	*	0.5	3.1	6.6	6.0	4.2	2.3	0.9
2006.....	2.5	*	*	*	*	0.3	2.9	5.8	5.8	4.1	2.9	1.0
2005.....	2.5	*	*	*	*	0.4	2.9	5.7	5.5	4.8	2.7	0.8
2004.....	2.4	*	*	*	*	0.3	2.8	5.9	5.6	4.3	2.2	1.0
2003.....	2.3	*	*	*	*	0.3	3.0	5.3	5.4	4.1	2.6	1.0
2002.....	2.4	*	*	*	*	0.5	3.1	5.2	5.4	4.8	3.0	0.9
2001.....	2.4	*	*	*	*	0.5	3.2	5.5	5.0	4.7	2.5	1.0
2000.....	2.3	*	*	*	*	0.4	3.1	4.8	5.9	4.5	2.5	1.0
1999.....	2.2	*	*	*	*	0.5	2.9	4.7	5.3	4.2	3.3	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Non-Hispanic white, both sexes

2007 5/.....	5.2	*	*	*	*	0.6	4.4	11.6	12.8	9.5	5.3	1.9
2006 5/.....	4.7	*	*	*	*	0.5	4.0	10.4	11.8	9.0	5.6	2.0
2005 5/.....	4.7	*	*	*	*	0.4	4.1	10.4	11.8	9.2	5.5	1.8
2004 5/.....	4.6	*	*	*	*	0.5	4.2	10.3	11.1	9.7	5.1	2.2
2003 5/.....	4.5	*	*	*	*	0.5	4.4	10.1	10.7	9.3	5.4	1.7
2002.....	4.4	*	*	*	*	0.5	4.3	9.7	10.8	9.5	5.5	1.9
2001.....	4.4	*	*	*	*	0.6	4.5	9.9	10.5	9.3	5.3	2.1
2000.....	4.3	*	*	*	*	0.6	4.5	9.4	11.1	9.5	5.6	2.2
1999.....	4.3	*	*	*	*	0.6	4.4	9.5	10.8	9.7	6.0	2.4

Non-Hispanic white, male

2007 5/.....	7.6	*	*	*	*	0.8	5.5	16.5	19.9	15.3	9.6	4.1
2006 5/.....	6.9	*	*	*	*	0.7	5.1	14.9	18.0	14.5	9.3	4.0
2005 5/.....	6.8	*	*	*	*	0.5	5.2	14.9	18.4	14.2	9.4	3.9
2004 5/.....	6.7	*	*	*	*	0.7	5.5	14.6	17.0	15.7	9.2	4.7
2003 5/.....	6.6	*	*	*	*	0.8	5.6	15.1	16.4	15.3	9.4	3.5
2002.....	6.4	*	*	*	*	0.7	5.4	14.4	16.7	14.9	9.4	4.1
2001.....	6.4	*	*	*	*	0.8	5.8	14.7	16.4	14.6	9.3	4.8
2000.....	6.5	*	*	*	*	0.8	6.1	14.5	17.0	15.4	10.6	5.1
1999.....	6.5	*	*	*	*	0.7	6.0	14.7	17.1	16.3	10.3	5.1

Non-Hispanic white, female

2007 5/.....	2.9	*	*	*	*	0.4	3.3	6.8	6.0	4.4	2.4	0.9
2006 5/.....	2.7	*	*	*	*	0.3	3.0	6.0	6.0	4.2	3.0	1.1
2005 5/.....	2.6	*	*	*	*	0.3	3.0	5.9	5.5	4.9	2.8	0.8
2004 5/.....	2.5	*	*	*	*	0.3	3.0	6.0	5.5	4.5	2.3	1.0
2003 5/.....	2.4	*	*	*	*	0.3	3.1	5.3	5.2	4.1	2.8	1.0
2002.....	2.4	*	*	*	*	0.4	3.2	5.1	5.2	4.9	3.0	1.0
2001.....	2.4	*	*	*	*	0.4	3.2	5.3	5.0	4.8	2.5	1.0
2000.....	2.2	*	*	*	*	0.4	3.0	4.4	5.5	4.6	2.4	1.0
1999.....	2.2	*	*	*	*	0.5	2.8	4.3	4.9	4.2	3.2	1.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Non-Hispanic black, both sexes

2007 5/.....	3.1	*	*	*	*	*	2.3	8.4	12.1	9.7	3.9	*
2006 5/.....	2.7	*	*	*	*	*	2.6	7.7	9.5	7.8	4.1	*
2005 5/.....	3.1	*	*	*	*	*	2.7	9.1	11.5	8.8	4.6	*
2004 5/.....	3.1	*	*	*	*	*	3.0	9.2	12.8	9.1	3.2	*
2003 5/.....	3.3	*	*	*	*	*	3.4	10.0	12.7	9.4	3.4	*
2002.....	3.2	*	*	*	*	0.5	3.0	10.2	12.3	8.7	5.1	*
2001.....	3.7	*	*	*	*	0.5	4.4	11.7	13.1	10.8	4.6	*
2000.....	3.9	*	*	*	*	0.7	5.1	12.3	13.5	10.4	5.6	*
1999.....	4.2	*	*	*	*	0.5	5.1	12.9	15.4	11.9	7.2	*

Non-Hispanic black, male

2007 5/.....	4.6	*	*	*	*	*	3.0	12.5	19.9	18.7	8.3	*
2006 5/.....	3.8	*	*	*	*	*	3.1	11.3	15.0	13.7	7.5	*
2005 5/.....	4.4	*	*	*	*	*	3.2	13.7	18.6	15.4	9.0	*
2004 5/.....	4.6	*	*	*	*	*	4.1	13.2	20.7	17.2	*	*
2003 5/.....	4.9	*	*	*	*	*	4.2	16.0	20.3	16.2	6.9	*
2002.....	4.6	*	*	*	*	*	3.9	15.1	18.6	15.1	8.8	*
2001.....	5.5	*	*	*	*	*	6.1	17.4	21.7	20.0	7.8	*
2000.....	5.5	*	*	*	*	*	7.1	17.9	19.6	18.4	10.2	*
1999.....	6.1	*	*	*	*	*	6.9	19.7	24.6	21.9	12.8	*

Non-Hispanic black, female

2007 5/.....	1.7	*	*	*	*	*	1.7	4.9	5.8	3.4	*	*
2006 5/.....	1.7	*	*	*	*	*	2.0	4.6	5.1	3.5	*	*
2005 5/.....	1.9	*	*	*	*	*	2.3	5.2	5.8	4.1	*	*
2004 5/.....	1.8	*	*	*	*	*	2.0	5.7	6.4	3.4	*	*
2003 5/.....	1.9	*	*	*	*	*	2.6	4.8	6.6	4.6	*	*
2002.....	2.1	*	*	*	*	*	2.2	6.1	7.2	4.2	*	*
2001.....	2.2	*	*	*	*	*	2.8	6.8	6.2	4.3	*	*
2000.....	2.5	*	*	*	*	*	3.4	7.5	8.6	4.8	*	*
1999.....	2.4	*	*	*	*	*	3.5	7.0	8.1	4.8	4.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/, both sexes

2007.....	4.9	*	*	*	*	0.2	1.7	7.0	11.3	16.6	22.6	17.7
2006.....	4.8	*	*	*	*	0.2	1.9	7.3	10.7	16.6	23.0	17.2
2005.....	4.9	*	*	*	*	0.2	2.0	7.1	11.2	17.3	23.2	17.8
2004.....	4.9	*	*	*	*	0.2	2.1	7.5	10.9	17.8	23.3	17.6
2003.....	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
2002.....	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001.....	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000.....	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999.....	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4

All origins 3/, male

2007.....	5.8	*	*	*	*	0.3	2.3	9.8	15.5	19.3	26.3	21.0
2006.....	5.7	*	*	*	*	0.2	2.5	10.4	14.6	19.0	25.7	21.4
2005.....	5.8	*	*	*	*	0.2	2.7	10.3	15.3	19.9	26.2	23.3
2004.....	5.8	*	*	*	*	0.3	2.9	10.8	14.6	20.5	26.6	21.0
2003.....	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
2002.....	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001.....	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000.....	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999.....	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0

All origins 3/, female

2007.....	4.0	*	*	*	*	0.2	1.1	4.4	7.4	14.3	20.1	16.1
2006.....	4.0	*	*	*	*	0.2	1.3	4.2	7.1	14.5	21.2	15.2
2005.....	4.0	*	*	*	*	0.2	1.3	4.1	7.4	15.1	21.2	15.3
2004.....	4.1	*	*	*	*	0.1	1.3	4.2	7.5	15.4	21.1	16.1
2003.....	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
2002.....	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001.....	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000.....	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999.....	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Hispanic, both sexes

2007.....	3.9	*	*	*	*	0.3	2.0	8.5	15.2	29.1	38.2	28.6
2006.....	3.8	*	*	*	*	*	2.0	8.6	14.9	26.0	40.0	32.9
2005.....	4.1	*	*	*	*	*	2.4	9.1	18.0	28.6	42.6	30.4
2004.....	4.1	*	*	*	*	*	2.3	9.6	17.4	31.2	45.6	24.3
2003.....	4.2	*	*	*	*	0.4	2.3	9.8	17.0	33.4	47.9	26.9
2002.....	4.5	*	*	*	*	0.4	3.2	10.8	18.5	35.3	43.0	38.8
2001.....	4.3	*	*	*	*	*	3.5	10.9	17.6	33.3	43.5	30.6
2000.....	4.3	*	*	*	*	0.4	2.9	10.5	18.0	35.4	48.4	44.5
1999.....	4.2	*	*	*	*	0.5	3.6	9.7	17.2	35.7	45.0	40.0

Hispanic, male

2007.....	4.3	*	*	*	*	*	3.0	12.4	19.9	31.1	33.0	25.4
2006.....	4.1	*	*	*	*	*	3.0	12.7	20.3	24.8	33.8	33.8
2005.....	4.8	*	*	*	*	*	3.4	13.9	25.0	31.0	45.1	*
2004.....	4.5	*	*	*	*	*	3.4	14.1	22.7	30.8	42.6	*
2003.....	4.7	*	*	*	*	0.6	3.5	14.7	23.8	32.3	44.9	*
2002.....	5.1	*	*	*	*	*	4.6	17.1	23.8	37.5	46.4	*
2001.....	5.0	*	*	*	*	*	5.1	17.1	25.0	35.1	45.6	*
2000.....	4.8	*	*	*	*	0.6	4.4	16.2	23.3	36.7	46.2	50.4
1999.....	4.9	*	*	*	*	0.6	5.7	15.1	22.6	39.5	43.8	45.3

Hispanic, female

2007.....	3.5	*	*	*	*	*	0.9	4.5	10.9	27.4	41.8	30.3
2006.....	3.4	*	*	*	*	*	0.9	4.4	10.0	27.0	44.3	32.4
2005.....	3.4	*	*	*	*	*	1.2	4.2	11.5	26.7	40.9	33.9
2004.....	3.7	*	*	*	*	*	1.1	5.0	12.5	31.5	47.7	24.5
2003.....	3.7	*	*	*	*	*	1.1	5.0	10.8	34.3	50.0	26.6
2002.....	3.8	*	*	*	*	*	1.7	4.6	13.8	33.6	40.8	42.3
2001.....	3.4	*	*	*	*	*	1.8	4.7	11.0	31.8	42.0	33.5
2000.....	3.7	*	*	*	*	*	1.3	4.8	13.3	34.3	49.9	41.5
1999.....	3.5	*	*	*	*	*	1.3	4.5	12.4	32.7	45.8	37.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 4/, both sexes

2007.....	5.1	*	*	*	*	0.2	1.6	6.8	10.9	15.6	21.6	17.1
2006.....	5.0	*	*	*	*	0.2	1.8	7.1	10.3	15.8	21.9	16.4
2005.....	5.1	*	*	*	*	0.2	1.9	6.9	10.6	16.4	22.0	17.2
2004.....	5.0	*	*	*	*	0.2	2.1	7.2	10.4	16.7	22.0	17.3
2003.....	5.4	*	*	*	*	0.2	2.4	7.5	10.9	18.7	23.0	17.9
2002.....	5.4	*	*	*	*	0.3	2.5	7.3	10.7	18.2	24.8	18.5
2001.....	5.3	*	*	*	*	0.3	2.5	7.5	10.7	18.9	23.6	19.4
2000.....	5.2	*	*	*	0.1	0.3	2.5	7.0	11.0	18.4	23.8	19.9
1999.....	5.2	*	*	*	*	0.3	2.3	6.7	11.2	19.2	24.6	19.6

Non-Hispanic 4/, male

2007.....	6.0	*	*	*	*	0.3	2.2	9.4	15.0	18.3	25.8	20.7
2006.....	6.0	*	*	*	*	0.2	2.4	10.1	14.1	18.6	25.1	20.8
2005.....	6.0	*	*	*	*	0.2	2.6	9.8	14.5	19.1	25.0	23.3
2004.....	6.0	*	*	*	*	0.3	2.8	10.4	13.9	19.7	25.7	20.7
2003.....	6.4	*	*	*	*	0.3	3.2	11.0	14.7	22.3	25.9	23.7
2002.....	6.2	*	*	*	*	0.4	3.3	10.6	14.0	20.9	27.7	23.4
2001.....	6.2	*	*	*	*	0.4	3.1	10.8	13.8	22.8	25.7	26.4
2000.....	6.1	*	*	*	*	0.3	3.3	10.3	14.7	21.7	26.0	28.0
1999.....	6.1	*	*	*	*	0.4	3.3	10.1	14.7	22.6	27.7	26.2

Non-Hispanic 4/, female

2007.....	4.1	*	*	*	*	0.1	1.1	4.4	7.1	13.2	18.7	15.4
2006.....	4.1	*	*	*	*	0.1	1.3	4.2	6.9	13.5	19.7	14.3
2005.....	4.1	*	*	*	*	0.2	1.3	4.1	7.0	14.1	20.0	14.4
2004.....	4.1	*	*	*	*	0.1	1.3	4.1	7.1	14.2	19.6	15.7
2003.....	4.4	*	*	*	*	0.2	1.6	4.2	7.3	15.7	21.1	15.3
2002.....	4.6	*	*	*	*	0.3	1.8	4.2	7.6	15.8	22.8	16.4
2001.....	4.5	*	*	*	*	0.2	1.9	4.3	7.7	15.7	22.2	16.5
2000.....	4.4	*	*	*	*	0.3	1.6	3.9	7.6	15.8	22.4	16.7
1999.....	4.4	*	*	*	*	0.3	1.4	3.5	8.0	16.3	22.6	17.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic white, both sexes

2007 5/.....	5.6	*	*	*	*	0.2	1.7	6.9	10.7	16.3	22.7	17.8
2006 5/.....	5.5	*	*	*	*	0.2	1.9	7.2	10.3	16.6	22.9	16.9
2005 5/.....	5.5	*	*	*	*	0.2	2.1	6.7	10.5	17.3	22.7	17.7
2004 5/.....	5.5	*	*	*	*	0.2	2.0	7.1	10.2	17.4	23.2	17.7
2003 5/.....	5.9	*	*	*	*	0.2	2.4	7.4	10.8	19.6	24.3	18.6
2002.....	5.9	*	*	*	*	0.3	2.6	7.1	10.7	19.0	26.0	19.3
2001.....	5.8	*	*	*	*	0.3	2.5	7.1	10.5	19.6	24.7	20.4
2000.....	5.7	*	*	*	*	0.3	2.4	6.7	11.0	18.9	25.0	20.6
1999.....	5.6	*	*	*	*	0.3	2.3	6.4	11.0	19.8	25.6	20.2

Non-Hispanic white, male

2007 5/.....	6.6	*	*	*	*	0.3	2.3	9.5	14.6	19.0	26.9	22.0
2006 5/.....	6.5	*	*	*	*	0.2	2.3	10.2	13.9	19.4	26.1	21.5
2005 5/.....	6.5	*	*	*	*	0.2	2.7	9.7	14.1	20.0	25.4	24.2
2004 5/.....	6.5	*	*	*	*	0.2	2.8	10.2	13.6	20.4	27.1	21.7
2003 5/.....	6.9	*	*	*	*	0.2	3.3	10.9	14.4	23.4	27.1	24.5
2002.....	6.7	*	*	*	*	0.3	3.4	10.3	13.7	21.7	29.3	24.7
2001.....	6.6	*	*	*	*	0.4	3.0	10.2	13.3	23.5	26.9	28.1
2000.....	6.6	*	*	*	*	0.3	3.4	9.9	14.6	22.3	27.2	29.0
1999.....	6.6	*	*	*	*	0.4	3.2	9.5	14.4	23.3	28.9	27.5

Non-Hispanic white, female

2007 5/.....	4.6	*	*	*	*	*	1.2	4.3	7.0	13.9	19.7	15.7
2006 5/.....	4.6	*	*	*	*	*	1.4	4.2	6.9	14.2	20.6	14.8
2005 5/.....	4.6	*	*	*	*	0.2	1.4	3.8	7.1	15.0	20.8	14.8
2004 5/.....	4.5	*	*	*	*	*	1.2	4.1	7.0	14.9	20.5	16.0
2003 5/.....	4.8	*	*	*	*	0.2	1.5	4.0	7.4	16.4	22.4	16.0
2002.....	5.0	*	*	*	*	0.2	1.8	3.9	7.7	16.6	23.8	17.0
2001.....	5.0	*	*	*	*	0.3	1.9	4.1	7.9	16.3	23.2	17.2
2000.....	4.8	*	*	*	*	0.3	1.5	3.6	7.6	16.0	23.5	17.2
1999.....	4.8	*	*	*	*	0.2	1.4	3.2	7.7	16.8	23.5	17.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic black, both sexes

2007 5/.....	3.5	*	*	*	*	*	1.4	8.1	14.2	12.6	11.6	8.4
2006 5/.....	3.4	*	*	*	*	*	2.1	8.1	12.9	11.2	12.2	10.1
2005 5/.....	3.6	*	*	*	*	*	1.6	9.0	13.8	11.7	14.7	10.7
2004 5/.....	3.6	*	*	*	*	*	2.6	9.5	13.1	12.9	11.2	11.2
2003 5/.....	3.8	*	*	*	*	0.4	2.5	10.1	13.4	14.7	10.6	11.0
2002.....	3.8	*	*	*	*	0.4	2.6	11.0	13.0	13.1	13.2	9.7
2001.....	3.9	*	*	*	*	*	3.2	11.2	13.2	14.5	13.6	7.8
2000.....	3.8	*	*	*	*	0.4	2.9	10.3	12.7	16.1	12.6	12.4
1999.....	3.9	*	*	*	*	0.5	3.1	10.1	14.3	15.0	14.5	12.3

Non-Hispanic black, male

2007 5/.....	4.5	*	*	*	*	*	1.8	10.9	21.3	17.7	16.5	*
2006 5/.....	4.4	*	*	*	*	*	2.8	11.6	18.9	15.6	17.0	*
2005 5/.....	4.7	*	*	*	*	*	2.3	12.8	21.3	16.1	20.5	*
2004 5/.....	4.8	*	*	*	*	*	3.5	14.5	19.2	17.2	15.1	*
2003 5/.....	4.9	*	*	*	*	*	3.0	14.9	20.9	17.5	13.3	*
2002.....	4.7	*	*	*	*	*	2.9	15.5	19.6	17.3	13.2	*
2001.....	5.3	*	*	*	*	*	4.3	17.6	20.3	19.7	14.7	*
2000.....	4.8	*	*	*	*	*	3.6	15.5	18.9	19.8	14.8	*
1999.....	4.9	*	*	*	*	*	4.5	16.0	19.3	19.1	17.2	*

Non-Hispanic black, female

2007 5/.....	2.6	*	*	*	*	*	1.0	5.6	8.5	8.9	8.8	9.0
2006 5/.....	2.5	*	*	*	*	*	1.4	5.1	8.1	8.1	9.5	9.0
2005 5/.....	2.5	*	*	*	*	*	0.9	5.7	7.7	8.6	11.4	10.0
2004 5/.....	2.6	*	*	*	*	*	1.8	5.3	8.1	9.8	9.0	10.8
2003 5/.....	2.8	*	*	*	*	*	2.0	6.0	7.4	12.6	9.1	8.3
2002.....	3.0	*	*	*	*	*	2.3	7.1	7.7	10.1	13.3	9.6
2001.....	2.7	*	*	*	*	*	2.1	5.7	7.5	10.8	13.0	*
2000.....	2.9	*	*	*	*	*	2.3	5.9	7.7	13.5	11.3	11.3
1999.....	2.9	*	*	*	*	*	1.9	5.0	10.4	12.1	13.0	12.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins 3/, both sexes

2007.....	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.5	7.8	23.3
2006.....	1.0	*	*	*	*	0.0	0.1	0.3	0.9	2.5	7.8	21.8
2005.....	1.0	*	*	*	*	*	0.1	0.3	0.7	2.4	8.1	22.9
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.8	23.8
2003.....	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002.....	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001.....	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9

All origins 3/, male

2007.....	1.0	*	*	*	*	*	0.1	0.3	0.8	3.0	9.5	23.8
2006.....	0.9	*	*	*	*	*	0.1	0.3	0.9	3.1	9.6	21.7
2005.....	0.9	*	*	*	*	*	*	0.3	0.8	3.2	9.4	24.3
2004.....	0.9	*	*	*	*	*	0.1	0.4	1.0	2.9	9.3	25.5
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000.....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999.....	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7

All origins 3/, female

2007.....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.5	23.1
2006.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.0	6.6	21.9
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.6	1.8	7.2	22.3
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	23.0
2003.....	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001.....	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999.....	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Hispanic, both sexes

2007.....	0.5	*	*	*	*	*	*	*	1.0	3.2	8.5	24.0
2006.....	0.4	*	*	*	*	*	*	*	0.9	2.5	8.6	20.0
2005.....	0.5	*	*	*	*	*	*	*	1.0	3.4	7.7	29.9
2004.....	0.5	*	*	*	*	*	*	*	1.0	2.8	9.6	27.1
2003.....	0.4	*	*	*	*	*	*	*	*	3.6	9.5	18.7
2002.....	0.5	*	*	*	*	*	*	*	1.3	3.6	9.2	24.4
2001.....	0.5	*	*	*	*	*	*	*	1.2	4.5	8.5	31.3
2000.....	0.5	*	*	*	*	*	*	*	*	3.8	10.1	29.2
1999.....	0.5	*	*	*	*	*	*	*	1.2	3.3	11.9	32.3

Hispanic, male

2007.....	0.5	*	*	*	*	*	*	*	*	4.2	9.7	25.4
2006.....	0.3	*	*	*	*	*	*	*	*	*	9.5	*
2005.....	0.5	*	*	*	*	*	*	*	*	4.1	10.4	28.6
2004.....	0.5	*	*	*	*	*	*	*	*	*	12.2	*
2003.....	0.5	*	*	*	*	*	*	*	*	4.4	13.8	*
2002.....	0.5	*	*	*	*	*	*	*	*	4.0	9.4	*
2001.....	0.5	*	*	*	*	*	*	*	*	5.0	11.2	*
2000.....	0.5	*	*	*	*	*	*	*	*	5.3	15.7	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	15.1	*

Hispanic, female

2007.....	0.5	*	*	*	*	*	*	*	*	*	7.6	23.2
2006.....	0.5	*	*	*	*	*	*	*	*	*	8.0	22.8
2005.....	0.6	*	*	*	*	*	*	*	*	2.9	5.8	30.6
2004.....	0.6	*	*	*	*	*	*	*	*	2.8	7.9	31.0
2003.....	0.4	*	*	*	*	*	*	*	*	2.9	6.6	19.5
2002.....	0.5	*	*	*	*	*	*	*	*	3.2	9.0	23.2
2001.....	0.6	*	*	*	*	*	*	*	*	4.1	6.6	30.7
2000.....	0.5	*	*	*	*	*	*	*	*	*	*	31.7
1999.....	0.6	*	*	*	*	*	*	*	*	*	9.7	32.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 4/, both sexes

2007.....	1.2	*	*	*	*	0.1	0.1	0.3	0.7	2.4	7.7	23.2
2006.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.5	7.8	21.9
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.7	2.3	8.1	22.6
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.4	7.7	23.6
2003.....	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	8.1	21.9
2002.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.7	22.8
2001.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.8	7.5	23.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.4	24.6
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.6	23.6

Non-Hispanic 4/, male

2007.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.9	9.5	23.6
2006.....	1.0	*	*	*	*	*	*	0.3	0.9	3.1	9.6	22.0
2005.....	1.0	*	*	*	*	*	*	0.3	0.8	3.1	9.3	24.0
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.9	9.1	25.8
2003.....	1.0	*	*	*	*	*	*	0.3	0.8	3.1	9.9	23.0
2002.....	1.0	*	*	*	*	*	0.1	0.4	0.9	3.0	9.8	23.9
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.2	26.6
2000.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	27.7
1999.....	1.0	*	*	*	*	*	0.1	0.3	1.0	3.0	9.0	27.5

Non-Hispanic 4/, female

2007.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	6.5	23.0
2006.....	1.2	*	*	*	*	*	*	0.3	0.8	1.9	6.5	21.9
2005.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.7	7.3	21.9
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.0	6.7	22.6
2003.....	1.2	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	2.4	6.3	22.4
2001.....	1.2	*	*	*	*	*	*	0.3	0.8	2.5	6.3	22.1
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.2	23.4
1999.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.6	22.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic white, both sexes

2007 5/.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.4	7.7	23.9
2006 5/.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.2	7.7	22.4
2005 5/.....	1.2	*	*	*	*	*	*	0.2	0.6	2.1	8.1	22.7
2004 5/.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.3	7.5	24.0
2003 5/.....	1.2	*	*	*	*	*	0.1	0.2	0.6	2.6	8.1	22.0
2002.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.6	7.5	23.4
2001.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.6	7.3	23.6
2000.....	1.2	*	*	*	*	*	0.1	0.2	0.7	2.2	7.3	25.3
1999.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.4	7.6	24.0

Non-Hispanic white, male

2007 5/.....	1.2	*	*	*	*	*	*	0.3	0.8	3.0	9.6	24.2
2006 5/.....	1.1	*	*	*	*	*	*	0.3	0.8	2.8	9.6	23.2
2005 5/.....	1.1	*	*	*	*	*	*	0.2	0.7	2.9	9.5	24.5
2004 5/.....	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.1	27.0
2003 5/.....	1.1	*	*	*	*	*	*	0.2	0.7	3.1	9.8	23.3
2002.....	1.1	*	*	*	*	*	*	0.3	0.9	3.0	9.8	24.5
2001.....	1.1	*	*	*	*	*	*	0.4	1.0	2.9	9.2	26.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.8	2.6	9.3	28.9
1999.....	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.3	27.8

Non-Hispanic white, female

2007 5/.....	1.4	*	*	*	*	*	*	0.2	0.5	1.9	6.3	23.8
2006 5/.....	1.3	*	*	*	*	*	*	0.2	0.7	1.6	6.4	22.0
2005 5/.....	1.3	*	*	*	*	*	*	0.2	0.5	1.4	7.1	21.8
2004 5/.....	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.5	22.7
2003 5/.....	1.3	*	*	*	*	*	*	0.2	0.5	2.1	6.9	21.4
2002.....	1.3	*	*	*	*	*	*	0.2	0.6	2.3	6.1	22.9
2001.....	1.3	*	*	*	*	*	*	0.2	0.7	2.2	6.1	22.4
2000.....	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.0	23.9
1999.....	1.3	*	*	*	*	*	*	0.3	0.6	1.9	6.5	22.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic black, both sexes

2007 5/.....	0.8	*	*	*	*	*	*	0.5	1.3	2.9	8.6	18.2
2006 5/.....	0.9	*	*	*	*	*	*	0.6	1.4	4.1	9.3	17.5
2005 5/.....	0.9	*	*	*	*	*	*	0.6	1.2	4.2	9.4	21.0
2004 5/.....	0.8	*	*	*	*	*	*	0.8	1.6	3.6	9.3	19.6
2003 5/.....	0.8	*	*	*	*	*	*	0.7	1.2	3.0	8.1	21.1
2002.....	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.6	15.1
2001.....	0.9	*	*	*	*	*	*	0.7	1.7	4.8	8.3	20.0
2000.....	0.8	*	*	*	*	*	*	0.7	1.2	3.8	8.3	18.1
1999.....	0.7	*	*	*	*	*	0.4	0.5	1.5	3.7	7.3	15.5

Non-Hispanic black, male

2007 5/.....	0.6	*	*	*	*	*	*	*	*	3.0	10.2	20.1
2006 5/.....	0.6	*	*	*	*	*	*	*	1.6	3.8	9.7	*
2005 5/.....	0.8	*	*	*	*	*	*	*	*	5.3	10.4	20.0
2004 5/.....	0.7	*	*	*	*	*	*	1.0	1.6	3.7	9.1	*
2003 5/.....	0.7	*	*	*	*	*	*	*	*	3.3	10.7	21.3
2002.....	0.6	*	*	*	*	*	*	*	*	*	10.8	*
2001.....	0.8	*	*	*	*	*	*	*	2.2	5.0	6.3	27.7
2000.....	0.6	*	*	*	*	*	*	*	*	3.4	8.0	*
1999.....	0.6	*	*	*	*	*	*	*	*	3.8	7.2	*

Non-Hispanic black, female

2007 5/.....	0.9	*	*	*	*	*	*	*	1.4	2.8	7.7	17.4
2006 5/.....	1.0	*	*	*	*	*	*	*	1.3	4.2	9.0	20.2
2005 5/.....	1.0	*	*	*	*	*	*	*	*	3.4	8.9	21.4
2004 5/.....	1.0	*	*	*	*	*	*	*	1.6	3.6	9.3	21.7
2003 5/.....	0.8	*	*	*	*	*	*	0.8	*	2.8	6.7	21.0
2002.....	0.8	*	*	*	*	*	*	*	*	3.8	9.0	16.3
2001.....	1.0	*	*	*	*	*	*	*	*	4.6	9.4	17.1
2000.....	1.0	*	*	*	*	*	*	*	*	4.0	8.5	19.6
1999.....	0.8	*	*	*	*	*	*	*	*	3.6	7.4	15.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes

2007.....	15.4	3.4	0.1	0.1	0.2	0.6	1.7	5.1	13.6	40.1	113.0	290.6
2006.....	15.1	3.9	*	*	0.2	0.7	1.8	5.2	13.8	39.4	111.4	290.5
2005.....	14.8	3.9	*	0.1	0.2	0.7	1.7	4.8	13.6	39.3	110.3	288.3
2004.....	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.6	38.6	108.4	286.6
2003.....	14.6	4.5	*	0.1	0.2	0.7	1.8	4.9	13.6	40.1	109.5	293.1
2002.....	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001.....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000.....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999.....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9

All origins 3/, male

2007.....	15.2	3.6	*	*	0.2	0.7	2.0	5.8	15.7	45.5	139.9	376.3
2006.....	15.0	4.8	*	*	0.3	0.8	2.1	6.3	16.0	45.1	136.8	377.1
2005.....	14.6	4.2	*	*	0.2	0.7	1.9	5.9	16.1	43.8	135.9	376.5
2004.....	14.1	5.3	*	*	0.2	0.6	2.0	5.9	15.1	43.6	134.3	372.2
2003.....	14.3	5.2	*	*	0.3	0.7	2.1	6.0	15.4	45.3	136.8	387.6
2002.....	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001.....	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000.....	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999.....	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1

All origins 3/, female

2007.....	15.6	3.2	*	*	0.2	0.6	1.5	4.4	11.6	35.4	94.4	249.9
2006.....	15.3	3.0	*	*	0.2	0.6	1.5	4.1	11.8	34.6	94.1	250.0
2005.....	15.0	3.5	*	*	0.2	0.7	1.5	3.7	11.4	35.5	92.9	247.8
2004.....	14.8	3.2	*	*	0.2	0.7	1.6	4.1	12.2	34.3	90.9	248.1
2003.....	14.9	3.8	*	*	0.2	0.7	1.5	3.9	12.0	35.6	91.3	251.4
2002.....	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001.....	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000.....	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999.....	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Hispanic, both sexes

2007.....	5.9	3.5	*	*	*	0.4	1.1	4.8	13.8	42.7	98.2	190.2
2006.....	5.8	3.7	*	*	*	0.5	1.0	4.9	14.4	40.3	99.5	206.2
2005.....	5.5	3.6	*	*	*	0.4	1.1	3.9	14.8	38.3	95.8	193.4
2004.....	5.3	4.1	*	*	*	0.3	1.2	4.5	14.4	41.2	91.9	183.6
2003.....	5.4	4.0	*	*	*	0.4	1.3	4.6	15.2	39.6	90.2	233.2
2002.....	4.9	4.2	*	*	*	0.5	1.1	4.3	13.5	36.2	88.7	201.2
2001.....	5.0	*	*	*	*	0.4	1.4	3.9	12.8	41.4	94.5	222.0
2000.....	4.7	3.2	*	*	*	0.4	1.0	3.5	13.6	37.5	84.7	229.6
1999.....	4.7	2.6	*	*	*	0.3	1.3	3.6	14.0	35.6	88.7	232.2

Hispanic, male

2007.....	5.8	4.4	*	*	*	*	1.2	6.0	16.0	48.4	118.2	223.6
2006.....	5.6	*	*	*	*	*	1.2	6.2	15.8	48.2	113.8	241.3
2005.....	5.4	*	*	*	*	*	1.1	5.2	17.7	41.3	112.9	259.0
2004.....	5.2	5.4	*	*	*	*	1.4	5.3	17.0	44.4	109.4	231.2
2003.....	5.2	*	*	*	*	*	1.6	5.6	16.2	42.4	107.4	297.1
2002.....	4.7	4.9	*	*	*	*	1.3	4.8	15.1	40.9	112.0	235.9
2001.....	4.7	*	*	*	*	*	1.6	3.8	15.2	43.3	112.2	281.9
2000.....	4.4	*	*	*	*	*	1.4	4.4	14.3	40.9	101.3	292.2
1999.....	4.7	*	*	*	*	*	1.2	4.4	17.8	39.1	110.5	336.4

Hispanic, female

2007.....	6.0	*	*	*	*	*	0.9	3.7	11.7	38.1	84.1	172.1
2006.....	6.1	*	*	*	*	*	0.8	3.6	13.2	33.9	89.4	187.5
2005.....	5.5	*	*	*	*	*	1.0	2.5	12.1	35.9	84.0	159.0
2004.....	5.5	*	*	*	*	*	0.9	3.7	11.9	38.6	79.8	159.2
2003.....	5.7	*	*	*	*	*	1.0	3.5	14.4	37.4	78.5	200.8
2002.....	5.1	*	*	*	*	0.6	1.0	3.8	12.1	32.4	72.8	184.0
2001.....	5.4	*	*	*	*	*	1.1	4.0	10.7	39.8	82.5	192.7
2000.....	4.9	*	*	*	*	*	*	2.5	12.9	34.8	73.6	198.8
1999.....	4.7	*	*	*	*	*	1.3	2.9	10.7	32.9	74.2	182.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 4/, both sexes

2007.....	17.1	3.3	0.2	0.1	0.2	0.7	1.9	5.1	13.5	39.8	113.9	295.6
2006.....	16.7	4.0	*	*	0.2	0.7	1.9	5.2	13.7	39.2	112.0	294.4
2005.....	16.4	3.9	*	*	0.2	0.8	1.8	4.9	13.5	39.3	111.1	292.5
2004.....	15.9	4.3	*	*	0.2	0.7	1.9	5.0	13.5	38.3	109.0	290.9
2003.....	16.0	4.6	*	*	0.2	0.8	1.9	4.9	13.5	40.0	110.2	294.9
2002.....	15.6	4.3	*	*	0.2	0.7	1.7	4.7	12.9	39.3	109.8	291.2
2001.....	15.1	3.5	*	*	0.2	0.6	1.7	4.7	12.9	40.0	104.3	289.6
2000.....	14.4	4.4	*	0.1	0.2	0.7	1.7	4.4	12.6	37.9	101.2	279.0
1999.....	13.8	4.8	*	0.1	0.2	0.7	1.6	4.1	11.8	37.1	97.7	269.2

Non-Hispanic 4/, male

2007.....	17.0	3.3	*	*	0.2	0.8	2.1	5.8	15.6	45.2	141.3	384.6
2006.....	16.7	5.1	*	*	0.3	0.9	2.3	6.3	16.0	44.8	138.0	384.0
2005.....	16.2	4.2	*	*	0.2	0.8	2.1	6.0	15.9	43.9	137.2	382.0
2004.....	15.6	5.2	*	*	0.2	0.6	2.2	5.9	15.0	43.4	135.3	378.6
2003.....	15.8	5.4	*	*	0.3	0.8	2.1	6.0	15.2	45.4	138.0	390.8
2002.....	15.4	5.4	*	*	0.2	0.7	1.9	5.6	14.2	45.9	136.9	391.0
2001.....	14.8	3.7	*	*	0.2	0.6	1.9	5.6	14.0	46.1	130.6	397.6
2000.....	14.1	5.4	*	*	0.2	0.7	2.0	5.4	14.1	42.9	128.2	384.0
1999.....	13.5	4.8	*	*	0.2	0.7	1.9	4.7	13.0	42.3	125.7	370.9

Non-Hispanic 4/, female

2007.....	17.2	3.4	*	*	0.2	0.6	1.6	4.4	11.5	35.2	95.0	253.6
2006.....	16.8	2.9	*	*	0.2	0.6	1.6	4.2	11.6	34.6	94.2	252.7
2005.....	16.5	3.5	*	*	0.2	0.7	1.5	3.8	11.3	35.4	93.4	251.6
2004.....	16.2	3.4	*	*	0.2	0.7	1.7	4.1	12.2	33.9	91.3	251.6
2003.....	16.2	3.6	*	*	0.2	0.7	1.6	3.9	11.8	35.5	91.7	252.7
2002.....	15.8	3.2	*	*	0.2	0.7	1.6	3.9	11.7	33.7	91.8	248.2
2001.....	15.4	3.3	*	*	0.2	0.6	1.6	3.8	12.0	34.8	87.1	244.6
2000.....	14.7	3.4	*	*	0.2	0.6	1.5	3.6	11.3	33.8	83.6	236.6
1999.....	14.1	4.8	*	*	0.2	0.7	1.3	3.5	10.7	32.9	79.5	228.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic white, both sexes

2007 5/.....	17.0	2.7	*	*	0.2	0.5	1.2	3.5	10.0	33.1	107.0	287.1
2006 5/.....	16.5	3.4	*	*	0.2	0.5	1.2	3.5	10.3	32.1	104.2	284.8
2005 5/.....	16.2	2.7	*	*	0.2	0.5	1.1	3.4	9.8	32.7	103.5	284.0
2004 5/.....	15.8	2.6	*	*	0.2	0.4	1.3	3.3	9.8	32.2	101.8	282.1
2003 5/.....	15.8	3.5	*	*	0.1	0.5	1.1	3.2	10.0	33.2	102.7	285.9
2002.....	15.4	3.3	*	*	0.1	0.5	1.1	3.1	9.5	33.0	102.5	281.1
2001.....	14.9	2.4	*	*	0.2	0.4	1.1	3.1	9.3	33.5	96.6	279.7
2000.....	14.2	2.7	*	*	0.2	0.4	1.1	2.9	9.3	31.4	94.1	269.4
1999.....	13.4	3.5	*	*	0.1	0.4	1.0	2.6	8.7	31.1	89.9	255.3

Non-Hispanic white, male

2007 5/.....	17.1	2.4	*	*	0.1	0.5	1.4	4.1	11.8	38.2	135.8	382.0
2006 5/.....	16.7	4.6	*	*	0.3	0.6	1.5	4.3	12.3	37.7	130.5	380.7
2005 5/.....	16.2	2.7	*	*	0.2	0.5	1.4	4.2	11.7	38.2	130.4	379.1
2004 5/.....	15.6	3.1	*	*	0.2	0.4	1.5	3.8	11.1	37.2	129.4	373.0
2003 5/.....	15.7	4.4	*	*	0.2	0.5	1.3	3.8	11.4	38.5	131.0	386.3
2002.....	15.3	4.1	*	*	*	0.5	1.2	3.6	10.3	39.7	130.8	382.6
2001.....	14.8	2.5	*	*	0.2	0.4	1.3	3.7	10.5	39.9	123.7	392.6
2000.....	14.1	3.5	*	*	0.2	0.5	1.3	3.5	10.6	37.3	121.7	378.9
1999.....	13.3	3.9	*	*	*	0.5	1.2	2.9	9.9	36.6	117.3	359.6

Non-Hispanic white, female

2007 5/.....	16.9	3.0	*	*	0.2	0.4	1.0	3.0	8.4	28.6	86.8	242.2
2006 5/.....	16.4	2.1	*	*	*	0.4	1.0	2.7	8.4	27.3	86.0	240.1
2005 5/.....	16.2	2.7	*	*	0.2	0.5	0.9	2.6	8.0	27.9	85.0	240.5
2004 5/.....	16.0	2.1	*	*	0.2	0.5	1.1	2.7	8.6	27.8	83.0	241.4
2003 5/.....	15.9	2.5	*	*	*	0.5	0.9	2.6	8.7	28.7	83.5	241.7
2002.....	15.6	2.5	*	*	*	0.4	1.0	2.6	8.7	27.3	83.6	237.3
2001.....	15.0	2.3	*	*	0.2	0.4	0.9	2.4	8.3	28.0	78.6	232.7
2000.....	14.2	1.8	*	*	0.2	0.4	1.0	2.3	8.0	26.5	76.0	225.2
1999.....	13.5	3.1	*	*	*	0.4	0.8	2.2	7.5	26.4	71.9	213.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic black, both sexes

2007 5/.....	21.8	6.7	*	*	0.4	2.0	6.1	15.9	42.2	102.1	204.0	443.9
2006 5/.....	22.1	6.8	*	*	0.5	2.0	6.1	16.7	42.7	104.5	213.8	449.4
2005 5/.....	21.5	9.1	*	*	0.5	2.3	5.7	15.2	45.1	101.6	211.9	429.8
2004 5/.....	21.0	10.4	*	*	0.5	2.0	5.8	17.1	45.6	96.6	205.1	437.2
2003 5/.....	21.4	9.5	*	*	0.6	2.3	6.4	16.8	43.7	105.6	209.9	446.5
2002.....	20.5	8.5	*	*	0.6	2.1	5.5	16.5	41.8	101.5	205.5	444.0
2001.....	20.2	8.5	*	*	0.5	1.8	5.5	16.2	42.5	104.2	204.0	432.9
2000.....	19.5	12.0	*	*	0.4	1.9	5.5	16.2	41.2	101.2	197.3	410.2
1999.....	19.2	10.2	*	*	0.5	2.0	5.6	15.6	37.8	96.6	196.4	436.8

Non-Hispanic black, male

2007 5/.....	20.5	7.7	*	*	*	2.3	6.9	18.2	49.7	116.2	235.7	500.9
2006 5/.....	21.0	7.4	*	*	*	2.2	7.1	20.6	51.0	117.5	256.1	496.6
2005 5/.....	20.3	11.2	*	*	*	2.4	6.4	19.4	56.3	104.8	249.8	487.2
2004 5/.....	19.9	12.4	*	*	*	2.0	6.4	21.5	51.1	109.9	235.9	520.3
2003 5/.....	20.8	9.9	*	*	0.7	2.6	7.4	21.8	52.5	123.3	248.7	526.3
2002.....	19.7	10.6	*	*	*	2.0	6.2	20.6	50.1	116.3	237.1	568.9
2001.....	18.6	8.9	*	*	*	2.2	5.3	19.9	46.3	116.0	235.3	517.4
2000.....	18.3	13.6	*	*	*	2.0	6.7	20.8	45.6	105.7	234.2	496.7
1999.....	18.1	8.7	*	*	*	1.9	6.7	19.4	41.2	103.9	251.1	532.4

Non-Hispanic black, female

2007 5/.....	23.0	*	*	*	*	1.8	5.3	14.0	36.1	92.0	186.0	420.3
2006 5/.....	23.0	*	*	*	*	1.8	5.3	13.3	36.1	95.3	189.8	430.1
2005 5/.....	22.5	6.8	*	*	*	2.2	5.1	11.6	36.1	99.3	190.5	406.7
2004 5/.....	22.0	8.4	*	*	*	1.9	5.2	13.2	41.2	87.2	187.7	404.4
2003 5/.....	21.9	9.0	*	*	*	2.0	5.6	12.5	36.6	93.0	187.9	415.6
2002.....	21.2	6.3	*	*	0.7	2.2	4.9	13.1	35.1	91.0	187.7	396.3
2001.....	21.7	8.1	*	*	*	1.4	5.6	13.0	39.4	95.8	186.5	401.7
2000.....	20.7	10.2	*	*	*	1.8	4.4	12.2	37.6	98.1	176.6	378.1
1999.....	20.2	11.8	*	*	*	2.0	4.7	12.3	35.1	91.5	165.8	401.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All origins 3/, both sexes

2007.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	0.9
2006.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.6
2005.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
2004.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8

All origins 3/, male

2007.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.2
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*

All origins 3/, female

2007.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
2006.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Non-Hispanic 4/, both sexes

2007.....	0.1	*	*	*	*	*	*	*	*	0.1	0.2	0.5	0.9
2006.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	0.7
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	0.3	0.7
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.2	0.4	0.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.2	0.5	0.8

Non-Hispanic 4/, male

2007.....	0.1	*	*	*	*	*	*	*	*	*	*	0.6	*
2006.....	0.1	*	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	0.4	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	0.5	1.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.7	*

Non-Hispanic 4/, female

2007.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	0.8
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	0.3	*
2005.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	0.7
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	0.3	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	0.5	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	0.4	0.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	0.4	0.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	0.4	0.7





Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins 3/, both sexes

2007.....	1.0	*	*	*	*	*	0.1	0.3	0.6	2.0	7.1	22.3
2006.....	0.6	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.4	13.4
2005.....	0.3	*	*	*	*	0.1	0.1	0.1	0.3	0.7	1.9	5.4
2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.3	2.8
2003.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.6	1.3	2.8
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2

All origins 3/, male

2007.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	8.7	29.9
2006.....	0.6	*	*	*	*	*	*	0.2	0.6	1.5	5.7	17.2
2005.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	2.4	7.5
2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.6	4.0
2003.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.6	4.1
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0

All origins 3/, female

2007.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.6	6.0	18.7
2006.....	0.6	*	*	*	*	*	*	0.2	0.4	1.2	3.5	11.6
2005.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.6	4.4
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic 4/, both sexes

2007.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.0	7.1	22.8
2006.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	4.4	13.6
2005.....	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	1.9	5.5
2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.3	2.8
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	2.8
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.7	3.2

Non-Hispanic 4/, male

2007.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.4	8.6	31.0
2006.....	0.7	*	*	*	*	*	*	0.2	0.6	1.5	5.7	17.4
2005.....	0.3	*	*	*	*	*	0.1	0.1	0.4	0.8	2.5	7.5
2004.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.7	3.8
2003.....	0.2	*	*	*	*	*	*	0.2	*	0.6	1.6	4.3
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.2
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.3	5.1

Non-Hispanic 4/, female

2007.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	6.0	18.9
2006.....	0.7	*	*	*	*	*	*	0.2	0.4	1.2	3.5	11.9
2005.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.5	4.5
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.1	2.2
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.2
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic white, both sexes

2007 5/.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	6.5	21.4
2006 5/.....	0.7	*	*	*	*	*	*	0.1	0.3	1.1	4.0	12.7
2005 5/.....	0.3	*	*	*	*	*	*	0.1	0.3	0.6	1.8	5.2
2004 5/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	2.7
2003 5/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.6	1.1	2.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9

Non-Hispanic white, male

2007 5/.....	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.1	29.7
2006 5/.....	0.7	*	*	*	*	*	*	0.1	0.4	1.3	5.2	16.2
2005 5/.....	0.3	*	*	*	*	*	*	*	0.3	0.7	2.4	7.3
2004 5/.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	3.6
2003 5/.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.4	4.0
2002.....	0.2	*	*	*	*	*	*	*	*	0.7	1.7	3.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.8	1.7	4.0
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.7
1999.....	0.2	*	*	*	*	*	*	*	*	0.5	2.2	4.6

Non-Hispanic white, female

2007 5/.....	1.1	*	*	*	*	*	*	0.2	0.4	1.3	5.3	17.5
2006 5/.....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.1	11.1
2005 5/.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	4.3
2004 5/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.3
2003 5/.....	0.2	*	*	*	*	*	*	*	*	0.5	1.0	2.1
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.0
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.1	1.9
1999.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Renal failure (N17-N19)

##### All origins 3/, both sexes

2007.....	14.3	3.2	*	*	0.2	0.6	1.6	4.8	12.9	37.9	105.3	267.3
2006.....	14.5	3.7	*	*	0.2	0.6	1.7	5.0	13.3	37.9	106.7	276.4
2005.....	14.5	3.7	*	*	0.2	0.7	1.6	4.6	13.3	38.4	108.0	282.1
2004.....	14.2	4.0	*	*	0.2	0.6	1.7	4.8	13.4	37.9	106.6	282.9
2003.....	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002.....	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001.....	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000.....	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999.....	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8

##### All origins 3/, male

2007.....	14.2	3.4	*	*	0.2	0.6	1.9	5.4	14.8	42.8	130.5	345.0
2006.....	14.3	4.4	*	*	0.2	0.7	2.0	6.0	15.4	43.4	130.7	358.8
2005.....	14.2	4.1	*	*	0.2	0.7	1.8	5.7	15.6	42.8	133.1	368.0
2004.....	13.8	5.1	*	*	0.2	0.5	2.0	5.7	14.8	42.9	132.1	367.4
2003.....	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002.....	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001.....	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000.....	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999.....	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6

##### All origins 3/, female

2007.....	14.5	3.1	*	*	0.2	0.6	1.4	4.1	11.0	33.7	88.0	230.4
2006.....	14.6	3.0	*	*	0.2	0.6	1.4	3.9	11.4	33.2	90.2	237.8
2005.....	14.7	3.2	*	*	0.2	0.6	1.4	3.6	11.0	34.8	91.0	242.6
2004.....	14.6	3.0	*	*	0.2	0.6	1.5	3.9	12.0	33.7	89.5	244.9
2003.....	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002.....	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001.....	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000.....	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999.....	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Renal failure (N17-N19)</b>												
<b>Hispanic, both sexes</b>												
2007.....	5.5	3.4	*	*	*	0.3	1.0	4.5	13.2	40.5	89.7	174.8
2006.....	5.6	3.7	*	*	*	0.4	0.9	4.7	13.8	38.7	95.0	197.2
2005.....	5.3	3.6	*	*	*	0.3	1.0	3.7	14.5	37.5	93.5	187.4
2004.....	5.2	3.9	*	*	*	0.3	1.2	4.4	14.0	40.5	91.0	179.3
2003.....	5.3	3.8	*	*	*	0.4	1.3	4.4	14.8	39.0	88.5	230.6
2002.....	4.8	4.2	*	*	*	0.4	1.1	4.2	13.2	35.7	86.9	198.4
2001.....	4.9	*	*	*	*	0.3	1.2	3.7	12.5	40.5	93.2	217.6
2000.....	4.5	2.9	*	*	*	0.4	1.0	3.3	13.3	36.9	83.0	225.6
1999.....	4.5	*	*	*	*	*	1.2	3.5	13.6	34.4	86.9	228.0
<b>Hispanic, male</b>												
2007.....	5.3	4.2	*	*	*	*	1.2	5.5	15.0	45.1	107.0	208.4
2006.....	5.4	*	*	*	*	*	1.1	6.0	15.1	46.6	108.8	227.8
2005.....	5.2	*	*	*	*	*	1.0	5.0	17.2	40.3	111.6	249.1
2004.....	5.0	5.2	*	*	*	*	1.4	5.2	16.4	43.5	108.4	224.2
2003.....	5.1	*	*	*	*	*	1.5	5.3	15.8	41.9	105.1	295.5
2002.....	4.6	4.9	*	*	*	*	1.3	4.8	14.4	40.7	109.2	232.6
2001.....	4.6	*	*	*	*	*	1.4	3.7	14.6	42.5	110.4	278.1
2000.....	4.3	*	*	*	*	*	1.3	4.2	13.8	40.5	99.3	280.1
1999.....	4.5	*	*	*	*	*	1.1	4.3	17.2	38.2	109.0	330.0
<b>Hispanic, female</b>												
2007.....	5.6	*	*	*	*	*	0.8	3.4	11.5	36.7	77.5	156.7
2006.....	5.8	*	*	*	*	*	0.7	3.5	12.6	32.4	85.4	180.9
2005.....	5.3	*	*	*	*	*	0.9	2.5	11.9	35.2	80.9	155.1
2004.....	5.4	*	*	*	*	*	0.9	3.6	11.9	38.0	79.0	156.3
2003.....	5.6	*	*	*	*	*	0.9	3.5	13.9	36.6	77.2	197.7
2002.....	5.0	*	*	*	*	0.6	0.8	3.6	12.0	31.8	71.7	181.5
2001.....	5.2	*	*	*	*	*	1.0	3.7	10.5	38.9	81.6	188.0
2000.....	4.8	*	*	*	*	*	*	2.4	12.8	34.0	72.0	198.8
1999.....	4.5	*	*	*	*	*	1.3	2.7	10.4	31.3	72.2	178.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Renal failure (N17-N19)

##### Non-Hispanic 4/, both sexes

2007.....	15.9	3.2	*	*	0.2	0.7	1.7	4.8	12.8	37.6	106.3	271.9
2006.....	16.0	3.7	*	*	0.2	0.7	1.8	5.0	13.2	37.8	107.2	280.0
2005.....	16.0	3.7	*	*	0.2	0.7	1.7	4.7	13.1	38.5	108.8	286.2
2004.....	15.7	4.1	*	*	0.2	0.6	1.8	4.8	13.3	37.6	107.2	287.2
2003.....	15.7	4.3	*	*	0.2	0.7	1.8	4.8	13.2	39.2	108.5	291.4
2002.....	15.3	4.1	*	*	0.2	0.7	1.7	4.6	12.6	38.6	107.9	287.4
2001.....	14.9	3.3	*	*	0.2	0.6	1.7	4.6	12.7	39.2	102.7	285.9
2000.....	14.1	4.2	*	*	0.2	0.6	1.6	4.3	12.3	37.1	99.3	274.9
1999.....	13.5	4.7	*	*	0.2	0.6	1.5	3.9	11.5	36.4	95.4	265.0

##### Non-Hispanic 4/, male

2007.....	15.8	3.2	*	*	0.2	0.7	2.0	5.4	14.7	42.5	132.0	352.4
2006.....	15.9	4.6	*	*	0.3	0.8	2.2	6.0	15.4	43.1	131.8	365.5
2005.....	15.8	4.1	*	*	0.2	0.8	1.9	5.8	15.5	42.9	134.3	373.6
2004.....	15.3	5.1	*	*	0.2	0.6	2.1	5.8	14.6	42.7	133.1	374.0
2003.....	15.5	5.0	*	*	0.2	0.8	2.1	5.8	15.0	44.6	136.0	386.2
2002.....	15.1	5.1	*	*	0.2	0.6	1.8	5.4	13.9	45.1	134.7	386.2
2001.....	14.6	3.5	*	*	0.2	0.6	1.8	5.4	13.7	45.3	128.5	392.7
2000.....	13.8	5.2	*	*	0.2	0.6	1.9	5.1	13.7	42.0	125.9	377.7
1999.....	13.2	4.6	*	*	0.1	0.6	1.8	4.5	12.7	41.5	122.6	364.4

##### Non-Hispanic 4/, female

2007.....	16.0	3.2	*	*	0.2	0.6	1.5	4.2	11.0	33.4	88.6	233.9
2006.....	16.1	2.8	*	*	0.2	0.6	1.5	4.0	11.2	33.2	90.4	240.2
2005.....	16.2	3.2	*	*	0.2	0.7	1.5	3.7	11.0	34.7	91.5	246.4
2004.....	16.0	3.1	*	*	0.2	0.6	1.6	4.0	12.0	33.3	89.8	248.5
2003.....	15.9	3.5	*	*	0.2	0.7	1.5	3.8	11.6	34.7	90.1	249.7
2002.....	15.5	2.9	*	*	0.2	0.7	1.5	3.8	11.4	33.2	90.2	244.8
2001.....	15.2	3.1	*	*	0.2	0.5	1.6	3.7	11.8	34.1	85.8	241.5
2000.....	14.4	3.1	*	*	0.2	0.5	1.4	3.5	11.0	33.2	82.0	233.4
1999.....	13.8	4.8	*	*	0.2	0.6	1.3	3.3	10.4	32.1	77.8	225.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Renal failure (N17-N19)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	15.8	2.5	*	*	0.1	0.4	1.1	3.3	9.4	31.3	100.1	264.8
2006 5/.....	15.8	3.1	*	*	0.2	0.5	1.2	3.4	9.9	30.9	99.8	271.3
2005 5/.....	15.8	2.5	*	*	0.2	0.5	1.1	3.3	9.5	32.0	101.3	278.1
2004 5/.....	15.5	2.5	*	*	0.1	0.4	1.3	3.2	9.6	31.5	100.0	278.6
2003 5/.....	15.5	3.2	*	*	0.1	0.5	1.1	3.1	9.8	32.5	101.1	282.6
2002.....	15.2	3.1	*	*	0.1	0.4	1.1	3.0	9.2	32.3	100.8	277.6
2001.....	14.6	2.2	*	*	0.2	0.4	1.1	3.0	9.2	32.7	95.1	276.1
2000.....	13.9	2.5	*	*	0.2	0.4	1.1	2.8	9.0	30.7	92.3	265.7
1999.....	13.1	3.5	*	*	0.1	0.4	0.9	2.4	8.4	30.4	87.8	251.6
<b>Non-Hispanic white, male</b>												
2007 5/.....	15.9	2.1	*	*	*	0.5	1.3	3.8	11.1	36.0	127.0	351.1
2006 5/.....	15.9	4.1	*	*	0.2	0.6	1.4	4.2	11.9	36.4	124.8	363.4
2005 5/.....	15.8	2.6	*	*	0.2	0.5	1.3	4.1	11.3	37.3	127.6	370.9
2004 5/.....	15.3	3.0	*	*	0.2	0.3	1.4	3.7	10.9	36.5	127.2	368.8
2003 5/.....	15.4	4.0	*	*	0.2	0.5	1.3	3.7	11.2	37.8	129.3	381.9
2002.....	15.0	3.9	*	*	*	0.4	1.2	3.5	10.0	38.8	128.8	377.8
2001.....	14.5	2.3	*	*	*	0.3	1.3	3.6	10.3	39.1	121.7	387.6
2000.....	13.7	3.3	*	*	0.2	0.4	1.2	3.3	10.3	36.5	119.4	373.4
1999.....	13.0	3.8	*	*	*	0.4	1.1	2.8	9.6	35.9	114.4	353.8
<b>Non-Hispanic white, female</b>												
2007 5/.....	15.8	2.9	*	*	*	0.4	0.9	2.8	7.9	27.2	81.2	223.9
2006 5/.....	15.7	2.1	*	*	*	0.4	0.9	2.6	8.1	26.2	82.5	228.4
2005 5/.....	15.9	2.3	*	*	*	0.5	0.9	2.5	7.7	27.4	83.2	235.7
2004 5/.....	15.7	2.0	*	*	*	0.4	1.1	2.6	8.4	27.2	81.5	238.2
2003 5/.....	15.6	2.3	*	*	*	0.5	0.9	2.5	8.5	28.0	82.0	238.9
2002.....	15.3	2.2	*	*	*	0.4	1.0	2.5	8.5	26.7	82.0	234.3
2001.....	14.7	2.1	*	*	0.2	0.4	0.9	2.4	8.1	27.3	77.5	229.7
2000.....	14.0	*	*	*	*	0.3	0.9	2.3	7.7	25.8	74.4	222.2
1999.....	13.2	3.1	*	*	*	0.4	0.8	2.1	7.3	25.8	70.3	210.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Renal failure (N17-N19)

##### Non-Hispanic black, both sexes

2007 5/.....	20.3	6.5	*	*	0.4	1.9	5.7	15.0	40.3	96.4	189.0	402.7
2006 5/.....	21.1	6.6	*	*	0.4	1.8	5.9	15.9	41.1	100.4	205.1	425.2
2005 5/.....	21.0	8.7	*	*	0.4	2.1	5.5	14.8	44.2	99.8	208.0	418.9
2004 5/.....	20.7	10.1	*	*	0.5	1.8	5.5	16.7	45.0	95.4	202.5	432.6
2003 5/.....	21.1	9.3	*	*	0.5	2.2	6.3	16.5	43.3	104.3	207.1	442.3
2002.....	20.2	8.0	*	*	0.6	2.0	5.4	16.0	41.4	100.9	202.7	436.5
2001.....	20.0	8.5	*	*	0.4	1.7	5.3	15.8	42.1	102.9	201.9	428.9
2000.....	19.2	11.4	*	*	0.4	1.8	5.3	15.6	40.7	99.9	194.6	404.2
1999.....	18.8	10.2	*	*	0.5	1.9	5.4	15.2	37.0	95.3	193.0	428.7

##### Non-Hispanic black, male

2007 5/.....	19.1	7.7	*	*	*	2.0	6.6	17.1	47.2	109.2	218.9	452.0
2006 5/.....	20.0	7.1	*	*	*	1.9	6.9	19.7	48.7	112.9	245.0	462.7
2005 5/.....	19.8	11.2	*	*	*	2.1	6.0	18.9	55.1	103.4	243.9	473.9
2004 5/.....	19.6	12.1	*	*	*	2.0	6.2	21.2	50.2	108.6	233.0	514.2
2003 5/.....	20.5	9.6	*	*	*	2.5	7.3	21.3	52.1	121.6	244.4	521.0
2002.....	19.4	10.0	*	*	*	1.8	6.1	20.1	49.8	115.4	233.3	562.3
2001.....	18.4	8.9	*	*	*	2.1	5.1	19.3	45.7	114.3	233.5	514.0
2000.....	17.9	13.6	*	*	*	1.9	6.5	19.9	45.0	104.2	231.7	487.3
1999.....	17.7	8.7	*	*	*	1.8	6.4	19.0	40.5	102.5	246.7	519.4

##### Non-Hispanic black, female

2007 5/.....	21.4	*	*	*	*	1.7	4.9	13.2	34.7	87.3	171.9	382.3
2006 5/.....	22.1	*	*	*	*	1.7	5.0	12.7	35.1	91.4	182.5	409.9
2005 5/.....	22.1	*	*	*	*	2.1	5.0	11.3	35.4	97.2	187.6	396.8
2004 5/.....	21.7	8.0	*	*	*	1.6	4.9	12.9	40.9	86.0	185.3	400.4
2003 5/.....	21.7	9.0	*	*	*	1.9	5.5	12.4	36.3	92.0	186.1	411.9
2002.....	20.9	*	*	*	*	2.1	4.8	12.5	34.7	90.6	185.4	388.4
2001.....	21.4	8.1	*	*	*	1.4	5.5	12.8	39.2	94.9	184.1	397.4
2000.....	20.3	9.2	*	*	*	1.7	4.1	11.8	37.3	96.9	173.7	373.3
1999.....	19.8	11.8	*	*	*	2.0	4.5	12.0	34.3	90.2	163.0	394.9











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All origins 3/, both sexes

2007.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.5
2006.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.7
2005.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.2
2004.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.6	1.9	5.9
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000.....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5

All origins 3/, male

2007.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	3.3
2006.....	0.1	*	*	*	*	*	*	*	0.1	0.5	0.8	3.7
2005.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	3.6
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8

All origins 3/, female

2007.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.5	3.6
2006.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.2
2005.....	0.4	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.9
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.5
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.1
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000.....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.5
2006.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.8
2005.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.2
2004.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.3
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.5	1.9	6.0
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	6.4
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.1
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.6
1999.....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.7	1.9	7.5

Non-Hispanic 4/, male

2007.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	3.5
2006.....	0.2	*	*	*	*	*	*	*	*	0.4	0.8	3.9
2005.....	0.2	*	*	*	*	*	*	*	*	0.4	1.5	3.5
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.6
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.8
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.9

Non-Hispanic 4/, female

2007.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.5	3.5
2006.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.2
2005.....	0.4	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.9
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.6
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.2
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.8
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.4	2.3	6.9
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999.....	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic white, both sexes

2007 5/.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	3.7
2006 5/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.3	5.0
2005 5/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.8	5.2
2004 5/.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.3
2003 5/.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	6.0
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.5
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.4	2.0	6.2
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999.....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	1.9	7.7

Non-Hispanic white, male

2007 5/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	3.7
2006 5/.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	4.0
2005 5/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	3.4
2004 5/.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.8
2003 5/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.3	5.8
2002.....	0.2	*	*	*	*	*	*	*	*	0.5	1.3	5.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.5	4.2
2000.....	0.2	*	*	*	*	*	*	*	*	*	1.5	6.0
1999.....	0.2	*	*	*	*	*	*	*	*	0.4	1.7	7.3

Non-Hispanic white, female

2007 5/.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	3.7
2006 5/.....	0.4	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.5
2005 5/.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.0	6.0
2004 5/.....	0.4	*	*	*	*	*	*	0.1	0.2	0.7	2.0	5.6
2003 5/.....	0.4	*	*	*	*	*	*	0.2	0.2	0.7	2.2	6.1
2002.....	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	6.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	7.1
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.7	2.1	8.2
1999.....	0.5	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.9























Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/, both sexes

2007.....	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2006.....	0.3	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005.....	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.4	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
1999.....	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*

All origins 3/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins 3/, female

2007.....	0.5	...	...	*	0.8	1.6	1.0	0.3	*	*	*	*
2006.....	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2004.....	0.5	...	...	*	0.8	1.5	0.7	0.4	*	*	*	*
2003.....	0.4	...	...	*	0.6	1.2	0.7	0.1	*	*	*	*
2002.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2001.....	0.3	...	...	*	0.6	1.0	0.5	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
1999.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

Hispanic, both sexes

2007.....	0.3	...	...	*	0.4	0.7	0.6	*	*	*	*	*
2006.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2005.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2004.....	0.3	...	...	*	0.4	0.6	0.4	*	*	*	*	*
2003.....	0.3	...	...	*	0.3	0.6	0.5	*	*	*	*	*
2002.....	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001.....	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000.....	0.2	...	...	*	0.4	0.6	0.4	*	*	*	*	*
1999.....	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*

Hispanic, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, female

2007.....	0.6	...	...	*	0.8	1.6	1.2	*	*	*	*	*
2006.....	0.7	...	...	*	1.1	1.9	1.1	*	*	*	*	*
2005.....	0.6	...	...	*	1.0	1.7	1.1	*	*	*	*	*
2004.....	0.6	...	...	*	0.9	1.4	0.9	*	*	*	*	*
2003.....	0.5	...	...	*	0.7	1.4	1.0	*	*	*	*	*
2002.....	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001.....	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000.....	0.5	...	...	*	0.8	1.2	0.9	*	*	*	*	*
1999.....	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 4/, both sexes

2007.....	0.2	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2006.....	0.2	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005.....	0.2	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*

Non-Hispanic 4/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 4/, female

2007.....	0.5	...	...	*	0.8	1.6	1.0	0.3	*	*	*	*
2006.....	0.5	...	...	*	0.8	1.6	0.8	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2004.....	0.4	...	...	*	0.8	1.5	0.7	0.4	*	*	*	*
2003.....	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.4	0.9	0.4	*	*	*	*	*
2001.....	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000.....	0.3	...	...	*	0.4	0.9	0.5	*	*	*	*	*
1999.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic white, both sexes

2007 5/.....	0.2	...	...	*	0.2	0.6	0.4	0.1	*	*	*	*
2006 5/.....	0.1	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2005 5/.....	0.2	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2004 5/.....	0.2	...	...	*	0.2	0.5	0.2	0.1	*	*	*	*
2003 5/.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2002.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.3	0.1	*	*	*	*	*
2000.....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*

Non-Hispanic white, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic white, female

2007 5/.....	0.3	...	...	*	0.5	1.1	0.7	0.2	*	*	*	*
2006 5/.....	0.3	...	...	*	0.6	0.9	0.5	0.2	*	*	*	*
2005 5/.....	0.3	...	...	*	0.6	1.0	0.6	0.2	*	*	*	*
2004 5/.....	0.3	...	...	*	0.5	1.0	0.5	0.3	*	*	*	*
2003 5/.....	0.2	...	...	*	0.3	0.7	0.4	*	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.6	0.3	*	*	*	*	*
2001.....	0.2	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000.....	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
1999.....	0.2	...	...	*	0.2	0.6	0.3	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic black, both sexes

2007 5/.....	0.7	...	...	*	1.1	2.0	1.1	*	*	*	*	*
2006 5/.....	0.7	...	...	*	0.9	2.2	1.1	0.4	*	*	*	*
2005 5/.....	0.7	...	...	*	1.1	2.2	1.2	*	*	*	*	*
2004 5/.....	0.7	...	...	*	1.2	2.0	0.8	0.5	*	*	*	*
2003 5/.....	0.5	...	...	*	0.7	1.6	1.1	*	*	*	*	*
2002.....	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*
2001.....	0.4	...	...	*	0.8	1.3	0.7	*	*	*	*	*
2000.....	0.4	...	...	*	0.6	1.3	0.7	*	*	*	*	*
1999.....	0.5	...	...	*	0.8	1.3	0.7	*	*	*	*	*

Non-Hispanic black, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic black, female

2007 5/.....	1.2	...	...	*	2.1	3.8	2.1	*	*	*	*	*
2006 5/.....	1.3	...	...	*	1.8	4.1	2.1	0.8	*	*	*	*
2005 5/.....	1.4	...	...	*	2.2	4.2	2.2	*	*	*	*	*
2004 5/.....	1.3	...	...	*	2.4	3.9	1.5	0.9	*	*	*	*
2003 5/.....	1.0	...	...	*	1.4	3.0	2.0	*	*	*	*	*
2002.....	0.8	...	...	*	1.5	2.4	1.4	*	*	*	*	*
2001.....	0.8	...	...	*	1.6	2.4	1.4	*	*	*	*	*
2000.....	0.8	...	...	*	1.2	2.4	1.3	*	*	*	*	*
1999.....	0.9	...	...	*	1.6	2.5	1.4	*	*	*	*	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All origins 3/, both sexes

2007.....	0.2	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2006.....	0.2	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005.....	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.5	0.4	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.2	0.4	0.3	*	*	*	*	*

All origins 3/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins 3/, female

2007.....	0.5	...	...	*	0.7	1.6	1.0	0.3	*	*	*	*
2006.....	0.5	...	...	*	0.8	1.6	0.9	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.9	1.5	0.9	0.2	*	*	*	*
2004.....	0.4	...	...	*	0.8	1.4	0.7	0.4	*	*	*	*
2003.....	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001.....	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
1999.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Hispanic, both sexes

2007.....	0.3	...	...	*	0.3	0.7	0.6	*	*	*	*	*
2006.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2005.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2004.....	0.3	...	...	*	0.4	0.6	0.4	*	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.5	0.5	*	*	*	*	*
2002.....	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001.....	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000.....	0.2	...	...	*	0.3	0.5	0.4	*	*	*	*	*
1999.....	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*

Hispanic, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, female

2007.....	0.6	...	...	*	0.7	1.5	1.2	*	*	*	*	*
2006.....	0.7	...	...	*	1.1	1.9	1.1	*	*	*	*	*
2005.....	0.6	...	...	*	1.0	1.7	1.1	*	*	*	*	*
2004.....	0.6	...	...	*	0.9	1.4	0.9	*	*	*	*	*
2003.....	0.5	...	...	*	0.7	1.2	1.0	*	*	*	*	*
2002.....	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001.....	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000.....	0.5	...	...	*	0.7	1.1	0.9	*	*	*	*	*
1999.....	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic 4/, both sexes

2007.....	0.2	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2006.....	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2005.....	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.2	0.5	0.3	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*

Non-Hispanic 4/, male

2007.....	...	...	...	...	...	...	...	...	...	...	...	...
2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 4/, female

2007.....	0.5	...	...	*	0.7	1.6	1.0	0.3	*	*	*	*
2006.....	0.4	...	...	*	0.8	1.5	0.8	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.8	1.5	0.9	0.3	*	*	*	*
2004.....	0.4	...	...	*	0.7	1.4	0.6	0.3	*	*	*	*
2003.....	0.3	...	...	*	0.5	1.0	0.7	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2000.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
1999.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic white, both sexes

2007 5/.....	0.2	...	...	*	0.2	0.5	0.3	0.1	*	*	*	*
2006 5/.....	0.1	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2005 5/.....	0.2	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2004 5/.....	0.1	...	...	*	0.2	0.5	0.2	0.1	*	*	*	*
2003 5/.....	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
2002.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2000.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
1999.....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*

Non-Hispanic white, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic white, female

2007 5/.....	0.3	...	...	*	0.5	1.1	0.7	0.2	*	*	*	*
2006 5/.....	0.3	...	...	*	0.6	0.9	0.5	0.2	*	*	*	*
2005 5/.....	0.3	...	...	*	0.6	1.0	0.6	0.2	*	*	*	*
2004 5/.....	0.3	...	...	*	0.4	1.0	0.5	0.3	*	*	*	*
2003 5/.....	0.2	...	...	*	0.3	0.7	0.4	*	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
1999.....	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic black, both sexes

2007 5/.....	0.6	...	...	*	1.0	1.8	1.1	*	*	*	*	*
2006 5/.....	0.6	...	...	*	0.9	2.0	1.1	0.4	*	*	*	*
2005 5/.....	0.7	...	...	*	1.1	2.0	1.1	*	*	*	*	*
2004 5/.....	0.6	...	...	*	1.1	1.9	0.7	0.5	*	*	*	*
2003 5/.....	0.5	...	...	*	0.6	1.5	1.0	*	*	*	*	*
2002.....	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2001.....	0.4	...	...	*	0.7	1.2	0.6	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	1.1	0.6	*	*	*	*	*
1999.....	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*

Non-Hispanic black, male

2007 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2006 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic black, female

2007 5/.....	1.2	...	...	*	2.0	3.5	2.0	*	*	*	*	*
2006 5/.....	1.2	...	...	*	1.8	3.9	2.0	0.8	*	*	*	*
2005 5/.....	1.3	...	...	*	2.2	3.9	2.1	*	*	*	*	*
2004 5/.....	1.2	...	...	*	2.1	3.7	1.3	0.9	*	*	*	*
2003 5/.....	0.9	...	...	*	1.1	2.8	1.8	*	*	*	*	*
2002.....	0.7	...	...	*	1.5	2.2	1.2	*	*	*	*	*
2001.....	0.7	...	...	*	1.3	2.2	1.2	*	*	*	*	*
2000.....	0.7	...	...	*	1.1	2.1	1.1	*	*	*	*	*
1999.....	0.8	...	...	*	1.5	2.3	1.3	*	*	*	*	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/, both sexes

2007.....	3.5	135.9	3.3	0.9	0.9	1.0	1.2	1.6	2.1	1.8	3.1	4.8
2006.....	3.5	140.9	3.2	0.9	1.1	1.1	1.1	1.5	2.3	1.9	3.0	5.6
2005.....	3.5	135.2	3.2	1.0	1.2	1.1	1.2	1.6	2.4	1.8	3.1	6.4
2004.....	3.6	137.9	3.6	1.0	1.2	1.0	1.1	1.7	2.2	2.3	3.6	6.1
2003.....	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002.....	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001.....	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000.....	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999.....	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8

All origins 3/, male

2007.....	3.7	140.7	3.4	0.9	1.1	1.2	1.2	1.7	2.4	1.9	3.4	5.0
2006.....	3.8	146.6	2.9	0.9	1.3	1.3	1.2	1.8	2.6	2.1	3.6	6.8
2005.....	3.7	141.1	3.2	0.9	1.5	1.2	1.2	1.7	2.5	1.8	3.6	6.9
2004.....	3.8	139.0	3.9	1.0	1.3	1.2	1.3	1.8	2.3	2.5	3.7	6.8
2003.....	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002.....	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001.....	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000.....	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999.....	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5

All origins 3/, female

2007.....	3.2	130.8	3.2	0.9	0.8	0.9	1.2	1.5	1.8	1.7	3.0	4.7
2006.....	3.2	134.9	3.4	0.8	0.9	0.9	0.9	1.3	2.0	1.8	2.6	5.1
2005.....	3.3	129.0	3.3	1.0	0.9	1.0	1.1	1.6	2.2	1.8	2.8	6.2
2004.....	3.4	136.8	3.2	0.9	1.0	0.9	1.0	1.5	2.1	2.2	3.6	5.8
2003.....	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002.....	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001.....	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000.....	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999.....	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Hispanic, both sexes

2007.....	4.4	147.1	3.4	0.9	0.7	0.7	0.6	0.9	1.1	*	*	*
2006.....	4.4	151.0	3.3	0.8	1.0	0.7	0.6	1.1	1.3	1.7	*	*
2005.....	4.4	148.3	3.3	1.1	1.0	0.7	0.7	1.2	1.7	*	3.2	*
2004.....	4.3	141.0	4.0	0.9	0.9	0.7	0.6	1.1	1.4	1.6	3.4	*
2003.....	4.5	153.0	3.7	0.9	0.8	0.8	0.7	1.0	1.7	*	*	*
2002.....	4.4	150.7	3.5	1.0	1.0	0.7	0.7	1.1	1.3	*	3.8	*
2001.....	4.4	146.2	3.4	0.8	1.1	0.7	0.8	1.1	1.3	1.9	4.3	*
2000.....	4.6	153.3	3.8	0.8	0.8	0.7	0.8	1.1	1.8	2.0	4.7	*
1999.....	4.4	141.8	3.5	0.9	0.9	0.9	0.7	1.4	1.6	2.9	*	*

Hispanic, male

2007.....	4.5	148.9	3.8	0.8	1.0	0.8	0.7	1.1	*	*	*	*
2006.....	4.6	153.4	3.6	0.8	1.2	0.9	0.7	1.0	*	*	*	*
2005.....	4.5	151.2	3.5	1.2	1.1	0.7	0.8	1.1	2.0	*	*	*
2004.....	4.2	134.3	4.6	1.3	1.1	0.6	0.7	1.3	*	*	*	*
2003.....	4.5	156.9	3.7	0.7	0.8	0.9	0.8	1.2	*	*	*	*
2002.....	4.5	153.1	3.8	0.9	1.1	0.9	*	1.3	*	*	*	*
2001.....	4.7	153.4	3.5	1.0	1.1	0.7	1.0	1.5	*	*	*	*
2000.....	4.7	153.3	4.0	0.8	1.1	0.7	1.0	1.4	*	*	*	*
1999.....	4.3	138.1	4.3	0.9	0.9	0.7	*	1.6	*	*	*	*

Hispanic, female

2007.....	4.2	145.2	2.9	0.9	*	*	*	*	*	*	*	*
2006.....	4.2	148.5	3.1	0.7	0.7	*	*	1.3	*	*	*	*
2005.....	4.4	145.2	3.1	1.0	0.9	0.6	0.7	1.3	*	*	*	*
2004.....	4.3	147.9	3.3	0.6	0.6	0.8	*	*	*	*	*	*
2003.....	4.4	148.9	3.7	1.0	0.8	0.6	*	*	*	*	*	*
2002.....	4.3	148.2	3.3	1.0	0.9	*	0.8	*	*	*	*	*
2001.....	4.1	138.8	3.3	0.6	1.1	*	*	*	*	*	*	*
2000.....	4.4	153.3	3.5	0.7	*	0.7	*	*	*	*	*	*
1999.....	4.5	145.8	2.7	0.9	0.9	1.0	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (000-099)

Non-Hispanic 4/, both sexes

2007.....	3.3	131.5	3.3	0.9	1.0	1.1	1.3	1.6	2.2	1.8	3.2	4.9
2006.....	3.3	136.8	3.1	0.9	1.1	1.2	1.2	1.6	2.4	1.9	3.1	5.8
2005.....	3.3	130.4	3.2	1.0	1.2	1.2	1.3	1.7	2.4	1.9	3.1	6.6
2004.....	3.4	135.9	3.4	1.0	1.2	1.1	1.2	1.7	2.2	2.4	3.7	6.2
2003.....	3.5	136.0	3.4	0.9	1.1	1.1	1.3	1.7	2.4	2.3	3.7	6.7
2002.....	3.6	134.9	3.3	1.0	1.2	1.3	1.4	1.8	2.5	2.5	3.8	6.7
2001.....	3.5	132.8	3.7	0.9	1.3	1.3	1.4	1.7	2.4	2.5	3.7	7.2
2000.....	3.6	148.8	3.1	1.0	1.2	1.3	1.3	1.7	2.4	2.6	3.9	7.5
1999.....	3.6	143.1	3.6	1.1	1.2	1.2	1.3	1.8	2.4	2.5	4.0	7.9

Non-Hispanic 4/, male

2007.....	3.5	137.2	3.2	1.0	1.1	1.3	1.3	1.7	2.5	1.9	3.5	5.0
2006.....	3.6	143.6	2.7	0.9	1.3	1.4	1.3	1.9	2.7	2.1	3.6	6.8
2005.....	3.6	137.1	3.1	0.9	1.6	1.3	1.3	1.8	2.6	1.9	3.6	7.2
2004.....	3.6	139.1	3.7	0.9	1.4	1.3	1.4	1.9	2.3	2.5	3.7	7.0
2003.....	3.7	144.2	3.3	1.0	1.4	1.2	1.5	1.8	2.5	2.4	3.5	6.7
2002.....	3.8	138.1	3.5	1.0	1.4	1.5	1.4	2.0	2.6	2.6	3.9	7.0
2001.....	3.7	139.4	3.6	1.0	1.6	1.4	1.5	1.7	2.3	2.7	4.0	7.6
2000.....	3.9	154.4	3.1	1.0	1.4	1.5	1.5	1.9	2.2	2.7	3.9	10.4
1999.....	3.8	148.4	3.9	1.1	1.4	1.3	1.4	1.9	2.6	2.4	4.2	8.5

Non-Hispanic 4/, female

2007.....	3.1	125.6	3.3	0.9	0.8	1.0	1.3	1.5	1.9	1.7	3.1	4.8
2006.....	3.0	129.7	3.5	0.9	0.9	1.0	1.0	1.3	2.1	1.8	2.7	5.3
2005.....	3.1	123.3	3.3	1.0	0.9	1.1	1.2	1.6	2.3	1.9	2.8	6.3
2004.....	3.2	132.6	3.2	1.0	1.1	1.0	1.0	1.6	2.1	2.2	3.6	5.9
2003.....	3.3	127.6	3.4	0.9	0.9	1.0	1.2	1.6	2.3	2.2	3.8	6.8
2002.....	3.4	131.5	3.2	1.0	1.1	1.1	1.3	1.7	2.4	2.3	3.8	6.6
2001.....	3.3	125.8	3.7	0.9	1.0	1.2	1.3	1.7	2.4	2.2	3.5	7.0
2000.....	3.4	142.8	3.0	1.0	1.0	1.1	1.2	1.4	2.5	2.4	3.9	6.4
1999.....	3.4	137.4	3.3	1.1	0.9	1.1	1.2	1.6	2.2	2.6	3.9	7.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (000-099)

Non-Hispanic white, both sexes

2007 5/.....	3.1	123.6	2.9	0.9	0.9	1.1	1.3	1.7	2.3	1.8	3.3	5.0
2006 5/.....	3.2	129.5	2.8	0.8	1.0	1.2	1.2	1.6	2.5	2.0	3.2	5.9
2005 5/.....	3.2	123.0	2.9	0.9	1.2	1.2	1.3	1.8	2.6	2.0	3.3	6.7
2004 5/.....	3.3	130.5	2.9	0.9	1.1	1.2	1.2	1.8	2.4	2.5	3.7	6.5
2003 5/.....	3.3	129.8	3.0	0.9	1.1	1.2	1.4	1.7	2.6	2.3	3.8	6.9
2002.....	3.4	131.2	3.0	1.0	1.2	1.3	1.4	1.9	2.5	2.5	3.9	6.9
2001.....	3.4	127.4	3.1	0.9	1.2	1.3	1.4	1.7	2.5	2.5	3.8	7.4
2000.....	3.4	141.2	2.6	0.9	1.1	1.3	1.3	1.7	2.4	2.6	4.0	7.8
1999.....	3.4	136.8	3.0	0.9	1.1	1.3	1.3	1.8	2.5	2.5	4.0	8.0

Non-Hispanic white, male

2007 5/.....	3.3	131.0	2.9	0.9	1.0	1.2	1.3	1.7	2.6	2.0	3.5	5.2
2006 5/.....	3.4	135.2	2.6	0.9	1.2	1.4	1.4	1.8	2.8	2.1	3.7	6.8
2005 5/.....	3.4	129.9	3.0	0.8	1.6	1.3	1.3	1.9	2.7	2.0	3.8	7.2
2004 5/.....	3.4	132.8	3.1	0.9	1.2	1.4	1.4	2.0	2.4	2.7	3.7	7.4
2003 5/.....	3.4	136.1	2.9	0.9	1.3	1.2	1.5	1.8	2.6	2.3	3.4	6.8
2002.....	3.6	137.2	3.2	1.0	1.4	1.6	1.4	2.0	2.6	2.7	3.8	7.3
2001.....	3.6	135.5	3.1	1.0	1.4	1.5	1.5	1.7	2.4	2.9	4.1	7.4
2000.....	3.6	147.9	2.6	0.9	1.3	1.6	1.4	1.9	2.3	2.6	4.1	10.8
1999.....	3.6	144.4	3.2	0.9	1.4	1.3	1.4	2.0	2.7	2.4	4.1	8.6

Non-Hispanic white, female

2007 5/.....	2.9	115.9	2.9	0.8	0.9	1.0	1.3	1.6	2.0	1.7	3.1	4.9
2006 5/.....	2.9	123.5	2.9	0.7	0.9	1.0	1.0	1.4	2.2	1.9	2.8	5.5
2005 5/.....	3.0	115.8	2.9	0.9	0.8	1.0	1.3	1.7	2.4	2.0	2.9	6.5
2004 5/.....	3.2	128.0	2.8	0.9	1.0	1.1	1.1	1.6	2.3	2.4	3.7	6.1
2003 5/.....	3.2	123.2	3.0	0.9	0.9	1.1	1.3	1.7	2.5	2.3	4.0	6.9
2002.....	3.3	125.0	2.9	1.0	1.1	1.1	1.3	1.8	2.5	2.4	3.9	6.7
2001.....	3.2	118.8	3.2	0.7	1.0	1.2	1.3	1.7	2.6	2.3	3.6	7.4
2000.....	3.2	134.2	2.6	1.0	1.0	1.1	1.2	1.6	2.5	2.6	4.0	6.6
1999.....	3.3	128.9	2.9	1.0	0.8	1.2	1.2	1.7	2.3	2.6	4.0	7.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (000-099)

Non-Hispanic black, both sexes

2007 5/.....	4.6	166.3	4.6	1.2	1.3	1.5	1.5	1.6	1.8	2.0	3.4	*
2006 5/.....	4.6	174.3	4.2	1.1	1.4	1.3	1.4	1.9	2.0	1.6	3.0	5.6
2005 5/.....	4.4	163.4	4.1	1.3	1.6	1.5	1.4	1.4	2.0	*	2.1	6.0
2004 5/.....	4.6	164.3	5.1	1.3	1.8	1.1	1.3	1.7	1.7	1.6	4.0	*
2003 5/.....	4.6	160.5	4.8	1.1	1.5	1.4	1.2	1.7	2.2	2.4	3.1	5.9
2002.....	4.7	156.9	4.7	1.3	1.7	1.5	1.6	1.8	2.7	2.4	3.4	*
2001.....	4.7	163.5	5.4	1.1	2.0	1.3	1.6	1.8	1.9	2.1	3.0	*
2000.....	5.0	184.1	5.2	1.4	1.6	1.4	1.6	1.6	2.4	2.7	2.8	*
1999.....	4.8	174.8	5.8	1.6	1.5	1.1	1.4	1.9	2.0	2.9	4.3	7.4

Non-Hispanic black, male

2007 5/.....	5.0	167.8	4.4	1.2	1.6	2.0	1.5	1.7	2.2	*	*	*
2006 5/.....	5.1	182.2	3.2	0.9	1.6	1.7	1.5	2.4	2.4	*	*	*
2005 5/.....	4.8	167.1	3.6	1.2	2.0	1.7	1.7	1.5	2.3	*	*	*
2004 5/.....	5.2	170.5	5.9	1.2	2.4	1.4	1.6	1.8	2.0	*	*	*
2003 5/.....	5.2	173.1	5.0	1.2	1.8	1.5	1.2	2.0	2.8	3.6	*	*
2002.....	5.2	153.3	4.8	1.4	2.0	1.8	1.7	2.3	3.5	3.1	*	*
2001.....	5.1	167.5	5.5	1.0	2.5	1.2	1.8	1.7	2.1	*	*	*
2000.....	5.5	186.0	5.6	1.5	2.1	1.7	1.9	2.2	*	3.9	*	*
1999.....	5.3	179.5	6.6	1.6	1.8	1.3	1.6	2.3	2.1	*	*	*

Non-Hispanic black, female

2007 5/.....	4.3	164.7	4.8	1.3	0.9	1.0	1.4	1.5	1.5	2.3	3.1	*
2006 5/.....	4.2	166.0	5.3	1.2	1.1	1.0	1.2	1.5	1.7	*	*	*
2005 5/.....	4.1	159.6	4.7	1.4	1.2	1.4	1.1	1.4	1.7	*	*	*
2004 5/.....	4.1	157.8	4.3	1.4	1.2	0.8	0.9	1.7	1.4	*	3.4	*
2003 5/.....	4.0	147.3	4.7	1.1	1.2	1.4	1.3	1.5	1.8	*	*	*
2002.....	4.4	160.6	4.7	1.1	1.3	1.2	1.6	1.5	2.0	*	*	*
2001.....	4.4	159.2	5.4	1.3	1.4	1.4	1.5	1.8	1.8	2.4	*	*
2000.....	4.5	182.2	4.8	1.2	1.2	1.1	1.4	1.2	3.1	*	*	*
1999.....	4.4	170.0	5.0	1.6	1.1	1.0	1.2	1.5	1.9	3.2	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes

2007.....	11.1	85.0	1.4	0.3	1.4	2.2	3.3	5.0	6.8	12.9	44.8	251.4
2006.....	10.6	83.8	1.5	0.3	1.6	2.5	3.7	5.2	6.5	12.2	41.7	236.1
2005.....	10.8	87.4	1.4	0.4	1.8	2.9	4.7	6.7	7.2	12.8	39.0	224.4
2004.....	10.3	82.4	1.2	0.3	1.6	2.7	4.3	5.8	7.0	12.6	39.8	221.5
2003.....	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002.....	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001.....	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000.....	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999.....	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7

All origins 3/, male

2007.....	9.6	96.3	1.6	0.3	1.8	2.8	4.1	6.3	8.8	15.5	45.8	197.6
2006.....	9.5	95.2	1.6	0.4	2.0	3.5	4.8	6.7	8.3	14.8	44.2	191.1
2005.....	10.0	101.8	1.6	0.5	2.5	4.0	5.8	8.7	9.3	15.5	40.7	183.9
2004.....	9.5	96.9	1.2	0.3	2.1	3.6	5.6	7.6	9.1	15.0	41.8	183.6
2003.....	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002.....	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001.....	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000.....	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999.....	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3

All origins 3/, female

2007.....	12.6	73.1	1.3	0.2	0.9	1.5	2.5	3.7	4.9	10.7	44.1	277.0
2006.....	11.7	71.9	1.3	0.3	1.1	1.5	2.7	3.8	4.8	10.0	39.9	257.2
2005.....	11.5	72.3	1.3	0.3	1.1	1.9	3.6	4.7	5.3	10.6	37.8	243.0
2004.....	11.0	67.3	1.1	0.3	1.0	1.8	3.0	4.0	5.1	10.6	38.4	238.5
2003.....	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002.....	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001.....	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000.....	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999.....	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, both sexes

2007.....	4.3	54.0	1.4	0.3	1.5	1.8	2.4	3.6	4.3	7.4	23.9	108.7
2006.....	4.0	52.5	1.2	*	1.3	1.9	2.5	3.0	3.8	6.9	18.7	117.0
2005.....	4.4	59.5	1.2	0.3	1.5	2.4	3.1	3.6	3.6	7.8	18.6	105.2
2004.....	3.9	50.4	1.1	*	1.6	2.1	3.1	3.6	3.9	6.2	16.4	99.4
2003.....	4.1	56.8	0.9	0.3	1.4	2.3	3.1	3.7	3.5	6.1	18.4	110.9
2002.....	4.0	56.6	1.3	*	1.3	2.0	3.0	3.9	4.8	7.0	18.7	91.5
2001.....	5.9	60.7	1.6	0.5	2.8	4.2	6.0	8.3	6.9	9.9	19.3	108.2
2000.....	5.9	65.6	1.7	0.5	2.1	4.0	6.7	8.0	6.7	9.9	20.5	102.2
1999.....	3.8	54.9	1.0	*	1.3	2.2	3.0	3.1	3.8	7.0	19.4	87.0

Hispanic, male

2007.....	4.8	61.3	1.4	*	2.4	2.6	3.5	5.0	5.7	9.5	26.3	91.5
2006.....	4.5	62.1	1.1	*	1.9	2.9	3.9	4.2	5.4	9.7	17.4	90.2
2005.....	5.1	68.9	1.3	*	2.0	3.6	4.3	5.5	5.1	10.2	21.0	99.6
2004.....	4.5	60.0	1.2	*	2.3	2.8	4.5	4.9	5.8	7.1	15.7	86.9
2003.....	4.8	69.9	*	*	1.7	3.4	4.4	5.7	5.6	8.1	18.8	95.4
2002.....	4.6	66.4	1.6	*	1.7	2.9	4.0	4.7	6.3	9.6	23.4	88.7
2001.....	7.4	67.2	1.8	*	3.9	6.0	8.5	12.6	10.6	14.2	25.0	87.6
2000.....	7.5	78.6	2.1	*	2.9	5.8	9.9	12.4	9.0	14.3	25.6	106.8
1999.....	4.4	61.3	*	*	1.9	2.9	4.3	4.7	5.5	8.3	24.0	77.6

Hispanic, female

2007.....	3.7	46.4	1.4	*	*	0.9	1.2	2.1	3.1	5.7	22.3	118.0
2006.....	3.4	42.5	1.3	*	0.7	0.7	1.0	1.7	2.3	4.6	19.7	131.2
2005.....	3.5	49.8	1.2	*	0.9	0.9	1.8	1.6	2.3	5.9	17.0	108.2
2004.....	3.3	40.4	*	*	0.8	1.3	1.6	2.4	2.2	5.5	16.8	105.9
2003.....	3.2	43.2	*	*	1.0	1.0	1.6	1.7	*	4.6	18.2	118.8
2002.....	3.3	46.4	*	*	0.9	1.0	2.0	3.2	3.5	5.0	15.5	92.8
2001.....	4.3	53.9	*	0.6	1.5	2.1	3.4	4.1	3.6	6.5	15.4	118.2
2000.....	4.1	52.1	1.4	*	1.3	2.0	3.2	3.7	4.6	6.5	17.2	99.9
1999.....	3.1	48.2	*	*	*	1.3	1.6	1.7	2.4	6.0	16.3	91.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 4/, both sexes

2007.....	12.3	94.5	1.4	0.3	1.3	2.2	3.4	5.2	7.0	13.3	46.1	258.9
2006.....	11.7	92.7	1.5	0.3	1.6	2.6	3.9	5.4	6.6	12.5	43.0	241.9
2005.....	11.8	94.5	1.5	0.4	1.9	3.1	4.9	6.9	7.5	13.1	39.9	229.6
2004.....	11.2	90.9	1.2	0.3	1.6	2.8	4.5	5.9	7.2	13.0	40.9	226.7
2003.....	11.8	89.4	1.8	0.5	2.1	3.5	5.7	7.3	7.7	14.3	39.9	228.4
2002.....	11.3	92.8	1.7	0.4	1.9	3.6	5.6	6.9	7.7	13.5	39.6	211.9
2001.....	12.1	92.6	1.8	0.4	2.3	3.7	7.0	8.9	9.2	15.4	40.7	209.6
2000.....	12.0	102.4	1.6	0.4	2.2	3.8	6.5	8.2	9.3	16.8	41.4	211.6
1999.....	10.3	101.9	1.6	0.4	1.6	2.8	4.7	6.0	8.0	15.0	36.7	190.9

Non-Hispanic 4/, male

2007.....	10.5	107.2	1.6	0.3	1.7	2.9	4.2	6.5	9.1	15.9	47.1	203.6
2006.....	10.3	104.3	1.7	0.4	2.0	3.6	4.9	7.0	8.4	15.1	45.9	196.3
2005.....	10.8	110.2	1.7	0.5	2.6	4.0	5.9	8.9	9.6	15.8	41.6	187.5
2004.....	10.2	106.6	1.2	0.3	2.0	3.8	5.7	7.8	9.2	15.5	43.1	188.0
2003.....	11.0	102.9	2.0	0.5	2.8	4.7	7.2	9.4	9.9	17.7	43.0	181.1
2002.....	10.9	106.2	1.9	0.4	2.5	4.8	7.3	9.1	10.1	16.3	44.6	178.6
2001.....	11.8	106.2	2.1	0.5	3.0	4.9	9.0	11.7	12.0	18.2	44.6	178.2
2000.....	11.9	118.0	1.7	0.5	2.9	5.1	8.6	11.2	12.3	20.7	46.6	185.4
1999.....	10.0	116.7	1.9	0.4	2.2	3.7	6.1	8.2	10.6	18.7	40.6	169.5

Non-Hispanic 4/, female

2007.....	14.0	81.2	1.3	0.2	0.9	1.6	2.7	3.9	5.1	11.1	45.5	285.0
2006.....	13.0	80.5	1.3	0.3	1.2	1.7	3.0	4.0	5.0	10.4	41.1	263.1
2005.....	12.7	78.0	1.3	0.3	1.1	2.1	3.9	4.9	5.5	10.8	38.8	248.9
2004.....	12.1	74.3	1.2	0.3	1.0	1.9	3.2	4.2	5.3	10.9	39.4	244.0
2003.....	12.6	75.3	1.6	0.5	1.4	2.4	4.1	5.3	5.6	11.4	37.8	249.2
2002.....	11.7	78.7	1.4	0.3	1.3	2.3	4.0	4.8	5.4	11.2	36.3	226.3
2001.....	12.3	78.3	1.5	0.4	1.5	2.6	5.1	6.2	6.6	13.1	38.0	222.7
2000.....	12.1	86.1	1.5	0.4	1.5	2.6	4.5	5.4	6.6	13.6	38.0	222.3
1999.....	10.5	86.3	1.4	0.3	1.0	2.0	3.3	3.9	5.7	11.9	34.2	199.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic white, both sexes

2007 5/.....	13.0	79.1	1.3	0.2	1.3	2.0	3.3	4.7	6.6	12.8	46.1	264.8
2006 5/.....	12.4	77.1	1.2	0.3	1.6	2.6	3.8	5.2	6.3	12.1	42.8	247.4
2005 5/.....	12.3	77.9	1.2	0.3	1.8	3.0	4.8	6.5	6.9	12.5	39.8	234.9
2004 5/.....	11.7	71.6	1.0	0.2	1.5	2.8	4.3	5.6	6.7	12.5	40.7	230.0
2003 5/.....	12.3	72.6	1.5	0.4	2.2	3.4	5.5	6.7	7.1	13.7	40.1	232.9
2002.....	11.6	76.0	1.3	0.3	1.8	3.4	5.4	6.1	7.1	13.1	39.4	215.8
2001.....	12.4	73.1	1.5	0.3	2.2	3.6	6.9	8.5	8.6	14.7	40.4	213.3
2000.....	12.2	79.7	1.4	0.3	2.1	3.6	6.3	7.6	8.6	15.9	40.9	215.8
1999.....	10.2	79.3	1.4	0.3	1.4	2.5	4.1	5.2	7.3	14.3	35.7	193.8

Non-Hispanic white, male

2007 5/.....	10.7	92.3	1.6	0.3	1.6	2.6	3.9	5.8	8.5	15.2	46.9	207.6
2006 5/.....	10.5	89.1	1.4	0.4	2.0	3.5	4.7	6.5	8.0	14.2	45.1	199.4
2005 5/.....	10.9	94.6	1.4	0.3	2.5	4.0	5.9	8.4	8.8	14.9	41.0	189.8
2004 5/.....	10.3	83.1	1.0	0.2	2.1	3.7	5.6	7.2	8.4	14.9	42.8	191.0
2003 5/.....	11.0	87.4	1.6	0.4	2.9	4.6	6.9	8.6	9.0	16.8	42.8	183.3
2002.....	10.7	88.2	1.4	0.3	2.5	4.7	6.9	7.9	9.3	15.7	43.6	181.5
2001.....	11.8	85.2	1.7	0.4	2.9	4.9	8.8	11.1	10.9	17.3	44.7	180.4
2000.....	11.8	91.8	1.7	0.4	2.8	4.8	8.5	10.3	11.1	19.3	46.0	189.6
1999.....	9.6	92.3	1.7	0.3	1.9	3.3	5.5	6.9	9.5	17.7	39.2	171.1

Non-Hispanic white, female

2007 5/.....	15.3	65.3	1.1	0.2	0.9	1.4	2.6	3.5	4.8	10.7	45.6	291.9
2006 5/.....	14.2	64.5	1.1	0.2	1.1	1.7	2.9	3.9	4.7	10.3	41.2	269.7
2005 5/.....	13.7	60.4	1.0	0.2	1.1	2.0	3.7	4.6	5.2	10.4	39.0	255.5
2004 5/.....	13.0	59.4	1.0	0.2	1.0	1.8	3.0	4.0	5.0	10.4	39.3	247.5
2003 5/.....	13.5	57.1	1.4	0.4	1.4	2.2	4.0	4.8	5.4	11.0	38.3	254.7
2002.....	12.5	63.2	1.3	0.2	1.1	2.2	3.8	4.3	5.1	10.9	36.6	230.6
2001.....	13.1	60.4	1.3	0.3	1.5	2.4	5.0	5.9	6.4	12.5	37.6	227.0
2000.....	12.6	66.9	1.2	0.3	1.3	2.3	4.1	4.9	6.2	13.0	37.5	226.4
1999.....	10.9	65.5	1.0	0.3	0.9	1.8	2.8	3.5	5.2	11.4	33.4	202.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic black, both sexes

2007 5/.....	12.0	161.6	1.9	0.4	1.8	3.9	5.5	9.5	12.1	21.2	59.3	245.2
2006 5/.....	11.4	161.4	2.8	0.6	1.9	3.5	5.8	8.9	10.9	19.6	56.9	226.9
2005 5/.....	12.4	170.4	2.5	0.9	2.5	4.2	7.0	11.7	13.9	22.1	51.8	213.3
2004 5/.....	11.9	174.5	2.0	0.6	1.8	4.1	6.6	9.8	13.4	20.8	53.1	223.3
2003 5/.....	12.7	161.6	3.3	0.8	2.5	5.1	8.6	13.6	14.3	24.0	45.7	214.5
2002.....	12.6	165.4	2.7	0.8	2.6	5.3	8.8	13.9	13.9	20.4	49.9	198.1
2001.....	12.9	176.9	3.0	0.6	2.7	5.2	9.3	14.0	15.1	24.6	50.3	194.2
2000.....	13.9	205.6	2.3	0.7	3.2	5.9	9.3	14.8	17.3	29.2	56.1	189.2
1999.....	13.0	202.0	2.9	0.5	2.7	5.0	9.1	13.2	16.0	24.6	54.8	182.0

Non-Hispanic black, male

2007 5/.....	12.2	173.4	1.9	*	2.4	5.0	7.1	12.6	17.3	28.2	64.9	187.6
2006 5/.....	12.0	172.0	3.1	0.7	2.3	4.7	7.3	12.5	14.0	28.0	70.4	196.1
2005 5/.....	13.3	187.0	2.7	0.9	3.4	5.4	8.1	15.2	19.6	29.5	62.5	196.0
2004 5/.....	12.8	208.3	2.2	*	2.3	5.3	8.1	13.8	18.9	25.4	59.3	182.9
2003 5/.....	13.8	171.9	3.6	0.8	3.0	7.0	11.2	17.9	20.7	32.2	52.8	197.1
2002.....	14.4	186.3	3.6	0.9	3.1	7.1	11.7	19.6	20.0	26.4	68.6	182.7
2001.....	14.2	195.9	3.7	0.8	3.8	6.1	11.4	18.2	23.8	30.7	52.2	186.0
2000.....	15.7	236.8	2.2	0.9	3.8	7.8	10.9	20.6	25.1	40.0	66.0	163.2
1999.....	14.9	228.0	2.9	0.6	3.6	6.9	11.6	19.4	23.0	32.8	66.9	180.7

Non-Hispanic black, female

2007 5/.....	11.8	149.2	1.9	*	1.2	2.8	4.1	6.9	7.9	16.1	56.2	269.1
2006 5/.....	10.8	150.2	2.4	*	1.4	2.3	4.4	5.8	8.4	13.6	49.2	239.5
2005 5/.....	11.5	153.1	2.3	0.8	1.5	3.1	5.9	8.6	9.4	16.9	45.8	220.2
2004 5/.....	11.0	139.4	1.7	*	1.3	3.1	5.3	6.3	8.9	17.6	49.6	239.3
2003 5/.....	11.6	150.9	3.0	0.8	1.9	3.5	6.3	10.0	9.3	18.2	41.7	221.2
2002.....	10.9	143.6	1.9	0.7	2.0	3.7	6.2	9.0	8.9	16.1	39.3	204.0
2001.....	11.7	156.8	2.2	*	1.7	4.3	7.4	10.3	8.1	20.3	49.2	197.2
2000.....	12.4	173.4	2.4	*	2.6	4.3	7.8	9.9	11.0	21.6	50.5	198.9
1999.....	11.2	175.3	2.8	*	1.9	3.2	7.0	7.9	10.3	18.8	48.0	182.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All origins 3/, both sexes												
2007.....	79.0	31.4	3.6	2.2	4.6	7.8	16.4	37.6	70.0	149.0	469.6	1,701.9
2006.....	79.3	28.4	3.4	2.0	5.0	8.3	18.4	40.1	71.6	151.6	470.5	1,724.2
2005.....	73.4	29.2	3.8	2.1	4.9	7.9	18.5	40.3	69.5	146.1	434.3	1,559.1
2004.....	68.8	29.2	3.8	2.2	4.7	8.1	18.3	38.9	67.0	141.4	409.6	1,461.6
2003.....	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
2002.....	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001.....	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000.....	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999.....	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7
All origins 3/, male												
2007.....	64.9	35.8	3.9	2.5	5.3	8.6	18.7	45.3	80.6	161.6	474.7	1,449.1
2006.....	65.2	32.5	3.6	2.1	5.8	9.5	21.4	48.3	82.7	164.8	474.4	1,455.0
2005.....	60.9	34.8	4.3	2.3	5.8	8.9	22.0	49.1	80.9	156.1	437.0	1,305.3
2004.....	57.2	33.8	4.0	2.3	5.4	9.2	21.5	47.8	77.1	151.9	410.5	1,236.6
2003.....	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
2002.....	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001.....	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000.....	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999.....	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4
All origins 3/, female												
2007.....	92.7	26.9	3.4	1.8	3.9	6.8	14.1	30.1	60.1	138.3	466.0	1,822.1
2006.....	93.0	24.1	3.1	1.8	4.2	7.1	15.3	32.2	61.3	140.5	467.9	1,850.2
2005.....	85.6	23.2	3.3	1.9	4.0	6.8	15.0	31.8	59.0	137.7	432.5	1,675.6
2004.....	79.9	24.5	3.6	2.0	3.9	7.0	15.1	30.4	57.6	132.5	408.9	1,562.8
2003.....	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
2002.....	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001.....	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000.....	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999.....	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

#### Hispanic, both sexes

2007.....	25.7	28.8	3.1	2.0	3.9	5.1	11.0	26.8	51.4	106.5	318.7	1,049.8
2006.....	26.4	24.6	3.3	1.8	4.4	6.1	13.8	28.9	52.9	114.4	326.2	1,067.7
2005.....	24.0	29.1	4.1	1.6	4.0	6.1	13.3	28.3	52.5	107.5	290.7	890.7
2004.....	22.3	26.1	3.3	1.9	3.7	5.4	13.1	27.8	49.1	101.0	276.7	880.5
2003.....	23.1	27.1	4.3	2.1	4.1	6.1	14.0	30.3	53.5	106.9	284.5	887.2
2002.....	21.6	25.1	3.8	2.1	3.7	5.6	13.0	27.5	48.5	111.2	272.2	870.8
2001.....	21.2	22.9	3.7	1.7	3.7	5.2	12.3	26.3	49.7	107.2	277.5	991.0
2000.....	20.7	25.4	3.7	1.7	3.3	5.9	13.8	27.5	47.5	103.4	284.0	913.0
1999.....	21.5	26.5	3.7	1.6	3.5	6.5	14.0	28.2	51.4	109.3	263.5	1,039.7

#### Hispanic, male

2007.....	23.5	33.3	3.1	2.2	4.1	5.6	13.2	33.9	60.0	118.5	319.2	913.8
2006.....	24.9	28.0	3.3	1.9	5.3	7.4	17.6	37.3	64.2	127.6	334.2	897.6
2005.....	23.2	34.6	4.2	1.6	4.8	7.2	16.6	36.7	64.9	115.5	300.0	789.5
2004.....	21.3	28.7	3.5	2.1	4.5	6.2	17.2	37.5	57.6	109.6	264.7	770.5
2003.....	22.2	32.6	4.5	2.3	4.4	7.3	16.7	38.4	64.6	116.4	301.8	757.3
2002.....	21.1	27.2	4.1	2.4	4.0	6.6	17.1	35.5	56.4	126.0	290.1	808.2
2001.....	20.2	25.4	4.3	1.8	4.1	6.0	15.9	34.7	59.1	117.2	273.7	939.0
2000.....	20.2	25.3	3.6	1.8	3.9	7.0	18.2	35.6	57.7	119.9	273.9	870.7
1999.....	21.0	32.1	3.4	1.6	3.8	8.1	18.8	37.0	62.6	123.9	276.3	923.1

#### Hispanic, female

2007.....	28.1	24.2	3.2	1.8	3.6	4.4	8.6	19.6	43.3	96.7	318.3	1,123.6
2006.....	28.1	21.1	3.4	1.8	3.5	4.6	9.5	20.4	42.4	103.7	320.7	1,158.1
2005.....	24.9	23.3	3.9	1.6	3.2	4.8	9.6	19.8	41.1	101.0	284.2	943.7
2004.....	23.4	23.5	3.1	1.8	2.8	4.4	8.6	17.9	41.3	94.1	285.0	937.0
2003.....	24.1	21.5	4.0	2.0	3.8	4.7	11.0	22.1	43.5	99.3	272.6	953.1
2002.....	22.1	22.8	3.5	1.7	3.2	4.4	8.6	19.6	41.4	99.5	260.0	901.9
2001.....	22.3	20.2	3.1	1.7	3.2	4.2	8.4	18.0	41.4	99.4	280.2	1,016.5
2000.....	21.4	25.5	3.8	1.5	2.5	4.5	9.0	19.6	38.5	90.4	290.9	933.8
1999.....	22.0	20.6	4.0	1.7	3.1	4.6	8.8	19.7	41.6	97.9	255.0	1,096.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 4/, both sexes

2007.....	88.3	32.1	3.8	2.2	4.8	8.4	17.3	38.8	71.4	152.2	479.0	1,735.2
2006.....	88.3	29.5	3.4	2.0	5.1	8.8	19.1	41.2	73.1	154.2	478.9	1,755.2
2005.....	81.6	28.9	3.7	2.2	5.1	8.3	19.3	41.5	70.8	148.8	442.3	1,589.3
2004.....	76.2	30.0	3.9	2.2	4.9	8.7	19.1	40.0	68.3	144.0	416.3	1,485.2
2003.....	76.5	30.1	4.0	2.3	5.0	8.8	19.6	39.6	68.0	146.7	417.4	1,531.7
2002.....	74.4	29.0	4.3	2.1	4.7	8.3	19.7	38.6	66.4	142.2	410.7	1,514.9
2001.....	71.8	29.0	4.3	2.2	4.6	8.3	18.6	37.0	64.2	142.7	399.8	1,491.3
2000.....	69.3	29.6	4.0	2.2	4.7	8.1	18.4	35.8	63.3	137.5	391.9	1,462.0
1999.....	66.8	30.8	4.3	2.4	4.8	7.8	18.1	34.1	60.9	137.8	376.6	1,420.1

Non-Hispanic 4/, male

2007.....	72.5	36.4	4.1	2.6	5.5	9.5	19.8	46.6	82.2	164.7	484.4	1,479.5
2006.....	72.4	33.8	3.7	2.2	5.9	10.0	22.1	49.4	84.1	167.2	482.6	1,484.0
2005.....	67.4	34.5	4.2	2.4	5.9	9.4	23.0	50.3	81.9	158.7	444.6	1,330.8
2004.....	63.3	35.0	4.1	2.4	5.6	9.9	22.2	48.6	78.4	154.6	418.1	1,257.5
2003.....	63.0	35.0	4.2	2.6	5.8	9.7	23.1	47.9	77.2	157.9	420.3	1,278.3
2002.....	61.2	34.9	4.6	2.3	5.4	9.7	23.1	47.1	75.6	153.8	411.5	1,280.6
2001.....	58.8	32.7	4.6	2.2	5.4	9.2	21.9	44.2	71.7	155.2	402.2	1,282.3
2000.....	56.9	33.6	4.4	2.3	5.5	9.4	21.7	43.0	71.3	149.4	395.5	1,279.8
1999.....	55.2	35.1	4.7	2.4	5.6	9.0	21.7	40.5	69.1	149.6	387.8	1,234.0

Non-Hispanic 4/, female

2007.....	103.4	27.6	3.5	1.8	3.9	7.4	15.0	31.3	61.4	141.5	475.2	1,856.0
2006.....	103.5	25.0	3.0	1.8	4.3	7.7	16.3	33.4	62.8	143.2	476.5	1,881.3
2005.....	95.1	23.0	3.1	2.0	4.1	7.2	15.8	33.1	60.5	140.4	440.7	1,707.2
2004.....	88.5	24.7	3.7	2.1	4.2	7.5	16.0	31.6	58.9	135.1	415.1	1,587.0
2003.....	89.4	25.1	3.7	1.9	4.2	7.9	16.2	31.7	59.5	137.3	415.5	1,643.1
2002.....	86.8	22.7	4.0	1.9	4.1	7.0	16.4	30.4	57.9	132.5	410.1	1,615.9
2001.....	84.2	25.2	4.0	2.1	3.9	7.4	15.4	30.0	57.3	132.2	398.3	1,578.3
2000.....	81.0	25.4	3.5	2.1	3.9	6.8	15.2	28.9	55.9	127.6	389.5	1,535.7
1999.....	77.9	26.2	3.9	2.3	4.0	6.7	14.4	28.0	53.3	128.2	369.3	1,494.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic white, both sexes

2007 5/.....	97.1	23.8	3.1	2.1	4.5	7.6	15.8	36.4	68.6	149.5	483.0	1,771.7
2006 5/.....	96.9	23.3	3.0	1.9	4.8	8.2	17.6	38.5	69.8	151.7	482.4	1,790.9
2005 5/.....	88.9	23.2	3.3	2.0	4.8	7.5	17.8	38.5	67.6	145.9	443.5	1,618.2
2004 5/.....	82.7	24.8	3.5	2.1	4.5	7.6	17.4	36.9	64.8	140.9	417.7	1,511.9
2003 5/.....	83.0	24.6	3.5	2.1	4.8	7.8	17.8	36.2	64.3	144.1	417.9	1,558.9
2002.....	80.5	24.7	3.8	2.0	4.3	7.4	17.8	35.3	62.7	139.1	410.5	1,540.5
2001.....	77.3	23.2	3.9	2.1	4.3	7.2	16.4	33.2	60.1	139.7	399.5	1,512.9
2000.....	74.3	24.0	3.4	2.1	4.5	7.0	16.2	32.0	59.0	133.9	390.1	1,481.3
1999.....	71.5	24.2	4.0	2.2	4.5	7.0	15.5	30.1	56.9	133.5	375.1	1,443.8

Non-Hispanic white, male

2007 5/.....	78.4	26.6	3.5	2.4	5.4	8.9	18.5	44.3	78.5	160.3	485.3	1,512.0
2006 5/.....	78.2	26.8	3.3	2.0	5.9	9.9	20.8	46.9	80.0	163.8	482.2	1,512.3
2005 5/.....	72.3	27.7	3.8	2.2	5.9	8.9	22.0	47.5	78.1	154.6	442.1	1,354.9
2004 5/.....	67.4	29.6	3.7	2.3	5.4	9.0	21.1	45.6	74.3	150.0	415.3	1,273.4
2003 5/.....	66.9	30.4	3.6	2.5	6.0	9.0	21.5	44.3	72.6	153.3	416.3	1,300.3
2002.....	64.8	31.4	4.0	2.2	5.2	8.9	21.8	43.5	71.1	148.8	408.2	1,293.4
2001.....	61.9	25.3	4.4	2.2	5.3	8.2	19.8	40.0	66.9	150.8	399.4	1,293.1
2000.....	59.6	26.7	3.8	2.3	5.4	8.5	19.4	39.2	66.0	144.1	390.9	1,288.3
1999.....	57.6	27.4	4.5	2.2	5.4	8.4	19.4	36.2	64.5	144.2	382.8	1,239.8

Non-Hispanic white, female

2007 5/.....	115.1	20.9	2.8	1.7	3.6	6.3	13.2	28.7	59.1	140.1	481.3	1,894.7
2006 5/.....	114.9	19.5	2.7	1.7	3.7	6.5	14.4	30.2	60.1	141.3	482.5	1,920.7
2005 5/.....	105.0	18.4	2.8	1.7	3.7	6.1	13.7	29.6	57.7	138.4	444.5	1,738.7
2004 5/.....	97.5	19.8	3.3	1.8	3.6	6.1	13.8	28.4	55.8	133.1	419.4	1,618.7
2003 5/.....	98.4	18.6	3.5	1.7	3.6	6.5	14.0	28.3	56.3	136.2	419.1	1,672.7
2002.....	95.5	17.7	3.6	1.8	3.4	5.8	13.9	27.2	54.8	130.8	412.1	1,647.2
2001.....	92.1	21.0	3.4	2.1	3.3	6.2	13.1	26.4	53.7	130.2	399.5	1,604.4
2000.....	88.3	21.2	2.9	1.8	3.5	5.5	12.9	25.0	52.5	125.2	389.6	1,559.1
1999.....	84.9	20.9	3.5	2.2	3.5	5.6	11.7	24.1	49.8	124.5	370.2	1,525.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic black, both sexes

2007 5/.....	68.5	61.9	6.0	3.0	6.1	14.1	30.0	63.0	112.4	209.1	548.7	1,656.9
2006 5/.....	68.7	53.8	4.6	2.6	6.7	14.1	31.1	68.4	116.8	209.8	546.6	1,660.0
2005 5/.....	65.9	51.0	4.9	3.4	6.7	13.8	31.8	69.9	113.4	204.5	523.9	1,525.4
2004 5/.....	63.1	50.5	5.4	2.8	7.0	16.0	31.9	69.4	113.2	204.2	489.7	1,425.6
2003 5/.....	63.0	52.6	5.5	2.8	6.7	15.7	34.0	71.3	113.6	200.3	487.6	1,451.6
2002.....	61.7	46.1	6.1	2.7	7.0	15.0	34.7	69.9	110.2	197.3	482.8	1,443.4
2001.....	61.1	53.8	5.6	2.5	6.7	14.7	34.8	71.7	109.1	198.5	476.3	1,431.5
2000.....	60.1	51.3	6.1	2.9	6.6	15.6	35.1	70.6	109.6	195.0	472.0	1,403.3
1999.....	58.0	58.4	5.5	3.1	6.9	13.5	36.2	69.3	104.9	202.1	444.8	1,295.3

Non-Hispanic black, male

2007 5/.....	61.4	71.1	6.1	3.7	6.4	14.4	32.0	73.6	135.9	247.9	601.3	1,389.1
2006 5/.....	61.5	61.0	5.0	3.0	6.8	13.4	33.3	78.7	140.9	243.3	603.7	1,437.6
2005 5/.....	58.9	63.8	5.8	3.7	7.0	13.9	33.4	80.6	135.0	232.6	573.0	1,261.7
2004 5/.....	58.1	56.9	5.8	2.7	7.1	16.8	33.1	81.5	135.3	238.7	548.8	1,316.9
2003 5/.....	57.6	56.8	6.3	3.0	6.0	15.6	36.7	84.1	134.7	237.5	552.6	1,243.4
2002.....	56.6	50.6	6.6	2.9	6.9	15.8	36.2	83.4	131.4	232.9	526.1	1,331.3
2001.....	56.4	61.3	5.2	2.5	6.8	15.6	39.3	85.6	128.3	232.9	530.2	1,329.4
2000.....	56.1	61.4	6.4	2.6	7.1	16.3	39.6	82.4	133.0	231.8	527.9	1,340.9
1999.....	54.4	66.5	5.6	3.6	7.3	13.9	40.3	81.9	122.9	228.2	509.7	1,303.8

Non-Hispanic black, female

2007 5/.....	75.0	52.3	5.9	2.3	5.9	13.9	28.1	53.9	93.4	181.4	518.8	1,767.9
2006 5/.....	75.2	46.3	4.1	2.2	6.6	14.7	29.2	59.6	97.4	186.0	514.2	1,750.7
2005 5/.....	72.3	37.6	4.0	3.1	6.3	13.8	30.4	60.7	96.1	184.5	496.1	1,631.5
2004 5/.....	67.6	43.8	5.0	3.0	6.9	15.2	30.8	59.0	95.4	179.6	456.3	1,468.5
2003 5/.....	67.9	48.4	4.6	2.6	7.3	15.7	31.7	60.3	96.6	173.9	450.9	1,532.3
2002.....	66.3	41.3	5.6	2.4	7.2	14.3	33.3	58.3	93.2	172.0	458.3	1,486.2
2001.....	65.2	46.0	6.0	2.6	6.6	13.9	30.8	59.7	93.6	174.3	446.1	1,469.3
2000.....	63.7	40.9	5.8	3.2	6.2	15.0	31.0	60.5	90.8	169.1	440.6	1,426.5
1999.....	61.3	50.0	5.4	2.6	6.5	13.1	32.7	58.5	90.5	183.7	408.5	1,292.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes

2007.....	41.0	30.2	9.6	5.5	37.4	36.9	39.2	46.3	37.3	45.2	105.5	286.7
2006.....	40.6	27.8	9.9	5.6	38.2	37.0	40.2	45.5	36.2	44.5	105.1	274.9
2005.....	39.7	26.4	10.3	6.0	37.4	34.9	38.6	43.2	35.8	46.3	106.1	279.5
2004.....	38.1	25.8	10.3	6.5	37.0	32.6	37.3	40.7	33.2	44.0	103.7	276.7
2003.....	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002.....	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001.....	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4

All origins 3/, male

2007.....	53.7	32.8	11.7	6.6	54.5	55.0	55.6	64.9	53.0	62.9	136.8	349.0
2006.....	53.5	31.6	11.1	6.7	56.0	55.8	56.9	65.0	51.6	60.1	135.8	337.1
2005.....	52.3	27.5	12.4	7.2	54.6	52.2	54.8	62.2	51.3	62.6	139.0	344.2
2004.....	49.8	29.0	11.8	8.0	53.3	48.5	53.0	58.6	46.9	59.2	134.7	345.5
2003.....	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002.....	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001.....	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000.....	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999.....	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4

All origins 3/, female

2007.....	28.7	27.4	7.5	4.3	19.3	18.0	22.8	28.3	22.6	30.2	83.9	257.0
2006.....	28.1	23.8	8.6	4.4	19.4	17.6	23.4	26.6	22.0	31.3	84.1	245.8
2005.....	27.5	25.2	8.0	4.7	19.2	16.9	22.3	24.7	21.3	32.5	83.8	249.8
2004.....	26.8	22.5	8.7	5.1	19.9	16.0	21.7	23.4	20.5	31.1	82.9	245.8
2003.....	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002.....	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001.....	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000.....	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999.....	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, both sexes

2007.....	25.8	16.2	9.0	4.7	33.8	28.4	28.6	34.9	29.5	36.8	76.8	152.0
2006.....	27.2	19.5	9.7	5.1	36.1	29.7	31.6	35.6	33.2	37.6	74.9	142.8
2005.....	26.9	18.5	10.6	4.7	37.1	29.4	30.0	34.1	30.3	39.4	79.7	153.6
2004.....	25.2	17.5	9.0	5.1	33.4	27.4	29.6	31.0	30.5	37.2	74.2	160.8
2003.....	26.1	15.1	9.9	5.8	33.8	29.4	31.7	32.8	30.6	38.1	70.9	151.8
2002.....	26.1	15.2	10.9	5.4	35.5	27.6	31.7	32.0	28.7	39.5	77.4	153.0
2001.....	25.8	18.8	10.3	5.4	32.6	28.2	32.3	31.7	30.5	40.0	73.8	166.3
2000.....	25.0	15.3	10.4	5.6	31.8	27.3	31.5	30.4	28.8	38.8	72.1	184.5
1999.....	25.5	17.7	11.9	5.6	30.8	28.4	32.6	31.2	28.5	42.5	69.6	182.4

Hispanic, male

2007.....	37.6	17.4	11.3	5.8	52.1	43.1	42.9	54.5	42.9	54.7	101.6	186.0
2006.....	39.7	20.8	11.3	6.3	55.5	44.9	48.0	54.0	51.2	54.2	95.8	174.8
2005.....	39.0	18.3	13.3	5.6	55.8	44.9	44.8	52.9	46.6	58.4	106.2	168.1
2004.....	36.5	18.8	10.5	6.2	50.8	41.6	44.9	47.9	44.8	52.0	99.4	193.3
2003.....	38.3	17.4	11.9	7.5	50.7	45.5	48.0	50.6	45.3	53.3	102.4	195.5
2002.....	38.5	19.5	13.5	6.6	53.5	43.0	48.5	48.7	43.5	58.7	105.9	224.2
2001.....	37.6	18.0	12.7	6.2	49.2	43.8	50.1	48.8	45.0	54.9	105.1	198.1
2000.....	36.9	17.0	11.6	6.1	48.2	42.9	49.6	48.4	43.2	57.2	93.9	251.9
1999.....	37.7	20.7	13.5	6.8	46.2	44.1	51.3	50.2	43.5	63.3	94.9	258.8

Hispanic, female

2007.....	13.1	14.9	6.5	3.6	13.5	10.4	12.5	14.7	17.1	22.2	59.5	133.5
2006.....	13.8	18.1	8.0	3.9	14.2	11.0	13.1	16.7	16.7	24.1	60.2	125.8
2005.....	13.8	18.6	7.7	3.9	15.9	10.5	13.5	14.8	15.3	23.9	61.2	146.0
2004.....	13.1	16.2	7.5	4.0	13.5	10.2	12.6	14.0	17.5	25.4	56.8	144.0
2003.....	13.1	12.7	7.8	3.9	14.3	10.1	13.8	15.1	17.3	26.0	49.2	129.7
2002.....	12.8	10.8	8.2	4.1	14.5	9.5	13.4	15.3	15.3	24.2	57.9	117.7
2001.....	13.2	19.7	7.8	4.5	13.3	10.1	13.0	14.9	17.7	28.2	52.6	150.8
2000.....	12.4	13.5	9.2	5.2	12.5	9.2	12.2	12.8	16.0	24.4	57.4	151.3
1999.....	12.6	14.6	10.3	4.2	12.8	10.1	12.6	12.8	15.2	26.2	52.7	145.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 4/, both sexes

2007.....	43.6	34.6	9.8	5.6	38.1	38.9	41.1	47.5	37.8	45.8	107.1	293.6
2006.....	42.8	30.2	9.8	5.7	38.6	38.7	41.5	46.4	36.4	44.9	106.7	281.3
2005.....	41.8	28.5	10.2	6.3	37.4	36.0	39.9	44.0	36.1	46.7	107.5	285.1
2004.....	40.1	28.1	10.6	6.8	37.7	33.6	38.5	41.5	33.3	44.3	105.2	281.5
2003.....	39.2	25.9	11.1	6.5	37.6	31.8	38.5	39.2	32.9	44.3	103.2	283.7
2002.....	38.5	25.5	10.4	6.9	38.3	32.2	37.8	36.8	31.4	44.4	102.1	279.8
2001.....	36.9	25.4	11.3	7.2	36.6	30.1	35.6	34.0	30.0	42.8	101.9	279.8
2000.....	36.0	25.0	12.2	7.5	36.7	29.8	34.2	32.5	30.9	41.9	95.8	276.1
1999.....	36.2	23.4	12.4	7.9	36.0	29.6	33.7	31.6	30.5	44.5	101.3	285.0

Non-Hispanic 4/, male

2007.....	56.5	37.5	11.8	6.7	54.9	58.2	58.0	65.9	53.6	63.4	138.8	358.0
2006.....	55.8	34.7	10.9	6.8	55.9	58.6	58.3	66.0	51.3	60.4	137.8	345.9
2005.....	54.5	30.0	12.1	7.6	54.3	53.9	56.4	62.9	51.5	62.7	140.7	352.9
2004.....	51.9	31.8	12.1	8.4	53.7	50.1	54.2	59.4	46.8	59.5	136.4	352.5
2003.....	50.9	28.7	13.2	8.0	53.7	47.3	54.9	56.0	47.0	59.0	134.1	358.9
2002.....	50.3	28.6	12.7	8.4	55.0	48.6	54.2	53.4	44.9	60.3	132.3	351.2
2001.....	48.5	28.0	13.4	9.0	53.5	45.6	51.7	50.4	42.5	57.1	134.3	364.5
2000.....	47.3	27.7	14.5	9.4	52.7	45.2	50.8	48.6	43.8	55.5	125.2	363.1
1999.....	47.4	26.6	14.5	9.9	51.4	44.8	49.7	47.3	43.5	59.7	133.2	378.7

Non-Hispanic 4/, female

2007.....	31.2	31.5	7.7	4.5	20.5	19.7	24.5	29.8	23.0	30.8	85.3	263.2
2006.....	30.4	25.5	8.7	4.5	20.4	19.0	25.0	27.5	22.4	31.8	85.4	251.3
2005.....	29.6	26.9	8.1	4.9	19.8	18.2	23.6	25.7	21.7	33.1	85.0	254.1
2004.....	28.8	24.2	9.0	5.3	21.0	17.2	23.1	24.3	20.7	31.5	84.2	249.8
2003.....	28.1	22.9	8.9	5.0	20.9	16.3	22.4	23.0	19.8	32.0	82.6	250.6
2002.....	27.3	22.3	8.1	5.2	21.1	15.9	21.8	20.9	18.8	31.0	82.2	249.0
2001.....	26.0	22.6	9.2	5.3	19.3	14.7	19.9	18.3	18.5	30.8	80.6	244.6
2000.....	25.2	22.2	9.7	5.6	20.3	14.5	18.1	17.0	18.9	30.6	76.7	240.9
1999.....	25.5	20.0	10.2	5.9	20.2	14.6	18.0	16.5	18.5	32.0	80.6	247.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic white, both sexes

2007 5/.....	47.0	27.6	9.2	5.3	41.9	42.8	43.7	48.4	37.3	45.8	111.2	309.9
2006 5/.....	45.7	24.3	8.9	5.3	42.5	41.8	43.6	46.5	35.4	45.1	110.2	295.9
2005 5/.....	44.4	24.5	9.8	5.9	40.7	38.6	41.7	43.9	35.2	46.5	109.4	297.2
2004 5/.....	42.8	24.1	10.2	6.4	41.5	35.7	40.3	41.5	32.5	44.3	107.9	295.2
2003 5/.....	41.6	21.6	10.7	6.2	41.1	33.5	39.8	38.7	32.2	43.9	105.5	296.5
2002.....	40.6	19.8	9.4	6.3	41.6	33.7	38.7	36.3	30.6	43.9	104.4	290.9
2001.....	38.5	20.9	10.9	6.6	39.0	31.1	35.9	32.8	29.0	42.4	103.7	289.3
2000.....	37.3	20.1	11.1	7.2	39.2	30.0	34.2	31.0	30.0	41.0	97.2	285.6
1999.....	37.2	19.0	11.1	7.4	38.1	29.9	33.5	30.0	28.9	43.3	101.7	293.3

Non-Hispanic white, male

2007 5/.....	60.0	30.7	11.1	6.3	60.2	63.3	60.7	66.2	52.2	62.6	142.7	376.3
2006 5/.....	58.6	27.6	9.8	6.2	61.3	62.5	60.4	64.8	49.1	59.9	141.2	362.3
2005 5/.....	57.0	25.6	11.8	7.1	58.9	57.1	58.4	61.9	49.1	61.6	141.9	366.3
2004 5/.....	54.6	28.3	11.8	7.9	58.9	52.5	56.2	58.7	44.8	59.0	138.8	366.8
2003 5/.....	53.1	24.1	12.9	7.5	58.2	49.5	56.3	54.7	45.4	58.2	135.7	372.9
2002.....	52.0	23.3	11.1	7.7	59.2	50.3	55.0	52.1	43.0	58.8	133.5	361.6
2001.....	49.7	22.9	13.2	8.1	56.6	46.7	51.8	48.0	40.3	56.1	135.2	374.7
2000.....	48.1	22.9	13.4	8.9	55.9	45.2	50.4	45.8	41.6	53.6	125.2	373.8
1999.....	47.8	22.0	12.9	9.1	53.6	44.9	49.0	44.2	40.7	57.3	132.7	387.1

Non-Hispanic white, female

2007 5/.....	34.5	24.4	7.1	4.2	22.6	22.0	26.6	31.0	23.2	31.2	89.1	278.4
2006 5/.....	33.3	20.8	8.0	4.3	22.7	20.9	26.7	28.3	22.4	32.2	88.7	264.9
2005 5/.....	32.3	23.4	7.6	4.6	21.6	19.8	25.0	26.1	22.0	33.4	87.1	265.6
2004 5/.....	31.4	19.7	8.5	4.9	23.2	18.6	24.2	24.6	20.9	31.6	86.8	263.1
2003 5/.....	30.5	19.0	8.4	4.8	23.2	17.2	23.2	22.9	19.6	31.7	85.1	262.8
2002.....	29.6	16.0	7.5	4.8	23.2	16.8	22.4	20.6	18.8	31.2	84.9	260.3
2001.....	27.8	18.7	8.5	5.0	20.7	15.2	19.9	17.9	18.3	30.7	82.8	253.8
2000.....	26.9	17.1	8.7	5.3	21.9	14.6	18.0	16.3	19.1	30.4	78.8	250.0
1999.....	27.0	15.9	9.3	5.6	22.0	14.7	17.9	15.9	17.7	31.5	81.5	255.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic black, both sexes

2007 5/.....	34.9	64.0	12.6	7.3	26.4	32.3	39.2	53.3	48.5	50.4	75.4	156.2
2006 5/.....	36.2	54.8	14.2	7.4	26.8	35.6	42.4	56.2	49.5	48.7	82.0	153.2
2005 5/.....	35.9	48.6	12.6	7.9	27.7	33.1	39.6	54.0	48.5	53.6	97.8	176.7
2004 5/.....	33.7	47.0	12.8	8.8	26.3	32.5	37.8	50.7	44.5	48.5	88.2	155.1
2003 5/.....	33.3	44.7	13.7	8.0	26.4	30.9	39.7	49.3	42.9	51.1	83.1	161.8
2002.....	34.0	50.1	14.6	9.4	28.4	32.5	39.9	47.6	41.9	49.7	84.6	175.4
2001.....	34.4	44.5	14.2	9.7	29.6	31.1	40.9	48.5	42.4	51.3	89.6	183.9
2000.....	34.4	46.5	16.8	9.6	29.6	33.5	40.6	47.8	41.3	53.0	85.3	183.2
1999.....	36.1	44.1	18.2	10.9	30.5	33.2	40.7	48.3	45.1	58.6	103.8	208.9

Non-Hispanic black, male

2007 5/.....	50.0	65.2	14.5	8.8	38.9	51.4	60.3	81.3	78.3	79.8	107.8	200.7
2006 5/.....	52.5	64.1	16.3	9.1	40.2	57.6	64.1	88.1	79.8	73.9	117.3	202.5
2005 5/.....	51.3	52.3	14.1	9.8	41.2	52.4	59.3	83.8	80.9	82.3	145.2	227.4
2004 5/.....	47.9	49.2	14.4	10.6	38.5	52.0	55.6	77.9	71.9	74.1	133.2	218.5
2003 5/.....	47.4	49.0	15.9	10.1	39.8	48.4	59.4	75.8	69.0	74.9	123.8	228.0
2002.....	49.0	52.1	18.9	11.9	42.6	52.2	59.7	73.3	69.2	77.2	131.3	262.5
2001.....	49.5	50.4	15.8	13.1	44.5	48.8	61.3	78.6	68.2	74.9	138.3	256.4
2000.....	50.3	49.5	19.4	12.3	44.2	53.5	63.3	76.7	69.4	80.4	133.3	277.1
1999.....	52.3	48.5	21.8	14.3	46.4	53.0	63.0	79.4	71.8	89.6	151.0	318.5

Non-Hispanic black, female

2007 5/.....	21.2	62.8	10.6	5.8	13.6	14.5	20.5	29.3	24.6	29.3	56.9	137.8
2006 5/.....	21.5	45.0	12.0	5.6	13.1	15.2	23.2	28.9	25.1	30.7	62.0	133.0
2005 5/.....	21.8	44.8	11.1	6.0	14.0	15.3	22.1	28.5	22.5	33.2	70.9	156.3
2004 5/.....	20.8	44.8	11.1	6.8	13.9	14.7	21.9	27.5	22.4	30.3	62.8	130.1
2003 5/.....	20.5	40.3	11.4	5.9	12.9	14.9	22.4	26.6	22.0	34.1	60.1	136.2
2002.....	20.3	48.0	10.1	6.8	14.2	14.7	22.5	25.6	20.0	30.2	58.2	142.1
2001.....	20.6	38.3	12.4	6.2	14.8	15.1	22.7	22.7	21.6	34.6	62.2	157.0
2000.....	20.1	43.3	14.2	6.7	15.1	15.5	20.5	23.0	18.7	33.7	58.4	148.3
1999.....	21.4	39.6	14.5	7.4	14.8	15.4	21.1	21.8	23.7	36.8	77.4	167.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes

2007.....	15.5	3.0	3.5	3.4	25.7	18.4	15.8	16.4	14.8	16.5	22.9	24.1
2006.....	16.2	3.4	3.8	3.6	26.8	19.2	16.6	16.8	15.8	16.8	23.6	24.4
2005.....	16.3	3.6	4.0	3.8	26.8	19.0	16.6	16.7	16.2	18.0	24.3	26.1
2004.....	16.4	3.6	4.1	4.3	27.3	18.8	16.4	16.7	15.5	17.5	25.5	27.2
2003.....	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002.....	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001.....	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000.....	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999.....	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0

All origins 3/, male

2007.....	22.5	2.7	3.9	4.0	36.3	27.6	23.5	24.8	21.9	24.1	33.8	41.7
2006.....	23.1	3.3	4.0	4.1	37.9	28.9	23.9	25.1	22.9	23.4	33.2	42.9
2005.....	23.3	3.5	4.5	4.4	37.9	28.6	24.2	24.9	23.5	25.3	34.8	46.0
2004.....	23.0	3.9	4.4	5.0	37.8	27.7	24.0	24.7	22.2	23.7	36.5	46.5
2003.....	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002.....	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001.....	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000.....	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999.....	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5

All origins 3/, female

2007.....	8.8	3.3	3.1	2.9	14.5	8.8	8.1	8.3	8.1	10.1	15.4	15.7
2006.....	9.4	3.6	3.6	3.1	15.1	9.1	9.2	8.7	9.2	11.2	17.1	15.7
2005.....	9.5	3.6	3.5	3.2	15.1	9.1	9.1	8.8	9.4	11.8	17.1	17.0
2004.....	9.9	3.2	3.8	3.6	16.1	9.5	8.8	9.0	9.3	12.3	18.1	18.5
2003.....	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002.....	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001.....	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000.....	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999.....	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transport accidents (V01-V99,Y85)

Hispanic, both sexes

2007.....	13.4	3.0	4.3	3.3	24.9	17.1	13.5	14.5	12.9	15.7	21.0	17.5
2006.....	14.8	4.6	4.7	3.6	27.8	17.8	15.3	15.7	15.4	17.4	20.7	11.0
2005.....	14.9	4.7	5.3	3.6	28.4	18.3	14.9	14.9	14.3	16.3	24.1	15.4
2004.....	14.5	4.4	4.7	3.9	26.3	18.1	14.9	14.3	15.9	17.0	21.9	20.0
2003.....	15.2	4.5	4.2	4.1	26.2	19.1	16.4	15.1	15.8	18.6	23.6	24.9
2002.....	15.3	3.1	5.8	3.8	27.6	18.3	16.0	15.2	15.2	18.4	26.2	25.5
2001.....	15.6	5.3	5.3	3.8	26.0	18.8	17.3	16.4	17.9	20.2	30.1	23.8
2000.....	14.8	5.4	5.2	4.1	24.6	17.3	16.2	16.2	15.7	20.4	23.9	25.2
1999.....	14.4	5.3	5.5	3.8	23.1	17.8	15.4	15.0	15.5	22.0	25.0	25.3

Hispanic, male

2007.....	19.7	*	4.8	4.0	37.3	25.3	19.1	22.1	18.6	24.6	30.5	29.5
2006.....	21.7	4.6	5.3	4.3	42.0	26.5	22.4	23.0	22.7	24.2	29.4	*
2005.....	21.7	4.4	6.5	4.2	41.5	27.4	21.2	22.2	20.9	25.7	35.4	*
2004.....	21.3	*	5.4	4.6	39.3	27.0	22.2	21.6	21.6	23.9	34.7	28.0
2003.....	22.2	5.7	4.6	5.5	38.2	29.3	23.9	22.4	21.4	26.2	35.3	46.2
2002.....	22.4	*	7.1	4.6	40.4	27.5	23.9	21.6	22.1	28.5	37.3	45.2
2001.....	22.8	*	6.0	4.3	38.4	28.8	25.9	23.2	24.6	28.2	51.0	43.8
2000.....	21.6	6.1	5.6	4.5	36.1	26.4	24.1	24.1	23.5	29.5	36.4	44.3
1999.....	20.8	5.7	5.8	4.5	33.3	26.7	23.2	22.6	21.0	31.6	40.1	*

Hispanic, female

2007.....	6.7	*	3.7	2.5	11.1	6.9	7.1	6.7	7.6	8.5	14.2	11.0
2006.....	7.4	4.6	4.0	2.9	11.9	7.1	7.4	8.1	8.8	11.9	14.6	*
2005.....	7.6	5.0	4.1	3.1	13.6	7.2	7.7	7.5	8.3	8.6	16.3	*
2004.....	7.3	5.0	4.1	3.2	11.3	7.5	6.7	6.9	10.7	11.5	13.1	15.8
2003.....	7.6	*	3.8	2.7	12.3	7.0	8.3	7.9	10.7	12.5	15.5	*
2002.....	7.7	*	4.5	2.9	12.6	7.3	7.5	8.7	8.9	10.3	18.5	*
2001.....	8.0	6.7	4.6	3.2	11.6	7.1	8.0	9.7	12.0	13.9	16.0	*
2000.....	7.6	*	4.8	3.7	11.0	6.8	7.8	8.5	8.7	13.3	15.5	*
1999.....	7.6	*	5.2	3.1	11.0	7.6	7.0	7.7	10.6	14.5	14.9	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transport accidents (V01-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	15.9	3.0	3.3	3.4	25.9	18.6	16.2	16.6	14.9	16.5	23.0	24.4
2006.....	16.4	3.1	3.5	3.6	26.6	19.5	16.8	16.8	15.8	16.7	23.7	25.0
2005.....	16.5	3.2	3.6	3.9	26.4	19.1	16.9	16.9	16.3	18.1	24.2	26.5
2004.....	16.6	3.3	3.9	4.4	27.4	18.9	16.6	16.9	15.4	17.5	25.6	27.5
2003.....	16.7	3.5	4.0	4.3	28.0	18.0	17.2	16.6	15.8	17.3	26.3	29.8
2002.....	16.9	3.2	3.7	4.3	29.3	18.8	17.2	16.4	15.6	18.3	26.9	28.9
2001.....	16.7	3.4	4.2	4.6	28.2	18.8	17.3	15.8	15.0	17.7	27.7	29.0
2000.....	16.8	4.3	4.3	4.8	28.8	18.7	17.0	15.7	15.9	17.8	27.1	31.7
1999.....	16.9	4.9	4.4	5.0	28.4	18.5	16.9	15.4	16.0	19.2	28.3	32.2

Non-Hispanic 4/, male

2007.....	22.9	2.6	3.6	3.9	36.1	28.1	24.3	25.0	22.1	24.0	33.9	42.4
2006.....	23.3	2.9	3.5	4.1	36.9	29.5	24.1	25.3	22.8	23.2	33.3	44.2
2005.....	23.6	3.3	3.9	4.4	37.1	28.8	24.6	25.1	23.6	25.2	34.6	47.1
2004.....	23.3	3.9	4.1	5.0	37.5	27.8	24.3	24.9	22.2	23.6	36.5	47.3
2003.....	23.4	3.5	4.5	5.1	38.5	26.7	24.9	24.4	23.0	23.3	36.6	52.0
2002.....	23.8	3.4	4.2	5.0	40.6	28.1	24.6	24.1	22.4	24.9	37.5	53.2
2001.....	23.5	3.2	4.7	5.5	39.7	28.1	24.8	23.2	21.3	24.0	39.2	52.9
2000.....	23.4	4.5	4.8	5.6	39.8	27.8	24.7	22.8	22.2	23.9	38.0	60.4
1999.....	23.3	5.1	4.9	6.0	39.0	27.2	24.0	22.2	22.4	25.6	40.4	64.3

Non-Hispanic 4/, female

2007.....	9.1	3.4	3.0	3.0	15.1	9.1	8.2	8.5	8.2	10.2	15.4	15.9
2006.....	9.8	3.2	3.5	3.1	15.7	9.5	9.5	8.8	9.2	11.1	17.2	16.1
2005.....	9.8	3.2	3.3	3.3	15.3	9.5	9.3	8.9	9.5	12.1	17.2	17.1
2004.....	10.3	2.6	3.8	3.7	17.0	9.9	9.1	9.2	9.1	12.3	18.4	18.6
2003.....	10.3	3.5	3.6	3.6	17.0	9.4	9.6	9.2	9.2	12.3	19.5	20.1
2002.....	10.4	2.9	3.1	3.6	17.7	9.5	10.0	9.0	9.3	12.7	19.8	18.4
2001.....	10.2	3.6	3.6	3.6	16.3	9.5	10.0	8.8	9.1	12.5	20.2	19.1
2000.....	10.5	4.2	3.8	3.9	17.4	9.7	9.4	8.9	10.1	12.8	20.0	20.1
1999.....	10.7	4.7	3.9	4.0	17.5	10.0	9.9	8.8	10.0	13.9	20.5	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transport accidents (V01-V99,Y85)

Non-Hispanic white, both sexes

2007 5/.....	16.4	2.7	2.9	3.5	27.8	19.1	16.5	16.8	15.0	16.6	23.3	24.9
2006 5/.....	16.9	2.6	3.0	3.5	28.8	19.8	17.2	17.0	15.5	16.9	24.0	26.0
2005 5/.....	17.1	2.5	3.3	3.8	28.4	19.6	17.3	17.0	16.4	18.2	24.6	27.5
2004 5/.....	17.2	2.9	3.7	4.4	29.9	19.1	17.0	16.8	15.4	17.6	26.0	28.8
2003 5/.....	17.2	3.2	3.8	4.4	30.2	18.2	17.3	16.6	15.6	17.3	26.6	30.6
2002.....	17.5	2.5	3.3	4.3	31.8	19.0	17.3	16.4	15.5	18.3	27.4	29.7
2001.....	17.0	3.1	3.8	4.4	29.9	19.0	17.4	15.6	14.7	17.6	28.3	29.9
2000.....	17.1	3.5	3.8	4.8	30.9	18.5	17.0	15.4	15.8	17.6	27.6	32.4
1999.....	17.1	4.3	3.7	5.0	30.3	18.6	16.7	15.0	15.6	19.1	28.6	32.5

Non-Hispanic white, male

2007 5/.....	23.6	2.7	3.3	3.9	38.6	28.5	24.4	25.1	22.0	23.9	33.9	42.7
2006 5/.....	23.8	2.6	2.9	3.9	39.7	29.6	24.5	25.2	22.4	23.2	33.5	45.4
2005 5/.....	24.2	2.8	3.5	4.4	39.6	29.0	24.9	25.3	23.5	25.0	34.8	48.7
2004 5/.....	23.9	3.3	3.8	5.2	40.6	27.6	24.6	24.5	21.8	23.5	36.6	49.6
2003 5/.....	23.9	3.3	4.4	5.1	41.1	26.8	24.9	24.1	22.6	23.3	36.6	52.8
2002.....	24.3	2.9	3.6	5.0	43.6	28.1	24.6	24.0	22.0	24.5	37.5	54.0
2001.....	23.8	2.9	4.4	5.2	41.9	28.2	24.6	22.7	20.8	23.7	39.6	54.3
2000.....	23.7	3.8	4.3	5.7	42.5	27.1	24.5	22.2	21.6	23.2	38.1	62.3
1999.....	23.5	4.9	4.0	5.9	41.2	26.9	23.6	21.6	21.7	25.2	40.3	64.6

Non-Hispanic white, female

2007 5/.....	9.5	2.7	2.5	3.0	16.4	9.6	8.6	8.7	8.2	10.2	16.0	16.5
2006 5/.....	10.2	2.6	3.2	3.1	17.2	9.8	9.9	8.9	9.0	11.3	17.5	17.0
2005 5/.....	10.3	2.3	3.2	3.2	16.6	10.1	9.7	8.9	9.6	12.3	17.5	17.8
2004 5/.....	10.7	2.4	3.6	3.6	18.6	10.4	9.3	9.2	9.3	12.5	18.8	19.5
2003 5/.....	10.7	3.0	3.2	3.7	18.8	9.5	9.7	9.2	8.9	12.1	19.8	20.8
2002.....	10.9	2.1	2.9	3.5	19.4	9.8	10.1	8.9	9.4	13.0	20.7	19.2
2001.....	10.6	3.3	3.2	3.5	17.4	9.7	10.2	8.6	9.1	12.5	20.8	19.8
2000.....	10.9	3.2	3.4	4.0	18.9	9.7	9.5	8.6	10.4	12.8	20.7	20.4
1999.....	11.0	3.6	3.4	4.0	19.1	10.2	9.9	8.6	9.9	13.9	20.9	19.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic black, both sexes												
2007 5/.....	15.1	4.2	4.6	3.8	19.9	20.2	18.3	18.0	16.6	16.3	18.8	17.9
2006 5/.....	15.6	4.9	5.2	4.0	19.6	22.4	17.9	18.2	19.1	15.6	20.1	15.9
2005 5/.....	15.4	5.7	4.7	4.1	20.7	20.4	17.8	18.2	17.3	18.3	20.0	16.9
2004 5/.....	15.6	5.0	5.2	4.5	20.0	21.6	17.4	19.9	17.6	16.4	21.7	15.0
2003 5/.....	15.6	4.4	5.1	4.3	20.3	20.0	18.8	18.6	18.7	18.2	22.6	21.4
2002.....	15.7	5.6	5.0	4.7	21.6	21.1	18.5	18.0	17.3	17.3	21.8	18.8
2001.....	16.3	*	5.4	5.5	22.8	20.5	19.2	19.1	18.4	18.8	21.4	17.8
2000.....	16.4	6.9	5.9	4.8	22.4	22.2	19.1	19.7	17.5	19.9	19.9	21.9
1999.....	16.7	7.7	7.1	5.6	22.7	20.6	19.5	19.2	18.5	19.5	26.0	27.8
Non-Hispanic black, male												
2007 5/.....	23.4	*	4.8	4.4	29.0	33.0	30.3	29.5	27.2	26.9	34.9	39.3
2006 5/.....	23.6	*	5.8	4.9	28.4	36.4	28.3	29.3	30.0	25.6	31.7	37.6
2005 5/.....	23.4	*	5.1	4.8	30.2	33.5	28.5	28.4	28.0	30.2	34.2	34.3
2004 5/.....	23.4	6.2	5.4	4.9	28.6	35.1	27.3	31.5	28.2	26.4	38.5	29.5
2003 5/.....	23.3	*	5.1	5.3	30.3	31.8	29.1	29.7	29.6	26.5	38.1	43.7
2002.....	23.6	*	6.5	5.4	31.0	34.1	28.0	28.0	28.6	28.7	40.7	47.0
2001.....	24.5	*	5.8	7.1	33.4	32.6	29.7	30.7	30.2	27.4	36.9	35.8
2000.....	24.8	7.2	6.4	5.8	32.5	35.6	30.5	30.1	29.6	32.6	36.4	45.8
1999.....	24.9	*	8.0	7.0	32.9	33.0	29.8	30.7	29.3	29.5	43.5	68.9
Non-Hispanic black, female												
2007 5/.....	7.5	*	4.5	3.1	10.6	8.2	7.6	8.1	8.1	8.8	9.7	9.0
2006 5/.....	8.2	*	4.6	3.1	10.5	9.4	8.7	8.7	10.3	8.5	13.6	*
2005 5/.....	8.1	6.5	4.3	3.4	11.1	8.2	8.3	9.4	8.8	9.9	11.9	10.0
2004 5/.....	8.5	*	5.0	4.0	11.3	9.3	8.7	9.9	9.2	9.2	12.2	9.2
2003 5/.....	8.5	*	5.2	3.3	10.1	9.3	9.7	9.0	9.9	12.3	13.9	12.8
2002.....	8.5	*	3.6	3.9	12.1	9.3	10.1	9.4	8.2	9.1	11.1	*
2001.....	8.9	*	5.0	3.8	12.3	9.6	9.9	9.2	9.0	12.8	12.6	11.1
2000.....	8.8	*	5.4	3.7	12.5	10.1	8.9	10.8	7.8	11.0	10.6	13.1
1999.....	9.4	9.3	6.2	4.1	12.6	9.4	10.4	9.5	9.8	12.5	16.3	12.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All origins 3/, both sexes												
2007.....	14.6	2.9	3.3	3.2	24.9	17.5	14.8	14.9	13.3	15.2	21.8	23.2
2006.....	15.1	3.4	3.6	3.3	26.0	18.2	15.3	15.3	14.3	15.4	22.3	23.4
2005.....	15.3	3.6	3.8	3.6	25.9	18.0	15.4	15.1	14.7	16.7	22.9	25.1
2004.....	15.3	3.5	4.0	4.1	26.3	17.6	15.1	15.1	14.1	16.1	24.5	26.1
2003.....	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002.....	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001.....	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000.....	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999.....	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
All origins 3/, male												
2007.....	20.9	2.6	3.7	3.7	35.1	26.2	21.7	22.2	19.4	21.6	31.9	39.8
2006.....	21.4	3.3	3.8	3.7	36.6	27.4	21.8	22.6	20.5	21.1	30.8	41.0
2005.....	21.7	3.5	4.2	4.1	36.5	26.9	22.2	22.2	20.9	23.2	32.4	44.1
2004.....	21.3	3.8	4.2	4.7	36.4	25.8	21.9	21.9	19.8	21.3	34.7	44.2
2003.....	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002.....	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001.....	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000.....	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999.....	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
All origins 3/, female												
2007.....	8.4	3.2	3.0	2.7	14.1	8.4	7.7	7.8	7.6	9.7	14.9	15.2
2006.....	9.0	3.5	3.4	2.9	14.7	8.7	8.8	8.2	8.5	10.6	16.5	15.2
2005.....	9.1	3.6	3.4	3.0	14.7	8.8	8.6	8.2	8.9	11.2	16.5	16.4
2004.....	9.5	3.2	3.7	3.4	15.7	9.1	8.3	8.5	8.8	11.7	17.7	17.9
2003.....	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002.....	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001.....	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000.....	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999.....	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Hispanic, both sexes												
2007.....	12.8	2.9	4.1	3.2	24.3	16.5	12.7	13.3	11.7	14.6	19.6	15.4
2006.....	14.2	4.6	4.6	3.4	27.2	17.2	14.3	14.6	14.7	16.5	19.4	10.6
2005.....	14.3	4.7	5.2	3.5	27.7	17.7	14.1	13.8	13.5	15.5	22.5	15.0
2004.....	13.9	4.4	4.6	3.8	25.6	17.2	13.9	13.2	14.9	16.1	21.3	19.5
2003.....	14.4	4.5	4.1	3.9	25.5	18.1	15.4	14.2	14.6	17.4	23.1	24.4
2002.....	14.6	3.1	5.6	3.6	26.6	17.6	15.2	13.9	14.3	17.9	24.4	24.9
2001.....	14.3	4.9	4.7	3.5	25.1	17.4	15.3	13.8	15.1	17.7	26.3	20.0
2000.....	13.9	5.2	4.9	3.8	23.7	16.4	14.7	15.1	14.4	18.4	22.3	22.6
1999.....	13.3	5.3	5.2	3.6	21.9	16.4	13.9	13.7	14.5	19.7	23.5	23.2
Hispanic, male												
2007.....	18.7	*	4.6	3.9	36.3	24.3	17.8	19.8	16.4	22.1	28.1	27.4
2006.....	20.7	4.6	5.3	4.0	41.1	25.6	20.6	21.3	21.5	22.9	26.9	*
2005.....	20.7	4.4	6.3	4.0	40.3	26.3	20.2	20.4	19.4	24.6	32.1	*
2004.....	20.2	*	5.3	4.4	38.3	25.4	20.8	19.6	19.7	22.3	33.3	28.0
2003.....	21.1	5.7	4.3	5.2	37.2	27.6	22.1	20.8	19.8	24.7	34.1	46.2
2002.....	21.3	*	6.8	4.4	38.9	26.4	22.6	19.6	20.5	27.5	34.5	43.5
2001.....	21.0	*	5.3	4.0	36.9	26.8	23.1	20.2	21.2	24.6	44.3	*
2000.....	20.1	6.1	5.2	4.1	34.7	24.9	21.6	22.0	21.2	26.0	33.4	*
1999.....	19.1	5.7	5.6	4.1	31.6	24.4	20.8	20.1	19.0	27.2	37.0	*
Hispanic, female												
2007.....	6.5	*	3.6	2.4	10.9	6.8	6.8	6.5	7.3	8.5	13.6	*
2006.....	7.2	4.6	3.9	2.9	11.6	6.8	7.2	7.8	8.5	11.3	14.2	*
2005.....	7.4	5.0	4.1	2.9	13.4	7.2	7.4	7.0	8.2	8.2	15.8	*
2004.....	7.1	5.0	3.9	3.2	11.1	7.3	6.4	6.7	10.5	11.2	13.1	15.1
2003.....	7.3	*	3.8	2.7	11.9	6.6	8.0	7.5	9.8	11.6	15.5	*
2002.....	7.4	*	4.4	2.7	12.4	7.2	7.3	8.2	8.7	10.2	17.4	*
2001.....	7.2	6.2	4.1	3.0	11.2	6.5	6.9	7.6	9.7	12.3	14.2	*
2000.....	7.2	*	4.6	3.6	10.6	6.5	7.3	8.3	8.4	12.4	14.9	*
1999.....	7.2	*	4.7	3.0	10.6	7.1	6.5	7.4	10.6	13.8	14.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic 4/, both sexes												
2007.....	14.8	2.9	3.1	3.2	24.9	17.6	15.1	15.0	13.4	15.2	21.9	23.6
2006.....	15.3	3.0	3.3	3.3	25.6	18.4	15.4	15.3	14.2	15.3	22.4	24.1
2005.....	15.4	3.2	3.4	3.6	25.5	18.0	15.5	15.1	14.7	16.7	22.9	25.5
2004.....	15.5	3.2	3.8	4.1	26.4	17.6	15.2	15.2	14.0	16.1	24.6	26.3
2003.....	15.5	3.4	3.9	4.0	26.8	16.9	15.7	14.9	14.1	16.0	25.0	28.9
2002.....	15.9	3.0	3.5	4.0	28.5	17.7	15.8	14.8	14.1	16.9	25.7	28.1
2001.....	15.5	3.2	3.9	4.2	27.1	17.5	15.7	14.2	13.5	16.2	26.3	28.0
2000.....	15.6	4.2	4.1	4.4	27.5	17.4	15.4	14.1	14.4	16.3	25.8	30.6
1999.....	15.4	4.7	4.0	4.5	26.9	16.9	15.0	13.5	14.0	17.6	26.5	30.5
Non-Hispanic 4/, male												
2007.....	21.3	2.5	3.4	3.6	34.7	26.6	22.5	22.4	19.6	21.6	32.0	40.6
2006.....	21.5	2.9	3.3	3.6	35.5	27.8	21.9	22.7	20.3	20.9	30.9	42.3
2005.....	21.8	3.3	3.6	4.1	35.6	27.0	22.5	22.3	20.9	23.0	32.3	45.2
2004.....	21.5	3.8	3.9	4.7	36.0	25.8	22.0	22.1	19.7	21.2	34.6	45.0
2003.....	21.4	3.4	4.3	4.7	36.7	24.9	22.5	21.4	19.9	20.9	34.2	49.9
2002.....	21.9	3.2	4.0	4.6	39.2	26.4	22.2	21.3	19.7	22.3	35.1	51.9
2001.....	21.5	3.1	4.3	5.1	38.1	25.9	22.2	20.4	18.8	21.3	36.9	51.6
2000.....	21.4	4.2	4.5	5.1	37.9	25.5	22.0	20.2	19.6	21.3	35.7	58.3
1999.....	21.0	4.8	4.4	5.2	36.7	24.6	21.1	19.2	19.1	22.9	37.7	60.7
Non-Hispanic 4/, female												
2007.....	8.7	3.3	2.8	2.7	14.7	8.8	7.9	8.0	7.7	9.8	15.0	15.6
2006.....	9.3	3.1	3.3	2.9	15.3	9.1	9.0	8.2	8.5	10.5	16.6	15.6
2005.....	9.4	3.2	3.2	3.1	14.9	9.1	8.7	8.3	8.9	11.4	16.5	16.5
2004.....	9.8	2.6	3.7	3.4	16.5	9.4	8.6	8.6	8.6	11.7	17.9	18.0
2003.....	9.8	3.3	3.4	3.3	16.5	8.9	9.0	8.6	8.7	11.9	18.8	19.6
2002.....	10.1	2.9	3.0	3.4	17.4	9.1	9.5	8.6	8.9	12.4	19.4	17.8
2001.....	9.8	3.4	3.5	3.3	15.8	9.1	9.4	8.2	8.7	12.0	19.3	18.2
2000.....	10.0	4.2	3.6	3.6	16.9	9.2	8.9	8.2	9.6	12.2	19.3	19.4
1999.....	10.0	4.7	3.6	3.7	16.8	9.2	9.1	8.1	9.3	13.2	19.2	18.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic white, both sexes												
2007 5/.....	15.4	2.7	2.7	3.2	26.8	18.1	15.4	15.2	13.5	15.3	22.3	24.1
2006 5/.....	15.7	2.5	2.8	3.2	27.7	18.6	15.8	15.4	13.9	15.4	22.6	25.1
2005 5/.....	16.0	2.5	3.1	3.6	27.4	18.5	15.9	15.4	14.7	16.8	23.2	26.6
2004 5/.....	16.0	2.8	3.5	4.1	28.8	17.7	15.5	15.1	13.9	16.1	25.0	27.6
2003 5/.....	15.9	3.1	3.6	4.0	28.9	17.0	15.7	14.8	13.7	15.9	25.2	29.6
2002.....	16.3	2.4	3.0	4.0	30.8	17.8	15.8	14.8	14.0	16.9	26.2	28.9
2001.....	15.8	3.0	3.5	4.1	28.7	17.6	15.7	13.9	13.3	16.1	26.9	29.0
2000.....	15.8	3.3	3.7	4.4	29.5	17.0	15.3	13.7	14.2	16.1	26.2	31.3
1999.....	15.6	4.1	3.4	4.4	28.7	16.8	14.8	13.1	13.7	17.5	26.8	30.9
Non-Hispanic white, male												
2007 5/.....	21.8	2.7	3.0	3.6	37.2	26.8	22.5	22.5	19.5	21.5	32.0	40.8
2006 5/.....	22.0	2.6	2.7	3.4	38.1	27.8	22.2	22.6	19.8	20.8	30.9	43.6
2005 5/.....	22.4	2.8	3.2	4.1	38.2	27.1	22.7	22.5	20.7	22.7	32.5	46.7
2004 5/.....	22.0	3.3	3.6	4.8	39.1	25.5	22.2	21.7	19.3	21.0	34.8	47.1
2003 5/.....	21.8	3.2	4.2	4.7	39.1	24.9	22.4	21.1	19.4	20.9	34.3	50.7
2002.....	22.3	2.7	3.3	4.6	42.1	26.1	22.0	21.2	19.4	21.8	35.2	52.6
2001.....	21.7	2.9	3.9	4.8	40.1	25.9	21.9	19.9	18.2	21.0	37.3	53.1
2000.....	21.5	3.4	4.1	5.1	40.3	24.7	21.6	19.6	18.9	20.7	35.6	60.0
1999.....	21.1	4.7	3.6	5.1	38.8	24.2	20.6	18.5	18.5	22.6	37.6	61.2
Non-Hispanic white, female												
2007 5/.....	9.1	2.6	2.3	2.7	16.0	9.3	8.2	8.1	7.7	9.8	15.5	16.2
2006 5/.....	9.7	2.5	3.0	2.9	16.7	9.4	9.3	8.3	8.3	10.7	16.9	16.5
2005 5/.....	9.8	2.2	3.1	3.0	16.1	9.7	9.1	8.4	9.0	11.7	16.9	17.3
2004 5/.....	10.2	2.3	3.5	3.3	18.0	9.8	8.7	8.6	8.7	11.9	18.4	18.9
2003 5/.....	10.2	2.9	3.0	3.3	18.3	9.0	9.1	8.5	8.4	11.6	19.1	20.3
2002.....	10.5	2.1	2.8	3.3	19.0	9.4	9.7	8.4	8.8	12.7	20.2	18.6
2001.....	10.1	3.1	3.1	3.3	17.0	9.3	9.6	8.1	8.7	11.9	20.0	19.0
2000.....	10.3	3.2	3.2	3.6	18.4	9.3	9.0	7.9	9.8	12.3	20.0	19.8
1999.....	10.3	3.5	3.1	3.7	18.3	9.4	9.0	7.9	9.2	13.2	19.7	18.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic black, both sexes												
2007 5/.....	14.2	4.1	4.5	3.5	19.1	19.4	17.3	16.4	15.1	15.1	17.9	17.6
2006 5/.....	14.7	4.7	5.0	3.8	19.1	21.3	16.7	16.7	17.8	14.8	19.0	14.3
2005 5/.....	14.4	5.7	4.5	3.9	19.9	19.4	16.7	16.1	16.3	17.2	18.9	16.1
2004 5/.....	14.6	4.9	5.0	4.3	19.1	20.5	16.3	18.1	16.1	15.4	20.4	15.0
2003 5/.....	14.7	4.4	5.0	4.1	19.4	19.1	17.6	16.9	17.3	17.1	21.1	21.1
2002.....	14.8	5.4	4.9	4.3	21.0	20.3	17.2	16.3	15.7	16.0	20.2	18.5
2001.....	15.2	*	5.3	5.0	22.2	19.6	17.6	17.2	16.6	17.4	19.9	16.9
2000.....	15.5	6.9	5.6	4.5	21.6	21.1	17.6	18.3	16.3	18.4	18.8	20.7
1999.....	15.5	7.6	6.5	5.1	21.7	19.2	17.9	17.4	16.7	18.1	24.0	25.5
Non-Hispanic black, male												
2007 5/.....	21.9	*	4.6	4.1	27.8	31.7	28.5	26.5	24.3	24.2	33.0	39.3
2006 5/.....	22.2	*	5.7	4.6	27.7	34.5	26.2	26.4	27.6	24.0	30.3	33.9
2005 5/.....	21.8	*	4.9	4.5	28.7	32.0	26.6	24.9	26.1	27.9	32.2	33.3
2004 5/.....	21.7	*	5.2	4.7	27.0	33.0	25.2	28.5	25.3	24.7	36.2	29.5
2003 5/.....	21.7	*	5.0	4.9	28.8	30.2	27.1	26.5	26.7	24.1	35.5	42.6
2002.....	22.0	*	6.3	5.0	30.1	32.7	25.7	24.8	25.4	25.6	36.6	45.9
2001.....	22.7	*	5.6	6.5	32.4	31.0	26.9	27.5	26.8	24.2	34.8	35.8
2000.....	23.1	7.2	6.0	5.4	31.1	33.6	27.9	27.6	27.2	29.8	34.6	44.6
1999.....	22.8	*	7.0	6.2	31.5	30.7	26.9	27.5	25.8	26.8	41.3	61.8
Non-Hispanic black, female												
2007 5/.....	7.2	*	4.4	3.0	10.1	7.9	7.3	7.7	7.7	8.6	9.4	8.7
2006 5/.....	7.9	*	4.3	3.1	10.3	9.1	8.3	8.4	10.0	8.3	12.6	*
2005 5/.....	7.7	6.5	4.1	3.4	10.9	7.9	7.9	8.5	8.4	9.6	11.3	9.2
2004 5/.....	8.1	*	4.9	3.8	11.1	9.1	8.3	9.2	8.6	8.8	11.4	9.2
2003 5/.....	8.3	*	4.9	3.2	9.9	9.0	9.3	8.6	9.7	12.1	13.0	12.8
2002.....	8.3	*	3.5	3.6	11.9	9.1	9.7	9.0	8.0	9.1	10.9	*
2001.....	8.5	*	4.9	3.4	12.1	9.2	9.4	8.4	8.4	12.6	11.5	9.8
2000.....	8.5	*	5.2	3.6	12.3	9.8	8.5	10.2	7.5	10.4	9.9	11.8
1999.....	8.9	9.3	5.9	4.0	12.1	8.9	9.9	8.8	9.3	12.0	14.4	12.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
All origins 3/, both sexes												
2007.....	0.4	*	*	0.1	0.4	0.3	0.4	0.5	0.4	0.4	0.5	*
2006.....	0.4	*	*	0.1	0.4	0.3	0.5	0.6	0.5	0.5	0.6	0.5
2005.....	0.4	*	0.2	0.1	0.5	0.4	0.5	0.6	0.5	0.4	0.6	0.5
2004.....	0.4	*	*	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6
2003.....	0.5	*	*	0.1	0.6	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002.....	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001.....	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000.....	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999.....	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3
All origins 3/, male												
2007.....	0.6	*	*	0.1	0.6	0.6	0.7	0.9	0.6	0.8	0.8	*
2006.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.8	1.0	*
2005.....	0.7	*	*	0.2	0.7	0.7	0.7	1.0	0.8	0.6	1.1	*
2004.....	0.6	*	*	0.2	0.7	0.7	0.8	0.9	0.7	0.7	0.9	*
2003.....	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.8	1.0	*
2002.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001.....	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000.....	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999.....	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9
All origins 3/, female												
2007.....	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	*	0.3	*
2006.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	*
2005.....	0.2	*	*	*	0.2	0.1	0.2	0.3	0.2	0.3	0.3	*
2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002.....	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000.....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999.....	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic 4/, both sexes												
2007.....	0.4	*	*	0.1	0.4	0.3	0.4	0.5	0.3	0.4	0.4	*
2006.....	0.4	*	*	0.1	0.4	0.3	0.5	0.5	0.5	0.5	0.6	0.5
2005.....	0.4	*	0.2	0.1	0.5	0.4	0.5	0.6	0.5	0.4	0.6	0.5
2004.....	0.4	*	*	0.1	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.6
2003.....	0.5	*	*	0.1	0.7	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002.....	0.4	*	*	0.2	0.3	0.4	0.5	0.5	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.2	0.5	0.4	0.6	0.6	0.5	0.7	0.7	0.6
2000.....	0.5	*	*	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999.....	0.7	*	0.3	0.3	0.9	0.7	0.9	0.8	0.8	0.7	1.3	1.3
Non-Hispanic 4/, male												
2007.....	0.6	*	*	0.1	0.6	0.6	0.7	0.9	0.6	0.7	0.7	*
2006.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.9	1.0	*
2005.....	0.7	*	*	0.2	0.7	0.7	0.7	1.0	0.7	0.6	1.0	*
2004.....	0.6	*	*	0.2	0.7	0.7	0.7	0.8	0.7	0.7	0.9	*
2003.....	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.7	1.0	*
2002.....	0.6	*	*	0.2	0.6	0.6	0.8	0.9	0.7	0.7	0.9	*
2001.....	0.7	*	*	0.3	0.7	0.7	1.0	0.9	0.8	1.0	1.0	*
2000.....	0.8	*	*	0.3	1.0	0.8	1.1	0.8	0.8	0.8	1.1	*
1999.....	1.1	*	0.4	0.4	1.4	1.1	1.4	1.2	1.2	1.0	1.8	3.0
Non-Hispanic 4/, female												
2007.....	0.2	*	*	0.1	0.2	*	0.2	0.2	0.1	*	*	*
2006.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.3	*
2005.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.2	0.3	0.3	*
2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.3	0.2	0.2	*	0.4	*
2002.....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.5	*
2000.....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.4	*
1999.....	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5	1.0	*





Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins 3/, both sexes

2007.....	0.6	*	*	0.1	0.4	0.6	0.6	1.0	1.1	0.9	0.6	0.6
2006.....	0.6	*	*	0.2	0.5	0.7	0.8	0.9	1.0	0.8	0.8	0.4
2005.....	0.6	*	*	0.1	0.4	0.6	0.8	1.0	1.1	0.9	0.7	0.5
2004.....	0.7	*	*	0.1	0.5	0.8	0.8	1.1	1.0	0.9	0.5	0.5
2003.....	0.7	*	*	0.2	0.5	0.6	0.9	1.1	1.2	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001.....	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000.....	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999.....	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*

All origins 3/, male

2007.....	1.0	*	*	0.1	0.7	0.9	1.1	1.7	1.9	1.6	1.2	1.1
2006.....	1.0	*	*	0.2	0.7	1.1	1.3	1.6	1.7	1.4	1.4	*
2005.....	1.0	*	*	0.1	0.7	1.0	1.2	1.7	1.8	1.5	1.3	*
2004.....	1.1	*	*	0.1	0.7	1.2	1.4	1.8	1.7	1.7	0.9	*
2003.....	1.1	*	*	0.2	0.8	0.9	1.5	1.8	2.2	1.6	1.3	*
2002.....	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001.....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000.....	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999.....	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*

All origins 3/, female

2007.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.4	0.3	*	*
2006.....	0.3	*	*	0.1	0.2	0.3	0.3	0.3	0.4	0.4	0.3	*
2005.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.3	0.3	*
2004.....	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003.....	0.2	*	*	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001.....	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999.....	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	0.7	*	*	0.1	0.5	0.6	0.7	1.0	1.1	0.9	0.6	0.5
2006.....	0.7	*	*	0.2	0.5	0.7	0.9	1.0	1.1	0.9	0.8	0.5
2005.....	0.7	*	*	0.1	0.5	0.7	0.8	1.1	1.1	0.9	0.7	0.5
2004.....	0.7	*	*	0.1	0.5	0.8	0.9	1.2	1.0	0.9	0.5	0.6
2003.....	0.7	*	*	0.2	0.5	0.6	0.9	1.2	1.2	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.1	0.9	0.8	*
2001.....	0.7	*	*	0.1	0.6	0.9	1.0	1.1	1.0	0.9	0.7	*
2000.....	0.7	*	*	0.1	0.6	0.8	1.0	1.0	1.1	1.0	0.6	*
1999.....	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.6	*

Non-Hispanic 4/, male

2007.....	1.1	*	*	0.1	0.7	1.0	1.2	1.7	1.9	1.7	1.2	*
2006.....	1.1	*	*	0.2	0.8	1.2	1.4	1.7	1.7	1.5	1.4	*
2005.....	1.1	*	*	0.1	0.8	1.1	1.4	1.8	1.9	1.5	1.3	*
2004.....	1.2	*	*	0.2	0.7	1.3	1.5	1.9	1.8	1.8	1.0	*
2003.....	1.2	*	*	0.2	0.8	1.0	1.6	2.0	2.3	1.7	1.3	*
2002.....	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	2.0	1.5	*
2001.....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.3	*
2000.....	1.2	*	*	0.2	1.0	1.5	1.7	1.8	1.9	1.7	1.3	*
1999.....	1.2	*	*	0.3	0.8	1.5	1.6	1.8	2.1	1.7	1.0	*

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	0.2	0.3	0.2	0.3	0.4	0.3	*	*
2006.....	0.3	*	*	*	0.2	0.3	0.4	0.3	0.4	0.4	0.4	*
2005.....	0.3	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.4	0.3	*
2004.....	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003.....	0.3	*	*	0.2	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001.....	0.2	*	*	*	0.2	0.3	0.3	0.3	0.2	0.2	0.4	*
2000.....	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	0.3	*	*
1999.....	0.3	*	*	0.1	0.3	0.3	0.4	0.4	0.3	0.3	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic white, both sexes

2007 5/.....	0.7	*	*	0.1	0.5	0.7	0.8	1.1	1.2	1.0	0.6	0.5
2006 5/.....	0.8	*	*	0.2	0.6	0.8	1.0	1.1	1.1	0.9	0.8	*
2005 5/.....	0.8	*	*	0.2	0.5	0.8	0.9	1.1	1.2	1.0	0.7	0.5
2004 5/.....	0.8	*	*	0.2	0.6	1.0	1.0	1.3	1.1	1.0	0.6	0.6
2003 5/.....	0.8	*	*	0.2	0.5	0.7	1.1	1.2	1.3	0.9	0.7	*
2002.....	0.8	*	*	0.1	0.6	0.8	1.0	1.2	1.2	1.0	0.8	*
2001.....	0.8	*	*	0.1	0.7	1.0	1.1	1.1	1.0	0.9	0.7	*
2000.....	0.8	*	*	0.2	0.7	1.0	1.1	1.1	1.2	1.0	0.7	*
1999.....	0.8	*	*	0.2	0.6	1.0	1.1	1.1	1.1	0.9	0.5	*

Non-Hispanic white, male

2007 5/.....	1.2	*	*	*	0.8	1.1	1.4	1.8	2.0	1.7	1.2	*
2006 5/.....	1.3	*	*	0.3	1.0	1.3	1.5	1.8	1.9	1.6	1.5	*
2005 5/.....	1.3	*	*	0.2	0.8	1.3	1.5	1.9	2.1	1.7	1.3	*
2004 5/.....	1.3	*	*	0.2	0.8	1.5	1.6	2.1	1.9	1.9	1.0	*
2003 5/.....	1.3	*	*	0.2	0.8	1.2	1.7	2.1	2.4	1.7	1.3	*
2002.....	1.3	*	*	*	0.9	1.3	1.8	2.0	2.1	2.0	1.4	*
2001.....	1.3	*	*	0.2	1.1	1.6	1.8	1.9	1.8	1.7	1.4	*
2000.....	1.4	*	*	0.2	1.1	1.7	1.8	1.9	2.0	1.8	1.3	*
1999.....	1.3	*	*	0.3	1.0	1.6	1.7	1.9	2.0	1.7	1.0	*

Non-Hispanic white, female

2007 5/.....	0.3	*	*	*	0.2	0.3	0.3	0.4	0.4	0.3	*	*
2006 5/.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.3	*
2005 5/.....	0.3	*	*	*	0.3	0.2	0.3	0.3	0.4	0.4	0.4	*
2004 5/.....	0.3	*	*	*	0.4	0.4	0.4	0.4	0.3	*	*	*
2003 5/.....	0.3	*	*	0.2	0.2	0.3	0.4	0.4	0.3	0.3	*	*
2002.....	0.3	*	*	*	0.3	0.2	0.3	0.3	0.4	*	0.3	*
2001.....	0.3	*	*	*	0.3	0.3	0.4	0.3	0.2	0.2	0.3	*
2000.....	0.3	*	*	*	0.3	0.3	0.3	0.3	0.4	0.3	*	*
1999.....	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	0.3	*	*



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes

2007.....	25.5	27.2	6.1	2.0	11.7	18.5	23.4	29.9	22.5	28.7	82.6	262.6
2006.....	24.4	24.3	6.1	2.0	11.4	17.8	23.6	28.7	20.5	27.8	81.4	250.5
2005.....	23.4	22.8	6.3	2.1	10.6	15.8	21.9	26.5	19.6	28.3	81.9	253.4
2004.....	21.8	22.2	6.1	2.2	9.8	13.8	20.9	24.0	17.7	26.5	78.2	249.5
2003.....	21.0	19.9	6.8	2.1	9.3	13.2	20.7	22.3	16.9	26.6	75.6	249.2
2002.....	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001.....	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000.....	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999.....	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3

All origins 3/, male

2007.....	31.2	30.2	7.8	2.6	18.1	27.4	32.1	40.1	31.0	38.9	103.0	307.3
2006.....	30.4	28.3	7.2	2.6	18.1	26.8	32.9	39.9	28.7	36.7	102.6	294.3
2005.....	29.0	24.0	7.9	2.8	16.7	23.6	30.6	37.3	27.8	37.4	104.2	298.3
2004.....	26.8	25.1	7.4	3.0	15.4	20.8	28.9	34.0	24.6	35.5	98.2	299.0
2003.....	26.0	22.4	8.5	2.8	14.7	19.9	29.4	31.6	24.1	35.3	96.3	300.5
2002.....	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001.....	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000.....	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999.....	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9

All origins 3/, female

2007.....	19.9	24.1	4.3	1.5	4.9	9.3	14.7	20.0	14.5	20.1	68.5	241.4
2006.....	18.6	20.2	5.0	1.4	4.3	8.5	14.1	17.8	12.8	20.1	67.0	230.1
2005.....	18.0	21.5	4.5	1.4	4.1	7.8	13.2	16.0	11.8	20.7	66.7	232.8
2004.....	16.9	19.3	4.8	1.5	3.8	6.5	12.9	14.4	11.2	18.9	64.8	227.3
2003.....	16.2	17.3	5.0	1.4	3.6	6.3	12.0	13.3	10.3	19.3	61.8	226.5
2002.....	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001.....	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000.....	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999.....	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Hispanic, both sexes												
2007.....	12.3	13.2	4.7	1.4	8.9	11.3	15.1	20.4	16.6	21.1	55.9	134.4
2006.....	12.4	14.8	5.0	1.5	8.3	11.9	16.2	20.0	17.8	20.2	54.2	131.9
2005.....	12.0	13.7	5.2	1.1	8.7	11.1	15.2	19.2	15.9	23.1	55.5	138.2
2004.....	10.6	13.1	4.3	1.2	7.1	9.3	14.7	16.7	14.6	20.2	52.3	140.8
2003.....	11.0	10.6	5.6	1.6	7.6	10.2	15.3	17.7	14.8	19.6	47.3	127.0
2002.....	10.8	12.1	5.1	1.5	7.9	9.4	15.7	16.8	13.5	21.1	51.2	127.5
2001.....	10.1	13.6	5.0	1.6	6.6	9.4	15.0	15.3	12.6	19.8	43.6	142.6
2000.....	10.2	9.9	5.2	1.5	7.3	10.0	15.3	14.2	13.1	18.4	48.2	159.2
1999.....	11.1	12.4	6.4	1.7	7.7	10.5	17.2	16.2	13.0	20.5	44.6	157.2
Hispanic, male												
2007.....	17.9	14.6	6.5	1.8	14.8	17.7	23.8	32.4	24.3	30.2	71.0	156.5
2006.....	18.0	16.1	6.0	2.0	13.5	18.4	25.5	31.0	28.5	30.0	66.4	155.6
2005.....	17.3	13.9	6.8	1.4	14.3	17.5	23.6	30.7	25.7	32.7	70.8	146.9
2004.....	15.2	14.9	5.1	1.5	11.4	14.7	22.6	26.3	23.2	28.1	64.7	165.3
2003.....	16.1	11.8	7.3	2.1	12.5	16.2	24.1	28.2	23.9	27.1	67.1	149.3
2002.....	16.1	15.7	6.4	1.9	13.1	15.5	24.6	27.1	21.4	30.2	68.5	179.0
2001.....	14.8	14.2	6.7	1.9	10.8	14.9	24.3	25.6	20.4	26.8	54.1	154.3
2000.....	15.3	10.9	5.9	1.6	12.1	16.5	25.6	24.3	19.7	27.6	57.5	207.6
1999.....	16.9	15.0	7.7	2.3	12.9	17.4	28.2	27.6	22.5	31.6	54.7	217.8
Hispanic, female												
2007.....	6.4	11.7	2.8	1.1	2.4	3.5	5.4	8.0	9.5	13.7	45.2	122.5
2006.....	6.4	13.5	4.0	1.0	2.3	3.9	5.7	8.6	7.9	12.3	45.6	119.2
2005.....	6.2	13.6	3.6	0.8	2.3	3.3	5.8	7.4	7.0	15.3	44.9	133.6
2004.....	5.8	11.3	3.4	0.8	2.2	2.8	6.0	7.1	6.9	13.9	43.7	128.2
2003.....	5.4	9.4	3.9	1.2	2.0	3.1	5.5	7.2	6.7	13.5	33.7	115.6
2002.....	5.2	8.3	3.7	1.2	1.9	2.1	5.9	6.6	6.3	13.8	39.3	102.0
2001.....	5.2	13.0	3.3	1.3	1.7	2.9	5.0	5.2	5.7	14.3	36.6	136.8
2000.....	4.9	8.8	4.4	1.5	1.6	2.4	4.4	4.3	7.3	11.1	41.9	135.5
1999.....	5.0	9.8	5.2	1.1	1.7	2.6	5.5	5.1	4.6	11.7	37.8	127.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nontransport accidents (W00-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	27.7	31.6	6.5	2.2	12.2	20.3	24.9	30.9	22.9	29.3	84.2	269.2
2006.....	26.4	27.1	6.3	2.1	12.0	19.2	24.8	29.5	20.6	28.3	82.9	256.3
2005.....	25.2	25.3	6.6	2.4	11.0	16.9	23.0	27.1	19.7	28.6	83.3	258.6
2004.....	23.5	24.8	6.6	2.5	10.3	14.8	21.8	24.6	17.8	26.8	79.5	254.1
2003.....	22.5	22.4	7.1	2.2	9.6	13.8	21.4	22.5	17.0	27.0	76.9	253.9
2002.....	21.6	22.3	6.7	2.6	9.0	13.4	20.6	20.4	15.8	26.1	75.2	250.9
2001.....	20.2	22.0	7.2	2.6	8.5	11.3	18.3	18.2	15.0	25.1	74.2	250.8
2000.....	19.2	20.7	7.9	2.8	8.0	11.1	17.3	16.8	14.9	24.1	68.7	244.4
1999.....	19.3	18.5	8.0	2.9	7.6	11.1	16.8	16.2	14.5	25.3	72.9	252.8

Non-Hispanic 4/, male

2007.....	33.6	34.9	8.2	2.8	18.8	30.1	33.7	40.9	31.5	39.4	104.8	315.6
2006.....	32.6	31.8	7.4	2.7	19.0	29.0	34.2	40.7	28.5	37.2	104.6	301.7
2005.....	30.9	26.8	8.3	3.2	17.2	25.1	31.8	37.8	27.8	37.5	106.0	305.9
2004.....	28.7	27.9	8.0	3.3	16.3	22.3	29.9	34.6	24.6	35.9	100.0	305.1
2003.....	27.5	25.2	8.7	2.9	15.2	20.7	30.0	31.7	24.0	35.7	97.6	306.9
2002.....	26.5	25.2	8.5	3.5	14.4	20.5	29.6	29.3	22.6	35.4	94.7	298.0
2001.....	25.0	24.9	8.7	3.5	13.7	17.5	26.9	27.2	21.2	33.2	95.1	311.6
2000.....	23.9	23.2	9.7	3.8	12.9	17.4	26.1	25.8	21.5	31.6	87.2	302.7
1999.....	24.0	21.5	9.6	3.9	12.4	17.6	25.7	25.0	21.1	34.1	92.8	314.3

Non-Hispanic 4/, female

2007.....	22.1	28.1	4.8	1.6	5.4	10.6	16.3	21.3	14.9	20.6	69.9	247.3
2006.....	20.6	22.2	5.2	1.5	4.7	9.5	15.5	18.8	13.2	20.7	68.2	235.2
2005.....	19.8	23.7	4.8	1.6	4.5	8.7	14.3	16.8	12.2	21.1	67.9	237.0
2004.....	18.6	21.6	5.2	1.6	4.1	7.3	13.9	15.1	11.5	19.2	65.8	231.2
2003.....	17.8	19.5	5.3	1.4	3.9	6.9	12.9	13.8	10.6	19.7	63.1	230.6
2002.....	16.9	19.4	4.9	1.6	3.4	6.4	11.8	11.9	9.5	18.4	62.3	230.6
2001.....	15.7	19.1	5.6	1.7	3.0	5.2	9.9	9.5	9.4	18.3	60.4	225.5
2000.....	14.7	18.0	6.0	1.7	2.9	4.8	8.6	8.1	8.8	17.8	56.7	220.8
1999.....	14.8	15.2	6.3	1.9	2.7	4.6	8.2	7.7	8.4	18.1	60.1	228.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Nontransport accidents (W00-X59,Y86)</b>												
<b>Non-Hispanic white, both sexes</b>												
2007 5/.....	30.6	24.9	6.3	1.8	14.1	23.7	27.2	31.6	22.4	29.2	87.8	285.0
2006 5/.....	28.8	21.7	5.9	1.8	13.7	22.1	26.4	29.5	19.9	28.2	86.2	269.8
2005 5/.....	27.3	21.9	6.4	2.0	12.3	19.0	24.4	26.9	18.8	28.3	84.9	269.7
2004 5/.....	25.6	21.2	6.5	2.0	11.6	16.6	23.3	24.7	17.1	26.7	81.8	266.4
2003 5/.....	24.4	18.4	6.9	1.8	10.9	15.3	22.4	22.1	16.6	26.6	78.9	265.9
2002.....	23.1	17.2	6.1	2.0	9.8	14.7	21.4	19.9	15.1	25.6	77.0	261.2
2001.....	21.5	17.8	7.1	2.2	9.1	12.1	18.5	17.2	14.2	24.7	75.4	259.4
2000.....	20.1	16.6	7.3	2.3	8.3	11.5	17.2	15.6	14.1	23.4	69.6	253.2
1999.....	20.0	14.7	7.4	2.4	7.8	11.3	16.7	14.9	13.3	24.2	73.2	260.8
<b>Non-Hispanic white, male</b>												
2007 5/.....	36.5	28.0	7.9	2.4	21.7	34.9	36.3	41.0	30.1	38.7	108.8	333.6
2006 5/.....	34.8	25.0	6.9	2.4	21.6	32.9	36.0	39.6	26.7	36.7	107.8	316.9
2005 5/.....	32.7	22.8	8.4	2.7	19.3	28.1	33.5	36.7	25.6	36.7	107.1	317.6
2004 5/.....	30.7	25.0	8.0	2.7	18.3	24.9	31.7	34.2	22.9	35.5	102.2	317.3
2003 5/.....	29.2	20.8	8.5	2.4	17.1	22.7	31.4	30.6	22.8	34.8	99.1	320.0
2002.....	27.8	20.4	7.6	2.7	15.6	22.2	30.5	28.1	21.0	34.3	96.0	307.6
2001.....	25.9	20.0	8.8	2.9	14.7	18.6	27.2	25.3	19.6	32.5	95.6	320.4
2000.....	24.5	19.1	9.1	3.2	13.5	18.1	26.0	23.6	19.9	30.4	87.2	311.5
1999.....	24.3	17.1	8.9	3.2	12.5	17.9	25.4	22.7	19.1	32.0	92.4	322.6
<b>Non-Hispanic white, female</b>												
2007 5/.....	24.9	21.6	4.6	1.3	6.1	12.3	18.1	22.3	15.0	21.0	73.2	261.9
2006 5/.....	23.1	18.2	4.8	1.2	5.5	11.1	16.8	19.5	13.4	20.9	71.2	247.9
2005 5/.....	22.0	21.1	4.4	1.3	4.9	9.8	15.3	17.2	12.4	21.1	69.6	247.8
2004 5/.....	20.7	17.3	4.9	1.3	4.6	8.3	14.9	15.4	11.6	19.1	68.0	243.7
2003 5/.....	19.7	16.0	5.1	1.1	4.4	7.7	13.5	13.8	10.7	19.6	65.3	242.0
2002.....	18.6	13.9	4.6	1.2	3.8	7.1	12.3	11.8	9.4	18.3	64.2	241.1
2001.....	17.2	15.5	5.3	1.5	3.3	5.5	9.8	9.3	9.2	18.1	62.0	234.0
2000.....	16.0	14.0	5.4	1.3	3.0	4.9	8.4	7.7	8.7	17.6	58.1	229.7
1999.....	16.0	12.3	5.9	1.5	2.9	4.6	8.0	7.3	7.9	17.6	60.6	236.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Nontransport accidents (W00-X59,Y86)</b>												
<b>Non-Hispanic black, both sexes</b>												
2007 5/.....	19.9	59.8	8.0	3.5	6.5	12.2	20.9	35.3	31.9	34.0	56.5	138.3
2006 5/.....	20.7	49.9	9.0	3.4	7.2	13.3	24.5	38.0	30.3	33.1	61.9	137.2
2005 5/.....	20.5	42.9	7.9	3.8	7.0	12.7	21.8	35.8	31.2	35.3	77.8	159.8
2004 5/.....	18.2	42.0	7.6	4.3	6.3	10.9	20.3	30.9	26.8	32.1	66.5	140.2
2003 5/.....	17.7	40.3	8.5	3.8	6.1	10.9	20.9	30.7	24.3	32.9	60.5	140.4
2002.....	18.3	44.5	9.5	4.7	6.9	11.4	21.4	29.6	24.6	32.4	62.8	156.7
2001.....	18.1	41.6	8.7	4.3	6.8	10.5	21.7	29.4	23.9	32.5	68.2	166.1
2000.....	18.0	39.5	10.9	4.8	7.1	11.3	21.5	28.1	23.8	33.1	65.4	161.2
1999.....	19.4	36.4	11.1	5.3	7.8	12.7	21.2	29.1	26.6	39.1	77.8	181.1
<b>Non-Hispanic black, male</b>												
2007 5/.....	26.6	61.9	9.8	4.4	9.9	18.4	30.1	51.8	51.0	52.9	72.9	161.4
2006 5/.....	28.8	59.5	10.5	4.2	11.7	21.3	35.7	58.8	49.7	48.2	85.6	164.9
2005 5/.....	27.9	47.3	9.0	5.0	11.0	18.9	30.8	55.4	52.9	52.1	111.0	193.2
2004 5/.....	24.6	43.0	9.1	5.7	9.9	16.9	28.3	46.4	43.7	47.7	94.7	189.0
2003 5/.....	24.1	45.3	10.8	4.8	9.5	16.6	30.2	46.1	39.3	48.4	85.7	184.3
2002.....	25.5	46.7	12.5	6.5	11.6	18.0	31.7	45.3	40.6	48.5	90.6	215.5
2001.....	25.0	48.2	10.0	6.0	11.0	16.2	31.6	47.9	38.0	47.5	101.4	220.6
2000.....	25.5	42.3	12.9	6.5	11.7	17.9	32.7	46.6	39.8	47.8	96.9	231.3
1999.....	27.5	42.3	13.7	7.3	13.5	20.0	33.3	48.7	42.5	60.1	107.6	249.6
<b>Non-Hispanic black, female</b>												
2007 5/.....	13.7	57.6	6.1	2.7	3.1	6.3	12.8	21.1	16.6	20.5	47.2	128.8
2006 5/.....	13.3	39.9	7.5	2.5	2.6	5.8	14.5	20.2	14.7	22.2	48.4	125.9
2005 5/.....	13.7	38.3	6.8	2.6	2.9	7.1	13.8	19.0	13.7	23.3	58.9	146.3
2004 5/.....	12.3	40.9	6.1	2.8	2.6	5.4	13.3	17.6	13.3	21.0	50.6	120.9
2003 5/.....	12.0	35.1	6.2	2.7	2.7	5.6	12.7	17.6	12.2	21.8	46.3	123.4
2002.....	11.8	42.3	6.5	2.9	2.1	5.4	12.3	16.2	11.7	21.1	47.1	134.2
2001.....	11.8	34.6	7.4	2.4	2.5	5.4	12.9	13.5	12.6	21.9	49.5	145.9
2000.....	11.3	36.6	8.8	3.0	2.6	5.4	11.6	12.2	10.9	22.8	47.8	135.2
1999.....	12.0	30.3	8.3	3.3	2.1	6.1	10.7	12.3	13.8	24.3	61.1	155.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Falls (W00-W19)

##### All origins 3/, both sexes

2007.....	7.5	0.6	0.2	0.1	0.5	0.8	1.4	3.0	5.3	13.4	50.3	166.7
2006.....	7.0	0.6	0.2	0.1	0.6	0.8	1.4	3.0	5.0	12.6	47.7	151.6
2005.....	6.6	*	0.2	0.1	0.6	0.7	1.4	2.8	4.8	12.4	45.6	147.7
2004.....	6.4	0.6	0.3	0.1	0.6	0.8	1.5	2.8	4.8	12.2	43.8	143.3
2003.....	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002.....	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001.....	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000.....	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999.....	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2

##### All origins 3/, male

2007.....	7.8	*	0.3	0.1	0.9	1.4	2.2	4.5	7.5	17.8	62.7	196.7
2006.....	7.3	*	0.3	0.1	1.0	1.3	2.3	4.7	7.3	16.5	59.4	181.1
2005.....	7.0	*	0.2	0.1	0.9	1.3	2.2	4.3	7.0	16.2	57.8	174.8
2004.....	6.8	*	0.4	0.1	1.0	1.3	2.4	4.5	7.1	16.5	54.8	174.4
2003.....	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002.....	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001.....	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000.....	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999.....	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5

##### All origins 3/, female

2007.....	7.2	*	*	*	0.2	0.2	0.5	1.4	3.3	9.7	41.8	152.4
2006.....	6.6	*	*	*	0.1	0.2	0.5	1.3	3.0	9.4	39.8	137.8
2005.....	6.3	*	*	*	0.2	0.2	0.5	1.3	2.7	9.2	37.4	135.2
2004.....	6.0	*	*	*	0.2	0.2	0.6	1.3	2.7	8.6	36.4	129.2
2003.....	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002.....	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001.....	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000.....	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999.....	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

Hispanic, both sexes

2007.....	2.7	*	*	*	0.7	1.1	1.4	2.9	4.9	9.9	35.0	92.6
2006.....	2.7	*	*	*	0.7	1.2	1.8	3.2	4.6	10.7	34.5	90.4
2005.....	2.5	*	*	*	0.7	0.9	1.3	2.7	4.7	11.8	34.8	91.6
2004.....	2.4	*	*	*	0.6	1.1	1.8	2.3	4.9	10.2	30.8	87.1
2003.....	2.1	*	*	*	0.5	0.8	1.4	2.6	5.0	9.2	25.4	80.8
2002.....	2.1	*	*	*	0.7	0.9	1.6	2.5	4.4	10.0	29.2	71.5
2001.....	2.0	*	*	*	0.7	1.1	1.8	2.6	4.2	9.1	23.6	70.0
2000.....	2.0	*	*	*	0.7	1.1	1.3	2.5	4.3	8.0	26.5	81.0
1999.....	1.9	*	*	*	0.7	1.1	1.4	2.2	3.8	8.1	24.2	86.3

Hispanic, male

2007.....	3.2	*	*	*	1.2	2.0	2.5	4.8	6.6	12.0	42.3	106.7
2006.....	3.5	*	*	*	1.3	2.1	3.1	5.5	7.7	15.1	41.4	102.6
2005.....	3.1	*	*	*	1.2	1.6	2.2	4.8	7.5	16.4	46.8	87.2
2004.....	3.1	*	*	*	1.1	1.8	3.0	4.2	8.1	14.0	38.3	106.5
2003.....	2.7	*	*	*	0.9	1.4	2.6	4.3	7.5	11.8	36.0	90.8
2002.....	2.8	*	*	*	1.3	1.6	2.7	3.8	7.0	13.8	36.9	97.0
2001.....	2.8	*	*	*	1.2	1.9	3.3	4.5	7.1	11.4	27.7	72.4
2000.....	2.7	*	*	*	1.2	2.0	2.3	4.4	6.6	12.7	30.5	104.8
1999.....	2.7	*	*	*	1.3	1.9	2.4	3.8	7.2	11.4	31.3	120.8

Hispanic, female

2007.....	2.0	*	*	*	*	*	*	1.0	3.4	8.1	29.9	84.9
2006.....	1.8	*	*	*	*	*	*	*	1.7	7.1	29.7	83.9
2005.....	1.8	*	*	*	*	*	*	*	2.2	8.1	26.5	93.8
2004.....	1.6	*	*	*	*	*	*	*	2.0	7.2	25.7	77.1
2003.....	1.5	*	*	*	*	*	*	*	2.8	7.2	18.2	75.8
2002.....	1.4	*	*	*	*	*	*	1.1	2.1	7.0	23.9	58.9
2001.....	1.3	*	*	*	*	*	*	*	*	7.2	20.9	68.9
2000.....	1.2	*	*	*	*	*	*	*	2.3	4.3	23.8	69.2
1999.....	1.1	*	*	*	*	*	*	*	*	5.5	19.4	69.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

#### Falls (W00-W19)

##### Non-Hispanic 4/, both sexes

2007.....	8.3	0.6	0.2	0.1	0.5	0.7	1.4	3.0	5.3	13.6	51.2	170.4
2006.....	7.7	*	0.2	0.1	0.5	0.7	1.4	2.9	5.1	12.8	48.5	154.6
2005.....	7.3	*	0.2	0.1	0.5	0.7	1.4	2.8	4.8	12.5	46.2	150.2
2004.....	7.0	*	0.3	0.1	0.6	0.7	1.4	2.9	4.8	12.3	44.5	145.5
2003.....	6.5	*	0.3	0.1	0.6	0.7	1.4	2.5	4.3	11.3	41.5	137.9
2002.....	6.2	*	0.2	0.1	0.6	0.7	1.5	2.5	4.0	10.8	38.6	132.6
2001.....	5.7	*	0.2	0.1	0.6	0.8	1.4	2.6	3.9	10.0	35.8	123.1
2000.....	5.1	*	0.2	0.1	0.6	0.7	1.3	2.3	3.9	9.1	31.2	113.5
1999.....	5.1	*	0.4	0.1	0.6	0.8	1.4	2.2	3.7	9.0	31.7	110.7

##### Non-Hispanic 4/, male

2007.....	8.6	*	*	*	0.8	1.2	2.1	4.5	7.5	18.2	63.9	201.5
2006.....	8.0	*	0.3	*	0.9	1.1	2.2	4.6	7.2	16.6	60.4	185.4
2005.....	7.6	*	*	0.1	0.9	1.2	2.2	4.2	6.9	16.2	58.4	179.2
2004.....	7.4	*	0.4	0.1	0.9	1.2	2.3	4.4	6.9	16.7	55.6	177.2
2003.....	6.8	*	0.5	0.1	0.9	1.1	2.3	3.9	6.0	14.6	52.3	168.3
2002.....	6.5	*	0.3	0.1	0.9	1.3	2.3	3.9	5.9	14.4	48.4	161.3
2001.....	6.2	*	0.3	0.2	1.1	1.4	2.3	4.2	5.8	13.1	46.4	159.0
2000.....	5.5	*	*	0.2	1.0	1.2	2.2	3.6	5.7	11.9	40.2	146.5
1999.....	5.5	*	0.5	0.2	1.0	1.3	2.3	3.6	5.5	12.4	40.6	142.9

##### Non-Hispanic 4/, female

2007.....	8.1	*	*	*	0.2	0.3	0.6	1.5	3.3	9.8	42.5	155.7
2006.....	7.4	*	*	*	0.2	0.3	0.6	1.4	3.1	9.5	40.3	140.2
2005.....	7.0	*	*	*	0.2	0.2	0.6	1.4	2.8	9.3	37.9	136.9
2004.....	6.7	*	*	*	0.2	0.2	0.6	1.4	2.8	8.7	37.0	131.3
2003.....	6.2	*	*	*	0.2	0.3	0.6	1.2	2.7	8.5	34.3	124.6
2002.....	5.9	*	*	*	0.2	0.2	0.7	1.2	2.3	7.7	32.1	120.2
2001.....	5.3	*	*	*	0.1	0.3	0.5	1.0	2.2	7.4	28.8	108.1
2000.....	4.7	*	*	*	0.2	0.2	0.5	1.0	2.1	6.8	25.3	100.1
1999.....	4.6	*	*	*	0.2	0.3	0.5	0.9	2.0	6.3	25.9	97.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

Non-Hispanic white, both sexes

2007 5/.....	9.8	*	*	0.1	0.6	0.9	1.4	3.1	5.5	14.2	54.7	182.9
2006 5/.....	9.0	*	*	0.1	0.6	0.8	1.5	3.0	5.1	13.3	51.4	164.5
2005 5/.....	8.5	*	*	0.1	0.6	0.7	1.5	2.8	4.8	13.0	48.5	159.6
2004 5/.....	8.2	*	0.3	0.1	0.7	0.8	1.5	2.9	4.8	12.8	47.2	155.1
2003 5/.....	7.5	*	0.3	0.1	0.6	0.8	1.5	2.5	4.3	11.6	43.6	146.8
2002.....	7.1	*	*	0.1	0.7	0.8	1.5	2.5	4.1	11.1	40.8	140.4
2001.....	6.6	*	0.2	0.2	0.8	0.9	1.4	2.6	3.9	10.3	37.6	129.9
2000.....	5.8	*	*	0.1	0.7	0.7	1.4	2.3	3.8	9.2	32.7	120.2
1999.....	5.7	*	0.4	0.2	0.7	0.8	1.5	2.1	3.5	9.1	33.1	116.6

Non-Hispanic white, male

2007 5/.....	10.0	*	*	*	0.9	1.4	2.2	4.6	7.7	18.7	67.6	215.6
2006 5/.....	9.2	*	*	*	1.0	1.2	2.3	4.6	7.2	17.1	63.3	196.1
2005 5/.....	8.7	*	*	*	1.0	1.2	2.4	4.2	6.9	16.8	60.9	188.7
2004 5/.....	8.5	*	*	*	1.1	1.3	2.4	4.5	6.7	17.3	58.6	186.9
2003 5/.....	7.7	*	0.5	0.2	1.0	1.2	2.4	3.8	6.0	14.8	54.6	178.7
2002.....	7.3	*	*	*	1.0	1.4	2.3	3.9	5.9	14.8	50.6	169.2
2001.....	7.0	*	*	0.2	1.3	1.5	2.3	4.1	5.7	13.5	48.2	166.0
2000.....	6.2	*	*	0.2	1.1	1.3	2.2	3.6	5.6	12.0	41.7	154.1
1999.....	6.1	*	0.5	0.2	1.1	1.4	2.3	3.4	5.1	12.2	42.0	149.7

Non-Hispanic white, female

2007 5/.....	9.7	*	*	*	0.2	0.3	0.6	1.6	3.4	10.4	45.6	167.4
2006 5/.....	8.8	*	*	*	0.2	0.3	0.6	1.4	3.2	10.1	43.1	149.8
2005 5/.....	8.3	*	*	*	0.2	0.2	0.6	1.5	2.8	9.8	40.1	146.3
2004 5/.....	7.9	*	*	*	0.2	0.2	0.5	1.4	2.9	9.0	39.4	140.9
2003 5/.....	7.3	*	*	*	0.2	0.3	0.6	1.2	2.7	8.9	36.2	132.8
2002.....	7.0	*	*	*	0.2	0.2	0.7	1.2	2.4	8.1	34.2	128.0
2001.....	6.2	*	*	*	0.2	0.3	0.5	1.1	2.2	7.7	30.6	114.8
2000.....	5.5	*	*	*	0.2	0.2	0.5	1.0	2.1	6.9	26.8	106.5
1999.....	5.4	*	*	*	0.2	0.3	0.6	0.8	1.9	6.4	27.2	103.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

Non-Hispanic black, both sexes

2007 5/.....	2.6	*	*	*	*	0.4	1.1	2.9	4.6	9.5	20.5	56.3
2006 5/.....	2.8	*	*	*	*	0.4	1.2	3.1	4.8	9.8	23.9	54.9
2005 5/.....	2.7	*	*	*	*	0.5	1.0	3.1	4.7	9.0	25.7	53.0
2004 5/.....	2.7	*	*	*	*	0.6	1.5	2.9	5.6	10.1	20.8	47.5
2003 5/.....	2.5	*	*	*	*	0.5	1.3	3.2	4.9	9.0	19.3	44.6
2002.....	2.3	*	*	*	*	0.4	1.3	2.8	4.3	8.2	16.3	49.3
2001.....	2.3	*	*	*	*	0.5	1.6	3.0	4.4	8.8	18.7	44.0
2000.....	2.1	*	*	*	*	0.4	1.3	2.7	4.8	8.3	15.2	39.1
1999.....	2.5	*	*	*	0.4	0.8	1.3	3.5	5.5	9.3	17.2	45.3

Non-Hispanic black, male

2007 5/.....	3.2	*	*	*	*	*	1.8	4.8	7.5	15.0	26.7	65.4
2006 5/.....	3.6	*	*	*	*	*	1.9	5.4	8.0	14.5	34.5	68.7
2005 5/.....	3.3	*	*	*	*	*	1.6	5.3	7.7	12.6	35.6	63.8
2004 5/.....	3.5	*	*	*	*	1.1	2.2	4.8	10.2	14.5	29.9	73.2
2003 5/.....	3.0	*	*	*	*	*	2.2	5.2	7.2	12.6	24.5	56.5
2002.....	3.0	*	*	*	*	0.8	2.2	4.7	6.8	11.9	24.9	76.6
2001.....	3.2	*	*	*	*	*	2.7	5.4	6.9	12.6	29.1	68.1
2000.....	2.8	*	*	*	*	*	2.1	4.6	8.2	11.4	21.6	61.1
1999.....	3.4	*	*	*	*	1.4	2.5	5.9	8.7	15.0	24.1	58.2

Non-Hispanic black, female

2007 5/.....	2.1	*	*	*	*	*	*	1.2	2.3	5.5	17.0	52.4
2006 5/.....	2.1	*	*	*	*	*	*	1.2	2.3	6.4	17.8	49.3
2005 5/.....	2.1	*	*	*	*	*	*	1.2	2.2	6.4	20.0	48.6
2004 5/.....	1.9	*	*	*	*	*	0.9	1.4	1.9	7.0	15.6	37.3
2003 5/.....	2.0	*	*	*	*	*	*	1.5	3.0	6.5	16.3	40.0
2002.....	1.6	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.9
2001.....	1.6	*	*	*	*	*	*	*	2.3	6.1	12.8	35.1
2000.....	1.5	*	*	*	*	*	*	1.0	2.0	6.1	11.6	31.0
1999.....	1.7	*	*	*	*	*	*	1.4	2.9	5.3	13.3	40.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

All origins 3/, both sexes

2007.....	0.2	*	*	0.1	0.4	0.2	0.2	0.2	0.2	0.2	0.2	*
2006.....	0.2	*	*	0.1	0.5	0.3	0.2	0.2	0.2	0.2	0.3	*
2005.....	0.3	*	0.1	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.3	*
2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

All origins 3/, male

2007.....	0.4	*	*	0.2	0.7	0.4	0.3	0.3	0.3	0.3	0.5	*
2006.....	0.4	*	*	0.2	0.9	0.5	0.3	0.3	0.3	0.4	0.5	*
2005.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2004.....	0.4	*	*	0.2	0.7	0.5	0.4	0.4	0.2	0.4	0.4	*
2003.....	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*

All origins 3/, female

2007.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	*	0.1	0.4	0.2	0.2	0.2	0.2	0.2	0.3	*
2006.....	0.2	*	*	0.1	0.5	0.3	0.2	0.2	0.2	0.2	0.3	*
2005.....	0.3	*	0.2	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2004.....	0.2	*	*	0.1	0.4	0.3	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.4	0.3	0.3	0.2	0.2	0.2	*
2001.....	0.3	*	*	0.2	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.4	0.4	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.7	0.4	0.3	0.3	0.2	0.2	0.2	*

Non-Hispanic 4/, male

2007.....	0.4	*	*	0.2	0.7	0.4	0.4	0.3	0.3	0.3	0.5	*
2006.....	0.4	*	*	0.2	0.9	0.5	0.3	0.3	0.3	0.4	0.6	*
2005.....	0.5	*	*	0.2	0.9	0.6	0.5	0.4	0.4	0.4	0.5	*
2004.....	0.4	*	*	0.2	0.8	0.5	0.4	0.5	0.2	0.4	0.5	*
2003.....	0.5	*	*	0.2	0.9	0.6	0.5	0.4	0.5	0.4	0.7	*
2002.....	0.5	*	*	0.2	1.0	0.7	0.5	0.5	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	1.0	0.6	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.2	0.6	0.5	0.5	0.4	0.4	0.4	*

Non-Hispanic 4/, female

2007.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

Non-Hispanic white, both sexes

2007 5/.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	0.2	0.3	*
2006 5/.....	0.2	*	*	0.1	0.3	0.3	0.2	0.2	0.2	0.2	0.3	*
2005 5/.....	0.3	*	*	0.1	0.4	0.2	0.3	0.3	0.3	0.3	0.3	*
2004 5/.....	0.2	*	*	0.1	0.4	0.2	0.3	0.3	0.1	0.2	0.2	*
2003 5/.....	0.3	*	*	0.1	0.4	0.3	0.3	0.3	0.3	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.2	*
2001.....	0.3	*	*	0.2	0.5	0.3	0.4	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.4	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

Non-Hispanic white, male

2007 5/.....	0.4	*	*	0.2	0.5	0.3	0.4	0.4	0.4	0.3	0.6	*
2006 5/.....	0.4	*	*	0.2	0.6	0.5	0.3	0.3	0.3	0.4	0.6	*
2005 5/.....	0.4	*	*	0.2	0.7	0.4	0.4	0.4	0.5	0.5	0.5	*
2004 5/.....	0.4	*	*	0.2	0.7	0.4	0.4	0.5	0.2	0.4	*	*
2003 5/.....	0.5	*	*	0.2	0.8	0.5	0.6	0.4	0.5	0.4	0.6	*
2002.....	0.5	*	*	0.2	0.8	0.6	0.5	0.5	0.4	0.5	0.5	*
2001.....	0.5	*	*	0.3	0.8	0.6	0.6	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.0	0.6	0.5	0.5	0.4	0.4	*	*

Non-Hispanic white, female

2007 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

All origins 3/, both sexes

2007.....	1.1	1.3	2.8	0.6	1.5	0.9	1.0	1.1	1.0	1.0	1.4	1.6
2006.....	1.2	1.2	2.8	0.6	1.5	1.1	1.1	1.1	1.0	1.1	1.5	1.4
2005.....	1.2	1.6	3.0	0.6	1.5	1.0	1.1	1.2	0.9	1.0	1.4	1.8
2004.....	1.1	1.5	2.7	0.7	1.4	1.0	1.0	1.1	0.9	1.0	1.2	1.6
2003.....	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002.....	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001.....	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999.....	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6

All origins 3/, male

2007.....	1.8	1.4	3.7	0.8	2.5	1.6	1.5	1.8	1.6	1.7	2.2	3.0
2006.....	1.9	1.3	3.5	0.9	2.5	1.9	1.7	1.8	1.5	1.7	2.4	2.4
2005.....	1.9	1.3	4.1	0.9	2.7	1.6	1.9	1.9	1.5	1.6	2.3	3.9
2004.....	1.8	1.8	3.2	0.9	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.9
2003.....	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002.....	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001.....	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000.....	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999.....	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3

All origins 3/, female

2007.....	0.5	1.3	1.8	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.9	0.9
2006.....	0.5	1.2	2.1	0.3	0.3	0.2	0.4	0.5	0.5	0.7	0.8	0.9
2005.....	0.5	1.8	2.0	0.3	0.3	0.3	0.4	0.5	0.3	0.6	0.8	0.9
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.0
2003.....	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002.....	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001.....	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999.....	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

Non-Hispanic 4/, both sexes

2007.....	1.2	1.4	3.0	0.6	1.4	0.9	1.0	1.1	1.0	1.0	1.4	1.6
2006.....	1.2	1.1	2.9	0.7	1.3	1.1	1.1	1.2	1.0	1.2	1.5	1.4
2005.....	1.2	1.5	3.1	0.7	1.4	0.9	1.2	1.2	0.9	1.1	1.4	1.8
2004.....	1.1	1.4	2.8	0.7	1.4	1.0	1.0	1.1	1.0	1.0	1.2	1.6
2003.....	1.1	1.5	3.0	0.7	1.3	0.8	1.1	1.0	1.0	1.1	1.3	1.8
2002.....	1.2	1.4	3.1	0.9	1.4	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001.....	1.1	1.6	3.1	0.9	1.5	0.9	1.0	0.9	0.8	0.9	1.3	1.7
2000.....	1.2	2.1	3.4	1.0	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.5
1999.....	1.2	1.7	3.3	0.9	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.7

Non-Hispanic 4/, male

2007.....	1.8	1.4	3.9	0.8	2.3	1.6	1.6	1.8	1.7	1.7	2.2	3.0
2006.....	1.9	*	3.5	1.0	2.3	1.9	1.8	1.9	1.6	1.7	2.5	2.4
2005.....	1.9	1.3	4.1	1.0	2.5	1.6	2.0	1.9	1.5	1.6	2.4	3.8
2004.....	1.8	1.8	3.4	1.0	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.7
2003.....	1.8	1.7	3.7	1.0	2.4	1.5	1.8	1.6	1.6	1.8	2.1	3.4
2002.....	1.9	1.6	4.1	1.3	2.6	1.8	2.0	1.5	1.3	1.6	2.0	3.6
2001.....	1.8	1.3	3.8	1.2	2.6	1.6	1.7	1.6	1.3	1.5	2.1	3.4
2000.....	2.0	2.4	4.3	1.4	2.7	1.8	1.9	1.6	1.6	1.6	1.8	2.8
1999.....	2.0	1.7	4.2	1.4	2.7	1.9	1.8	1.7	1.5	1.6	2.3	2.4

Non-Hispanic 4/, female

2007.....	0.5	1.3	2.0	0.4	0.4	0.3	0.4	0.5	0.4	0.5	0.9	0.9
2006.....	0.6	*	2.2	0.3	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.9
2005.....	0.5	1.8	2.0	0.4	0.3	0.3	0.4	0.5	0.3	0.6	0.8	0.9
2004.....	0.5	*	2.2	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.1
2003.....	0.5	*	2.3	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	1.1
2002.....	0.5	*	2.1	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001.....	0.5	1.9	2.4	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.7	1.0
2000.....	0.5	1.9	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999.....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.4	0.6	0.6	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

Non-Hispanic white, both sexes

2007 5/.....	1.1	1.2	3.3	0.4	1.2	0.8	1.0	1.2	1.0	1.0	1.4	1.6
2006 5/.....	1.2	1.3	3.1	0.5	1.1	1.0	1.2	1.2	1.0	1.1	1.5	1.4
2005 5/.....	1.2	1.8	3.4	0.5	1.2	0.9	1.2	1.2	0.9	1.0	1.3	1.8
2004 5/.....	1.1	1.5	3.2	0.5	1.2	0.9	1.0	1.1	0.9	1.0	1.2	1.5
2003 5/.....	1.1	1.4	3.3	0.5	1.2	0.8	1.1	1.0	0.9	1.1	1.3	1.7
2002.....	1.1	1.4	3.3	0.5	1.2	1.0	1.2	0.9	0.8	0.9	1.2	1.6
2001.....	1.1	1.5	3.5	0.5	1.3	0.8	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.1	1.7	3.7	0.7	1.3	1.0	1.1	0.9	0.9	1.0	1.3	1.4
1999.....	1.1	1.3	3.6	0.5	1.3	1.0	1.1	1.0	0.9	1.0	1.2	1.6

Non-Hispanic white, male

2007 5/.....	1.7	*	4.2	0.6	2.0	1.4	1.5	1.8	1.6	1.7	2.3	3.1
2006 5/.....	1.8	*	3.6	0.7	1.9	1.8	1.8	1.9	1.5	1.8	2.5	2.6
2005 5/.....	1.8	*	4.6	0.6	2.1	1.5	1.8	1.9	1.4	1.5	2.3	3.9
2004 5/.....	1.7	1.8	3.8	0.7	2.1	1.5	1.6	1.7	1.4	1.6	2.2	2.6
2003 5/.....	1.7	*	4.0	0.7	2.1	1.3	1.8	1.5	1.5	1.8	2.1	3.4
2002.....	1.7	*	4.3	0.7	2.2	1.7	1.9	1.4	1.2	1.5	2.0	3.7
2001.....	1.6	*	4.3	0.7	2.2	1.4	1.5	1.5	1.2	1.5	2.1	3.5
2000.....	1.8	1.9	4.7	1.0	2.2	1.7	1.8	1.5	1.6	1.5	1.8	2.7
1999.....	1.8	*	4.5	0.7	2.2	1.6	1.8	1.7	1.5	1.4	2.2	2.4

Non-Hispanic white, female

2007 5/.....	0.5	*	2.5	0.2	0.4	0.3	0.5	0.5	0.5	0.4	0.9	0.9
2006 5/.....	0.6	*	2.5	0.3	0.4	0.2	0.5	0.5	0.5	0.6	0.8	0.9
2005 5/.....	0.5	2.3	2.3	0.3	0.2	0.3	0.5	0.5	0.3	0.6	0.7	0.8
2004 5/.....	0.5	*	2.5	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.5	1.0
2003 5/.....	0.5	*	2.7	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	0.9
2002.....	0.5	*	2.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.7	0.7
2001.....	0.5	1.7	2.6	0.4	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	*	2.6	0.4	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
1999.....	0.5	*	2.6	0.3	0.3	0.3	0.4	0.4	0.4	0.6	0.6	1.3



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes

2007.....	1.1	0.9	1.2	0.5	0.5	0.5	0.7	1.1	1.5	2.2	3.3	4.9
2006.....	1.0	0.7	1.2	0.4	0.4	0.5	0.7	1.1	1.5	2.1	3.4	4.3
2005.....	1.1	0.8	1.3	0.5	0.4	0.6	0.8	1.2	1.3	2.1	3.6	5.0
2004.....	1.1	0.6	1.4	0.6	0.4	0.6	0.7	1.2	1.4	1.9	3.5	5.6
2003.....	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002.....	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001.....	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000.....	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999.....	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9

All origins 3/, male

2007.....	1.3	0.9	1.5	0.6	0.6	0.7	0.9	1.4	2.1	2.8	4.0	6.5
2006.....	1.2	*	1.4	0.4	0.5	0.6	0.8	1.4	1.9	2.6	4.5	6.3
2005.....	1.3	*	1.5	0.6	0.5	0.7	1.0	1.6	1.7	2.6	4.6	7.0
2004.....	1.3	*	1.5	0.7	0.5	0.8	0.9	1.6	1.8	2.5	4.5	7.2
2003.....	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002.....	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001.....	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000.....	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999.....	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3

All origins 3/, female

2007.....	0.9	*	0.9	0.5	0.3	0.4	0.5	0.8	1.0	1.6	2.9	4.2
2006.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.8	1.0	1.7	2.7	3.4
2005.....	0.9	*	1.0	0.5	0.3	0.4	0.5	0.8	1.0	1.6	2.9	4.0
2004.....	0.9	*	1.3	0.5	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003.....	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002.....	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001.....	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000.....	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999.....	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 4/, both sexes

2007.....	1.2	1.0	1.3	0.6	0.5	0.6	0.8	1.2	1.6	2.3	3.3	5.0
2006.....	1.1	0.8	1.3	0.5	0.4	0.5	0.7	1.2	1.5	2.2	3.6	4.5
2005.....	1.2	0.8	1.4	0.6	0.4	0.6	0.8	1.2	1.4	2.1	3.6	5.1
2004.....	1.2	0.7	1.6	0.7	0.5	0.7	0.8	1.3	1.5	1.9	3.5	5.6
2003.....	1.3	0.8	1.6	0.6	0.5	0.7	1.0	1.3	1.5	2.4	3.6	5.4
2002.....	1.2	1.0	1.5	0.7	0.5	0.8	0.9	1.2	1.3	2.1	3.4	4.8
2001.....	1.3	1.3	1.7	0.7	0.6	0.7	1.0	1.1	1.5	2.3	3.4	5.7
2000.....	1.3	1.0	2.2	0.7	0.5	0.7	0.9	1.2	1.5	2.2	3.8	6.3
1999.....	1.3	1.2	2.1	0.7	0.5	0.8	1.0	1.0	1.4	2.3	4.0	6.0

Non-Hispanic 4/, male

2007.....	1.4	*	1.6	0.6	0.6	0.7	1.0	1.5	2.2	3.0	3.9	6.6
2006.....	1.3	*	1.5	0.4	0.5	0.7	0.9	1.5	2.0	2.7	4.6	6.5
2005.....	1.4	*	1.6	0.7	0.5	0.8	1.1	1.6	1.7	2.7	4.7	7.2
2004.....	1.4	*	1.7	0.8	0.6	0.9	1.0	1.7	1.9	2.5	4.5	7.3
2003.....	1.5	*	1.9	0.6	0.6	0.8	1.3	1.7	1.9	3.0	4.5	7.3
2002.....	1.5	*	1.7	0.8	0.7	1.0	1.2	1.5	1.7	2.9	4.6	7.0
2001.....	1.6	1.5	2.0	0.8	0.8	0.9	1.5	1.6	2.0	2.8	4.2	8.3
2000.....	1.6	*	2.6	0.8	0.7	0.9	1.3	1.7	2.1	2.8	5.0	9.3
1999.....	1.6	*	2.3	0.8	0.7	1.1	1.4	1.4	1.7	3.0	5.1	9.2

Non-Hispanic 4/, female

2007.....	1.0	*	1.0	0.5	0.4	0.4	0.6	0.9	1.0	1.7	3.0	4.2
2006.....	0.9	*	1.2	0.5	0.4	0.4	0.6	0.8	1.1	1.8	2.8	3.5
2005.....	0.9	*	1.1	0.5	0.4	0.5	0.5	0.8	1.0	1.7	2.8	4.1
2004.....	1.0	*	1.5	0.6	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003.....	1.0	1.3	1.3	0.5	0.4	0.6	0.7	1.0	1.0	1.8	3.1	4.5
2002.....	0.9	*	1.2	0.6	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.8
2001.....	1.0	*	1.3	0.5	0.4	0.5	0.6	0.7	1.1	1.9	2.9	4.6
2000.....	1.0	*	1.7	0.7	0.4	0.5	0.6	0.7	1.0	1.7	3.1	5.0
1999.....	1.0	*	1.8	0.6	0.4	0.5	0.6	0.6	1.1	1.8	3.3	4.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic white, both sexes

2007 5/.....	1.1	*	0.8	0.4	0.5	0.5	0.7	1.1	1.5	2.0	3.0	4.1
2006 5/.....	1.1	*	0.9	0.3	0.4	0.5	0.7	1.1	1.4	2.1	3.1	3.9
2005 5/.....	1.1	*	1.0	0.5	0.4	0.6	0.8	1.2	1.1	1.9	3.1	4.3
2004 5/.....	1.1	*	1.3	0.5	0.5	0.6	0.7	1.1	1.3	1.6	3.0	4.9
2003 5/.....	1.2	*	1.3	0.4	0.5	0.7	1.0	1.2	1.3	2.1	3.1	4.7
2002.....	1.0	*	1.0	0.4	0.4	0.7	0.8	1.1	1.1	1.8	3.0	3.9
2001.....	1.1	1.1	1.3	0.5	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.7
2000.....	1.1	*	1.5	0.5	0.5	0.6	0.8	1.1	1.3	1.8	3.1	5.1
1999.....	1.1	0.9	1.5	0.5	0.5	0.8	0.9	0.9	1.2	1.9	3.2	4.7

Non-Hispanic white, male

2007 5/.....	1.3	*	1.1	0.4	0.6	0.7	0.9	1.4	2.0	2.5	3.5	5.3
2006 5/.....	1.3	*	1.2	0.3	0.5	0.6	0.8	1.4	1.8	2.4	4.0	6.2
2005 5/.....	1.3	*	1.4	0.5	0.5	0.7	1.1	1.5	1.4	2.3	4.1	6.1
2004 5/.....	1.3	*	1.4	0.6	0.6	0.9	0.9	1.5	1.6	2.0	3.9	6.5
2003 5/.....	1.4	*	1.6	0.4	0.7	0.8	1.3	1.5	1.7	2.6	3.6	6.5
2002.....	1.3	*	1.1	0.6	0.6	0.9	1.1	1.4	1.4	2.4	4.2	5.7
2001.....	1.4	*	1.7	0.6	0.7	0.8	1.3	1.3	1.7	2.3	3.5	6.9
2000.....	1.4	*	1.8	0.6	0.6	0.8	1.1	1.6	1.7	2.3	4.1	8.1
1999.....	1.4	*	1.7	0.6	0.7	1.1	1.3	1.2	1.5	2.3	4.2	7.8

Non-Hispanic white, female

2007 5/.....	0.9	*	0.5	0.4	0.4	0.4	0.5	0.8	1.0	1.6	2.6	3.5
2006 5/.....	0.9	*	0.7	0.4	0.3	0.4	0.6	0.8	1.0	1.7	2.5	2.8
2005 5/.....	0.9	*	0.7	0.4	0.3	0.4	0.5	0.8	0.9	1.6	2.4	3.5
2004 5/.....	0.9	*	1.2	0.4	0.4	0.4	0.5	0.8	1.0	1.2	2.4	4.2
2003 5/.....	1.0	*	1.0	0.4	0.4	0.6	0.6	0.9	0.9	1.6	2.8	3.9
2002.....	0.8	*	0.8	0.3	0.3	0.5	0.5	0.7	0.8	1.3	2.2	3.2
2001.....	0.9	*	0.9	0.4	0.3	0.4	0.6	0.6	1.0	1.7	2.5	3.8
2000.....	0.9	*	1.1	0.5	0.3	0.4	0.5	0.7	0.8	1.5	2.4	3.9
1999.....	0.8	*	1.3	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.6	3.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic black, both sexes

2007 5/.....	2.0	*	3.2	1.4	0.5	1.0	1.3	2.0	2.8	5.0	8.0	15.6
2006 5/.....	1.8	*	3.1	1.1	0.6	0.9	1.1	2.0	2.8	4.2	9.6	12.5
2005 5/.....	2.0	*	2.8	1.3	0.6	1.0	1.2	2.0	3.4	4.4	9.4	14.5
2004 5/.....	2.1	*	2.9	1.7	0.6	1.0	1.2	2.5	3.5	5.0	9.7	15.3
2003 5/.....	2.1	*	3.1	1.5	0.6	0.9	1.2	2.6	2.9	5.4	10.3	14.9
2002.....	2.1	*	3.7	1.5	0.8	1.2	1.1	2.2	3.3	5.0	8.7	16.0
2001.....	2.3	*	3.4	1.3	0.7	1.2	1.8	2.4	3.6	6.1	9.9	17.8
2000.....	2.6	*	5.1	1.6	0.7	1.2	1.7	2.4	4.2	6.1	13.0	20.7
1999.....	2.6	*	4.7	1.6	0.7	1.0	1.8	2.2	3.4	7.0	13.5	22.0

Non-Hispanic black, male

2007 5/.....	2.5	*	3.5	1.6	*	1.1	2.0	2.7	4.5	8.1	10.2	23.6
2006 5/.....	2.1	*	3.1	0.9	*	1.2	1.4	2.6	4.0	5.6	13.9	*
2005 5/.....	2.4	*	2.8	1.4	0.6	1.1	1.8	3.1	5.2	6.4	13.5	21.9
2004 5/.....	2.7	*	3.0	2.0	0.7	1.3	1.7	3.9	4.5	7.7	13.4	*
2003 5/.....	2.5	*	3.4	1.5	*	1.1	1.5	3.4	4.4	7.7	16.2	*
2002.....	2.6	*	4.3	1.6	1.0	1.4	1.7	3.1	4.6	7.8	11.1	26.3
2001.....	3.0	*	3.7	1.6	0.8	1.4	2.8	3.9	5.1	9.0	13.2	26.6
2000.....	3.2	*	5.8	1.7	0.9	1.6	2.4	3.5	6.3	9.0	17.9	28.2
1999.....	3.1	*	5.3	1.8	0.9	1.2	2.7	3.3	4.3	10.4	18.4	29.7

Non-Hispanic black, female

2007 5/.....	1.6	*	2.8	1.3	*	0.8	0.8	1.3	1.4	2.9	6.7	12.3
2006 5/.....	1.6	*	3.1	1.2	*	*	0.7	1.4	1.7	3.1	7.1	12.0
2005 5/.....	1.5	*	2.7	1.2	*	0.9	*	1.1	2.0	2.9	7.2	11.5
2004 5/.....	1.7	*	2.9	1.3	*	*	0.7	1.3	2.6	3.0	7.6	13.7
2003 5/.....	1.8	*	2.8	1.4	0.7	*	0.9	2.0	1.7	3.8	7.0	13.2
2002.....	1.6	*	3.0	1.3	*	1.0	*	1.4	2.2	3.1	7.3	12.1
2001.....	1.8	*	3.1	1.0	*	1.1	1.0	1.1	2.4	4.0	8.1	14.5
2000.....	2.0	*	4.3	1.4	*	0.9	1.2	1.4	2.5	4.0	10.2	17.9
1999.....	2.1	*	4.1	1.4	*	0.8	1.1	1.3	2.8	4.7	10.7	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes

2007.....	9.9	*	0.2	0.2	7.4	14.0	17.6	20.5	9.5	3.1	2.7	3.5
2006.....	9.2	*	0.2	0.1	6.9	13.0	17.3	19.0	7.6	2.6	2.6	3.8
2005.....	8.0	0.5	0.1	0.1	5.9	10.9	15.3	16.4	6.6	2.3	2.4	3.5
2004.....	7.1	*	*	0.1	5.4	9.1	14.6	14.5	5.4	2.3	2.4	3.4
2003.....	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002.....	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001.....	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000.....	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999.....	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4

All origins 3/, male

2007.....	13.2	*	*	0.2	11.1	20.2	22.9	25.6	12.1	4.0	2.9	3.7
2006.....	12.6	*	*	0.2	10.6	18.9	23.0	24.9	9.9	2.9	2.9	4.6
2005.....	10.9	*	*	0.1	9.0	15.6	20.2	21.7	8.7	2.7	3.0	4.6
2004.....	9.6	*	*	0.1	8.2	13.1	19.0	19.1	6.6	2.6	2.7	4.2
2003.....	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002.....	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001.....	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000.....	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999.....	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4

All origins 3/, female

2007.....	6.7	*	*	0.2	3.5	7.7	12.2	15.6	7.2	2.4	2.6	3.4
2006.....	5.9	*	*	0.1	3.0	6.9	11.6	13.4	5.5	2.3	2.4	3.4
2005.....	5.1	*	*	0.1	2.7	6.0	10.5	11.3	4.7	2.0	2.0	3.0
2004.....	4.7	*	*	0.1	2.4	4.9	10.3	10.0	4.3	2.0	2.2	3.0
2003.....	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002.....	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001.....	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000.....	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999.....	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Hispanic, both sexes

2007.....	5.4	*	*	*	4.1	6.8	10.1	13.2	6.5	2.8	*	*
2006.....	5.2	*	*	*	3.3	7.0	10.6	12.3	7.0	1.5	*	*
2005.....	4.7	*	*	*	3.4	6.2	9.9	10.9	5.5	2.2	*	*
2004.....	4.2	*	*	*	2.8	4.9	9.2	9.7	5.4	2.4	*	*
2003.....	4.5	*	*	*	3.1	5.7	10.3	10.5	4.4	1.8	*	*
2002.....	4.3	*	*	*	2.9	5.0	10.1	10.1	4.0	2.2	*	*
2001.....	3.6	*	*	*	2.2	4.9	8.7	7.7	3.5	*	*	*
2000.....	3.7	*	*	*	2.0	5.3	9.6	8.0	3.1	*	*	*
1999.....	4.4	*	*	*	2.3	6.1	11.9	9.8	3.5	*	*	*

Hispanic, male

2007.....	8.3	*	*	*	6.5	10.3	15.4	20.6	9.3	4.8	*	*
2006.....	7.8	*	*	*	4.8	10.3	15.9	18.1	11.7	*	*	*
2005.....	7.3	*	*	*	5.2	9.3	15.0	16.8	9.3	*	*	*
2004.....	6.2	*	*	*	4.2	7.6	13.5	14.2	8.4	*	*	*
2003.....	6.9	*	*	*	4.8	8.6	15.6	16.4	7.4	*	*	*
2002.....	6.6	*	*	*	4.4	7.8	15.4	15.8	6.4	*	*	*
2001.....	5.6	*	*	*	3.1	7.4	13.6	12.9	5.8	*	*	*
2000.....	6.0	*	*	*	3.2	8.4	15.6	13.6	4.4	*	*	*
1999.....	7.3	*	*	*	3.6	9.8	19.4	16.4	6.6	*	*	*

Hispanic, female

2007.....	2.2	*	*	*	1.5	2.5	4.2	5.6	3.9	*	*	*
2006.....	2.4	*	*	*	1.5	2.9	4.8	6.4	2.8	*	*	*
2005.....	2.0	*	*	*	1.4	2.3	4.3	5.0	1.9	*	*	*
2004.....	1.9	*	*	*	1.3	1.6	4.4	5.1	2.6	*	*	*
2003.....	1.9	*	*	*	1.2	2.2	4.5	4.6	*	*	*	*
2002.....	1.7	*	*	*	1.1	1.6	4.3	4.4	*	*	*	*
2001.....	1.5	*	*	*	1.1	1.9	3.4	2.6	*	*	*	*
2000.....	1.3	*	*	*	*	1.6	3.1	2.5	*	*	*	*
1999.....	1.4	*	*	*	0.8	1.7	3.9	3.3	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 4/, both sexes

2007.....	10.7	*	0.2	0.2	8.1	15.8	18.9	21.3	9.7	3.1	2.8	3.6
2006.....	9.8	*	0.2	0.1	7.7	14.5	18.4	19.7	7.6	2.6	2.6	3.9
2005.....	8.5	*	*	0.1	6.4	12.0	16.2	16.9	6.7	2.3	2.5	3.5
2004.....	7.6	*	*	0.1	5.9	10.1	15.4	14.9	5.4	2.3	2.4	3.4
2003.....	7.0	*	0.3	0.2	5.2	9.2	14.5	13.5	4.9	2.2	2.2	3.5
2002.....	6.3	0.6	0.2	0.1	4.4	8.4	13.8	11.8	4.1	2.1	2.3	4.1
2001.....	5.1	*	0.2	0.1	3.6	6.6	11.4	9.1	3.1	1.6	2.3	3.6
2000.....	4.6	*	0.2	0.1	3.1	6.0	10.3	8.0	2.8	1.5	2.0	4.4
1999.....	4.3	*	0.2	0.1	2.5	5.8	9.7	7.5	2.7	1.6	2.2	3.5

Non-Hispanic 4/, male

2007.....	14.1	*	*	0.3	12.1	22.9	24.3	26.1	12.2	3.9	3.0	3.7
2006.....	13.4	*	*	0.2	11.9	21.3	24.2	25.5	9.7	2.9	2.9	4.7
2005.....	11.5	*	*	0.2	9.8	17.3	21.0	22.1	8.5	2.6	3.1	4.7
2004.....	10.2	*	*	0.1	9.1	14.5	19.8	19.5	6.4	2.6	2.7	4.2
2003.....	9.5	*	0.3	0.2	7.9	13.3	19.2	17.9	6.1	2.6	2.8	4.0
2002.....	8.8	*	*	0.1	6.6	12.3	18.7	15.9	5.2	2.4	2.6	5.3
2001.....	7.2	*	*	0.1	5.7	9.6	15.9	12.9	3.9	2.0	2.8	3.9
2000.....	6.6	*	*	0.1	4.9	9.1	14.8	11.7	3.3	1.7	2.1	6.2
1999.....	6.3	*	*	0.1	3.8	8.8	14.2	11.1	3.6	2.0	2.5	4.4

Non-Hispanic 4/, female

2007.....	7.4	*	*	0.1	3.9	8.8	13.6	16.7	7.4	2.5	2.7	3.5
2006.....	6.4	*	*	*	3.3	7.8	12.7	14.1	5.8	2.4	2.4	3.5
2005.....	5.6	*	*	*	2.9	6.8	11.5	12.0	4.9	2.1	2.0	3.0
2004.....	5.1	*	*	0.1	2.7	5.6	11.1	10.5	4.4	2.0	2.2	3.1
2003.....	4.6	*	*	0.2	2.4	5.2	9.9	9.2	3.8	1.9	1.8	3.3
2002.....	4.0	*	*	*	2.0	4.5	8.9	7.9	3.0	1.8	2.1	3.5
2001.....	3.1	*	*	*	1.5	3.5	7.1	5.4	2.3	1.2	1.9	3.4
2000.....	2.7	*	*	*	1.4	3.0	5.9	4.4	2.3	1.3	2.0	3.6
1999.....	2.4	*	*	*	1.2	2.8	5.4	4.0	1.8	1.3	2.0	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic white, both sexes

2007 5/.....	11.7	*	*	0.2	10.0	19.1	21.2	22.0	9.3	3.0	2.8	3.7
2006 5/.....	10.5	*	*	0.1	9.5	17.2	19.8	19.7	7.1	2.4	2.6	3.8
2005 5/.....	9.0	*	*	0.1	7.9	14.0	17.6	16.9	6.1	2.1	2.4	3.6
2004 5/.....	8.2	*	*	0.1	7.3	11.8	16.8	15.0	5.0	2.2	2.4	3.5
2003 5/.....	7.4	*	*	0.1	6.4	10.6	15.5	13.4	4.8	2.0	2.3	3.5
2002.....	6.6	*	*	0.1	5.3	9.6	14.5	11.4	3.7	2.0	2.3	4.1
2001.....	5.2	*	*	0.1	4.3	7.3	11.7	8.3	2.7	1.5	2.1	3.4
2000.....	4.6	*	*	0.1	3.7	6.5	10.4	7.2	2.5	1.4	2.1	4.1
1999.....	4.2	*	*	0.1	2.8	6.0	9.6	6.7	2.4	1.5	2.3	3.3

Non-Hispanic white, male

2007 5/.....	15.4	*	*	0.2	15.0	27.6	27.2	26.4	11.1	3.5	2.9	3.9
2006 5/.....	14.2	*	*	0.2	14.7	24.9	25.8	24.8	8.2	2.4	2.9	4.8
2005 5/.....	12.2	*	*	0.2	12.1	20.1	22.7	21.5	7.3	2.1	3.0	4.9
2004 5/.....	10.9	*	*	0.2	11.2	17.1	21.4	19.3	5.4	2.3	2.5	4.3
2003 5/.....	10.1	*	*	*	9.8	15.3	20.3	17.5	5.6	2.1	2.9	3.6
2002.....	9.1	*	*	0.2	8.0	13.9	19.6	15.0	4.3	2.2	2.5	5.4
2001.....	7.3	*	*	*	6.7	10.7	16.3	11.5	3.1	1.9	2.5	3.8
2000.....	6.6	*	*	*	5.8	9.8	14.9	10.3	2.7	1.5	2.2	5.8
1999.....	6.1	*	*	*	4.3	9.2	14.0	9.6	2.9	1.8	2.5	4.6

Non-Hispanic white, female

2007 5/.....	8.1	*	*	*	4.7	10.6	15.3	17.7	7.6	2.5	2.8	3.6
2006 5/.....	7.0	*	*	*	4.0	9.4	13.8	14.7	5.9	2.4	2.5	3.3
2005 5/.....	6.1	*	*	*	3.5	7.9	12.4	12.4	5.0	2.2	2.1	2.9
2004 5/.....	5.5	*	*	*	3.2	6.5	12.1	10.8	4.5	2.1	2.3	3.1
2003 5/.....	4.9	*	*	*	2.9	5.9	10.6	9.4	4.0	1.9	1.9	3.5
2002.....	4.3	*	*	*	2.5	5.2	9.3	7.8	3.1	1.9	2.2	3.5
2001.....	3.1	*	*	*	1.7	3.8	7.1	5.2	2.4	1.2	1.9	3.2
2000.....	2.7	*	*	*	1.5	3.2	5.8	4.2	2.3	1.4	2.0	3.4
1999.....	2.4	*	*	*	1.3	2.8	5.1	3.8	1.9	1.2	2.1	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic black, both sexes

2007 5/.....	8.4	*	*	*	2.3	7.1	14.0	24.0	16.1	5.3	3.2	*
2006 5/.....	9.2	*	*	*	2.2	8.3	17.9	26.1	14.6	4.7	2.8	5.6
2005 5/.....	7.8	*	*	*	1.8	7.3	14.6	22.2	13.1	4.5	2.9	*
2004 5/.....	6.7	*	*	*	1.8	5.5	13.4	19.2	10.0	3.8	3.4	*
2003 5/.....	6.4	*	*	*	1.4	5.8	13.9	18.3	7.6	4.1	*	*
2002.....	6.4	*	*	*	1.5	5.8	14.1	18.3	8.6	2.5	2.8	*
2001.....	6.0	*	*	*	1.6	5.3	13.4	17.0	6.4	2.4	3.7	6.6
2000.....	5.7	*	*	*	1.4	5.7	13.2	15.7	5.5	2.5	*	8.3
1999.....	5.8	*	*	*	1.7	6.3	13.2	15.6	6.0	3.0	*	*

Non-Hispanic black, male

2007 5/.....	11.6	*	*	*	3.1	10.3	18.4	33.4	25.9	9.0	*	*
2006 5/.....	13.3	*	*	*	3.1	13.0	24.9	38.8	25.0	7.8	*	*
2005 5/.....	11.0	*	*	*	2.4	10.4	19.0	32.7	22.4	8.2	*	*
2004 5/.....	9.1	*	*	*	2.6	7.8	17.0	27.5	16.5	6.4	*	*
2003 5/.....	9.1	*	*	*	2.1	8.3	19.2	26.6	12.9	7.5	*	*
2002.....	9.3	*	*	*	2.2	8.7	19.8	27.1	15.2	4.5	*	*
2001.....	8.6	*	*	*	2.3	7.6	18.2	27.0	11.0	3.7	6.0	*
2000.....	8.4	*	*	*	1.8	8.5	19.0	25.3	9.5	4.3	*	*
1999.....	8.8	*	*	*	2.8	9.3	19.5	25.9	11.0	4.4	*	*

Non-Hispanic black, female

2007 5/.....	5.6	*	*	*	1.4	4.1	10.0	15.9	8.3	2.6	*	*
2006 5/.....	5.4	*	*	*	1.3	3.8	11.8	15.2	6.2	2.4	*	*
2005 5/.....	4.9	*	*	*	1.2	4.5	10.8	13.3	5.7	*	*	*
2004 5/.....	4.4	*	*	*	1.1	3.4	10.1	12.0	4.7	*	*	*
2003 5/.....	4.1	*	*	*	0.7	3.6	9.3	11.1	3.4	*	*	*
2002.....	3.8	*	*	*	0.7	3.2	9.1	10.8	3.3	*	*	*
2001.....	3.6	*	*	*	0.9	3.2	9.2	8.4	2.6	*	*	*
2000.....	3.2	*	*	*	1.0	3.1	8.0	7.5	2.4	*	*	*
1999.....	3.1	*	*	*	*	3.5	7.7	6.7	1.9	2.1	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes

2007.....	5.7	23.9	1.6	0.6	1.4	2.0	2.6	4.0	5.0	8.9	24.5	85.8
2006.....	5.8	21.5	1.6	0.6	1.6	2.1	2.9	4.2	5.2	9.1	25.9	89.3
2005.....	6.3	19.5	1.5	0.6	1.7	2.3	3.1	4.7	5.8	10.2	28.6	95.1
2004.....	5.8	19.2	1.6	0.6	1.5	2.1	2.9	4.2	5.0	8.8	27.0	95.5
2003.....	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
2002.....	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001.....	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000.....	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999.....	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9

All origins 3/, male

2007.....	6.7	26.5	1.9	0.7	2.3	3.2	4.2	6.4	7.4	12.3	30.6	97.1
2006.....	7.0	25.2	1.7	0.8	2.6	3.5	4.8	6.8	7.7	12.7	32.9	99.4
2005.....	7.4	21.1	1.8	0.9	2.8	3.9	4.9	7.4	8.6	13.8	35.9	107.1
2004.....	6.8	21.7	2.0	0.9	2.6	3.4	4.6	6.6	7.5	11.8	33.6	109.8
2003.....	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
2002.....	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001.....	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000.....	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999.....	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7

All origins 3/, female

2007.....	4.6	21.2	1.2	0.4	0.4	0.6	1.0	1.6	2.7	5.9	20.3	80.4
2006.....	4.7	17.6	1.5	0.3	0.5	0.7	1.0	1.8	2.8	6.1	21.2	84.5
2005.....	5.1	17.9	1.2	0.4	0.6	0.7	1.2	2.0	3.1	7.1	23.6	89.6
2004.....	4.8	16.5	1.1	0.3	0.4	0.7	1.2	1.8	2.7	6.4	22.6	89.1
2003.....	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
2002.....	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001.....	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000.....	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999.....	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Hispanic, both sexes

2007.....	2.6	10.6	1.2	0.4	1.5	1.9	2.4	3.1	3.8	6.7	15.2	35.0
2006.....	2.7	12.4	1.4	0.4	1.5	2.0	2.5	3.1	4.4	6.4	14.7	37.2
2005.....	2.8	11.0	1.2	0.4	1.7	2.4	2.6	3.8	4.1	6.8	15.1	38.1
2004.....	2.5	9.8	1.2	0.4	1.7	1.9	2.4	3.3	3.5	5.2	14.6	44.2
2003.....	2.5	8.2	1.6	0.4	1.6	2.0	2.2	3.0	3.9	6.3	16.9	37.3
2002.....	2.5	8.2	1.3	0.5	1.5	1.9	2.5	2.8	3.4	6.2	15.3	48.2
2001.....	2.7	9.5	1.4	0.3	1.5	1.8	2.7	3.6	3.8	7.7	16.0	63.8
2000.....	2.7	7.5	1.4	0.4	1.6	2.2	2.9	2.7	4.0	6.2	16.6	70.3
1999.....	2.6	8.6	1.9	0.5	1.6	1.6	2.4	2.8	4.2	8.7	15.0	65.9

Hispanic, male

2007.....	3.7	11.7	1.6	0.5	2.5	3.2	4.0	5.0	6.4	10.9	20.0	38.6
2006.....	3.9	13.7	1.4	0.6	2.5	3.2	4.6	5.4	6.6	9.7	18.3	45.1
2005.....	4.0	11.5	1.5	0.5	2.9	4.1	4.4	6.4	6.4	9.7	18.0	46.1
2004.....	3.5	11.7	1.6	0.6	2.8	2.9	3.9	5.6	5.1	6.9	17.5	43.4
2003.....	3.6	9.1	1.9	0.5	2.9	3.4	3.7	4.7	6.4	9.0	24.9	44.6
2002.....	3.6	10.1	1.8	0.6	2.5	3.3	4.2	4.8	5.6	8.4	21.3	70.3
2001.....	3.7	9.6	1.8	*	2.6	3.0	4.6	6.1	5.6	10.8	20.6	68.6
2000.....	3.8	7.9	1.5	*	2.9	3.7	5.2	4.5	6.6	8.7	19.7	88.7
1999.....	3.6	9.0	2.5	*	2.6	2.7	4.0	4.8	6.2	13.2	15.6	84.1

Hispanic, female

2007.....	1.4	9.3	*	*	*	*	*	1.0	1.4	3.3	11.9	33.1
2006.....	1.5	11.0	1.5	*	*	0.6	*	*	2.4	3.7	12.2	33.0
2005.....	1.5	10.5	*	*	*	*	*	1.3	2.0	4.5	13.0	33.9
2004.....	1.5	7.9	*	*	*	0.6	*	*	2.0	3.8	12.6	44.7
2003.....	1.3	7.3	1.3	*	*	*	*	1.3	*	4.1	11.3	33.6
2002.....	1.2	6.1	*	*	*	*	*	*	*	4.4	11.2	37.3
2001.....	1.6	9.5	*	*	*	*	*	1.2	2.1	5.4	13.0	61.4
2000.....	1.4	7.2	*	*	*	*	*	*	*	4.3	14.5	61.3
1999.....	1.6	8.1	*	*	*	*	*	*	2.4	5.2	14.6	57.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	6.2	28.1	1.7	0.6	1.4	2.0	2.6	4.1	5.0	9.0	25.1	88.5
2006.....	6.4	24.2	1.6	0.6	1.6	2.2	3.0	4.3	5.2	9.3	26.6	91.8
2005.....	6.8	21.9	1.6	0.7	1.7	2.3	3.1	4.7	5.8	10.4	29.3	97.7
2004.....	6.3	21.9	1.7	0.7	1.5	2.1	3.0	4.2	5.1	9.1	27.7	97.7
2003.....	6.4	19.2	1.8	0.6	1.5	2.0	3.1	4.0	5.2	9.8	28.0	105.1
2002.....	6.5	18.9	1.6	0.7	1.6	2.1	3.1	3.8	5.3	10.0	29.4	107.6
2001.....	6.7	18.3	1.9	0.7	1.6	2.0	3.1	4.2	5.4	10.1	31.1	116.6
2000.....	6.7	16.9	1.8	0.7	1.7	2.2	3.2	4.1	5.7	10.1	30.2	118.5
1999.....	7.1	15.0	1.9	0.8	1.7	2.3	3.3	4.2	5.6	11.1	33.6	130.7

Non-Hispanic 4/, male

2007.....	7.3	31.0	2.0	0.8	2.3	3.2	4.2	6.6	7.5	12.4	31.3	100.5
2006.....	7.5	28.6	1.7	0.8	2.6	3.6	4.8	6.9	7.8	12.8	33.6	102.2
2005.....	8.0	23.8	1.9	1.0	2.8	3.8	5.0	7.5	8.7	14.0	36.9	110.2
2004.....	7.4	24.6	2.1	1.0	2.5	3.5	4.7	6.7	7.6	12.1	34.5	113.1
2003.....	7.5	22.2	2.3	0.8	2.5	3.4	4.9	6.1	7.9	13.4	35.3	123.5
2002.....	7.5	21.0	2.0	0.9	2.6	3.4	4.9	6.1	8.1	13.7	36.6	120.4
2001.....	7.7	21.1	2.1	1.0	2.6	3.5	5.1	6.6	7.8	13.3	39.0	136.5
2000.....	7.7	19.0	2.2	1.0	2.7	3.8	5.3	6.7	8.4	13.2	37.5	136.9
1999.....	8.1	17.9	2.3	1.1	2.9	3.9	5.4	6.6	8.4	14.8	41.8	154.8

Non-Hispanic 4/, female

2007.....	5.1	25.0	1.3	0.4	0.4	0.7	1.0	1.7	2.8	6.1	20.8	82.8
2006.....	5.2	19.6	1.5	0.3	0.5	0.8	1.2	1.9	2.8	6.3	21.7	87.0
2005.....	5.7	19.9	1.3	0.4	0.7	0.8	1.3	2.0	3.2	7.3	24.2	92.0
2004.....	5.3	19.0	1.2	0.4	0.5	0.7	1.3	1.9	2.8	6.6	23.1	90.9
2003.....	5.4	16.0	1.3	0.3	0.6	0.7	1.3	1.9	2.7	6.8	23.1	97.0
2002.....	5.6	16.8	1.3	0.4	0.5	0.7	1.2	1.6	2.8	7.0	24.6	102.2
2001.....	5.8	15.4	1.6	0.4	0.6	0.6	1.2	2.0	3.2	7.4	25.9	108.4
2000.....	5.7	14.7	1.4	0.4	0.5	0.7	1.1	1.6	3.1	7.5	25.4	111.1
1999.....	6.2	12.0	1.5	0.5	0.5	0.7	1.2	1.8	3.1	8.1	28.2	121.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic white, both sexes

2007 5/.....	6.7	22.1	1.7	0.6	1.6	2.1	2.6	4.0	4.9	8.8	25.6	92.5
2006 5/.....	6.9	18.9	1.5	0.6	1.7	2.3	3.1	4.3	5.2	9.1	27.3	96.0
2005 5/.....	7.2	18.6	1.6	0.7	1.8	2.5	3.2	4.5	5.6	10.0	29.1	100.2
2004 5/.....	6.9	18.4	1.6	0.7	1.7	2.3	3.1	4.3	5.0	8.9	27.9	101.3
2003 5/.....	6.9	15.5	1.7	0.6	1.6	2.2	3.2	3.8	5.0	9.6	28.3	109.1
2002.....	7.0	14.5	1.5	0.7	1.7	2.3	3.1	3.8	5.2	9.6	29.4	110.9
2001.....	7.2	14.5	1.7	0.7	1.8	2.1	3.1	4.1	5.2	9.8	31.2	119.5
2000.....	7.1	13.7	1.7	0.7	1.7	2.3	3.2	3.9	5.4	9.7	30.3	122.0
1999.....	7.6	12.1	1.7	0.9	1.9	2.4	3.4	3.9	5.2	10.6	33.1	134.3

Non-Hispanic white, male

2007 5/.....	7.7	25.0	2.1	0.8	2.6	3.5	4.1	6.4	7.4	12.1	31.9	105.2
2006 5/.....	8.0	22.0	1.7	0.9	2.9	3.9	4.9	6.7	7.7	12.6	34.4	106.7
2005 5/.....	8.4	20.1	2.0	1.0	2.9	4.1	5.1	7.1	8.1	13.5	36.3	113.3
2004 5/.....	7.9	22.0	2.2	0.9	2.8	3.8	4.8	6.7	7.5	11.9	34.6	116.4
2003 5/.....	7.9	18.0	2.2	0.8	2.7	3.6	5.0	5.9	7.5	13.1	35.3	127.5
2002.....	7.9	17.0	1.8	0.9	2.9	3.7	5.1	6.0	7.7	13.0	36.3	123.1
2001.....	8.1	16.8	2.0	1.0	2.9	3.6	5.1	6.3	7.4	13.0	38.6	139.7
2000.....	8.0	16.0	2.1	1.1	2.9	4.0	5.3	6.3	7.9	12.6	36.8	139.7
1999.....	8.4	14.2	2.0	1.2	3.1	4.0	5.5	6.2	7.7	13.9	41.0	157.3

Non-Hispanic white, female

2007 5/.....	5.6	19.0	1.3	0.4	0.4	0.7	1.1	1.6	2.5	6.0	21.2	86.5
2006 5/.....	5.8	15.6	1.4	0.3	0.5	0.7	1.2	1.9	2.8	6.0	22.3	91.0
2005 5/.....	6.2	17.0	1.2	0.4	0.6	0.8	1.2	1.9	3.2	7.0	24.2	94.2
2004 5/.....	5.9	14.7	0.9	0.4	0.5	0.7	1.4	2.0	2.7	6.4	23.3	94.5
2003 5/.....	6.0	12.9	1.2	0.3	0.5	0.7	1.3	1.8	2.6	6.7	23.6	101.0
2002.....	6.1	11.9	1.2	0.4	0.5	0.8	1.2	1.5	2.7	6.7	24.8	105.7
2001.....	6.4	12.1	1.5	0.4	0.7	0.6	1.1	1.9	3.1	7.1	26.3	111.2
2000.....	6.3	11.2	1.3	0.4	0.5	0.7	1.1	1.6	3.1	7.3	26.0	114.9
1999.....	6.8	9.8	1.5	0.5	0.6	0.7	1.3	1.7	2.8	7.9	28.0	125.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic black, both sexes

2007 5/.....	5.2	53.1	1.8	0.5	0.8	1.8	3.3	5.3	7.3	13.0	24.1	61.6
2006 5/.....	5.1	46.4	2.2	0.7	1.1	1.9	3.1	5.5	6.8	12.9	24.9	62.6
2005 5/.....	6.2	38.8	1.7	0.7	1.6	2.1	3.4	7.1	8.8	15.8	38.4	87.4
2004 5/.....	4.9	38.0	1.9	0.6	1.0	2.0	3.0	4.8	6.6	11.9	31.3	71.4
2003 5/.....	5.1	35.7	2.1	0.5	1.3	1.8	3.4	5.5	7.5	13.1	27.8	74.1
2002.....	5.4	38.6	2.5	0.9	1.2	1.7	3.2	4.8	7.4	15.6	32.9	84.4
2001.....	5.5	35.5	2.6	0.6	1.1	2.0	3.3	5.7	8.0	14.2	34.2	96.5
2000.....	5.6	31.7	2.3	0.7	1.6	2.3	3.6	5.8	8.4	15.1	34.1	90.9
1999.....	6.3	29.5	2.6	0.7	1.3	2.5	3.5	6.7	10.2	18.2	43.1	106.7

Non-Hispanic black, male

2007 5/.....	6.4	54.1	2.2	*	1.0	2.8	5.6	8.8	10.9	18.5	31.1	68.1
2006 5/.....	6.6	55.5	2.1	0.9	1.7	2.8	5.4	9.6	10.3	18.2	32.0	76.0
2005 5/.....	7.9	42.6	1.8	0.8	2.4	3.1	5.3	11.7	15.2	21.8	54.4	99.0
2004 5/.....	6.3	38.7	1.7	1.1	1.5	3.5	4.9	7.8	10.5	16.7	43.4	89.4
2003 5/.....	6.6	41.6	2.9	0.7	1.8	3.0	5.3	8.7	12.2	18.8	39.8	92.7
2002.....	6.8	40.0	3.3	1.2	1.8	2.9	4.8	7.6	12.1	22.1	46.6	107.2
2001.....	7.0	41.8	2.9	0.9	1.6	3.3	5.2	9.2	12.2	19.9	49.5	115.5
2000.....	7.5	32.8	2.7	1.0	2.6	3.8	6.1	10.3	14.2	20.8	53.7	124.5
1999.....	8.3	35.0	3.5	1.0	2.4	4.1	5.9	11.2	16.0	27.4	57.8	155.7

Non-Hispanic black, female

2007 5/.....	4.0	52.0	*	*	*	0.8	1.2	2.4	4.4	9.0	20.2	59.0
2006 5/.....	3.7	37.0	2.3	*	*	1.0	1.1	2.1	4.0	9.1	20.8	57.2
2005 5/.....	4.7	34.7	1.6	*	0.7	1.1	1.8	3.2	3.7	11.6	29.4	82.7
2004 5/.....	3.7	37.3	2.0	*	*	*	1.3	2.2	3.5	8.4	24.5	64.3
2003 5/.....	3.7	29.7	*	*	0.8	0.8	1.6	2.8	3.7	9.0	21.0	66.9
2002.....	4.2	37.2	*	*	*	*	1.8	2.4	3.6	11.0	25.2	75.7
2001.....	4.2	28.9	2.3	*	*	0.7	1.6	2.7	4.7	10.3	25.6	89.4
2000.....	3.9	30.7	2.0	*	*	0.9	1.3	1.9	3.7	11.1	23.1	78.5
1999.....	4.5	23.9	*	*	*	1.1	1.4	2.8	5.5	11.8	34.8	88.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

All origins 3/, both sexes

2007.....	11.5	...	...	0.5	9.7	13.0	15.6	17.7	15.5	12.6	16.3	15.6
2006.....	11.1	...	...	0.5	9.9	12.3	15.1	17.2	14.5	12.6	15.9	15.9
2005.....	11.0	...	...	0.7	10.0	12.4	14.9	16.5	13.9	12.6	16.9	16.9
2004.....	11.0	...	...	0.7	10.3	12.7	15.0	16.6	13.8	12.3	16.3	16.4
2003.....	10.8	...	...	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002.....	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 6/.....	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000.....	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999.....	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3

All origins 3/, male

2007.....	18.3	...	...	0.6	15.9	20.7	23.8	27.0	24.3	22.5	34.3	41.8
2006.....	17.8	...	...	0.7	16.2	19.7	23.2	26.2	22.7	22.7	33.3	43.2
2005.....	17.7	...	...	1.0	16.2	19.9	23.1	25.2	22.2	22.7	35.8	45.0
2004.....	17.7	...	...	0.9	16.8	20.4	23.0	24.8	22.1	22.6	34.8	45.0
2003.....	17.6	...	...	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002.....	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 6/.....	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000.....	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999.....	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1

All origins 3/, female

2007.....	4.8	...	...	0.3	3.2	5.0	7.3	8.8	7.3	4.2	3.8	3.1
2006.....	4.6	...	...	0.3	3.2	4.7	7.0	8.4	6.8	4.1	4.0	3.1
2005.....	4.5	...	...	0.3	3.5	4.7	6.8	8.0	6.1	4.0	4.0	4.0
2004.....	4.6	...	...	0.5	3.6	4.7	7.1	8.6	6.1	3.8	3.9	3.6
2003.....	4.3	...	...	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002.....	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 6/.....	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000.....	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999.....	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Hispanic, both sexes

2007.....	5.4	...	...	*	7.1	7.9	7.3	8.2	7.1	6.9	8.5	7.9
2006.....	4.9	...	...	0.3	7.4	7.1	6.6	6.5	6.0	5.2	7.6	*
2005.....	5.1	...	...	0.5	7.7	6.8	7.4	6.6	6.0	5.9	8.8	9.0
2004.....	5.3	...	...	0.5	8.0	7.2	6.8	7.5	7.0	6.8	9.1	*
2003.....	5.0	...	...	0.6	7.0	6.8	6.8	7.2	6.9	6.5	8.0	10.9
2002.....	5.0	...	...	0.5	6.7	6.6	7.0	7.1	7.2	7.5	9.2	12.7
2001 6/.....	5.0	...	...	0.5	6.2	7.1	7.3	7.0	6.2	7.6	8.8	13.8
2000.....	5.1	...	...	0.4	6.8	6.5	7.5	7.8	6.0	7.8	11.7	*
1999.....	5.0	...	...	0.5	6.2	6.7	7.5	7.1	6.8	7.0	13.1	15.4

Hispanic, male

2007.....	8.8	...	...	*	11.5	12.2	11.5	13.1	12.5	13.7	19.3	*
2006.....	7.9	...	...	*	11.6	10.9	10.6	10.1	10.5	9.1	16.1	*
2005.....	8.3	...	...	0.7	12.1	10.7	11.8	11.0	10.2	10.7	18.7	*
2004.....	8.6	...	...	0.6	12.8	11.2	10.8	12.0	11.3	13.6	19.7	*
2003.....	8.3	...	...	0.8	11.2	10.8	11.0	11.5	13.0	13.1	18.0	*
2002.....	8.3	...	...	0.6	10.6	10.7	11.2	11.3	13.2	14.7	19.3	35.1
2001 6/.....	8.3	...	...	0.7	9.5	11.3	12.4	11.4	11.5	15.4	20.1	41.9
2000.....	8.4	...	...	0.6	10.9	10.6	12.1	12.5	11.1	15.6	26.1	*
1999.....	8.2	...	...	0.6	9.9	10.7	12.1	11.6	12.2	13.8	28.2	*

Hispanic, female

2007.....	1.8	...	...	*	2.2	2.8	2.6	3.2	2.1	*	*	*
2006.....	1.7	...	...	*	2.6	2.4	2.2	2.8	1.9	*	*	*
2005.....	1.7	...	...	*	2.7	2.1	2.4	2.2	2.1	*	*	*
2004.....	1.8	...	...	*	2.5	2.3	2.4	3.1	3.1	*	*	*
2003.....	1.5	...	...	*	2.2	1.9	2.2	2.9	*	*	*	*
2002.....	1.6	...	...	*	2.1	1.7	2.4	2.9	*	*	*	*
2001 6/.....	1.5	...	...	*	2.3	2.2	1.8	2.7	*	*	*	*
2000.....	1.5	...	...	*	2.0	1.7	2.6	3.2	*	*	*	*
1999.....	1.6	...	...	*	1.8	2.0	2.6	2.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007.....	12.5	...	...	0.5	10.3	14.3	17.1	18.8	16.2	13.0	16.7	15.9
2006.....	12.2	...	...	0.6	10.4	13.6	16.6	18.3	15.2	13.1	16.4	16.3
2005.....	12.0	...	...	0.7	10.5	13.8	16.2	17.5	14.5	13.1	17.3	17.2
2004.....	11.9	...	...	0.7	10.8	13.9	16.3	17.5	14.3	12.7	16.7	16.8
2003.....	11.7	...	...	0.6	10.2	14.0	16.0	16.7	14.3	13.1	16.9	17.1
2002.....	11.8	...	...	0.7	10.5	13.9	16.4	16.5	14.0	13.8	18.1	18.1
2001 6/.....	11.6	...	...	0.7	10.7	13.9	15.7	15.9	13.6	13.6	17.8	17.5
2000.....	11.1	...	...	0.8	10.8	13.0	15.4	15.0	12.5	12.7	17.9	19.9
1999.....	11.2	...	...	0.6	10.8	13.8	15.1	14.4	12.5	13.7	18.3	19.3

Non-Hispanic 4/, male

2007.....	20.1	...	...	0.7	16.9	23.0	26.3	28.7	25.2	23.1	35.2	43.1
2006.....	19.6	...	...	0.8	17.1	22.1	25.5	28.0	23.8	23.7	34.3	44.4
2005.....	19.3	...	...	1.1	17.0	22.3	25.0	26.8	23.1	23.6	36.6	45.9
2004.....	19.2	...	...	1.0	17.5	22.7	25.0	26.1	22.8	23.1	35.6	46.2
2003.....	19.1	...	...	0.9	16.9	22.9	25.1	25.6	22.9	24.1	36.0	48.6
2002.....	19.4	...	...	1.0	17.6	22.7	25.5	25.6	22.8	25.2	39.0	50.9
2001 6/.....	19.0	...	...	1.1	18.1	23.1	24.6	24.5	21.7	25.0	38.5	51.3
2000.....	18.3	...	...	1.3	18.3	21.4	24.1	23.2	19.9	23.1	39.1	58.4
1999.....	18.4	...	...	1.0	18.2	22.7	23.6	22.2	20.2	25.2	39.1	57.6

Non-Hispanic 4/, female

2007.....	5.3	...	...	0.3	3.4	5.5	8.1	9.4	7.8	4.4	4.0	3.1
2006.....	5.1	...	...	0.4	3.3	5.2	7.8	9.0	7.3	4.2	4.2	3.2
2005.....	4.9	...	...	0.4	3.6	5.3	7.5	8.6	6.4	4.1	4.1	4.1
2004.....	5.0	...	...	0.5	3.7	5.2	7.8	9.2	6.4	3.9	4.0	3.6
2003.....	4.6	...	...	0.3	3.2	5.2	7.2	8.1	6.3	4.0	4.1	3.3
2002.....	4.6	...	...	0.3	3.1	5.2	7.4	7.8	5.9	4.3	4.3	3.9
2001 6/.....	4.5	...	...	0.3	3.0	4.9	7.0	7.6	6.1	4.1	4.2	3.5
2000.....	4.3	...	...	0.3	3.1	4.8	6.8	7.0	5.7	4.1	4.1	4.3
1999.....	4.3	...	...	0.2	3.2	5.0	6.8	6.9	5.4	4.3	4.8	4.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Non-Hispanic white, both sexes

2007 5/.....	14.4	...	...	0.5	11.1	16.1	19.7	21.7	18.2	14.5	18.1	17.3
2006 5/.....	13.9	...	...	0.6	11.2	15.3	19.1	21.0	17.1	14.5	17.8	17.4
2005 5/.....	13.6	...	...	0.7	11.4	15.3	18.5	20.0	16.2	14.4	18.7	18.3
2004 5/.....	13.5	...	...	0.7	11.6	15.4	18.7	19.9	15.8	13.8	17.9	17.9
2003 5/.....	13.3	...	...	0.6	10.9	15.5	18.3	19.1	15.8	14.3	18.2	18.4
2002.....	13.4	...	...	0.7	11.5	15.4	18.7	18.7	15.6	15.1	19.4	19.2
2001 6/.....	13.0	...	...	0.7	11.6	15.4	17.7	17.9	15.0	14.7	19.0	18.7
2000.....	12.5	...	...	0.9	11.6	14.2	17.2	16.7	13.9	13.8	19.0	21.3
1999.....	12.4	...	...	0.7	11.5	15.0	16.9	16.0	13.7	14.8	19.5	20.8

Non-Hispanic white, male

2007 5/.....	22.9	...	...	0.6	18.2	25.8	29.9	32.6	28.1	25.4	38.0	46.9
2006 5/.....	22.3	...	...	0.8	18.5	24.4	29.0	31.7	26.3	25.9	37.2	47.6
2005 5/.....	22.0	...	...	1.1	18.4	24.5	28.3	30.2	25.7	26.0	39.5	49.4
2004 5/.....	21.6	...	...	0.9	19.0	24.9	28.4	29.3	24.9	24.9	38.0	49.8
2003 5/.....	21.6	...	...	1.0	18.2	25.0	28.2	29.0	25.1	26.0	38.6	52.5
2002.....	21.9	...	...	1.1	19.3	24.6	28.7	28.6	25.0	27.5	41.7	54.3
2001 6/.....	21.4	...	...	1.1	19.6	25.2	27.4	27.4	23.6	26.9	41.1	55.2
2000.....	20.4	...	...	1.4	19.5	23.1	26.6	25.5	21.8	24.7	41.7	62.7
1999.....	20.3	...	...	1.1	19.2	24.3	26.0	24.4	21.9	27.0	41.6	62.2

Non-Hispanic white, female

2007 5/.....	6.1	...	...	0.3	3.7	6.2	9.5	10.9	8.9	4.9	4.1	3.3
2006 5/.....	5.9	...	...	0.4	3.5	6.1	9.2	10.5	8.3	4.5	4.4	3.4
2005 5/.....	5.6	...	...	0.3	3.9	5.9	8.7	9.9	7.2	4.4	4.3	4.0
2004 5/.....	5.7	...	...	0.5	4.0	5.6	9.0	10.6	7.1	4.3	4.2	3.6
2003 5/.....	5.3	...	...	0.2	3.3	5.8	8.3	9.3	7.0	4.3	4.4	3.4
2002.....	5.3	...	...	0.3	3.4	6.1	8.6	9.0	6.6	4.6	4.5	4.1
2001 6/.....	5.0	...	...	0.3	3.3	5.5	8.1	8.6	6.8	4.4	4.4	3.5
2000.....	4.9	...	...	0.4	3.3	5.2	7.8	7.9	6.4	4.6	4.2	4.6
1999.....	4.9	...	...	0.3	3.4	5.5	7.8	7.8	6.0	4.6	5.0	4.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Non-Hispanic black, both sexes

2007 5/.....	5.0	...	...	0.7	6.2	8.7	7.7	6.2	4.7	4.0	4.6	*
2006 5/.....	5.1	...	...	0.6	6.4	8.7	7.5	6.0	4.9	4.2	4.4	5.8
2005 5/.....	5.2	...	...	0.7	6.8	9.0	7.6	6.3	4.7	4.5	5.4	*
2004 5/.....	5.3	...	...	0.8	7.3	9.7	6.7	5.9	5.8	4.5	6.2	*
2003 5/.....	5.3	...	...	0.7	7.3	9.6	7.4	5.7	4.9	4.6	4.5	*
2002.....	5.2	...	...	0.5	6.6	9.6	7.7	5.9	4.9	4.9	5.4	*
2001 6/.....	5.3	...	...	0.6	7.3	9.2	7.5	6.2	5.1	5.5	5.6	*
2000.....	5.5	...	...	0.6	8.4	9.0	7.7	6.5	4.2	5.3	5.9	*
1999.....	5.5	...	...	0.5	8.3	9.5	7.8	5.7	5.0	5.8	5.8	*

Non-Hispanic black, male

2007 5/.....	8.6	...	...	0.9	10.5	15.3	13.0	10.2	8.3	8.6	11.0	*
2006 5/.....	9.1	...	...	0.9	10.9	16.1	13.4	10.6	9.1	8.8	11.7	*
2005 5/.....	8.9	...	...	0.8	11.9	15.5	13.0	10.5	8.0	8.4	12.6	*
2004 5/.....	9.2	...	...	0.9	12.4	16.8	11.4	9.7	11.2	10.1	15.4	*
2003 5/.....	9.0	...	...	1.0	12.5	17.1	12.5	9.3	8.5	8.5	11.3	*
2002.....	9.3	...	...	0.9	11.6	18.0	13.2	10.2	8.5	9.7	13.5	*
2001 6/.....	9.3	...	...	0.9	13.4	16.6	12.5	9.8	9.3	10.8	13.2	*
2000.....	9.6	...	...	1.0	14.6	16.1	13.3	11.1	7.5	11.4	12.3	*
1999.....	9.9	...	...	0.7	14.8	17.5	13.7	10.1	9.2	11.7	14.1	*

Non-Hispanic black, female

2007 5/.....	1.7	...	...	*	1.7	2.6	3.0	2.7	1.8	*	*	*
2006 5/.....	1.4	...	...	*	1.8	1.9	2.2	2.2	1.6	*	*	*
2005 5/.....	1.8	...	...	*	1.7	2.9	2.8	2.7	2.1	*	*	*
2004 5/.....	1.8	...	...	*	2.2	3.3	2.6	2.5	1.5	*	*	*
2003 5/.....	1.8	...	...	*	2.0	2.7	2.9	2.7	2.0	*	*	*
2002.....	1.6	...	...	*	1.7	2.1	2.9	2.2	2.0	*	*	*
2001 6/.....	1.7	...	...	*	1.3	2.4	3.0	3.1	1.8	*	*	*
2000.....	1.7	...	...	*	2.3	2.6	2.7	2.5	1.6	*	*	*
1999.....	1.6	...	...	*	2.0	2.4	2.6	2.0	1.6	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes

2007.....	5.8	...	...	0.1	4.5	5.7	6.7	8.0	8.5	8.8	12.2	11.0
2006.....	5.6	...	...	0.2	4.7	5.3	6.5	8.0	8.1	8.9	12.0	10.9
2005.....	5.7	...	...	0.2	4.7	5.7	6.5	8.2	8.1	9.0	12.6	11.2
2004.....	5.7	...	...	0.1	5.0	5.7	6.5	8.0	8.0	8.8	12.2	11.1
2003.....	5.8	...	...	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002.....	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001.....	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000.....	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999.....	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6

All origins 3/, male

2007.....	10.2	...	...	0.2	7.9	9.8	11.1	13.8	15.2	17.3	27.9	32.9
2006.....	10.0	...	...	0.3	8.2	9.2	10.8	13.7	14.3	17.6	27.7	32.7
2005.....	10.2	...	...	0.3	8.2	9.7	11.0	14.1	14.6	17.6	29.3	33.8
2004.....	10.0	...	...	0.2	8.8	9.7	10.8	13.5	14.2	17.5	28.3	34.4
2003.....	10.4	...	...	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002.....	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001.....	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000.....	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999.....	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6

All origins 3/, female

2007.....	1.4	...	...	*	0.8	1.4	2.3	2.5	2.3	1.5	1.4	0.6
2006.....	1.4	...	...	*	0.9	1.3	2.1	2.6	2.3	1.6	1.2	0.7
2005.....	1.4	...	...	*	0.9	1.5	2.0	2.4	2.1	1.7	1.3	0.8
2004.....	1.5	...	...	*	1.1	1.5	2.2	2.8	2.2	1.6	1.4	*
2003.....	1.4	...	...	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002.....	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001.....	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000.....	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999.....	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Hispanic, both sexes

2007.....	2.0	...	...	*	2.8	2.9	2.5	2.9	2.6	3.5	5.4	*
2006.....	1.8	...	...	*	3.1	2.6	2.2	2.1	2.2	2.0	4.3	*
2005.....	1.9	...	...	*	3.1	2.6	2.6	2.3	2.0	2.5	4.5	*
2004.....	2.1	...	...	*	3.7	2.7	2.4	3.0	2.7	3.5	5.0	*
2003.....	2.1	...	...	*	3.3	2.8	2.5	3.0	2.7	3.1	4.1	*
2002.....	2.2	...	...	*	3.3	2.7	2.5	2.7	3.4	4.4	4.7	*
2001.....	2.2	...	...	*	2.9	3.3	2.5	3.0	3.2	4.0	4.7	*
2000.....	2.3	...	...	*	3.4	2.8	2.8	4.0	2.7	4.6	5.3	*
1999.....	2.3	...	...	*	3.5	2.9	3.0	3.2	2.9	3.6	8.1	*

Hispanic, male

2007.....	3.6	...	...	*	5.0	4.9	4.2	4.9	5.2	7.5	12.7	*
2006.....	3.3	...	...	*	5.5	4.2	3.9	3.5	4.3	4.4	10.1	*
2005.....	3.4	...	...	*	5.4	4.3	4.4	4.3	4.1	5.2	11.0	*
2004.....	3.7	...	...	*	6.5	4.3	4.1	5.3	5.0	7.1	11.4	*
2003.....	3.8	...	...	*	5.6	4.8	4.3	5.3	5.6	7.0	9.6	*
2002.....	3.8	...	...	*	5.7	4.6	4.2	4.8	6.6	9.4	11.5	*
2001.....	3.8	...	...	*	4.8	5.5	4.5	5.3	6.6	8.8	11.6	*
2000.....	4.1	...	...	*	5.8	4.7	4.7	6.8	5.7	9.7	12.8	*
1999.....	4.0	...	...	*	5.8	4.8	5.0	5.5	5.9	7.7	19.8	*

Hispanic, female

2007.....	0.3	...	...	*	*	*	*	*	*	*	*	*
2006.....	0.3	...	...	*	*	0.6	*	*	*	*	*	*
2005.....	0.3	...	...	*	*	0.6	*	*	*	*	*	*
2004.....	0.4	...	...	*	0.6	0.7	*	*	*	*	*	*
2003.....	0.3	...	...	*	*	*	*	*	*	*	*	*
2002.....	0.4	...	...	*	*	*	*	*	*	*	*	*
2001.....	0.4	...	...	*	0.7	0.7	*	*	*	*	*	*
2000.....	0.4	...	...	*	*	*	*	*	*	*	*	*
1999.....	0.5	...	...	*	0.8	*	0.9	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic 4/, both sexes

2007.....	6.4	...	...	0.2	4.8	6.4	7.4	8.7	9.0	9.2	12.6	11.3
2006.....	6.3	...	...	0.2	5.0	6.0	7.2	8.7	8.6	9.4	12.4	11.2
2005.....	6.4	...	...	0.2	5.0	6.4	7.2	8.8	8.6	9.4	13.1	11.4
2004.....	6.3	...	...	0.2	5.3	6.4	7.1	8.6	8.4	9.2	12.6	11.4
2003.....	6.4	...	...	0.2	5.4	6.7	7.2	8.5	8.7	9.7	12.8	12.1
2002.....	6.5	...	...	0.2	5.5	6.7	7.7	8.3	8.8	10.0	13.6	12.1
2001.....	6.5	...	...	0.2	5.8	7.1	7.3	8.1	8.6	9.9	13.5	12.4
2000.....	6.4	...	...	0.3	6.2	6.8	7.6	7.9	7.9	9.5	13.4	13.4
1999.....	6.4	...	...	0.3	6.5	7.2	7.3	7.7	8.2	10.1	13.3	12.7

Non-Hispanic 4/, male

2007.....	11.4	...	...	0.3	8.5	11.2	12.4	14.9	16.0	18.0	28.9	34.1
2006.....	11.2	...	...	0.3	8.8	10.5	12.1	14.8	15.2	18.5	28.8	33.8
2005.....	11.4	...	...	0.4	8.8	11.1	12.2	15.2	15.4	18.5	30.3	34.7
2004.....	11.1	...	...	0.3	9.3	11.1	11.9	14.4	15.0	18.2	29.3	35.6
2003.....	11.5	...	...	0.3	9.6	11.7	12.2	14.7	15.6	19.2	29.7	37.5
2002.....	11.7	...	...	0.4	9.8	11.6	13.1	14.4	15.7	20.1	31.9	38.4
2001.....	11.6	...	...	0.4	10.4	12.5	12.3	14.0	15.1	19.8	31.5	39.7
2000.....	11.4	...	...	0.5	11.1	11.7	12.7	13.5	14.0	18.9	31.6	43.8
1999.....	11.5	...	...	0.4	11.5	12.5	12.3	13.1	14.5	20.1	31.5	42.2

Non-Hispanic 4/, female

2007.....	1.6	...	...	*	0.9	1.6	2.6	2.7	2.5	1.6	1.5	0.6
2006.....	1.6	...	...	*	1.0	1.5	2.4	2.8	2.5	1.7	1.2	0.7
2005.....	1.6	...	...	*	1.0	1.6	2.2	2.7	2.3	1.8	1.4	0.8
2004.....	1.7	...	...	*	1.1	1.7	2.5	3.0	2.3	1.6	1.5	*
2003.....	1.6	...	...	*	1.0	1.6	2.3	2.6	2.3	1.7	1.6	0.9
2002.....	1.6	...	...	*	1.1	1.9	2.4	2.4	2.3	1.6	1.4	0.8
2001.....	1.6	...	...	0.1	1.1	1.8	2.4	2.5	2.5	1.8	1.6	1.0
2000.....	1.6	...	...	0.1	1.3	1.9	2.6	2.4	2.3	1.8	1.5	1.1
1999.....	1.6	...	...	*	1.3	1.9	2.4	2.4	2.3	1.8	1.5	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic white, both sexes

2007 5/.....	7.5	...	...	0.2	5.2	7.3	8.8	10.1	10.3	10.4	13.9	12.5
2006 5/.....	7.3	...	...	0.2	5.4	6.7	8.4	10.2	9.8	10.6	13.7	12.0
2005 5/.....	7.4	...	...	0.3	5.4	7.2	8.3	10.2	9.8	10.6	14.3	12.5
2004 5/.....	7.3	...	...	0.2	5.8	7.1	8.3	9.9	9.5	10.2	13.7	12.5
2003 5/.....	7.4	...	...	0.2	5.8	7.4	8.4	9.9	9.9	10.8	14.0	13.2
2002.....	7.5	...	...	0.2	5.9	7.5	8.8	9.5	9.9	11.2	14.7	13.1
2001.....	7.4	...	...	0.3	6.2	8.0	8.3	9.3	9.6	10.9	14.5	13.4
2000.....	7.3	...	...	0.4	6.7	7.4	8.6	8.9	8.7	10.4	14.5	14.5
1999.....	7.3	...	...	0.3	6.8	7.9	8.3	8.6	9.0	11.0	14.3	13.7

Non-Hispanic white, male

2007 5/.....	13.3	...	...	0.3	9.0	12.6	14.5	17.1	18.0	20.1	31.5	37.7
2006 5/.....	13.0	...	...	0.3	9.5	11.6	13.9	17.1	17.0	20.6	31.4	36.3
2005 5/.....	13.2	...	...	0.5	9.4	12.3	13.9	17.3	17.4	20.6	32.9	37.9
2004 5/.....	12.8	...	...	0.3	10.1	12.1	13.7	16.5	16.7	19.9	31.5	39.0
2003 5/.....	13.2	...	...	0.4	10.3	12.7	14.1	16.8	17.4	21.2	32.1	41.0
2002.....	13.4	...	...	0.4	10.5	12.6	14.9	16.3	17.4	22.2	34.3	41.3
2001.....	13.2	...	...	0.4	11.0	13.8	13.9	15.9	16.7	21.6	33.8	43.0
2000.....	12.9	...	...	0.6	11.8	12.7	14.3	15.1	15.3	20.4	34.2	47.6
1999.....	12.9	...	...	0.5	12.0	13.4	13.7	14.6	15.9	21.8	33.8	45.7

Non-Hispanic white, female

2007 5/.....	1.9	...	...	*	1.0	1.9	3.1	3.2	2.9	1.9	1.6	0.6
2006 5/.....	1.9	...	...	*	1.1	1.7	3.0	3.4	2.9	2.0	1.4	0.7
2005 5/.....	1.8	...	...	*	1.1	2.0	2.7	3.2	2.6	1.9	1.5	0.9
2004 5/.....	1.9	...	...	*	1.3	2.0	3.0	3.5	2.7	1.9	1.6	*
2003 5/.....	1.8	...	...	*	1.1	2.0	2.8	3.1	2.7	1.9	1.7	1.0
2002.....	1.8	...	...	*	1.2	2.3	2.8	2.8	2.7	1.8	1.6	0.9
2001.....	1.9	...	...	*	1.2	2.1	2.7	2.9	2.9	1.9	1.7	1.1
2000.....	1.9	...	...	*	1.4	2.1	3.0	2.8	2.6	2.0	1.5	1.2
1999.....	1.9	...	...	*	1.4	2.3	2.9	2.7	2.6	2.0	1.6	1.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic black, both sexes

2007 5/.....	2.6	...	...	*	3.6	4.6	3.3	2.8	2.5	2.6	3.8	*
2006 5/.....	2.6	...	...	*	3.8	4.8	3.3	2.5	2.5	2.9	3.8	*
2005 5/.....	2.7	...	...	*	3.9	4.7	3.4	2.8	2.4	3.3	4.0	*
2004 5/.....	2.7	...	...	*	4.1	5.0	2.8	2.4	3.1	3.6	4.9	*
2003 5/.....	2.7	...	...	*	4.4	5.4	2.7	2.5	2.8	3.4	4.0	*
2002.....	2.9	...	...	*	4.2	5.3	3.6	2.9	2.8	3.3	4.3	*
2001.....	3.0	...	...	*	5.1	5.0	3.5	2.8	2.7	3.8	5.1	*
2000.....	3.1	...	...	*	5.4	5.2	3.5	3.2	3.0	3.8	3.6	*
1999.....	3.2	...	...	*	5.8	5.1	3.6	3.0	3.5	4.0	4.8	*

Non-Hispanic black, male

2007 5/.....	4.8	...	...	*	6.7	8.4	6.0	5.0	5.1	5.9	9.4	*
2006 5/.....	5.0	...	...	*	6.9	9.2	6.3	4.8	5.1	6.4	10.3	*
2005 5/.....	5.1	...	...	*	7.2	8.8	6.6	5.4	4.6	6.4	10.4	*
2004 5/.....	5.1	...	...	*	7.2	9.5	5.3	4.5	6.0	8.6	13.1	*
2003 5/.....	5.1	...	...	*	8.0	10.5	5.1	4.8	5.3	6.8	9.8	*
2002.....	5.5	...	...	*	7.7	10.5	6.5	5.5	5.5	7.0	11.4	*
2001.....	5.6	...	...	*	9.6	9.5	6.4	4.9	5.5	7.9	12.0	*
2000.....	5.7	...	...	*	9.9	9.6	6.4	5.6	5.7	9.0	8.3	*
1999.....	6.1	...	...	*	10.6	10.0	6.9	5.6	7.1	8.9	12.2	*

Non-Hispanic black, female

2007 5/.....	0.5	...	...	*	*	1.0	0.8	0.8	*	*	*	*
2006 5/.....	0.5	...	...	*	0.7	0.8	*	*	*	*	*	*
2005 5/.....	0.5	...	...	*	*	0.8	*	*	*	*	*	*
2004 5/.....	0.5	...	...	*	0.9	1.0	*	*	*	*	*	*
2003 5/.....	0.5	...	...	*	0.7	*	*	*	*	*	*	*
2002.....	0.5	...	...	*	0.7	*	1.0	*	*	*	*	*
2001.....	0.6	...	...	*	*	0.9	1.0	1.0	*	*	*	*
2000.....	0.7	...	...	*	1.0	1.1	0.8	1.1	*	*	*	*
1999.....	0.6	...	...	*	1.0	0.8	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes

2007.....	5.7	...	...	0.3	5.3	7.3	8.9	9.7	7.0	3.8	4.1	4.6
2006.....	5.5	...	...	0.4	5.2	7.0	8.6	9.1	6.4	3.7	3.9	5.0
2005.....	5.3	...	...	0.5	5.3	6.8	8.4	8.3	5.7	3.6	4.2	5.7
2004.....	5.3	...	...	0.6	5.3	7.0	8.5	8.5	5.8	3.5	4.1	5.4
2003.....	5.0	...	...	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
2002.....	5.0	...	...	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 6/.....	4.8	...	...	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000.....	4.5	...	...	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999.....	4.5	...	...	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7

All origins 3/, male

2007.....	8.1	...	...	0.4	8.0	10.9	12.8	13.2	9.1	5.2	6.4	8.8
2006.....	7.8	...	...	0.5	7.9	10.5	12.4	12.5	8.4	5.1	5.6	10.5
2005.....	7.5	...	...	0.6	8.0	10.2	12.1	11.1	7.6	5.1	6.5	11.2
2004.....	7.6	...	...	0.7	8.0	10.6	12.2	11.3	7.8	5.1	6.5	10.7
2003.....	7.3	...	...	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
2002.....	7.3	...	...	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 6/.....	7.1	...	...	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000.....	6.6	...	...	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999.....	6.6	...	...	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5

All origins 3/, female

2007.....	3.4	...	...	0.2	2.4	3.6	5.0	6.3	5.0	2.7	2.5	2.5
2006.....	3.2	...	...	0.3	2.3	3.4	4.9	5.8	4.5	2.4	2.8	2.4
2005.....	3.1	...	...	0.3	2.6	3.2	4.8	5.5	4.0	2.3	2.7	3.1
2004.....	3.1	...	...	0.4	2.5	3.2	4.9	5.9	3.9	2.2	2.5	3.0
2003.....	2.8	...	...	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
2002.....	2.9	...	...	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 6/.....	2.6	...	...	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000.....	2.5	...	...	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999.....	2.5	...	...	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Hispanic, both sexes

2007.....	3.4	...	...	*	4.2	5.1	4.8	5.4	4.5	3.4	3.1	*
2006.....	3.1	...	...	*	4.3	4.5	4.4	4.4	3.8	3.1	3.3	*
2005.....	3.2	...	...	0.4	4.6	4.2	4.8	4.3	4.0	3.4	4.3	*
2004.....	3.2	...	...	0.5	4.3	4.5	4.4	4.5	4.3	3.3	4.1	*
2003.....	2.9	...	...	0.5	3.8	3.9	4.3	4.2	4.2	3.4	3.9	*
2002.....	2.9	...	...	0.4	3.4	3.9	4.5	4.4	3.8	3.1	4.5	*
2001 6/.....	2.8	...	...	0.5	3.3	3.9	4.8	4.0	3.0	3.6	4.1	*
2000.....	2.8	...	...	0.3	3.4	3.7	4.7	3.8	3.2	3.3	6.3	*
1999.....	2.7	...	...	*	2.7	3.8	4.5	3.8	3.9	3.5	5.0	*

Hispanic, male

2007.....	5.2	...	...	*	6.6	7.3	7.3	8.2	7.3	6.2	6.7	*
2006.....	4.6	...	...	*	6.0	6.7	6.6	6.6	6.2	4.7	*	*
2005.....	4.9	...	...	0.6	6.7	6.5	7.4	6.7	6.1	5.4	7.7	*
2004.....	4.9	...	...	0.6	6.4	6.9	6.8	6.6	6.3	6.5	8.2	*
2003.....	4.5	...	...	0.6	5.6	6.0	6.7	6.2	7.5	6.1	8.4	*
2002.....	4.4	...	...	*	4.9	6.1	7.0	6.5	6.5	5.4	*	*
2001 6/.....	4.4	...	...	0.6	4.7	5.8	7.9	6.1	4.9	6.6	*	*
2000.....	4.3	...	...	*	5.1	5.9	7.3	5.7	5.4	5.9	13.3	*
1999.....	4.2	...	...	*	4.2	5.9	7.1	6.0	6.4	6.2	*	*

Hispanic, female

2007.....	1.4	...	...	*	1.7	2.3	2.0	2.4	1.8	*	*	*
2006.....	1.4	...	...	*	2.2	1.9	1.9	2.2	1.6	*	*	*
2005.....	1.4	...	...	*	2.2	1.5	1.9	1.8	2.0	*	*	*
2004.....	1.4	...	...	*	1.9	1.5	1.9	2.4	2.5	*	*	*
2003.....	1.2	...	...	*	1.6	1.5	1.7	2.3	*	*	*	*
2002.....	1.2	...	...	*	1.6	1.3	1.8	2.4	*	*	*	*
2001 6/.....	1.2	...	...	*	1.6	1.6	1.4	2.0	*	*	*	*
2000.....	1.1	...	...	*	1.4	1.1	1.9	2.0	*	*	*	*
1999.....	1.1	...	...	*	1.0	1.4	1.7	1.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007.....	6.1	...	...	0.4	5.5	7.9	9.7	10.2	7.2	3.9	4.1	4.6
2006.....	5.9	...	...	0.4	5.4	7.6	9.4	9.6	6.6	3.7	4.0	5.1
2005.....	5.6	...	...	0.5	5.5	7.4	9.0	8.7	5.8	3.6	4.2	5.8
2004.....	5.7	...	...	0.6	5.5	7.5	9.2	8.9	5.9	3.5	4.1	5.4
2003.....	5.3	...	...	0.4	4.8	7.3	8.8	8.2	5.5	3.4	4.0	5.1
2002.....	5.3	...	...	0.4	5.0	7.2	8.7	8.2	5.3	3.8	4.5	5.9
2001 6/.....	5.1	...	...	0.5	4.8	6.8	8.4	7.7	5.0	3.7	4.3	5.1
2000.....	4.8	...	...	0.5	4.6	6.2	7.8	7.1	4.6	3.2	4.5	6.5
1999.....	4.7	...	...	0.4	4.4	6.6	7.8	6.8	4.4	3.7	5.0	6.6

Non-Hispanic 4/, male

2007.....	8.7	...	...	0.5	8.3	11.9	13.8	13.8	9.2	5.1	6.4	9.0
2006.....	8.4	...	...	0.5	8.3	11.6	13.5	13.2	8.6	5.2	5.5	10.6
2005.....	8.0	...	...	0.7	8.2	11.2	12.9	11.6	7.7	5.1	6.3	11.3
2004.....	8.1	...	...	0.7	8.3	11.6	13.1	11.8	7.9	4.9	6.4	10.6
2003.....	7.7	...	...	0.6	7.4	11.2	12.8	11.0	7.3	4.9	6.3	11.2
2002.....	7.7	...	...	0.7	7.8	11.1	12.4	11.1	7.0	5.1	7.1	12.5
2001 6/.....	7.5	...	...	0.7	7.7	10.6	12.2	10.5	6.6	5.3	7.0	11.6
2000.....	6.9	...	...	0.8	7.2	9.6	11.4	9.7	6.0	4.2	7.5	14.6
1999.....	6.9	...	...	0.6	6.7	10.2	11.3	9.1	5.7	5.1	7.6	15.4

Non-Hispanic 4/, female

2007.....	3.7	...	...	0.3	2.5	3.9	5.5	6.7	5.3	2.8	2.6	2.5
2006.....	3.5	...	...	0.3	2.3	3.7	5.4	6.2	4.8	2.5	2.9	2.5
2005.....	3.4	...	...	0.3	2.6	3.6	5.2	5.9	4.1	2.4	2.7	3.2
2004.....	3.4	...	...	0.5	2.6	3.5	5.3	6.2	4.0	2.3	2.5	3.0
2003.....	3.1	...	...	0.2	2.1	3.5	4.9	5.5	3.9	2.2	2.6	2.4
2002.....	3.1	...	...	0.2	2.0	3.3	5.1	5.4	3.6	2.6	2.9	3.1
2001 6/.....	2.9	...	...	0.2	1.9	3.1	4.7	5.1	3.5	2.3	2.6	2.5
2000.....	2.7	...	...	0.2	1.9	2.9	4.3	4.6	3.4	2.4	2.6	3.2
1999.....	2.7	...	...	0.1	1.9	3.0	4.4	4.5	3.1	2.5	3.3	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic white, both sexes

2007 5/.....	6.9	...	...	0.3	6.0	8.8	10.9	11.6	8.0	4.1	4.2	4.8
2006 5/.....	6.6	...	...	0.4	5.8	8.6	10.7	10.8	7.3	3.8	4.1	5.4
2005 5/.....	6.2	...	...	0.4	6.0	8.1	10.2	9.8	6.4	3.8	4.3	5.7
2004 5/.....	6.3	...	...	0.5	5.9	8.3	10.4	9.9	6.3	3.6	4.2	5.4
2003 5/.....	5.9	...	...	0.4	5.1	8.1	9.9	9.2	6.0	3.5	4.2	5.2
2002.....	6.0	...	...	0.5	5.6	7.9	9.8	9.2	5.7	3.9	4.7	6.1
2001 6/.....	5.6	...	...	0.4	5.4	7.5	9.4	8.6	5.3	3.8	4.5	5.3
2000.....	5.2	...	...	0.5	4.9	6.8	8.6	7.8	5.1	3.3	4.5	6.8
1999.....	5.2	...	...	0.4	4.6	7.1	8.6	7.4	4.6	3.8	5.1	7.0

Non-Hispanic white, male

2007 5/.....	9.6	...	...	0.4	9.2	13.2	15.5	15.4	10.0	5.3	6.6	9.2
2006 5/.....	9.3	...	...	0.5	9.0	12.8	15.1	14.6	9.2	5.4	5.8	11.2
2005 5/.....	8.8	...	...	0.6	9.0	12.2	14.4	12.8	8.3	5.4	6.6	11.5
2004 5/.....	8.9	...	...	0.6	8.9	12.8	14.8	12.8	8.2	5.0	6.5	10.8
2003 5/.....	8.4	...	...	0.6	7.9	12.3	14.1	12.2	7.7	4.9	6.5	11.6
2002.....	8.5	...	...	0.7	8.8	12.0	13.8	12.3	7.6	5.3	7.3	13.0
2001 6/.....	8.2	...	...	0.7	8.6	11.5	13.5	11.5	6.9	5.3	7.3	12.3
2000.....	7.5	...	...	0.8	7.7	10.3	12.4	10.4	6.5	4.3	7.5	15.2
1999.....	7.4	...	...	0.6	7.2	10.8	12.3	9.8	6.0	5.3	7.8	16.5

Non-Hispanic white, female

2007 5/.....	4.2	...	...	0.2	2.6	4.4	6.3	7.8	6.0	3.1	2.6	2.7
2006 5/.....	4.0	...	...	0.3	2.4	4.3	6.3	7.1	5.4	2.5	3.0	2.7
2005 5/.....	3.7	...	...	0.2	2.8	4.0	6.0	6.7	4.6	2.5	2.8	3.1
2004 5/.....	3.7	...	...	0.4	2.7	3.6	6.1	7.1	4.5	2.5	2.6	3.0
2003 5/.....	3.4	...	...	0.2	2.2	3.9	5.6	6.2	4.3	2.4	2.7	2.4
2002.....	3.5	...	...	0.2	2.2	3.8	5.8	6.1	3.9	2.7	2.9	3.2
2001 6/.....	3.2	...	...	0.2	2.0	3.4	5.3	5.7	3.9	2.4	2.7	2.4
2000.....	3.0	...	...	0.2	1.9	3.1	4.8	5.2	3.8	2.5	2.6	3.4
1999.....	3.0	...	...	*	2.0	3.3	4.9	5.1	3.4	2.6	3.4	3.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic black, both sexes

2007 5/.....	2.5	...	...	0.5	2.5	4.1	4.4	3.4	2.2	1.4	*	*
2006 5/.....	2.4	...	...	0.5	2.6	3.9	4.2	3.5	2.4	1.3	*	*
2005 5/.....	2.6	...	...	0.6	2.9	4.3	4.2	3.5	2.3	1.2	*	*
2004 5/.....	2.6	...	...	0.6	3.2	4.7	3.9	3.5	2.7	*	*	*
2003 5/.....	2.5	...	...	0.6	2.9	4.2	4.7	3.2	2.1	1.3	*	*
2002.....	2.4	...	...	0.4	2.4	4.4	4.2	3.0	2.1	1.6	*	*
2001 6/.....	2.3	...	...	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000.....	2.4	...	...	0.5	3.0	3.9	4.2	3.3	1.3	1.4	2.3	*
1999.....	2.3	...	...	0.4	2.5	4.4	4.2	2.7	1.5	1.8	*	*

Non-Hispanic black, male

2007 5/.....	3.9	...	...	0.7	3.8	6.9	7.0	5.3	3.2	2.7	*	*
2006 5/.....	4.0	...	...	0.8	4.0	6.9	7.0	5.8	4.0	*	*	*
2005 5/.....	3.9	...	...	0.7	4.7	6.7	6.4	5.1	3.4	*	*	*
2004 5/.....	4.1	...	...	0.7	5.1	7.3	6.1	5.3	5.2	*	*	*
2003 5/.....	3.9	...	...	0.9	4.4	6.6	7.4	4.5	3.3	*	*	*
2002.....	3.8	...	...	0.6	3.8	7.5	6.7	4.7	3.0	*	*	*
2001 6/.....	3.7	...	...	0.7	3.8	7.1	6.1	4.9	3.7	3.0	*	*
2000.....	3.9	...	...	0.9	4.7	6.5	6.8	5.4	*	*	*	*
1999.....	3.8	...	...	*	4.1	7.5	6.8	4.5	2.1	*	*	*

Non-Hispanic black, female

2007 5/.....	1.2	...	...	*	1.2	1.6	2.2	1.9	1.3	*	*	*
2006 5/.....	1.0	...	...	*	1.1	1.0	1.7	1.6	1.2	*	*	*
2005 5/.....	1.4	...	...	*	1.2	2.1	2.2	2.1	1.4	*	*	*
2004 5/.....	1.3	...	...	*	1.3	2.3	2.0	1.9	*	*	*	*
2003 5/.....	1.3	...	...	*	1.3	2.0	2.3	2.1	*	*	*	*
2002.....	1.1	...	...	*	1.0	1.6	2.0	1.6	1.4	*	*	*
2001 6/.....	1.1	...	...	*	0.7	1.5	2.0	2.1	*	*	*	*
2000.....	1.1	...	...	*	1.3	1.5	1.9	1.5	*	*	*	*
1999.....	1.0	...	...	*	0.9	1.6	2.0	1.3	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

All origins 3/, both sexes

2007.....	6.1	8.3	2.4	0.9	13.1	11.7	7.1	4.9	3.0	2.1	2.1	1.5
2006.....	6.2	8.1	2.2	1.0	13.5	11.7	6.9	5.1	3.2	2.1	2.1	1.9
2005.....	6.1	7.5	2.3	0.8	13.0	11.8	7.1	4.8	2.8	2.4	2.2	2.1
2004.....	5.9	8.0	2.4	0.8	12.2	11.2	6.8	4.8	3.0	2.4	2.2	2.1
2003.....	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002.....	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 6/.....	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000.....	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999.....	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4

All origins 3/, male

2007.....	9.8	9.5	2.5	1.0	22.1	19.4	10.6	7.3	4.6	3.1	2.8	1.5
2006.....	10.0	9.4	2.5	1.2	22.8	19.4	10.4	7.6	4.8	3.0	2.7	2.3
2005.....	9.8	8.2	2.6	1.0	22.0	19.6	10.6	7.6	4.3	3.3	2.6	2.7
2004.....	9.4	8.0	2.5	1.0	20.4	18.5	10.1	7.3	4.6	3.6	2.6	2.7
2003.....	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002.....	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 6/.....	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000.....	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999.....	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4

All origins 3/, female

2007.....	2.5	7.0	2.3	0.7	3.5	3.7	3.5	2.5	1.5	1.3	1.5	1.4
2006.....	2.5	6.8	2.0	0.7	3.5	3.7	3.4	2.6	1.7	1.3	1.8	1.8
2005.....	2.5	6.6	2.0	0.7	3.4	3.8	3.6	2.2	1.4	1.6	1.9	1.9
2004.....	2.5	7.9	2.2	0.6	3.5	3.7	3.4	2.4	1.6	1.4	1.9	1.9
2003.....	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002.....	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 6/.....	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000.....	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999.....	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Hispanic, both sexes

2007.....	7.6	7.8	2.0	0.9	17.4	12.6	7.3	5.3	3.5	2.4	*	*
2006.....	8.0	8.3	1.7	1.2	18.2	12.3	8.1	5.8	3.7	2.3	*	*
2005.....	8.2	6.4	2.3	0.8	18.2	13.9	8.1	6.2	3.0	2.7	3.2	*
2004.....	7.9	8.1	2.3	0.9	17.4	12.6	7.7	5.9	3.8	2.3	*	*
2003.....	8.4	6.9	2.5	0.7	18.4	13.2	8.5	6.4	3.6	3.8	3.6	*
2002.....	8.1	6.2	2.6	0.9	17.6	12.3	8.0	5.9	4.4	2.5	*	*
2001 6/.....	9.0	6.7	2.3	0.8	18.3	14.0	10.4	7.3	4.4	4.0	*	*
2000.....	8.3	7.0	1.9	1.1	17.1	12.4	8.7	6.7	4.5	3.3	*	*
1999.....	8.4	7.0	1.8	1.4	17.7	12.7	8.8	6.6	4.2	3.2	*	*

Hispanic, male

2007.....	12.4	8.3	1.5	1.1	30.0	19.9	11.2	8.7	5.8	4.5	*	*
2006.....	13.1	10.1	1.8	1.5	31.0	19.7	12.5	9.1	6.7	3.2	*	*
2005.....	13.6	6.3	2.5	1.0	31.0	22.3	12.7	10.1	4.7	4.4	*	*
2004.....	13.0	9.1	2.5	1.0	29.6	20.3	11.8	9.5	6.5	4.1	*	*
2003.....	13.6	7.9	2.1	0.9	30.3	21.2	12.9	10.2	6.0	6.8	*	*
2002.....	13.2	6.6	3.1	1.0	29.6	19.8	12.1	9.1	7.7	4.8	*	*
2001 6/.....	14.5	7.9	2.5	1.1	30.5	21.6	15.9	11.3	6.8	6.6	*	*
2000.....	13.4	6.6	2.0	1.6	28.5	19.9	13.5	10.2	7.1	4.6	*	*
1999.....	13.8	6.7	1.7	1.8	29.1	20.6	13.9	10.3	7.3	5.7	*	*

Hispanic, female

2007.....	2.5	7.3	2.4	0.7	3.5	3.5	3.0	1.9	1.4	*	*	*
2006.....	2.4	6.5	1.5	0.8	3.8	3.2	3.1	2.4	*	*	*	*
2005.....	2.5	6.6	2.0	0.6	3.6	3.7	3.1	2.2	*	*	*	*
2004.....	2.4	7.0	2.1	0.7	3.3	3.4	3.2	2.2	*	*	*	*
2003.....	2.8	5.9	2.9	*	4.5	3.5	3.6	2.7	*	*	*	*
2002.....	2.6	5.9	2.1	0.8	3.8	3.4	3.5	2.7	*	*	*	*
2001 6/.....	3.2	5.5	2.1	*	4.0	5.1	4.4	3.4	2.4	*	*	*
2000.....	2.8	7.4	1.9	0.6	3.7	3.7	3.6	3.3	2.2	*	*	*
1999.....	2.8	7.3	1.9	0.9	4.4	3.5	3.3	3.1	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	5.8	8.3	2.5	0.8	12.1	11.5	7.0	4.8	2.9	2.1	2.0	1.5
2006.....	5.9	8.0	2.4	0.9	12.4	11.5	6.7	5.0	3.1	2.1	2.2	1.9
2005.....	5.7	7.7	2.3	0.9	11.8	11.3	6.9	4.7	2.8	2.3	2.1	2.2
2004.....	5.5	7.8	2.4	0.8	11.1	10.8	6.6	4.6	3.0	2.4	2.3	2.0
2003.....	5.7	8.9	2.3	0.8	11.8	10.8	6.7	4.8	2.7	2.2	2.4	2.1
2002.....	5.7	7.7	2.7	0.9	11.8	10.9	7.0	4.6	3.0	2.3	2.2	2.1
2001 6/.....	6.7	8.4	2.8	0.8	12.1	12.8	9.2	6.1	3.9	2.8	2.5	2.4
2000.....	5.6	9.6	2.4	0.9	11.6	10.0	6.9	4.4	2.9	2.3	2.4	2.3
1999.....	5.7	8.9	2.6	1.0	11.9	10.0	6.8	4.3	2.9	2.5	2.5	2.4

Non-Hispanic 4/, male

2007.....	9.2	9.6	2.8	1.0	20.3	19.2	10.4	7.1	4.5	3.0	2.8	1.4
2006.....	9.3	9.1	2.7	1.1	21.0	19.2	9.9	7.4	4.6	3.0	2.7	2.1
2005.....	9.1	8.8	2.6	1.0	19.9	18.8	10.1	7.2	4.3	3.2	2.4	2.8
2004.....	8.7	7.5	2.5	0.9	18.4	17.9	9.7	6.9	4.4	3.5	2.7	2.4
2003.....	9.0	10.5	2.6	1.0	19.8	17.7	10.0	7.1	4.0	3.1	3.2	3.0
2002.....	8.9	8.0	2.9	0.9	19.6	17.6	10.4	6.8	4.5	3.2	3.1	2.9
2001 6/.....	10.4	9.8	3.2	0.9	20.1	20.3	13.4	9.2	5.8	3.8	3.1	3.1
2000.....	8.6	11.2	2.6	1.0	19.1	15.9	9.8	6.6	4.4	3.2	3.1	3.1
1999.....	8.6	10.1	2.6	1.1	19.4	15.6	9.7	6.4	4.5	3.5	3.3	3.4

Non-Hispanic 4/, female

2007.....	2.5	6.9	2.2	0.7	3.5	3.7	3.6	2.6	1.5	1.3	1.5	1.5
2006.....	2.5	6.8	2.1	0.7	3.5	3.8	3.5	2.6	1.7	1.3	1.8	1.8
2005.....	2.5	6.6	2.0	0.7	3.4	3.8	3.7	2.2	1.4	1.6	1.9	1.9
2004.....	2.5	8.1	2.3	0.6	3.5	3.7	3.5	2.4	1.6	1.5	2.0	1.8
2003.....	2.5	7.2	2.1	0.6	3.5	4.0	3.6	2.5	1.5	1.5	1.8	1.8
2002.....	2.7	7.4	2.6	0.8	3.7	4.3	3.8	2.5	1.7	1.5	1.7	1.7
2001 6/.....	3.2	6.9	2.4	0.7	3.8	5.3	5.1	3.1	2.2	2.0	2.1	2.1
2000.....	2.7	7.9	2.2	0.7	3.8	4.1	4.0	2.3	1.5	1.6	1.9	2.0
1999.....	2.9	7.6	2.5	0.9	4.2	4.6	4.0	2.3	1.4	1.7	2.0	2.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Non-Hispanic white, both sexes

2007 5/.....	2.7	6.5	1.6	0.5	3.6	4.4	3.8	3.2	2.1	1.7	1.7	1.2
2006 5/.....	2.7	6.3	1.5	0.6	3.4	3.9	3.6	3.2	2.3	1.6	2.0	1.7
2005 5/.....	2.7	6.0	1.5	0.5	3.4	4.0	3.7	3.0	2.1	1.8	1.8	1.9
2004 5/.....	2.7	5.9	1.4	0.5	3.5	4.0	3.8	3.0	2.2	2.0	1.9	1.8
2003 5/.....	2.7	6.8	1.3	0.4	3.5	3.9	3.8	3.1	2.1	1.7	2.0	2.0
2002.....	2.8	5.0	1.8	0.5	3.7	3.9	4.1	3.0	2.4	1.9	1.8	1.8
2001 6/.....	4.0	5.8	1.7	0.6	4.2	6.5	6.6	4.6	3.3	2.3	2.1	2.2
2000.....	2.8	6.2	1.6	0.6	3.5	4.0	4.0	2.9	2.2	1.8	2.0	2.1
1999.....	2.9	6.3	1.7	0.7	3.9	4.3	4.1	2.9	2.2	2.1	2.0	2.1

Non-Hispanic white, male

2007 5/.....	3.7	7.2	1.9	0.5	4.9	6.2	5.1	4.4	2.9	2.2	2.0	*
2006 5/.....	3.6	6.5	1.6	0.6	4.7	5.5	4.9	4.5	3.2	2.1	2.2	1.8
2005 5/.....	3.6	6.9	1.6	0.5	4.7	5.4	4.8	4.3	3.1	2.2	1.9	2.0
2004 5/.....	3.6	6.1	1.6	0.6	4.7	5.5	5.0	4.3	3.1	2.6	2.1	1.9
2003 5/.....	3.6	8.0	1.3	0.6	5.0	5.2	4.9	4.3	2.8	2.3	2.6	2.9
2002.....	3.8	5.8	1.9	0.5	5.2	5.3	5.6	3.9	3.3	2.4	2.3	2.7
2001 6/.....	5.6	6.8	2.0	0.6	5.7	9.3	9.5	6.7	4.7	2.9	2.6	2.8
2000.....	3.6	8.3	1.8	0.7	4.7	5.2	5.2	3.9	3.1	2.2	2.4	2.6
1999.....	3.9	8.0	1.7	0.7	5.1	5.7	5.4	4.2	3.2	2.7	2.6	2.6

Non-Hispanic white, female

2007 5/.....	1.8	5.7	1.4	0.5	2.2	2.5	2.6	2.0	1.3	1.1	1.4	1.4
2006 5/.....	1.8	6.2	1.3	0.6	2.0	2.2	2.4	2.0	1.5	1.3	1.8	1.7
2005 5/.....	1.8	5.0	1.4	0.4	2.0	2.6	2.7	1.6	1.2	1.4	1.8	1.9
2004 5/.....	1.8	5.8	1.3	0.4	2.3	2.5	2.5	1.8	1.4	1.4	1.8	1.7
2003 5/.....	1.8	5.5	1.4	0.3	1.9	2.5	2.6	2.0	1.4	1.3	1.6	1.6
2002.....	1.9	4.1	1.8	0.6	2.2	2.6	2.5	2.1	1.5	1.4	1.5	1.5
2001 6/.....	2.4	4.7	1.5	0.7	2.7	3.7	3.7	2.6	1.9	1.8	1.8	2.0
2000.....	1.9	4.1	1.3	0.6	2.3	2.7	2.7	1.9	1.3	1.4	1.7	1.9
1999.....	2.0	4.4	1.6	0.7	2.6	2.9	2.8	1.7	1.3	1.6	1.6	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Non-Hispanic black, both sexes

2007 5/.....	22.9	16.1	6.3	2.2	49.2	46.7	25.1	15.6	9.7	5.9	6.6	*
2006 5/.....	23.6	15.9	6.0	2.4	51.2	49.3	24.5	16.2	9.7	6.1	4.5	*
2005 5/.....	22.8	14.5	5.5	2.3	48.5	47.9	24.8	16.0	8.2	7.2	5.9	*
2004 5/.....	21.7	15.8	6.2	2.0	45.1	45.9	23.3	15.4	8.8	6.2	6.4	*
2003 5/.....	22.7	16.8	6.4	2.3	49.1	47.1	24.2	15.8	8.1	6.3	6.6	*
2002.....	22.5	17.6	6.5	2.0	48.3	47.0	24.3	15.8	8.4	5.8	7.3	*
2001 6/.....	22.7	19.5	7.1	1.4	48.7	45.3	25.5	16.2	8.9	7.2	6.8	*
2000.....	22.1	24.1	6.1	1.9	49.3	42.2	24.5	15.0	8.4	7.6	7.1	*
1999.....	21.7	20.0	6.4	2.3	49.5	40.6	23.7	14.1	8.4	6.4	7.5	*

Non-Hispanic black, male

2007 5/.....	41.1	19.9	6.9	3.0	88.2	86.2	43.0	26.1	18.3	10.5	13.5	*
2006 5/.....	42.0	20.3	6.4	3.3	91.6	90.2	41.2	27.7	17.2	11.5	8.3	*
2005 5/.....	40.9	15.9	6.9	2.9	87.0	89.4	41.9	27.6	14.6	13.0	10.4	*
2004 5/.....	38.3	13.9	6.3	2.5	80.7	85.1	39.0	26.1	15.2	11.8	10.3	*
2003 5/.....	40.1	18.6	7.8	3.0	87.3	86.1	41.6	26.9	14.9	11.3	11.3	*
2002.....	39.4	16.4	7.3	2.6	85.8	85.3	39.8	27.5	15.8	10.5	13.8	*
2001 6/.....	39.3	21.7	8.4	2.0	88.3	81.7	39.5	27.2	15.0	12.6	10.2	*
2000.....	38.3	24.9	6.1	2.1	88.1	76.5	39.4	26.3	15.0	13.9	12.0	*
1999.....	37.2	20.1	6.2	2.7	88.1	71.5	37.9	23.7	15.4	11.9	12.2	*

Non-Hispanic black, female

2007 5/.....	6.4	12.1	5.8	1.4	9.2	9.8	9.3	6.5	2.8	2.6	*	*
2006 5/.....	6.8	11.3	5.6	1.4	9.8	11.5	9.7	6.4	3.7	2.2	*	*
2005 5/.....	6.3	13.0	4.1	1.7	9.0	9.7	9.5	6.1	3.1	3.1	3.3	*
2004 5/.....	6.6	17.7	6.1	1.6	9.1	10.1	9.3	6.2	3.7	2.1	4.2	*
2003 5/.....	6.8	14.8	4.9	1.5	10.4	11.6	8.8	6.2	2.7	2.8	3.9	*
2002.....	7.2	18.9	5.7	1.5	10.6	12.3	10.6	5.7	2.5	2.4	3.6	*
2001 6/.....	7.6	17.1	5.7	0.8	9.2	12.5	13.2	6.9	4.0	3.3	4.9	*
2000.....	7.4	23.3	6.0	1.6	11.0	11.4	11.3	5.3	3.2	3.2	4.3	*
1999.....	7.7	20.0	6.7	1.8	11.4	12.8	11.1	5.9	2.7	2.5	4.9	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

All origins 3/, both sexes

2007.....	4.2	*	0.3	0.5	11.0	9.2	4.7	2.6	1.4	1.0	0.7	0.4
2006.....	4.3	*	0.3	0.6	11.4	9.3	4.6	2.7	1.5	0.8	0.6	0.4
2005.....	4.2	*	0.2	0.5	10.7	9.4	4.6	2.6	1.3	1.1	0.7	0.5
2004.....	4.0	*	0.2	0.5	9.9	8.8	4.3	2.6	1.6	1.0	0.8	0.6
2003.....	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002.....	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001.....	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000.....	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999.....	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6

All origins 3/, male

2007.....	7.2	*	0.3	0.7	19.2	16.1	7.6	4.2	2.2	1.4	0.9	*
2006.....	7.4	*	0.3	0.8	20.0	16.2	7.3	4.3	2.2	1.2	0.8	*
2005.....	7.2	*	0.3	0.6	18.9	16.4	7.3	4.1	2.1	1.5	0.8	*
2004.....	6.9	*	*	0.6	17.3	15.3	7.1	4.0	2.5	1.6	0.8	*
2003.....	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002.....	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001.....	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000.....	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999.....	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*

All origins 3/, female

2007.....	1.2	*	0.3	0.3	2.3	2.1	1.8	1.2	0.6	0.6	0.5	0.5
2006.....	1.3	*	*	0.4	2.2	2.2	1.8	1.3	0.8	0.5	0.6	*
2005.....	1.2	*	*	0.3	2.0	2.2	1.9	1.1	0.6	0.7	0.7	*
2004.....	1.1	*	*	0.3	2.0	2.0	1.5	1.1	0.8	0.5	0.8	*
2003.....	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002.....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001.....	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000.....	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999.....	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hispanic, both sexes												
2007.....	5.2	*	*	0.5	14.1	9.2	4.7	3.0	1.4	*	*	*
2006.....	5.6	*	*	0.8	15.1	9.3	5.2	3.0	1.6	*	*	*
2005.....	5.7	*	*	0.5	14.5	10.6	5.3	3.3	1.3	1.5	*	*
2004.....	5.4	*	*	0.5	14.0	9.4	5.0	2.7	2.0	*	*	*
2003.....	5.8	*	*	0.4	14.9	9.9	5.3	3.6	1.2	*	*	*
2002.....	5.6	*	*	0.6	14.3	9.3	5.1	3.2	1.6	*	*	*
2001.....	5.7	*	*	0.4	14.6	9.5	5.6	3.0	1.1	*	*	*
2000.....	5.5	*	*	0.6	13.7	8.9	4.9	4.0	2.0	*	*	*
1999.....	5.7	*	*	0.8	14.4	8.7	5.2	3.8	2.1	*	*	*
Hispanic, male												
2007.....	9.1	*	*	0.8	24.8	15.1	7.5	5.1	2.2	*	*	*
2006.....	9.7	*	*	1.1	26.4	15.5	8.2	4.9	2.8	*	*	*
2005.....	10.0	*	*	0.8	25.5	17.6	8.3	5.5	2.3	*	*	*
2004.....	9.5	*	*	0.7	24.4	15.9	8.3	4.5	3.5	*	*	*
2003.....	10.0	*	*	0.6	25.5	16.8	8.4	5.9	2.2	*	*	*
2002.....	9.7	*	*	0.8	24.9	15.6	8.1	5.0	2.8	*	*	*
2001.....	9.9	*	*	0.6	24.9	16.1	9.1	4.8	*	*	*	*
2000.....	9.6	*	*	1.0	23.5	15.0	7.8	6.4	3.2	*	*	*
1999.....	9.8	*	*	1.2	24.2	14.7	8.6	5.7	3.5	*	*	*
Hispanic, female												
2007.....	1.1	*	*	*	2.4	1.9	1.5	*	*	*	*	*
2006.....	1.2	*	*	*	2.2	1.8	1.7	1.1	*	*	*	*
2005.....	1.2	*	*	*	1.9	2.1	1.9	1.2	*	*	*	*
2004.....	1.1	*	*	*	2.0	1.6	1.4	*	*	*	*	*
2003.....	1.3	*	*	*	2.7	1.7	1.9	1.4	*	*	*	*
2002.....	1.2	*	*	*	2.1	1.8	1.8	1.4	*	*	*	*
2001.....	1.3	*	*	*	2.5	1.8	1.9	1.2	*	*	*	*
2000.....	1.3	*	*	*	2.2	1.9	1.7	1.7	*	*	*	*
1999.....	1.4	*	*	*	2.9	1.7	1.7	1.9	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Non-Hispanic 4/, both sexes

2007.....	4.0	*	0.3	0.5	10.3	9.2	4.7	2.6	1.3	1.0	0.7	0.4
2006.....	4.0	*	0.3	0.5	10.5	9.3	4.4	2.7	1.5	0.9	0.7	0.4
2005.....	3.9	*	0.2	0.5	9.9	9.1	4.4	2.5	1.3	1.0	0.8	0.5
2004.....	3.7	*	0.2	0.4	9.0	8.5	4.1	2.5	1.6	1.1	0.8	0.5
2003.....	3.8	*	0.2	0.5	9.8	8.6	4.2	2.6	1.4	0.8	0.7	0.5
2002.....	3.8	*	0.3	0.5	9.8	8.5	4.4	2.5	1.5	0.9	0.9	*
2001.....	3.7	*	0.3	0.4	9.6	8.0	4.2	2.3	1.5	1.0	0.8	*
2000.....	3.6	*	0.2	0.4	9.3	7.4	4.1	2.4	1.5	1.0	0.7	0.7
1999.....	3.6	*	0.4	0.5	9.6	7.3	4.0	2.3	1.5	1.1	0.8	0.5

Non-Hispanic 4/, male

2007.....	6.9	*	0.3	0.6	17.9	16.2	7.6	4.1	2.1	1.4	0.9	*
2006.....	6.9	*	*	0.7	18.5	16.3	7.0	4.2	2.2	1.2	0.8	*
2005.....	6.7	*	*	0.6	17.4	15.9	7.0	3.9	2.1	1.5	0.8	*
2004.....	6.4	*	*	0.6	15.8	15.1	6.8	3.9	2.4	1.6	0.8	*
2003.....	6.5	*	*	0.7	17.2	15.0	6.7	4.1	2.1	1.3	0.7	*
2002.....	6.5	*	0.3	0.6	17.0	14.7	7.1	3.7	2.3	1.2	1.2	*
2001.....	6.3	*	0.3	0.6	17.0	13.8	6.5	3.7	2.2	1.4	1.1	*
2000.....	6.0	*	*	0.6	16.3	12.7	6.4	3.7	2.2	1.3	0.8	*
1999.....	6.0	*	0.4	0.7	16.7	12.4	6.0	3.6	2.5	1.5	0.9	*

Non-Hispanic 4/, female

2007.....	1.2	*	0.3	0.3	2.3	2.2	1.9	1.2	0.6	0.6	0.5	0.6
2006.....	1.3	*	*	0.4	2.2	2.3	1.9	1.3	0.8	0.5	0.6	*
2005.....	1.2	*	*	0.3	2.0	2.2	1.9	1.1	0.6	0.7	0.7	*
2004.....	1.2	*	*	0.2	2.0	2.1	1.5	1.2	0.8	0.6	0.8	*
2003.....	1.2	*	*	0.3	2.0	2.2	1.8	1.2	0.7	0.5	0.7	*
2002.....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001.....	1.2	*	0.3	0.3	2.0	2.3	2.0	1.0	0.8	0.7	0.6	*
2000.....	1.2	*	*	0.2	2.2	2.2	1.9	1.1	0.8	0.7	0.6	*
1999.....	1.3	*	0.4	0.3	2.4	2.3	2.0	1.1	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Non-Hispanic white, both sexes

2007 5/.....	1.5	*	*	0.2	2.5	3.0	2.3	1.6	0.9	0.8	0.6	*
2006 5/.....	1.4	*	*	0.3	2.2	2.5	2.2	1.6	1.1	0.6	0.6	*
2005 5/.....	1.4	*	*	0.2	2.1	2.6	2.2	1.6	1.0	0.8	0.7	*
2004 5/.....	1.5	*	*	0.2	2.1	2.6	2.2	1.6	1.2	0.9	0.7	*
2003 5/.....	1.4	*	*	0.2	2.4	2.5	2.1	1.7	1.1	0.7	0.6	0.5
2002.....	1.5	*	0.2	0.3	2.5	2.6	2.4	1.6	1.2	0.7	0.8	*
2001.....	1.6	*	0.2	0.4	2.4	2.7	2.4	1.6	1.2	0.8	0.7	*
2000.....	1.4	*	*	0.3	2.2	2.4	2.3	1.6	1.2	0.8	0.6	0.6
1999.....	1.5	*	0.2	0.3	2.4	2.6	2.3	1.5	1.1	0.9	0.6	0.6

Non-Hispanic white, male

2007 5/.....	2.2	*	*	0.2	3.5	4.5	3.4	2.3	1.3	1.0	0.7	*
2006 5/.....	2.0	*	*	0.3	3.4	3.8	3.1	2.2	1.5	0.8	0.5	*
2005 5/.....	2.1	*	*	0.3	3.2	3.8	3.0	2.3	1.5	1.0	0.7	*
2004 5/.....	2.1	*	*	0.3	3.1	4.0	3.1	2.3	1.7	1.2	0.7	*
2003 5/.....	2.1	*	*	0.3	3.7	3.7	2.9	2.4	1.6	1.0	0.5	*
2002.....	2.2	*	*	0.3	3.7	3.7	3.4	2.2	1.7	0.9	1.0	*
2001.....	2.2	*	*	0.4	3.6	3.9	3.4	2.3	1.6	1.0	0.9	*
2000.....	2.0	*	*	0.4	3.2	3.4	3.2	2.2	1.6	0.9	0.6	*
1999.....	2.2	*	*	0.4	3.5	3.9	3.2	2.2	1.7	1.2	0.6	*

Non-Hispanic white, female

2007 5/.....	0.9	*	*	0.2	1.3	1.5	1.3	1.0	0.5	0.6	0.5	*
2006 5/.....	0.8	*	*	0.3	1.1	1.2	1.2	1.1	0.8	0.5	0.7	*
2005 5/.....	0.8	*	*	*	1.0	1.4	1.4	0.9	0.6	0.7	0.7	*
2004 5/.....	0.8	*	*	*	1.1	1.2	1.2	0.9	0.7	0.7	0.8	*
2003 5/.....	0.8	*	*	*	1.0	1.3	1.3	1.0	0.7	0.4	0.6	*
2002.....	0.9	*	*	0.3	1.2	1.4	1.3	1.1	0.7	0.6	0.7	*
2001.....	0.9	*	*	0.3	1.2	1.5	1.4	0.8	0.8	0.7	0.6	*
2000.....	0.9	*	*	0.2	1.2	1.4	1.4	0.9	0.8	0.7	0.6	*
1999.....	0.9	*	*	0.2	1.2	1.4	1.5	0.8	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hispanic black, both sexes												
2007 5/.....	18.0	*	*	1.5	44.7	40.4	18.5	9.1	5.0	2.6	2.0	*
2006 5/.....	18.6	*	*	1.6	46.4	43.1	17.7	9.6	4.3	2.9	*	*
2005 5/.....	17.7	*	*	1.4	43.6	41.8	17.4	8.7	3.6	3.1	*	*
2004 5/.....	16.6	*	*	1.3	40.3	39.4	16.0	8.6	4.7	2.4	*	*
2003 5/.....	17.3	*	*	1.5	43.3	40.5	16.5	8.9	4.0	2.5	*	*
2002.....	17.1	*	*	1.2	43.1	39.7	16.7	8.6	4.2	2.4	2.1	*
2001.....	16.3	*	*	0.9	43.0	36.4	15.5	7.7	3.7	2.6	*	*
2000.....	16.0	*	*	1.0	42.6	34.5	15.4	8.5	3.6	2.8	*	*
1999.....	15.9	*	*	1.3	43.6	32.4	14.4	8.1	4.5	2.4	2.4	*
Non-Hispanic black, male												
2007 5/.....	34.0	*	*	2.2	81.9	77.2	33.5	16.5	9.6	4.8	*	*
2006 5/.....	35.0	*	*	2.3	84.7	81.2	31.8	17.9	8.0	5.6	*	*
2005 5/.....	33.6	*	*	2.0	80.3	80.3	31.8	15.9	6.9	6.1	*	*
2004 5/.....	31.4	*	*	1.9	74.1	75.4	29.8	15.5	8.5	5.6	*	*
2003 5/.....	32.6	*	*	2.1	79.4	77.4	30.1	16.4	7.4	4.8	*	*
2002.....	31.9	*	*	1.6	78.6	75.6	30.0	15.3	7.7	4.7	*	*
2001.....	30.6	*	*	1.3	80.6	69.4	26.7	14.0	7.1	5.5	*	*
2000.....	30.1	*	*	1.6	79.1	65.7	27.4	15.3	6.5	5.5	*	*
1999.....	29.3	*	*	1.8	80.0	61.1	24.6	14.1	9.1	4.7	*	*
Non-Hispanic black, female												
2007 5/.....	3.4	*	*	0.8	6.5	6.2	5.2	2.7	1.2	*	*	*
2006 5/.....	3.7	*	*	0.8	7.2	7.8	5.2	2.5	1.3	*	*	*
2005 5/.....	3.2	*	*	0.7	6.1	6.4	4.6	2.5	*	*	*	*
2004 5/.....	3.1	*	*	*	6.0	6.4	3.8	2.6	1.6	*	*	*
2003 5/.....	3.4	*	*	0.9	6.7	6.9	4.4	2.6	*	*	*	*
2002.....	3.7	*	*	0.8	7.5	7.2	5.0	2.8	*	*	*	*
2001.....	3.2	*	*	*	5.6	6.5	5.5	2.2	*	*	*	*
2000.....	3.3	*	*	*	6.7	6.4	4.7	2.6	*	*	*	*
1999.....	3.7	*	*	0.7	7.8	6.6	5.5	3.0	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes

2007.....	1.9	7.9	2.1	0.4	2.1	2.5	2.3	2.2	1.6	1.2	1.4	1.0
2006.....	1.9	8.0	2.0	0.4	2.1	2.4	2.4	2.4	1.7	1.3	1.5	1.5
2005.....	1.9	7.3	2.1	0.4	2.3	2.4	2.5	2.3	1.5	1.3	1.4	1.7
2004.....	2.0	7.8	2.1	0.4	2.3	2.5	2.5	2.3	1.4	1.4	1.5	1.5
2003.....	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
2002.....	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 6/.....	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000.....	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999.....	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8

All origins 3/, male

2007.....	2.5	8.9	2.3	0.3	2.9	3.4	3.0	3.1	2.4	1.8	1.9	1.3
2006.....	2.6	9.3	2.2	0.4	2.8	3.2	3.1	3.4	2.6	1.8	1.9	1.8
2005.....	2.6	7.9	2.3	0.4	3.1	3.3	3.3	3.4	2.2	1.8	1.8	2.1
2004.....	2.5	7.9	2.3	0.3	3.1	3.2	3.0	3.3	2.1	1.9	1.9	1.7
2003.....	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
2002.....	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 6/.....	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000.....	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999.....	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6

All origins 3/, female

2007.....	1.3	6.8	2.0	0.4	1.2	1.6	1.7	1.4	0.9	0.6	1.0	0.9
2006.....	1.3	6.6	1.8	0.4	1.3	1.5	1.6	1.4	0.9	0.8	1.2	1.4
2005.....	1.3	6.6	1.8	0.4	1.5	1.6	1.7	1.1	0.8	0.9	1.2	1.5
2004.....	1.4	7.7	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.5
2003.....	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
2002.....	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 6/.....	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000.....	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999.....	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Hispanic, both sexes

2007.....	2.4	7.3	1.8	0.4	3.3	3.4	2.6	2.3	2.1	1.7	*	*
2006.....	2.4	8.2	1.4	0.4	3.2	3.0	2.9	2.8	2.1	1.7	*	*
2005.....	2.5	6.3	2.1	0.3	3.7	3.3	2.8	2.8	1.7	*	*	*
2004.....	2.5	7.7	2.1	0.4	3.4	3.2	2.7	3.2	1.8	*	*	*
2003.....	2.6	6.8	2.2	0.3	3.5	3.3	3.2	2.8	2.3	2.4	3.1	*
2002.....	2.5	6.0	2.2	0.3	3.3	3.1	2.9	2.7	2.8	1.7	*	*
2001 6/.....	3.3	6.2	1.9	0.4	3.7	4.5	4.8	4.3	3.4	2.7	*	*
2000.....	2.7	6.9	1.8	0.5	3.4	3.5	3.9	2.7	2.5	2.0	*	*
1999.....	2.7	6.7	1.6	0.6	3.3	4.0	3.5	2.9	2.1	2.3	*	*

Hispanic, male

2007.....	3.4	7.4	1.4	*	5.2	4.8	3.6	3.6	3.6	3.1	*	*
2006.....	3.4	10.1	1.5	*	4.6	4.3	4.3	4.2	3.9	*	*	*
2005.....	3.6	6.1	2.3	*	5.5	4.7	4.4	4.6	2.4	*	*	*
2004.....	3.5	8.9	2.3	*	5.3	4.4	3.6	5.0	3.0	*	*	*
2003.....	3.6	7.7	1.8	*	4.8	4.5	4.6	4.3	3.7	4.4	*	*
2002.....	3.5	6.6	2.6	*	4.7	4.3	4.0	4.1	4.9	*	*	*
2001 6/.....	4.5	7.2	2.2	*	5.6	5.5	6.9	6.5	5.1	4.6	*	*
2000.....	3.8	6.6	1.8	0.6	5.0	5.0	5.7	3.8	3.9	*	*	*
1999.....	4.0	6.2	1.5	0.6	4.9	5.9	5.3	4.5	3.8	4.4	*	*

Hispanic, female

2007.....	1.3	7.1	2.2	*	1.2	1.6	1.5	1.1	*	*	*	*
2006.....	1.3	6.3	1.3	*	1.6	1.3	1.4	1.3	*	*	*	*
2005.....	1.3	6.6	1.9	*	1.7	1.6	1.2	1.0	*	*	*	*
2004.....	1.4	6.5	1.9	*	1.3	1.8	1.7	1.3	*	*	*	*
2003.....	1.5	5.9	2.6	*	1.9	1.8	1.7	1.4	*	*	*	*
2002.....	1.4	5.4	1.8	*	1.7	1.6	1.7	1.4	*	*	*	*
2001 6/.....	1.9	5.2	1.6	*	1.5	3.2	2.6	2.2	*	*	*	*
2000.....	1.6	7.2	1.7	*	1.5	1.9	1.9	1.6	*	*	*	*
1999.....	1.4	7.3	1.8	*	1.5	1.8	1.6	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	1.8	8.0	2.2	0.4	1.8	2.3	2.3	2.2	1.6	1.1	1.4	1.1
2006.....	1.8	7.8	2.2	0.4	1.9	2.2	2.3	2.3	1.7	1.2	1.5	1.5
2005.....	1.8	7.6	2.1	0.4	2.0	2.2	2.4	2.2	1.5	1.3	1.4	1.7
2004.....	1.8	7.7	2.1	0.4	2.0	2.3	2.4	2.1	1.4	1.4	1.5	1.5
2003.....	1.9	8.6	2.1	0.3	2.1	2.2	2.5	2.1	1.3	1.4	1.7	1.6
2002.....	1.9	7.5	2.4	0.4	2.0	2.4	2.6	2.1	1.5	1.4	1.3	1.7
2001 6/.....	3.1	8.2	2.4	0.4	2.5	4.7	5.0	3.8	2.5	1.8	1.7	2.1
2000.....	2.0	9.2	2.2	0.4	2.3	2.6	2.8	2.0	1.4	1.4	1.7	1.6
1999.....	2.1	8.7	2.2	0.5	2.3	2.7	2.9	2.0	1.4	1.5	1.7	1.8

Non-Hispanic 4/, male

2007.....	2.4	9.3	2.5	0.3	2.4	2.9	2.8	3.0	2.3	1.6	1.8	1.3
2006.....	2.4	9.0	2.4	0.4	2.4	2.9	2.9	3.3	2.5	1.7	1.9	1.7
2005.....	2.4	8.5	2.3	0.4	2.5	2.9	3.0	3.3	2.2	1.8	1.6	2.2
2004.....	2.3	7.4	2.3	0.3	2.6	2.9	2.9	3.0	2.0	1.9	1.9	1.7
2003.....	2.4	10.1	2.4	0.4	2.6	2.7	3.3	3.0	1.9	1.8	2.5	2.3
2002.....	2.5	7.8	2.5	0.3	2.6	2.9	3.3	3.1	2.2	2.0	1.8	2.3
2001 6/.....	4.2	9.5	2.8	0.3	3.2	6.5	7.0	5.5	3.6	2.4	2.0	2.7
2000.....	2.6	10.6	2.4	0.4	2.9	3.2	3.4	2.9	2.2	2.0	2.3	2.4
1999.....	2.6	10.0	2.3	0.4	2.8	3.2	3.8	2.9	2.0	2.0	2.4	2.6

Non-Hispanic 4/, female

2007.....	1.3	6.7	1.9	0.4	1.2	1.6	1.7	1.4	0.9	0.7	1.0	0.9
2006.....	1.3	6.6	1.9	0.4	1.3	1.5	1.6	1.4	0.9	0.8	1.2	1.4
2005.....	1.3	6.6	1.8	0.4	1.4	1.6	1.8	1.1	0.8	0.9	1.2	1.4
2004.....	1.4	8.0	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.4
2003.....	1.4	7.1	1.8	0.3	1.5	1.8	1.8	1.3	0.8	1.0	1.1	1.3
2002.....	1.4	7.1	2.3	0.4	1.4	1.9	1.9	1.2	0.9	0.9	1.0	1.4
2001 6/.....	2.0	6.8	2.0	0.4	1.9	2.9	3.1	2.1	1.4	1.3	1.5	1.8
2000.....	1.5	7.8	2.0	0.5	1.7	1.9	2.1	1.2	0.7	0.9	1.3	1.3
1999.....	1.6	7.3	2.2	0.6	1.8	2.3	2.0	1.2	0.8	1.0	1.3	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic white, both sexes

2007 5/.....	1.2	6.3	1.4	0.3	1.1	1.4	1.5	1.6	1.1	0.9	1.1	0.9
2006 5/.....	1.3	6.3	1.3	0.3	1.2	1.4	1.4	1.6	1.2	1.0	1.4	1.4
2005 5/.....	1.2	5.9	1.3	0.2	1.3	1.4	1.5	1.4	1.1	0.9	1.1	1.5
2004 5/.....	1.3	5.9	1.4	0.3	1.4	1.4	1.6	1.4	1.0	1.1	1.2	1.3
2003 5/.....	1.2	6.8	1.2	0.2	1.1	1.4	1.6	1.4	1.0	1.1	1.4	1.5
2002.....	1.3	5.0	1.6	0.3	1.2	1.4	1.7	1.4	1.2	1.1	1.0	1.4
2001 6/.....	2.4	5.6	1.5	0.3	1.7	3.8	4.2	3.1	2.1	1.5	1.4	1.9
2000.....	1.3	6.0	1.4	0.4	1.3	1.6	1.7	1.4	1.0	1.0	1.4	1.5
1999.....	1.4	6.2	1.4	0.4	1.5	1.7	1.8	1.4	1.1	1.2	1.4	1.5

Non-Hispanic white, male

2007 5/.....	1.5	7.0	1.6	0.2	1.4	1.7	1.7	2.2	1.5	1.2	1.3	*
2006 5/.....	1.6	6.5	1.5	0.2	1.4	1.7	1.8	2.3	1.7	1.3	1.7	1.6
2005 5/.....	1.5	6.7	1.4	0.2	1.5	1.7	1.8	2.0	1.6	1.2	1.2	1.6
2004 5/.....	1.5	6.1	1.5	0.2	1.6	1.5	1.9	1.9	1.4	1.5	1.4	*
2003 5/.....	1.5	8.0	1.2	0.3	1.3	1.5	2.0	1.9	1.3	1.3	2.1	2.2
2002.....	1.6	5.8	1.6	0.2	1.5	1.6	2.2	1.8	1.6	1.6	1.3	2.0
2001 6/.....	3.4	6.7	1.8	0.2	2.0	5.4	6.1	4.5	3.1	1.9	1.7	2.3
2000.....	1.6	8.1	1.7	0.3	1.5	1.8	2.0	1.8	1.5	1.3	1.8	1.9
1999.....	1.7	7.8	1.5	0.3	1.6	1.8	2.3	2.0	1.5	1.5	2.0	*

Non-Hispanic white, female

2007 5/.....	0.9	5.6	1.2	0.3	0.8	1.0	1.3	1.0	0.8	0.5	0.9	0.8
2006 5/.....	1.0	6.1	1.1	0.3	1.0	1.0	1.1	0.9	0.7	0.7	1.1	1.3
2005 5/.....	0.9	5.0	1.2	0.3	1.0	1.1	1.2	0.8	0.7	0.7	1.1	1.4
2004 5/.....	1.0	5.8	1.2	0.3	1.2	1.3	1.3	0.9	0.6	0.8	1.0	1.3
2003 5/.....	1.0	5.4	1.2	0.2	1.0	1.2	1.3	1.0	0.7	0.9	1.0	1.2
2002.....	1.0	4.1	1.6	0.3	0.9	1.2	1.2	1.0	0.8	0.8	0.8	1.2
2001 6/.....	1.5	4.5	1.2	0.4	1.4	2.2	2.3	1.7	1.2	1.1	1.2	1.7
2000.....	1.0	3.9	1.2	0.4	1.2	1.3	1.4	1.0	0.5	0.7	1.1	1.3
1999.....	1.1	4.4	1.4	0.4	1.4	1.6	1.3	0.9	0.7	0.9	1.0	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic black, both sexes

2007 5/.....	4.9	15.3	5.6	0.8	4.5	6.3	6.6	6.5	4.7	3.3	4.6	*
2006 5/.....	5.0	15.3	5.2	0.8	4.7	6.3	6.7	6.6	5.5	3.2	3.1	*
2005 5/.....	5.1	14.1	5.0	0.9	4.9	6.1	7.4	7.3	4.6	4.1	4.4	*
2004 5/.....	5.1	15.3	5.4	0.8	4.8	6.5	7.2	6.8	4.1	3.7	4.9	*
2003 5/.....	5.4	15.8	5.6	0.8	5.8	6.6	7.7	6.8	4.2	3.8	4.9	*
2002.....	5.4	16.7	5.9	0.8	5.2	7.3	7.6	7.2	4.2	3.4	5.2	*
2001 6/.....	6.4	19.1	6.3	0.6	5.7	9.0	10.1	8.6	5.1	4.5	4.9	*
2000.....	6.1	22.9	5.7	0.8	6.7	7.8	9.1	6.5	4.9	4.8	5.2	*
1999.....	5.8	19.7	5.7	1.0	5.9	8.2	9.2	6.0	3.9	4.0	5.2	*

Non-Hispanic black, male

2007 5/.....	7.1	19.0	6.2	0.8	6.3	9.0	9.5	9.6	8.7	5.6	8.8	*
2006 5/.....	7.1	19.7	5.6	0.9	6.8	9.0	9.4	9.8	9.3	5.9	*	*
2005 5/.....	7.4	15.2	6.2	0.9	6.7	9.1	10.1	11.7	7.7	6.8	7.0	*
2004 5/.....	6.9	13.6	5.6	0.6	6.6	9.6	9.2	10.6	6.6	6.3	8.3	*
2003 5/.....	7.6	16.8	6.9	0.9	7.9	8.7	11.5	10.5	7.4	6.5	8.7	*
2002.....	7.6	15.8	6.6	0.9	7.3	9.7	9.9	12.3	8.1	5.9	9.4	*
2001 6/.....	8.7	21.4	7.7	0.7	7.7	12.3	12.8	13.2	7.9	7.1	*	*
2000.....	8.2	22.5	5.7	*	9.1	10.8	12.0	11.0	8.5	8.4	8.3	*
1999.....	7.9	19.7	5.6	0.9	8.1	10.5	13.3	9.6	6.3	7.2	7.2	*

Non-Hispanic black, female

2007 5/.....	3.0	11.5	4.9	0.7	2.7	3.7	4.0	3.9	1.5	*	*	*
2006 5/.....	3.1	10.6	4.9	*	2.6	3.7	4.4	3.9	2.4	*	*	*
2005 5/.....	3.1	13.0	3.7	0.9	2.9	3.3	4.9	3.6	2.1	2.1	*	*
2004 5/.....	3.5	17.1	5.3	1.0	3.1	3.7	5.5	3.5	2.1	*	*	*
2003 5/.....	3.4	14.8	4.3	0.7	3.7	4.8	4.4	3.6	1.5	*	*	*
2002.....	3.5	17.7	5.2	0.7	3.1	5.1	5.7	2.9	*	*	*	*
2001 6/.....	4.4	16.8	5.0	*	3.6	5.9	7.7	4.6	2.9	2.7	4.4	*
2000.....	4.1	23.3	5.8	1.1	4.3	5.0	6.5	2.7	2.0	2.3	3.5	*
1999.....	4.0	19.6	5.7	1.1	3.6	6.1	5.6	2.9	1.9	*	4.0	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	1.8	2.1	0.3	0.1	1.4	2.1	2.9	3.4	1.8	0.9	1.0	1.3
2006.....	1.7	2.2	0.3	0.1	1.4	2.0	2.9	3.1	1.6	0.9	1.0	1.4
2005.....	1.6	2.5	0.4	0.2	1.2	1.8	2.7	3.1	1.4	0.8	0.8	1.5
2004.....	1.7	2.6	0.3	0.2	1.2	2.0	3.1	3.1	1.4	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002.....	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001.....	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999.....	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4

All origins 3/, male

2007.....	2.3	2.5	0.3	0.2	2.0	2.8	3.6	4.0	2.3	1.2	1.5	1.6
2006.....	2.2	2.7	0.3	0.2	2.1	2.8	3.5	3.9	1.9	1.1	1.4	2.2
2005.....	2.0	3.2	0.5	0.2	1.6	2.3	3.2	3.6	1.7	1.1	0.9	1.6
2004.....	2.2	3.1	0.3	0.2	1.8	2.7	4.0	4.0	1.7	1.1	1.0	1.9
2003.....	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002.....	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001.....	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000.....	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999.....	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1

All origins 3/, female

2007.....	1.3	1.6	0.3	*	0.6	1.5	2.2	2.9	1.4	0.7	0.7	1.1
2006.....	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005.....	1.3	1.8	0.3	0.1	0.6	1.3	2.3	2.6	1.1	0.7	0.7	1.4
2004.....	1.2	2.0	0.3	0.1	0.6	1.3	2.3	2.3	1.1	0.6	0.9	1.6
2003.....	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002.....	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000.....	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999.....	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	2.0	2.1	0.3	0.1	1.5	2.5	3.2	3.7	1.9	1.0	1.0	1.3
2006.....	1.9	2.3	0.3	0.2	1.6	2.4	3.2	3.3	1.6	0.9	1.1	1.3
2005.....	1.7	2.7	0.5	0.2	1.2	2.1	3.0	3.2	1.4	0.9	0.8	1.5
2004.....	1.8	2.8	0.3	0.2	1.3	2.2	3.4	3.3	1.4	0.9	0.9	1.6
2003.....	1.9	2.2	0.2	0.1	1.5	2.3	3.9	3.2	1.3	0.7	0.7	1.1
2002.....	1.8	2.7	0.4	0.2	1.3	2.3	3.6	3.0	1.2	0.7	0.8	1.4
2001.....	1.6	2.4	0.3	0.1	1.1	2.0	3.3	2.6	0.9	0.5	0.9	0.9
2000.....	1.5	1.9	0.3	0.2	1.0	1.8	3.0	2.4	0.9	0.6	0.6	1.4
1999.....	1.5	1.6	0.3	0.1	1.2	2.0	3.1	2.3	0.9	0.6	0.8	1.4

Non-Hispanic 4/, male

2007.....	2.5	2.8	0.3	0.2	2.2	3.2	4.1	4.3	2.3	1.2	1.5	1.6
2006.....	2.4	2.7	0.4	0.2	2.3	3.2	3.8	4.1	1.9	1.1	1.5	2.2
2005.....	2.1	3.6	0.6	0.2	1.8	2.7	3.5	3.8	1.7	1.1	0.9	1.7
2004.....	2.4	3.3	0.3	0.2	1.9	3.1	4.3	4.1	1.7	1.1	1.0	1.7
2003.....	2.5	3.0	*	0.2	2.2	3.1	4.9	4.2	1.7	0.8	0.9	*
2002.....	2.3	2.8	0.3	0.2	1.8	3.2	4.7	3.6	1.6	1.0	0.9	2.1
2001.....	2.1	2.7	*	0.2	1.7	2.8	4.4	3.5	1.2	0.7	1.2	*
2000.....	2.0	1.6	0.4	0.3	1.5	2.6	4.2	3.2	1.3	0.8	1.0	1.7
1999.....	2.1	1.9	*	0.2	1.8	2.7	4.2	3.1	1.3	0.9	1.1	2.1

Non-Hispanic 4/, female

2007.....	1.5	1.3	*	*	0.7	1.8	2.4	3.1	1.5	0.8	0.7	1.1
2006.....	1.4	1.9	0.3	*	0.8	1.5	2.6	2.6	1.3	0.7	0.8	1.0
2005.....	1.4	1.9	0.4	*	0.6	1.5	2.5	2.7	1.2	0.7	0.7	1.4
2004.....	1.3	2.3	0.3	*	0.6	1.4	2.6	2.5	1.1	0.7	0.9	1.5
2003.....	1.3	1.3	*	*	0.8	1.5	2.8	2.3	1.0	0.7	0.6	1.0
2002.....	1.3	2.6	0.4	0.2	0.7	1.4	2.6	2.3	0.9	0.5	0.7	1.1
2001.....	1.0	2.0	0.4	*	0.5	1.2	2.2	1.7	0.6	0.4	0.7	0.8
2000.....	0.9	2.2	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999.....	0.9	*	*	*	0.5	1.2	2.0	1.5	0.6	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic white, both sexes

2007 5/.....	2.0	1.5	0.3	0.1	1.6	2.9	3.5	3.8	1.8	0.9	0.9	1.3
2006 5/.....	1.9	2.0	0.2	0.2	1.8	2.7	3.3	3.3	1.5	0.9	1.0	1.1
2005 5/.....	1.8	2.2	0.4	0.2	1.3	2.4	3.1	3.3	1.4	0.9	0.7	1.5
2004 5/.....	1.9	2.3	0.3	0.1	1.5	2.5	3.6	3.4	1.4	0.8	0.9	1.6
2003 5/.....	2.0	2.0	*	0.1	1.7	2.6	4.1	3.3	1.3	0.7	0.7	1.2
2002.....	1.8	1.8	0.4	0.2	1.4	2.4	3.7	2.9	1.2	0.7	0.7	1.5
2001.....	1.6	1.7	0.3	0.1	1.3	2.2	3.3	2.5	0.9	0.5	0.9	1.0
2000.....	1.4	1.5	0.2	0.1	1.1	1.9	3.0	2.3	0.8	0.5	0.7	1.3
1999.....	1.4	1.3	*	0.1	1.2	2.0	3.0	2.1	0.9	0.6	0.8	1.4

Non-Hispanic white, male

2007 5/.....	2.5	2.2	*	*	2.3	3.7	4.2	4.2	2.1	1.1	1.3	1.7
2006 5/.....	2.4	2.4	*	0.2	2.6	3.6	3.8	3.9	1.7	0.9	1.2	1.9
2005 5/.....	2.1	3.0	0.4	0.2	1.9	2.9	3.5	3.7	1.6	1.0	0.7	1.8
2004 5/.....	2.4	2.7	*	0.2	2.2	3.4	4.5	4.1	1.6	1.0	0.9	1.9
2003 5/.....	2.5	2.4	*	0.2	2.4	3.4	5.1	4.0	1.5	0.7	0.9	*
2002.....	2.3	1.9	*	0.2	2.0	3.4	4.8	3.4	1.5	0.8	0.9	2.2
2001.....	2.1	2.0	*	0.2	1.8	3.0	4.3	3.3	1.1	0.7	1.2	*
2000.....	2.0	*	*	0.2	1.6	2.6	4.1	3.0	1.1	0.7	1.0	*
1999.....	2.0	*	*	0.2	1.8	2.7	4.0	2.6	1.1	0.8	1.1	2.1

Non-Hispanic white, female

2007 5/.....	1.6	*	*	*	0.8	2.1	2.7	3.4	1.5	0.8	0.7	1.1
2006 5/.....	1.5	*	*	*	0.9	1.7	2.9	2.7	1.3	0.8	0.8	0.7
2005 5/.....	1.5	*	*	*	0.7	1.8	2.6	2.9	1.3	0.7	0.7	1.4
2004 5/.....	1.4	2.0	*	*	0.7	1.6	2.8	2.7	1.2	0.6	0.9	1.5
2003 5/.....	1.4	*	*	*	1.0	1.7	3.1	2.5	1.1	0.6	0.6	1.1
2002.....	1.3	1.8	*	0.2	0.8	1.5	2.7	2.4	0.9	0.6	0.6	1.1
2001.....	1.1	*	*	*	0.6	1.4	2.4	1.8	0.7	0.4	0.6	0.8
2000.....	0.9	*	*	*	0.6	1.2	1.9	1.5	0.6	0.3	0.4	1.2
1999.....	1.0	*	*	*	0.5	1.3	1.9	1.5	0.7	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic black, both sexes

2007 5/.....	2.0	4.2	*	*	1.1	1.6	3.1	4.5	3.0	1.5	*	*
2006 5/.....	1.9	3.5	0.8	*	0.9	1.6	3.4	4.1	2.4	1.3	*	*
2005 5/.....	1.8	4.9	0.9	*	0.9	1.7	3.4	3.8	1.7	*	*	*
2004 5/.....	1.9	4.9	*	0.4	0.9	1.9	3.4	4.0	1.8	1.3	*	*
2003 5/.....	1.8	3.3	*	*	0.8	1.8	3.9	4.1	2.0	1.3	*	*
2002.....	2.0	5.7	*	*	0.9	2.3	4.3	4.3	1.5	*	*	*
2001.....	1.8	5.9	*	*	0.7	1.7	4.0	3.8	1.2	*	*	*
2000.....	1.9	*	*	0.3	0.9	2.0	4.1	3.7	1.8	*	*	*
1999.....	2.0	*	*	*	1.3	2.3	4.3	4.2	1.2	*	*	*

Non-Hispanic black, male

2007 5/.....	2.8	*	*	*	1.8	2.1	4.5	6.3	4.8	*	*	*
2006 5/.....	2.8	*	*	*	1.4	2.6	4.9	6.4	3.7	*	*	*
2005 5/.....	2.5	*	*	*	1.3	2.4	4.4	5.7	2.9	*	*	*
2004 5/.....	2.7	*	*	*	1.4	2.8	4.8	6.0	3.0	*	*	*
2003 5/.....	2.8	*	*	*	1.4	3.0	5.4	6.6	3.3	*	*	*
2002.....	2.9	*	*	*	1.3	3.4	6.2	6.5	2.6	*	*	*
2001.....	2.8	6.4	*	*	1.3	2.7	6.3	6.3	2.4	*	*	*
2000.....	2.9	*	*	*	1.6	3.2	6.1	6.0	3.0	*	*	*
1999.....	3.1	*	*	*	2.0	3.7	6.6	7.3	2.6	*	*	*

Non-Hispanic black, female

2007 5/.....	1.2	*	*	*	*	1.1	1.9	2.9	1.5	*	*	*
2006 5/.....	1.1	*	*	*	*	0.8	2.0	2.1	1.3	*	*	*
2005 5/.....	1.2	*	*	*	*	1.0	2.6	2.2	*	*	*	*
2004 5/.....	1.1	*	*	*	*	1.0	2.2	2.3	*	*	*	*
2003 5/.....	1.0	*	*	*	*	*	2.5	2.0	*	*	*	*
2002.....	1.2	*	*	*	*	1.3	2.7	2.4	*	*	*	*
2001.....	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000.....	1.0	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	1.0	2.4	1.6	*	*	*	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	1.7	2.1	0.3	0.1	1.2	2.0	2.8	3.3	1.7	0.9	1.0	1.3
2006.....	1.6	2.2	0.3	0.1	1.2	1.9	2.8	3.1	1.5	0.8	1.0	1.4
2005.....	1.5	2.5	0.4	0.1	1.0	1.8	2.7	3.0	1.3	0.8	0.8	1.4
2004.....	1.6	2.6	0.3	0.1	1.1	1.9	3.0	3.1	1.3	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.2	2.0	3.5	3.0	1.2	0.7	0.7	1.1
2002.....	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001.....	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000.....	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999.....	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3

All origins 3/, male

2007.....	2.1	2.5	0.3	0.1	1.7	2.6	3.5	3.8	2.1	1.1	1.4	1.5
2006.....	2.1	2.7	0.3	0.2	1.8	2.6	3.4	3.8	1.8	1.0	1.4	2.2
2005.....	1.8	3.2	0.5	0.2	1.3	2.2	3.1	3.5	1.6	1.0	0.9	1.5
2004.....	2.1	3.1	0.3	0.2	1.5	2.6	3.8	3.8	1.6	0.9	0.9	1.8
2003.....	2.2	2.4	*	0.1	1.7	2.7	4.5	3.9	1.5	0.7	0.8	1.4
2002.....	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001.....	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000.....	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999.....	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8

All origins 3/, female

2007.....	1.3	1.6	0.3	*	0.6	1.5	2.1	2.8	1.4	0.7	0.7	1.1
2006.....	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005.....	1.2	1.8	0.3	*	0.6	1.3	2.2	2.5	1.1	0.6	0.7	1.4
2004.....	1.2	2.0	0.3	*	0.6	1.2	2.3	2.3	1.1	0.6	0.8	1.5
2003.....	1.2	1.3	*	*	0.7	1.3	2.5	2.2	1.0	0.7	0.6	0.9
2002.....	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000.....	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999.....	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	1.9	2.1	0.3	0.1	1.3	2.4	3.1	3.6	1.8	0.9	1.0	1.3
2006.....	1.8	2.3	0.3	0.2	1.4	2.2	3.1	3.3	1.5	0.8	1.1	1.3
2005.....	1.7	2.7	0.4	0.1	1.0	2.0	2.9	3.2	1.4	0.8	0.8	1.5
2004.....	1.7	2.8	0.3	0.2	1.1	2.1	3.3	3.2	1.3	0.8	0.9	1.5
2003.....	1.8	2.2	0.2	0.1	1.3	2.2	3.8	3.2	1.3	0.7	0.7	1.1
2002.....	1.7	2.7	0.3	0.2	1.1	2.2	3.5	2.9	1.2	0.7	0.7	1.4
2001.....	1.5	2.4	0.3	0.1	1.0	1.9	3.2	2.5	0.8	0.5	0.8	0.9
2000.....	1.4	1.9	0.3	0.1	0.9	1.7	2.9	2.3	0.9	0.5	0.6	1.3
1999.....	1.4	1.6	0.3	0.1	0.8	1.8	2.9	2.2	0.9	0.5	0.7	1.3

Non-Hispanic 4/, male

2007.....	2.3	2.8	0.3	0.1	1.9	3.0	3.9	4.1	2.2	1.1	1.4	1.5
2006.....	2.3	2.7	0.3	0.2	2.0	3.1	3.7	4.1	1.8	1.0	1.4	2.2
2005.....	2.0	3.6	0.5	0.2	1.5	2.5	3.4	3.7	1.6	1.0	0.8	1.6
2004.....	2.2	3.3	0.3	0.2	1.7	3.0	4.2	4.0	1.6	0.9	0.9	1.7
2003.....	2.3	3.0	*	0.1	1.9	3.0	4.8	4.1	1.5	0.7	0.8	*
2002.....	2.2	2.8	0.3	0.2	1.5	3.0	4.5	3.5	1.5	0.9	0.8	2.1
2001.....	2.0	2.7	*	0.2	1.4	2.6	4.2	3.4	1.1	0.6	1.0	*
2000.....	1.9	1.6	0.4	0.2	1.2	2.4	4.1	3.1	1.2	0.7	0.9	*
1999.....	1.9	1.9	*	0.1	1.2	2.5	4.0	2.9	1.2	0.7	0.9	1.7

Non-Hispanic 4/, female

2007.....	1.4	1.3	*	*	0.6	1.7	2.4	3.1	1.5	0.7	0.7	1.1
2006.....	1.3	1.9	0.3	*	0.8	1.4	2.6	2.5	1.3	0.7	0.8	1.0
2005.....	1.3	1.9	0.3	*	0.6	1.5	2.5	2.7	1.1	0.7	0.7	1.4
2004.....	1.3	2.3	0.3	*	0.6	1.3	2.5	2.4	1.1	0.7	0.9	1.5
2003.....	1.3	1.3	*	*	0.8	1.4	2.7	2.3	1.0	0.7	0.6	1.0
2002.....	1.2	2.6	0.4	0.2	0.7	1.3	2.5	2.3	0.9	0.5	0.6	1.1
2001.....	1.0	2.0	0.4	*	0.5	1.2	2.2	1.6	0.6	0.3	0.7	0.8
2000.....	0.9	2.2	*	*	0.4	1.0	1.8	1.5	0.6	0.3	0.4	1.2
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	0.6	0.4	0.5	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic white, both sexes

2007 5/.....	2.0	1.5	0.3	0.1	1.5	2.8	3.4	3.6	1.7	0.8	0.9	1.3
2006 5/.....	1.9	2.0	*	0.1	1.6	2.6	3.3	3.3	1.4	0.8	0.9	1.1
2005 5/.....	1.7	2.2	0.4	0.1	1.2	2.3	3.0	3.2	1.4	0.8	0.7	1.5
2004 5/.....	1.8	2.3	0.3	0.1	1.3	2.4	3.5	3.3	1.3	0.7	0.9	1.6
2003 5/.....	1.9	2.0	*	0.1	1.6	2.5	4.0	3.2	1.2	0.6	0.7	1.2
2002.....	1.7	1.8	0.4	0.2	1.3	2.3	3.6	2.8	1.2	0.6	0.7	1.5
2001.....	1.5	1.7	0.3	0.1	1.1	2.0	3.2	2.5	0.8	0.4	0.8	0.9
2000.....	1.4	1.5	0.2	0.1	0.9	1.8	2.9	2.2	0.8	0.5	0.6	1.2
1999.....	1.3	1.3	*	0.1	0.9	1.9	2.8	2.0	0.9	0.5	0.7	1.3

Non-Hispanic white, male

2007 5/.....	2.4	2.2	*	*	2.2	3.5	4.1	4.0	1.9	0.9	1.2	1.7
2006 5/.....	2.3	2.4	*	0.2	2.3	3.4	3.7	3.8	1.6	0.9	1.1	1.9
2005 5/.....	2.0	3.0	0.4	0.2	1.7	2.8	3.4	3.6	1.5	0.9	0.7	1.7
2004 5/.....	2.3	2.7	*	*	1.9	3.3	4.3	4.0	1.5	0.8	0.8	1.8
2003 5/.....	2.4	2.4	*	*	2.2	3.3	5.0	3.9	1.3	0.6	0.8	*
2002.....	2.2	1.9	*	0.2	1.7	3.2	4.6	3.3	1.4	0.7	0.8	2.2
2001.....	2.0	2.0	*	0.2	1.6	2.7	4.1	3.2	1.0	0.6	1.0	*
2000.....	1.8	*	*	0.2	1.4	2.4	4.0	2.9	1.0	0.6	0.9	*
1999.....	1.8	*	*	0.2	1.4	2.5	3.7	2.5	1.0	0.7	0.9	*

Non-Hispanic white, female

2007 5/.....	1.6	*	*	*	0.7	2.0	2.6	3.3	1.5	0.8	0.6	1.1
2006 5/.....	1.4	*	*	*	0.8	1.7	2.8	2.7	1.3	0.7	0.8	0.7
2005 5/.....	1.4	*	*	*	0.7	1.7	2.6	2.9	1.3	0.7	0.7	1.4
2004 5/.....	1.4	2.0	*	*	0.7	1.5	2.7	2.6	1.2	0.6	0.9	1.5
2003 5/.....	1.4	*	*	*	0.9	1.7	3.0	2.5	1.1	0.6	0.6	1.1
2002.....	1.3	1.8	*	*	0.8	1.5	2.7	2.4	0.9	0.5	0.6	1.1
2001.....	1.1	*	*	*	0.6	1.3	2.3	1.7	0.7	0.3	0.6	0.8
2000.....	0.9	*	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999.....	0.9	*	*	*	0.4	1.3	1.9	1.5	0.7	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic black, both sexes												
2007 5/.....	1.8	4.2	*	*	0.6	1.4	3.1	4.3	2.9	1.4	*	*
2006 5/.....	1.8	3.5	0.8	*	0.5	1.4	3.2	4.0	2.4	1.3	*	*
2005 5/.....	1.7	4.9	0.8	*	0.6	1.6	3.4	3.8	1.6	*	*	*
2004 5/.....	1.8	4.9	*	0.3	0.7	1.7	3.4	3.9	1.7	1.2	*	*
2003 5/.....	1.8	3.3	*	*	0.5	1.7	3.8	4.1	1.9	1.3	*	*
2002.....	1.9	5.7	*	*	0.6	2.1	4.2	4.3	1.5	*	*	*
2001.....	1.7	5.9	*	*	0.5	1.7	3.9	3.7	1.1	*	*	*
2000.....	1.7	*	*	*	0.6	1.7	4.0	3.7	1.6	*	*	*
1999.....	1.8	*	*	*	0.6	2.0	4.2	4.1	1.2	*	*	*
Non-Hispanic black, male												
2007 5/.....	2.6	*	*	*	1.0	1.8	4.4	6.1	4.7	*	*	*
2006 5/.....	2.5	*	*	*	0.7	2.1	4.7	6.3	3.7	*	*	*
2005 5/.....	2.4	*	*	*	0.8	2.2	4.3	5.7	2.9	*	*	*
2004 5/.....	2.5	*	*	*	1.0	2.6	4.7	5.8	3.0	*	*	*
2003 5/.....	2.6	*	*	*	0.9	2.7	5.4	6.5	3.3	*	*	*
2002.....	2.7	*	*	*	0.8	3.0	6.0	6.5	2.6	*	*	*
2001.....	2.6	6.4	*	*	0.8	2.5	6.3	6.2	2.1	*	*	*
2000.....	2.6	*	*	*	1.0	2.7	5.9	5.9	2.7	*	*	*
1999.....	2.8	*	*	*	0.9	3.2	6.4	7.1	2.5	*	*	*
Non-Hispanic black, female												
2007 5/.....	1.1	*	*	*	*	1.0	1.9	2.8	1.5	*	*	*
2006 5/.....	1.0	*	*	*	*	0.7	2.0	2.0	1.3	*	*	*
2005 5/.....	1.1	*	*	*	*	1.0	2.5	2.2	*	*	*	*
2004 5/.....	1.1	*	*	*	*	0.9	2.2	2.3	*	*	*	*
2003 5/.....	1.0	*	*	*	*	*	2.4	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.3	2.6	2.4	*	*	*	*
2001.....	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000.....	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	0.9	2.3	1.6	*	*	*	*











Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/, both sexes

2007.....	0.9	0.5	0.1	0.1	0.1	0.2	0.3	0.6	1.2	2.5	5.5	7.9
2006.....	0.8	0.6	0.1	*	0.1	0.2	0.3	0.6	1.1	2.5	5.3	8.3
2005.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.2	2.6	5.8	9.1
2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.1	10.6
2003.....	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002.....	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001.....	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000.....	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999.....	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7

All origins 3/, male

2007.....	0.8	*	*	*	*	0.1	0.3	0.6	1.3	2.7	6.4	9.8
2006.....	0.8	*	*	*	0.1	0.1	0.2	0.7	1.1	2.4	6.0	8.8
2005.....	0.8	*	*	*	*	0.1	0.3	0.6	1.3	2.9	6.3	10.8
2004.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.6	3.3	7.4	11.9
2003.....	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002.....	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001.....	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000.....	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999.....	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5

All origins 3/, female

2007.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.3	4.9	6.9
2006.....	0.9	*	*	*	0.1	0.2	0.3	0.6	1.1	2.6	4.8	8.1
2005.....	1.0	*	*	*	*	0.2	0.4	0.7	1.1	2.3	5.5	8.3
2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.2	9.9
2003.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002.....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000.....	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999.....	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2



Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 4/, both sexes

2007.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.3	2.6	5.6	8.0
2006.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.2	2.5	5.4	8.6
2005.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.6	5.8	9.2
2004.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.1	6.1	10.8
2003.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	2.9	6.3	11.4
2002.....	1.1	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.6
2001.....	1.2	0.7	*	*	0.1	0.2	0.4	0.7	1.6	3.1	6.9	12.9
2000.....	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.6	3.5	6.7	13.7
1999.....	1.1	1.6	0.2	*	0.1	0.2	0.4	0.6	1.3	3.3	6.8	12.8

Non-Hispanic 4/, male

2007.....	0.9	*	*	*	*	0.2	0.3	0.6	1.3	2.8	6.5	10.1
2006.....	0.9	*	*	*	0.1	0.1	0.3	0.7	1.1	2.4	6.2	9.1
2005.....	0.9	*	*	*	*	0.1	0.3	0.7	1.3	3.0	6.4	11.2
2004.....	1.0	*	*	*	0.1	0.2	0.3	0.7	1.6	3.4	7.4	12.5
2003.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	7.0	13.2
2002.....	1.0	*	*	0.1	*	0.3	0.4	0.7	1.4	3.1	6.3	12.9
2001.....	1.1	*	*	*	*	0.2	0.4	0.8	1.7	3.3	7.7	14.5
2000.....	1.1	*	*	*	*	0.3	0.4	0.8	1.6	4.0	8.2	16.1
1999.....	1.0	1.8	*	*	*	0.2	0.4	0.6	1.5	3.5	7.9	16.8

Non-Hispanic 4/, female

2007.....	1.0	*	*	*	*	0.2	0.4	0.6	1.2	2.3	5.0	7.1
2006.....	1.0	*	*	*	*	0.2	0.3	0.6	1.2	2.6	4.9	8.3
2005.....	1.0	*	*	*	*	0.2	0.4	0.7	1.1	2.3	5.4	8.4
2004.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.3	10.1
2003.....	1.1	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.8	10.6
2002.....	1.2	*	*	*	*	0.2	0.4	0.7	1.3	3.1	6.2	11.1
2001.....	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.0	6.4	12.2
2000.....	1.2	*	*	*	*	0.2	0.3	0.8	1.6	3.1	5.7	12.7
1999.....	1.2	1.4	*	*	*	0.2	0.4	0.6	1.2	3.1	6.1	11.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic white, both sexes

2007 5/.....	1.0	*	*	*	*	0.2	0.3	0.5	1.2	2.4	5.4	7.7
2006 5/.....	1.0	*	*	*	0.1	0.2	0.3	0.6	1.0	2.4	5.2	8.5
2005 5/.....	1.0	*	*	*	0.1	0.2	0.3	0.6	1.1	2.3	5.7	9.4
2004 5/.....	1.1	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	5.9	10.6
2003 5/.....	1.1	*	*	*	0.1	0.2	0.4	0.6	1.3	2.8	6.2	11.2
2002.....	1.1	*	*	*	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.5
2001.....	1.2	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.2
2000.....	1.2	*	*	*	*	0.2	0.3	0.8	1.4	3.3	6.5	13.2
1999.....	1.1	1.3	*	*	0.1	0.2	0.3	0.5	1.2	3.1	6.5	12.1

Non-Hispanic white, male

2007 5/.....	1.0	*	*	*	*	0.2	0.3	0.5	1.2	2.6	6.5	9.8
2006 5/.....	0.9	*	*	*	*	*	0.3	0.6	1.0	2.3	6.1	9.1
2005 5/.....	1.0	*	*	*	*	0.2	0.3	0.6	1.2	2.8	6.3	11.6
2004 5/.....	1.1	*	*	*	*	0.2	0.3	0.7	1.5	3.3	7.4	12.3
2003 5/.....	1.1	*	*	*	*	0.2	0.4	0.6	1.4	3.1	7.1	13.1
2002.....	1.0	*	*	*	*	0.3	0.4	0.6	1.3	2.9	6.3	12.4
2001.....	1.0	*	*	*	*	0.2	0.3	0.6	1.4	2.9	7.8	13.5
2000.....	1.2	*	*	*	*	0.2	0.4	0.8	1.4	3.7	8.2	15.9
1999.....	1.1	*	*	*	*	0.2	0.3	0.5	1.3	3.3	7.7	15.8

Non-Hispanic white, female

2007 5/.....	1.0	*	*	*	*	0.2	0.4	0.5	1.1	2.2	4.7	6.8
2006 5/.....	1.0	*	*	*	*	0.2	0.3	0.5	1.0	2.5	4.6	8.2
2005 5/.....	1.0	*	*	*	*	0.2	0.4	0.6	1.1	1.9	5.3	8.3
2004 5/.....	1.1	*	*	*	*	0.2	0.3	0.6	1.1	2.6	5.0	9.9
2003 5/.....	1.1	*	*	*	*	*	0.3	0.6	1.1	2.5	5.5	10.4
2002.....	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	6.0	11.1
2001.....	1.3	*	*	*	*	0.2	0.3	0.6	1.4	2.7	6.2	11.6
2000.....	1.2	*	*	*	*	0.2	0.3	0.7	1.4	3.0	5.4	12.0
1999.....	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	5.7	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic black, both sexes

2007 5/.....	1.1	*	*	*	*	*	0.6	1.2	2.1	4.7	9.0	12.5
2006 5/.....	1.1	*	*	*	*	*	0.6	1.4	2.7	4.1	9.0	10.4
2005 5/.....	1.1	*	*	*	*	*	0.5	1.6	2.5	5.5	9.0	9.3
2004 5/.....	1.2	*	*	*	*	*	0.8	1.3	2.6	5.1	9.5	14.1
2003 5/.....	1.2	*	*	*	*	*	0.6	1.4	2.8	4.6	9.0	15.2
2002.....	1.2	*	*	*	*	*	0.9	1.7	2.7	4.7	8.5	14.2
2001.....	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.4	9.3	23.7
2000.....	1.4	*	*	*	*	0.4	0.6	1.4	3.6	6.2	10.6	21.3
1999.....	1.3	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.4

Non-Hispanic black, male

2007 5/.....	1.0	*	*	*	*	*	*	0.9	2.5	5.5	10.2	*
2006 5/.....	1.0	*	*	*	*	*	*	1.4	2.7	4.0	9.7	*
2005 5/.....	1.0	*	*	*	*	*	*	1.5	3.1	5.0	9.5	*
2004 5/.....	1.0	*	*	*	*	*	*	1.0	3.0	4.7	10.3	*
2003 5/.....	1.0	*	*	*	*	*	*	1.9	2.9	3.9	7.8	*
2002.....	1.1	*	*	*	*	*	*	1.4	2.6	5.4	7.3	*
2001.....	1.5	*	*	*	*	*	1.0	1.9	4.3	7.9	9.0	30.0
2000.....	1.2	*	*	*	*	*	*	1.3	3.2	7.8	10.2	*
1999.....	1.2	*	*	*	*	*	*	1.4	2.9	6.0	11.9	23.8

Non-Hispanic black, female

2007 5/.....	1.2	*	*	*	*	*	0.7	1.5	1.7	4.1	8.3	10.9
2006 5/.....	1.2	*	*	*	*	*	0.8	1.3	2.8	4.2	8.5	10.1
2005 5/.....	1.3	*	*	*	*	*	0.7	1.6	2.1	5.8	8.7	9.2
2004 5/.....	1.4	*	*	*	*	*	1.0	1.4	2.4	5.4	9.0	13.7
2003 5/.....	1.3	*	*	*	*	*	0.8	1.0	2.6	5.0	9.6	15.3
2002.....	1.4	*	*	*	*	*	1.1	1.9	2.8	4.2	9.1	11.7
2001.....	1.4	*	*	*	*	*	0.7	1.4	2.7	5.3	9.4	21.4
2000.....	1.6	*	*	*	*	*	*	1.5	3.9	5.2	10.8	22.2
1999.....	1.4	*	*	*	*	*	*	1.2	2.4	4.7	9.8	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

All origins 3/, both sexes

2007.....	7.7	*	*	*	0.5	1.8	7.3	18.3	20.2	15.5	9.7	4.6
2006.....	7.4	*	*	*	0.4	1.6	7.4	17.6	19.4	14.7	9.7	4.9
2005.....	7.3	*	*	*	0.4	1.3	7.5	17.6	19.5	15.1	9.2	4.6
2004.....	7.2	*	*	*	0.3	1.5	7.6	17.3	18.7	15.7	9.3	4.3
2003.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
2002.....	7.0	*	*	*	0.3	1.5	8.1	16.8	18.4	15.5	9.3	4.4
2001.....	7.1	*	*	*	0.3	1.6	8.3	17.2	18.2	15.5	9.6	5.0
2000.....	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4
1999.....	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5

All origins 3/, male

2007.....	11.7	*	*	*	0.7	2.7	10.2	27.4	32.5	26.4	17.8	9.6
2006.....	11.2	*	*	*	0.6	2.4	10.3	26.5	31.1	24.9	17.1	9.5
2005.....	11.1	*	*	*	0.6	2.0	10.6	26.7	31.6	25.0	17.0	9.3
2004.....	11.0	*	*	*	0.5	2.3	11.0	26.4	30.2	26.5	17.1	9.0
2003.....	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
2002.....	10.8	*	*	*	0.4	2.3	11.5	26.3	30.0	26.0	16.6	9.7
2001.....	10.8	*	*	*	0.4	2.3	11.9	26.6	29.7	26.3	17.2	11.6
2000.....	10.9	*	*	*	0.4	2.5	12.5	25.9	30.0	27.1	18.8	11.9
1999.....	10.9	*	*	*	0.4	2.3	12.6	26.2	30.5	27.4	19.4	11.8

All origins 3/, female

2007.....	3.8	*	*	*	0.2	0.9	4.3	9.5	8.6	6.2	4.1	2.2
2006.....	3.7	*	*	*	0.1	0.7	4.5	8.9	8.5	6.1	4.7	2.7
2005.....	3.6	*	*	*	0.1	0.7	4.3	8.8	8.4	6.8	3.9	2.5
2004.....	3.5	*	*	*	0.2	0.7	4.2	8.6	8.1	6.6	3.9	2.3
2003.....	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
2002.....	3.4	*	*	*	*	0.7	4.7	7.7	7.6	6.8	4.5	2.1
2001.....	3.4	*	*	*	*	0.9	4.8	8.1	7.5	6.6	4.6	2.3
2000.....	3.2	*	*	*	*	0.7	4.5	7.0	8.3	6.6	4.1	2.7
1999.....	3.2	*	*	*	0.1	0.9	4.5	7.0	7.7	6.5	4.8	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Hispanic, both sexes

2007.....	6.5	*	*	*	0.6	2.4	7.5	21.2	26.1	24.5	17.6	8.9
2006.....	6.3	*	*	*	0.5	2.3	8.0	20.2	26.3	21.7	19.0	9.4
2005.....	6.2	*	*	*	0.6	2.0	8.4	20.0	25.0	25.9	14.4	*
2004.....	5.8	*	*	*	0.6	1.9	8.1	20.6	22.5	20.7	17.1	*
2003.....	6.1	*	*	*	0.6	1.8	8.3	20.2	27.6	23.0	17.8	*
2002.....	6.2	*	*	*	0.5	2.0	8.7	21.2	26.4	27.0	14.5	*
2001.....	6.4	*	*	*	0.5	2.1	9.0	22.5	28.3	26.1	20.4	*
2000.....	6.6	*	*	*	0.5	2.1	10.4	22.3	29.5	28.9	19.0	*
1999.....	6.4	*	*	*	0.4	2.1	10.2	22.5	28.7	26.2	18.7	14.0

Hispanic, male

2007.....	10.8	*	*	*	1.0	3.8	12.1	35.3	46.5	47.1	33.3	23.4
2006.....	10.2	*	*	*	0.8	3.7	12.3	33.5	46.1	41.4	34.5	*
2005.....	10.3	*	*	*	1.1	3.2	13.1	33.7	45.0	49.8	29.7	*
2004.....	9.6	*	*	*	1.0	3.2	13.1	36.0	39.3	37.7	33.3	*
2003.....	9.9	*	*	*	0.9	2.8	13.2	35.2	50.5	40.4	34.9	*
2002.....	10.3	*	*	*	0.8	3.4	14.2	36.3	48.6	52.8	26.7	*
2001.....	10.7	*	*	*	0.7	3.3	14.7	38.7	51.8	52.3	38.0	*
2000.....	11.1	*	*	*	0.8	3.7	16.8	39.1	54.8	58.7	42.3	*
1999.....	10.7	*	*	*	0.7	3.4	16.7	39.3	52.8	51.2	36.0	*

Hispanic, female

2007.....	2.0	*	*	*	*	0.6	2.2	6.6	7.2	6.1	6.6	*
2006.....	2.2	*	*	*	*	*	3.2	6.6	8.0	5.7	8.2	*
2005.....	1.9	*	*	*	*	0.6	3.1	6.0	6.7	6.6	*	*
2004.....	1.8	*	*	*	*	*	2.5	5.0	7.3	7.1	5.9	*
2003.....	1.9	*	*	*	*	0.6	2.9	5.2	6.9	9.1	6.1	*
2002.....	1.8	*	*	*	*	*	2.7	6.2	6.4	6.5	6.2	*
2001.....	2.0	*	*	*	*	0.8	2.9	6.7	7.4	5.4	8.5	*
2000.....	1.7	*	*	*	*	*	3.5	5.8	7.1	5.5	*	*
1999.....	1.9	*	*	*	*	*	3.3	6.2	7.4	6.5	7.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic 4/, both sexes

2007.....	7.9	*	*	*	0.4	1.7	7.2	17.8	19.5	14.7	9.1	4.4
2006.....	7.5	*	*	*	0.3	1.4	7.3	17.2	18.7	14.1	9.1	4.6
2005.....	7.4	*	*	*	0.3	1.2	7.3	17.2	19.0	14.2	8.9	4.4
2004.....	7.4	*	*	*	0.3	1.4	7.5	16.9	18.3	15.2	8.8	4.4
2003.....	7.2	*	*	*	0.2	1.4	8.0	16.9	17.7	14.5	8.7	4.1
2002.....	7.1	*	*	*	0.2	1.4	7.9	16.3	17.6	14.6	9.0	4.1
2001.....	7.1	*	*	*	0.2	1.5	8.2	16.5	17.2	14.7	9.0	5.0
2000.....	7.0	*	*	*	0.2	1.5	8.2	15.6	17.8	14.9	9.4	5.3
1999.....	7.0	*	*	*	0.2	1.5	8.2	15.7	17.8	15.2	10.2	5.2

Non-Hispanic 4/, male

2007.....	11.8	*	*	*	0.6	2.3	9.8	26.2	31.1	24.7	16.7	8.8
2006.....	11.3	*	*	*	0.5	2.0	9.8	25.5	29.7	23.5	15.9	8.8
2005.....	11.2	*	*	*	0.5	1.7	10.1	25.7	30.2	23.0	16.2	8.6
2004.....	11.2	*	*	*	0.4	2.1	10.6	25.1	29.2	25.6	16.2	8.9
2003.....	11.0	*	*	*	0.4	2.2	11.1	26.0	27.9	24.2	15.8	7.8
2002.....	10.7	*	*	*	0.3	2.0	11.0	25.0	28.3	24.0	16.0	9.1
2001.....	10.8	*	*	*	0.4	2.1	11.4	25.2	27.7	24.4	16.2	11.4
2000.....	10.7	*	*	*	0.3	2.2	11.8	24.4	27.9	25.0	17.6	11.7
1999.....	10.8	*	*	*	0.3	2.0	11.9	24.8	28.7	25.8	18.7	11.1

Non-Hispanic 4/, female

2007.....	4.1	*	*	*	0.2	1.0	4.7	9.8	8.7	6.1	3.9	2.3
2006.....	3.9	*	*	*	0.1	0.7	4.8	9.2	8.5	6.2	4.5	2.7
2005.....	3.8	*	*	*	*	0.7	4.5	9.0	8.5	6.8	3.9	2.5
2004.....	3.7	*	*	*	0.2	0.8	4.4	8.9	8.1	6.5	3.8	2.3
2003.....	3.6	*	*	*	*	0.6	4.9	8.2	8.2	6.3	4.0	2.5
2002.....	3.6	*	*	*	*	0.8	5.0	7.8	7.7	6.8	4.4	1.9
2001.....	3.6	*	*	*	*	0.9	5.0	8.2	7.5	6.7	4.4	2.3
2000.....	3.4	*	*	*	*	0.8	4.7	7.1	8.4	6.6	4.2	2.6
1999.....	3.4	*	*	*	*	1.0	4.6	7.0	7.7	6.5	4.7	2.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic white, both sexes

2007 5/.....	8.4	*	*	*	0.5	1.7	7.6	18.5	19.7	14.8	9.2	4.3
2006 5/.....	8.0	*	*	*	0.3	1.3	7.5	17.8	18.8	14.1	9.4	4.7
2005 5/.....	7.9	*	*	*	0.3	1.1	7.5	17.5	19.0	14.2	8.9	4.4
2004 5/.....	7.7	*	*	*	0.3	1.4	7.8	17.2	17.9	14.8	8.9	4.5
2003 5/.....	7.5	*	*	*	0.2	1.4	8.1	16.9	17.1	14.4	8.8	4.1
2002.....	7.3	*	*	*	0.2	1.3	8.1	16.1	16.8	14.4	9.0	4.0
2001.....	7.2	*	*	*	0.2	1.4	8.0	15.9	16.4	14.3	9.0	4.7
2000.....	7.0	*	*	*	0.2	1.4	7.9	14.9	16.8	14.5	9.3	5.2
1999.....	6.9	*	*	*	0.2	1.4	7.7	14.8	16.6	14.5	9.8	5.0

Non-Hispanic white, male

2007 5/.....	12.6	*	*	*	0.7	2.4	10.2	26.9	31.2	24.3	16.6	8.8
2006 5/.....	12.0	*	*	*	0.5	2.0	10.2	26.2	29.4	23.1	16.3	8.9
2005 5/.....	11.8	*	*	*	0.5	1.6	10.5	25.8	30.0	22.5	15.9	8.7
2004 5/.....	11.7	*	*	*	0.4	2.0	10.9	25.3	28.3	24.4	16.3	9.3
2003 5/.....	11.4	*	*	*	0.4	2.2	11.3	25.5	26.7	24.0	15.7	7.7
2002.....	11.1	*	*	*	0.3	1.9	11.1	24.8	26.9	23.0	15.8	8.7
2001.....	10.8	*	*	*	0.3	2.0	11.0	24.2	26.0	23.2	16.0	10.8
2000.....	10.8	*	*	*	0.3	2.1	11.4	23.3	26.2	23.9	17.3	11.4
1999.....	10.7	*	*	*	0.3	2.0	11.3	23.4	26.6	24.1	18.0	10.5

Non-Hispanic white, female

2007 5/.....	4.4	*	*	*	0.2	0.9	5.0	10.2	8.8	6.5	4.1	2.2
2006 5/.....	4.2	*	*	*	*	0.7	4.8	9.5	8.7	6.4	4.6	2.7
2005 5/.....	4.0	*	*	*	*	0.7	4.5	9.3	8.5	7.0	4.0	2.5
2004 5/.....	3.9	*	*	*	*	0.8	4.6	9.1	8.0	6.6	3.9	2.4
2003 5/.....	3.8	*	*	*	*	0.7	4.9	8.3	8.0	6.3	4.2	2.5
2002.....	3.7	*	*	*	*	0.7	5.0	7.7	7.3	7.0	4.4	2.1
2001.....	3.6	*	*	*	*	0.9	5.0	7.6	7.4	6.8	4.4	2.2
2000.....	3.4	*	*	*	*	0.8	4.4	6.5	7.8	6.5	4.1	2.7
1999.....	3.3	*	*	*	*	0.9	4.2	6.4	7.2	6.4	4.5	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic black, both sexes

2007 5/.....	5.8	*	*	*	*	1.0	4.8	15.1	21.2	17.2	9.1	5.6
2006 5/.....	5.6	*	*	*	*	1.0	5.4	14.9	20.7	15.8	8.2	*
2005 5/.....	6.1	*	*	*	*	0.9	5.8	17.3	22.0	16.7	9.9	5.5
2004 5/.....	6.3	*	*	*	*	1.0	5.8	17.4	23.9	20.3	7.8	*
2003 5/.....	6.5	*	*	*	*	1.2	7.0	18.7	24.5	17.1	9.1	*
2002.....	6.6	*	*	*	*	1.4	6.9	18.9	25.4	18.4	11.0	*
2001.....	7.5	*	*	*	*	1.4	8.9	22.8	26.2	20.7	11.2	*
2000.....	7.6	*	*	*	*	1.8	9.3	23.1	26.9	21.2	11.6	6.4
1999.....	8.0	*	*	*	*	1.4	10.1	23.7	29.6	23.4	14.8	7.8

Non-Hispanic black, male

2007 5/.....	9.2	*	*	*	*	1.5	6.8	24.0	37.2	34.8	20.9	*
2006 5/.....	8.7	*	*	*	*	1.3	7.2	23.4	36.4	30.2	15.6	*
2005 5/.....	9.4	*	*	*	*	1.2	7.8	27.7	37.2	31.0	21.0	*
2004 5/.....	10.0	*	*	*	*	1.7	9.0	27.5	42.2	40.0	14.8	*
2003 5/.....	10.3	*	*	*	*	1.9	10.0	31.6	42.4	30.9	19.3	*
2002.....	10.3	*	*	*	*	2.1	9.6	30.0	43.7	36.0	21.7	*
2001.....	11.9	*	*	*	*	1.9	13.5	36.2	46.8	40.5	22.2	*
2000.....	11.7	*	*	*	*	2.6	13.7	36.6	44.7	40.3	24.4	*
1999.....	12.5	*	*	*	*	1.8	14.7	38.0	51.6	45.7	28.8	*

Non-Hispanic black, female

2007 5/.....	2.7	*	*	*	*	*	3.0	7.5	8.4	4.6	*	*
2006 5/.....	2.9	*	*	*	*	0.7	3.9	7.7	8.1	5.5	3.9	*
2005 5/.....	3.1	*	*	*	*	*	3.9	8.4	9.7	6.4	3.7	*
2004 5/.....	2.9	*	*	*	*	*	3.0	8.8	9.3	6.3	3.9	*
2003 5/.....	3.0	*	*	*	*	*	4.4	7.6	10.1	7.3	3.3	*
2002.....	3.3	*	*	*	*	*	4.5	9.5	10.8	5.9	5.0	*
2001.....	3.5	*	*	*	*	0.9	4.8	11.2	9.5	6.7	5.1	*
2000.....	3.9	*	*	*	*	1.1	5.4	11.5	12.5	7.8	4.3	*
1999.....	4.0	*	*	*	*	1.0	6.1	11.4	11.9	7.7	7.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

All origins 3/, both sexes

2007.....	12.7	0.8	0.3	0.2	8.6	17.1	22.4	26.9	13.6	4.7	3.9	4.8
2006.....	12.8	1.1	0.2	0.1	8.6	16.8	23.3	26.7	12.2	5.3	6.0	8.1
2005.....	11.3	0.9	0.2	0.1	7.4	14.3	21.3	23.6	10.7	4.8	5.4	7.6
2004.....	10.5	0.7	0.2	0.2	7.0	12.6	20.9	21.7	9.1	4.3	4.8	6.3
2003.....	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
2002.....	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.6	3.8	5.7
2001.....	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000.....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7
1999.....	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8

All origins 3/, male

2007.....	16.1	*	0.2	0.2	12.6	23.5	27.9	31.7	16.1	5.3	3.8	5.5
2006.....	16.6	1.1	0.2	0.2	12.7	23.4	29.4	33.3	14.9	5.8	6.9	10.7
2005.....	14.5	*	0.3	0.1	10.7	19.4	26.6	29.4	13.1	5.3	6.7	9.9
2004.....	13.4	*	*	0.1	10.2	17.5	26.0	27.0	10.7	4.7	5.3	8.6
2003.....	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
2002.....	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001.....	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.7
2000.....	9.5	*	*	0.1	5.8	12.5	21.8	18.2	5.4	2.8	3.8	7.5
1999.....	9.4	*	*	*	5.0	12.7	21.9	17.5	6.0	3.4	3.8	6.2

All origins 3/, female

2007.....	9.5	*	0.3	0.2	4.4	10.5	16.9	22.2	11.3	4.1	3.9	4.4
2006.....	9.1	1.0	*	0.1	4.2	10.0	17.1	20.3	9.8	4.9	5.4	6.8
2005.....	8.2	*	*	0.1	3.8	8.9	16.0	18.1	8.4	4.3	4.5	6.6
2004.....	7.6	*	*	0.2	3.6	7.7	15.8	16.5	7.7	3.9	4.4	5.2
2003.....	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
2002.....	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.9
2001.....	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.6	2.6	3.3	4.9
2000.....	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.9
1999.....	4.4	*	*	*	2.0	5.0	9.7	8.0	3.8	2.7	3.8	4.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Hispanic, both sexes

2007.....	6.0	*	*	*	4.1	7.2	11.4	15.1	8.8	3.1	*	*
2006.....	6.5	*	*	*	3.6	8.1	13.2	16.3	9.0	3.2	3.5	*
2005.....	6.1	*	*	*	3.9	7.4	12.5	15.1	7.8	3.7	4.0	*
2004.....	5.5	*	*	*	3.3	6.3	12.1	13.1	8.3	3.1	3.5	*
2003.....	5.9	*	*	*	3.5	7.2	13.3	14.9	6.5	3.1	3.7	*
2002.....	5.5	*	*	*	3.2	6.2	13.4	13.8	5.6	3.0	3.3	*
2001.....	4.7	*	*	*	2.4	5.9	11.8	11.0	4.8	2.6	*	*
2000.....	4.8	*	*	*	2.2	6.9	12.7	10.6	4.6	1.9	*	*
1999.....	5.8	*	*	*	2.7	8.1	15.0	13.4	5.1	2.7	*	*

Hispanic, male

2007.....	8.7	*	*	*	6.0	10.3	16.4	22.3	12.4	5.6	*	*
2006.....	9.3	*	*	*	5.1	11.5	18.7	23.2	14.5	4.4	*	*
2005.....	8.9	*	*	*	5.8	10.8	17.9	22.2	13.0	4.9	*	*
2004.....	7.8	*	*	*	4.6	9.3	17.2	18.3	12.5	4.8	*	*
2003.....	8.7	*	*	*	5.2	10.5	19.5	22.1	10.5	5.2	*	*
2002.....	8.2	*	*	*	4.7	9.3	19.9	21.1	8.7	*	*	*
2001.....	7.0	*	*	*	3.3	8.7	17.8	17.3	7.9	*	*	*
2000.....	7.4	*	*	*	3.4	10.5	19.6	17.1	6.5	*	*	*
1999.....	9.2	*	*	*	4.2	12.7	23.8	21.9	9.0	4.6	*	*

Hispanic, female

2007.....	3.1	*	*	*	2.1	3.5	5.9	7.7	5.5	*	*	*
2006.....	3.4	*	*	*	2.0	4.0	6.9	9.2	4.0	*	*	*
2005.....	3.0	*	*	*	1.9	3.3	6.4	7.9	3.1	2.7	*	*
2004.....	2.9	*	*	*	1.8	2.8	6.5	7.8	4.6	*	*	*
2003.....	2.9	*	*	*	1.5	3.3	6.5	7.8	2.8	*	*	*
2002.....	2.6	*	*	*	1.4	2.5	6.3	6.6	2.9	*	*	*
2001.....	2.2	*	*	*	1.4	2.8	5.3	4.8	*	*	*	*
2000.....	2.1	*	*	*	0.9	2.7	5.2	4.2	3.0	*	*	*
1999.....	2.2	*	*	*	1.0	2.8	5.7	5.2	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic 4/, both sexes

2007.....	13.9	0.8	0.3	0.2	9.6	19.6	24.4	28.2	14.0	4.8	4.0	4.9
2006.....	13.9	1.1	0.3	0.2	9.6	18.9	25.0	27.8	12.5	5.5	6.2	8.2
2005.....	12.1	0.9	0.2	0.1	8.0	15.9	22.7	24.4	10.9	4.8	5.5	7.7
2004.....	11.2	0.9	0.2	0.2	7.7	14.1	22.3	22.5	9.2	4.3	4.8	6.3
2003.....	10.4	0.7	0.2	0.1	6.9	13.2	21.6	20.4	8.1	4.2	4.2	6.0
2002.....	9.5	0.8	0.2	0.1	5.9	12.2	20.5	18.2	6.9	3.6	3.8	5.8
2001.....	8.0	0.7	0.2	0.1	4.9	10.1	17.6	15.0	5.3	3.0	3.5	5.2
2000.....	7.2	*	*	0.1	4.3	9.1	16.3	13.3	4.8	2.7	3.6	5.7
1999.....	6.9	*	0.2	0.1	3.6	9.0	15.6	12.4	4.8	3.0	3.9	4.9

Non-Hispanic 4/, male

2007.....	17.4	*	0.3	0.2	14.0	27.1	30.1	32.8	16.3	5.3	3.9	5.7
2006.....	17.9	*	*	0.2	14.4	26.6	31.3	34.3	14.8	5.9	7.1	10.8
2005.....	15.4	*	0.3	0.1	11.7	21.7	28.0	30.0	13.0	5.3	6.8	10.0
2004.....	14.3	*	*	0.1	11.4	19.5	27.4	27.8	10.5	4.6	5.3	8.6
2003.....	13.5	*	*	*	10.1	18.0	27.5	25.9	9.2	4.5	4.9	7.8
2002.....	12.3	*	*	0.1	8.4	17.0	26.3	23.0	8.0	3.7	4.1	7.5
2001.....	10.6	*	*	0.1	7.3	14.1	23.2	19.7	6.0	3.5	3.9	5.8
2000.....	9.7	*	*	0.1	6.3	12.8	21.9	17.9	5.2	2.8	3.9	7.5
1999.....	9.3	*	*	*	5.1	12.7	21.4	16.8	5.6	3.3	4.0	6.3

Non-Hispanic 4/, female

2007.....	10.5	*	0.3	0.2	4.9	12.0	18.8	23.8	11.8	4.4	4.1	4.6
2006.....	10.1	*	*	0.1	4.7	11.3	18.8	21.5	10.3	5.1	5.5	7.0
2005.....	9.0	*	*	0.2	4.2	10.1	17.4	19.1	8.9	4.4	4.6	6.7
2004.....	8.3	*	*	0.2	3.9	8.7	17.2	17.4	7.9	4.1	4.5	5.3
2003.....	7.6	*	*	*	3.7	8.4	15.9	15.2	7.0	3.9	3.7	5.3
2002.....	6.9	*	*	*	3.3	7.5	14.9	13.6	5.9	3.5	3.6	5.1
2001.....	5.5	*	*	*	2.6	6.1	12.2	10.5	4.8	2.6	3.3	5.0
2000.....	4.9	*	*	*	2.3	5.5	10.8	8.7	4.5	2.5	3.3	5.0
1999.....	4.6	*	*	*	2.1	5.4	10.0	8.2	3.9	2.8	3.9	4.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic white, both sexes

2007 5/.....	15.3	*	*	0.2	11.8	23.8	27.3	29.3	13.6	4.8	4.1	5.2
2006 5/.....	14.9	*	0.2	0.2	11.9	22.5	27.0	27.9	11.8	5.4	6.4	8.4
2005 5/.....	13.1	0.9	*	0.1	9.9	18.5	24.6	24.5	10.5	4.9	5.7	8.1
2004 5/.....	12.1	*	*	0.2	9.5	16.5	24.2	22.6	8.8	4.4	5.0	6.4
2003 5/.....	11.2	*	*	0.1	8.5	15.2	23.0	20.3	7.9	4.2	4.5	6.3
2002.....	10.0	*	*	0.1	7.1	13.8	21.6	17.8	6.6	3.7	3.9	5.9
2001.....	8.3	*	*	0.1	5.9	11.3	18.1	14.2	5.1	3.1	3.7	5.2
2000.....	7.4	*	*	0.1	5.1	9.9	16.4	12.3	4.7	2.7	3.7	5.7
1999.....	6.9	*	*	0.1	4.1	9.5	15.4	11.4	4.5	3.0	4.1	4.9

Non-Hispanic white, male

2007 5/.....	19.0	*	*	0.2	17.4	32.8	33.4	33.1	15.0	4.9	4.1	6.0
2006 5/.....	18.9	*	*	0.2	17.9	31.2	33.3	33.2	12.8	5.3	7.3	11.3
2005 5/.....	16.4	*	*	*	14.5	25.3	30.2	29.1	11.6	5.1	6.9	10.7
2004 5/.....	15.3	*	*	0.2	14.0	22.8	29.5	27.3	9.2	4.3	5.4	8.7
2003 5/.....	14.3	*	*	*	12.5	20.8	29.0	25.1	8.4	4.2	5.1	7.8
2002.....	12.8	*	*	*	10.1	19.2	27.5	21.7	7.0	3.5	4.1	7.8
2001.....	10.8	*	*	*	8.6	15.8	23.7	17.8	5.1	3.4	3.8	5.7
2000.....	9.8	*	*	0.2	7.5	13.8	21.9	16.1	4.6	2.7	4.1	7.2
1999.....	9.2	*	*	*	5.8	13.2	21.0	14.7	4.7	3.2	4.0	6.6

Non-Hispanic white, female

2007 5/.....	11.7	*	*	0.2	5.9	14.6	21.2	25.6	12.4	4.7	4.2	4.8
2006 5/.....	11.1	*	*	*	5.7	13.7	20.6	22.6	10.9	5.5	5.7	7.1
2005 5/.....	9.8	*	*	*	5.0	11.7	19.0	20.0	9.4	4.8	4.8	7.0
2004 5/.....	9.1	*	*	0.2	4.7	10.1	18.8	18.0	8.4	4.5	4.7	5.4
2003 5/.....	8.2	*	*	*	4.4	9.5	17.0	15.7	7.4	4.1	4.0	5.7
2002.....	7.4	*	*	*	3.9	8.4	15.7	13.9	6.3	3.8	3.8	5.1
2001.....	5.8	*	*	*	3.0	6.8	12.5	10.6	5.1	2.8	3.6	4.9
2000.....	5.1	*	*	*	2.6	6.0	10.8	8.6	4.8	2.7	3.5	5.1
1999.....	4.8	*	*	*	2.4	5.7	9.9	8.1	4.2	2.9	4.1	4.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic black, both sexes

2007 5/.....	10.8	*	*	*	2.5	8.8	18.9	31.2	20.9	6.1	3.7	*
2006 5/.....	12.5	*	*	*	2.6	10.5	23.8	35.9	21.5	7.1	4.4	6.6
2005 5/.....	10.8	*	*	*	2.2	9.7	20.5	31.5	17.7	5.2	4.3	*
2004 5/.....	9.7	*	*	*	2.2	8.2	19.4	28.7	14.7	4.7	3.8	*
2003 5/.....	9.5	*	*	*	1.9	8.3	21.4	27.5	11.7	4.9	2.2	*
2002.....	9.4	*	*	*	2.1	8.7	21.2	27.3	11.7	2.9	2.6	*
2001.....	8.7	*	*	*	1.9	7.8	20.4	25.6	8.7	3.2	2.7	6.2
2000.....	8.5	*	*	*	2.0	8.5	20.6	24.1	7.5	3.0	2.4	7.6
1999.....	8.7	*	*	*	2.2	9.3	21.2	23.9	8.2	3.4	2.2	*

Non-Hispanic black, male

2007 5/.....	14.6	*	*	*	3.5	12.0	24.4	42.8	33.2	11.0	*	*
2006 5/.....	17.8	*	*	*	3.5	16.1	32.6	52.7	35.8	12.9	*	*
2005 5/.....	14.7	*	*	*	2.9	13.2	25.7	45.3	29.6	8.9	6.4	*
2004 5/.....	13.1	*	*	*	3.1	11.4	24.6	39.9	24.7	8.7	6.6	*
2003 5/.....	13.0	*	*	*	2.5	10.9	28.4	39.7	19.0	8.4	*	*
2002.....	13.2	*	*	*	3.0	12.2	28.3	40.1	20.2	5.0	*	*
2001.....	12.5	*	*	*	2.9	11.0	27.5	40.1	14.8	5.2	*	*
2000.....	12.3	*	*	*	2.7	12.3	28.7	38.1	12.3	5.1	*	*
1999.....	12.9	*	*	*	3.4	13.8	30.4	38.3	15.0	5.3	*	*

Non-Hispanic black, female

2007 5/.....	7.3	*	*	*	1.5	5.7	13.9	21.2	10.9	2.6	3.9	*
2006 5/.....	7.6	*	*	*	1.8	5.2	16.1	21.4	10.0	2.9	3.9	7.5
2005 5/.....	7.2	*	*	*	1.4	6.5	15.9	19.7	8.1	2.5	*	*
2004 5/.....	6.6	*	*	*	1.3	5.2	14.7	19.1	6.7	*	*	*
2003 5/.....	6.3	*	*	*	1.2	5.9	15.2	17.1	5.9	2.5	*	*
2002.....	6.0	*	*	*	1.3	5.6	14.9	16.3	4.9	*	*	*
2001.....	5.3	*	*	*	0.9	5.0	14.1	13.1	3.7	*	*	*
2000.....	5.0	*	*	*	1.3	5.0	13.4	12.1	3.6	*	*	*
1999.....	4.9	*	*	*	1.1	5.3	13.1	11.5	2.7	2.1	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

All origins 3/, both sexes

2007.....	10.4	*	0.4	0.8	16.2	15.5	11.9	11.1	10.2	10.0	13.2	11.6
2006.....	10.3	*	0.3	0.9	16.9	15.3	11.5	11.2	9.8	10.0	12.9	11.5
2005.....	10.4	*	0.4	0.8	16.2	15.7	11.6	11.2	9.8	10.3	13.7	12.0
2004.....	10.1	*	0.3	0.7	15.7	15.0	11.3	11.0	9.8	10.2	13.3	11.9
2003.....	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
2002.....	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001.....	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000.....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999.....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5

All origins 3/, male

2007.....	18.2	*	0.5	1.2	28.5	27.0	19.5	18.6	17.9	19.1	29.5	33.5
2006.....	18.1	*	0.5	1.2	29.8	26.6	18.9	18.5	17.0	19.2	29.1	33.6
2005.....	18.3	*	0.5	1.2	28.7	27.2	19.2	18.9	17.4	19.7	30.8	35.4
2004.....	17.6	*	0.4	1.1	27.5	26.1	18.7	18.3	17.1	19.7	29.8	35.9
2003.....	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
2002.....	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001.....	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000.....	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999.....	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3

All origins 3/, female

2007.....	2.7	*	0.4	0.4	3.2	3.6	4.2	3.7	3.0	2.3	2.0	1.1
2006.....	2.8	*	*	0.4	3.2	3.7	4.1	4.0	3.2	2.2	1.8	1.1
2005.....	2.7	*	0.3	0.4	3.0	3.8	4.0	3.7	2.8	2.5	2.1	1.3
2004.....	2.7	*	0.3	0.3	3.2	3.7	3.9	4.0	3.1	2.2	2.3	1.0
2003.....	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
2002.....	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001.....	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000.....	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999.....	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Hispanic, both sexes

2007.....	7.7	*	*	0.7	17.9	12.7	7.5	6.2	4.3	4.3	5.9	*
2006.....	7.8	*	*	0.9	19.1	12.6	7.8	5.4	4.0	2.5	4.7	*
2005.....	8.1	*	*	0.6	18.8	13.8	8.3	6.1	3.4	4.2	5.3	*
2004.....	7.9	*	*	0.6	18.5	12.7	7.7	6.2	4.8	4.4	5.4	*
2003.....	8.3	*	*	0.7	19.2	13.3	8.1	6.8	4.1	4.7	4.7	*
2002.....	8.1	*	*	0.8	18.5	12.5	7.9	6.0	5.0	5.3	5.5	*
2001.....	8.3	*	*	0.6	18.4	13.5	8.6	6.3	4.4	5.3	5.6	*
2000.....	8.2	*	*	0.8	18.0	12.1	8.1	8.1	4.9	5.9	5.5	*
1999.....	8.5	*	*	1.1	18.8	12.2	8.7	7.4	5.3	4.9	9.2	*

Hispanic, male

2007.....	13.4	*	*	1.0	31.4	21.1	12.4	10.6	7.7	9.0	13.0	*
2006.....	13.7	*	*	1.3	33.6	20.9	12.9	9.1	7.5	5.0	11.1	*
2005.....	14.2	*	*	0.9	33.0	22.9	13.4	10.4	6.7	8.1	12.0	*
2004.....	13.9	*	*	0.8	32.4	21.3	12.9	10.6	8.6	8.8	11.8	*
2003.....	14.6	*	*	1.0	32.8	22.8	13.3	11.5	8.1	10.0	10.7	*
2002.....	14.2	*	*	1.0	32.1	21.2	12.9	10.1	9.5	10.9	13.1	*
2001.....	14.6	*	*	0.9	31.4	22.7	14.4	10.7	8.6	11.0	12.5	*
2000.....	14.2	*	*	1.3	30.8	20.3	13.2	13.5	9.1	11.4	12.8	*
1999.....	14.6	*	*	1.5	31.6	20.6	14.3	11.8	9.9	10.1	21.9	*

Hispanic, female

2007.....	1.5	*	*	*	2.9	2.5	2.0	1.6	*	*	*	*
2006.....	1.5	*	*	*	2.7	2.4	2.1	1.7	*	*	*	*
2005.....	1.6	*	*	*	2.6	2.8	2.5	1.7	*	*	*	*
2004.....	1.5	*	*	*	2.6	2.4	2.0	1.7	*	*	*	*
2003.....	1.7	*	*	*	3.5	2.1	2.4	2.0	*	*	*	*
2002.....	1.6	*	*	*	2.8	2.3	2.5	1.9	*	*	*	*
2001.....	1.7	*	*	*	3.3	2.7	2.3	2.0	*	*	*	*
2000.....	1.8	*	*	*	2.9	2.6	2.5	2.8	*	*	*	*
1999.....	2.0	*	*	*	3.8	2.4	2.6	3.0	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic 4/, both sexes

2007.....	10.8	*	0.4	0.8	15.8	16.2	12.6	11.7	10.7	10.4	13.6	11.9
2006.....	10.7	*	0.4	0.8	16.3	15.9	12.1	11.8	10.3	10.6	13.4	11.8
2005.....	10.7	*	0.4	0.9	15.6	16.1	12.1	11.7	10.3	10.8	14.1	12.3
2004.....	10.4	*	0.4	0.8	15.0	15.5	11.8	11.5	10.2	10.6	13.8	12.1
2003.....	10.6	*	0.3	0.8	16.0	16.0	12.0	11.6	10.5	10.8	13.9	12.7
2002.....	10.8	*	0.4	0.9	16.2	15.9	12.7	11.2	10.6	11.2	14.8	12.7
2001.....	10.6	*	0.5	0.9	16.3	15.8	12.1	10.9	10.4	11.2	14.7	12.9
2000.....	10.4	*	0.3	0.9	16.4	14.9	12.3	10.7	9.7	10.8	14.3	14.4
1999.....	10.6	*	0.5	1.0	17.3	15.3	11.9	10.4	10.0	11.4	14.4	13.6

Non-Hispanic 4/, male

2007.....	19.0	*	0.5	1.2	27.8	28.5	20.8	19.6	18.8	19.9	30.5	34.7
2006.....	18.8	*	0.5	1.2	28.8	28.0	19.9	19.6	17.8	20.3	30.2	34.7
2005.....	18.9	*	0.6	1.3	27.6	28.2	20.2	19.8	18.2	20.5	31.8	36.3
2004.....	18.2	*	0.4	1.2	26.4	27.2	19.7	19.1	17.7	20.5	30.8	37.0
2003.....	18.8	*	0.4	1.3	28.3	28.0	19.9	19.5	18.4	21.0	31.2	38.7
2002.....	19.0	*	0.5	1.2	28.4	27.6	21.2	18.9	18.5	21.9	33.8	39.5
2001.....	18.7	*	0.5	1.2	29.0	27.5	19.9	18.4	17.9	21.7	33.6	40.8
2000.....	18.3	*	0.4	1.5	28.9	25.7	20.1	17.9	16.7	20.7	33.0	45.8
1999.....	18.5	*	0.5	1.5	30.3	26.3	19.3	17.4	17.6	22.1	33.2	43.9

Non-Hispanic 4/, female

2007.....	2.9	*	0.4	0.4	3.3	3.9	4.6	4.0	3.2	2.4	2.0	1.2
2006.....	2.9	*	*	0.5	3.2	3.9	4.4	4.2	3.4	2.3	1.9	1.2
2005.....	2.8	*	*	0.5	3.1	4.0	4.2	3.9	3.0	2.6	2.1	1.3
2004.....	2.9	*	*	0.3	3.3	3.9	4.2	4.3	3.2	2.3	2.3	1.0
2003.....	2.9	*	*	0.3	3.2	3.9	4.2	4.0	3.1	2.3	2.3	1.3
2002.....	3.0	*	*	0.5	3.6	4.3	4.4	3.8	3.2	2.4	2.2	1.2
2001.....	3.0	*	0.4	0.5	3.1	4.3	4.6	3.7	3.5	2.5	2.3	1.4
2000.....	3.0	*	*	0.4	3.6	4.2	4.6	3.7	3.1	2.6	2.1	1.7
1999.....	3.0	*	0.4	0.5	3.9	4.4	4.6	3.6	3.0	2.5	2.2	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic white, both sexes

2007 5/.....	9.4	*	0.3	0.6	8.1	10.8	11.6	12.1	11.5	11.5	14.8	13.1
2006 5/.....	9.1	*	*	0.6	8.2	9.7	11.1	12.2	11.2	11.6	14.6	12.6
2005 5/.....	9.2	*	0.3	0.7	8.2	10.3	11.1	12.2	11.3	11.8	15.3	13.3
2004 5/.....	9.1	*	*	0.6	8.5	10.2	11.1	12.0	10.9	11.5	14.8	13.1
2003 5/.....	9.3	*	*	0.6	8.8	10.4	11.2	12.0	11.3	11.8	14.9	13.8
2002.....	9.4	*	0.3	0.6	9.2	10.7	11.8	11.5	11.4	12.3	15.8	13.7
2001.....	9.4	*	0.3	0.8	9.4	11.4	11.4	11.4	11.2	12.1	15.7	14.0
2000.....	9.1	*	0.2	0.8	9.6	10.5	11.5	10.8	10.2	11.5	15.4	15.5
1999.....	9.3	*	0.3	0.9	10.1	11.2	11.2	10.5	10.5	12.2	15.3	14.7

Non-Hispanic white, male

2007 5/.....	16.1	*	*	0.8	13.4	18.0	18.6	20.1	20.0	21.6	32.9	38.3
2006 5/.....	15.6	*	*	0.9	13.9	16.3	17.8	19.9	19.0	21.9	32.6	37.0
2005 5/.....	15.9	*	*	1.0	13.9	16.9	17.8	20.3	19.6	22.2	34.2	39.2
2004 5/.....	15.6	*	*	0.9	14.3	16.9	17.8	19.6	18.7	21.8	32.9	40.1
2003 5/.....	16.0	*	*	0.9	15.2	17.3	18.1	19.9	19.7	22.7	33.5	42.1
2002.....	16.3	*	*	0.9	15.6	17.4	19.3	19.2	19.7	23.7	35.9	42.5
2001.....	16.3	*	*	1.2	16.0	18.9	18.4	19.0	19.0	23.1	35.7	44.0
2000.....	15.7	*	*	1.3	16.2	17.2	18.4	18.0	17.5	21.9	35.5	49.6
1999.....	15.9	*	*	1.3	17.2	18.6	17.8	17.6	18.1	23.5	35.1	47.5

Non-Hispanic white, female

2007 5/.....	2.9	*	*	0.3	2.5	3.4	4.6	4.3	3.5	2.6	2.2	1.2
2006 5/.....	2.8	*	*	0.3	2.2	3.1	4.4	4.6	3.8	2.6	2.1	1.1
2005 5/.....	2.8	*	*	0.3	2.2	3.5	4.3	4.2	3.3	2.7	2.4	1.5
2004 5/.....	2.9	*	*	0.3	2.5	3.3	4.3	4.6	3.5	2.6	2.5	1.0
2003 5/.....	2.7	*	*	0.2	2.2	3.4	4.3	4.2	3.4	2.4	2.4	1.4
2002.....	2.8	*	*	0.4	2.5	3.8	4.3	4.0	3.5	2.6	2.3	1.2
2001.....	2.9	*	*	0.4	2.5	3.7	4.4	3.8	3.9	2.7	2.4	1.5
2000.....	2.9	*	*	0.4	2.7	3.7	4.5	3.8	3.4	2.8	2.2	1.8
1999.....	2.9	*	*	0.5	2.8	3.8	4.5	3.6	3.3	2.7	2.3	1.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic black, both sexes

2007 5/.....	21.3	*	1.0	1.8	50.3	46.5	22.3	12.2	7.7	5.4	5.9	*
2006 5/.....	22.0	*	1.1	1.8	52.1	49.4	21.6	12.6	6.9	5.9	5.3	6.4
2005 5/.....	21.1	*	1.1	1.7	48.9	47.9	21.6	11.9	6.3	6.6	5.7	*
2004 5/.....	19.9	*	1.1	1.6	45.7	45.7	19.4	11.3	8.1	6.4	6.8	*
2003 5/.....	20.7	*	0.9	1.8	49.3	47.4	19.7	11.7	6.9	6.2	6.0	*
2002.....	20.7	*	*	1.6	49.2	46.4	21.0	11.9	7.2	5.8	6.9	*
2001.....	19.8	*	1.0	1.2	49.6	42.2	19.5	10.7	6.9	6.6	7.4	*
2000.....	19.8	*	*	1.4	49.8	40.9	19.7	12.1	6.8	7.0	5.6	*
1999.....	20.0	*	1.1	1.6	51.8	38.9	18.8	11.6	8.4	6.8	7.6	*

Non-Hispanic black, male

2007 5/.....	40.3	*	*	2.8	92.2	88.5	40.6	22.1	15.2	11.2	14.3	*
2006 5/.....	41.4	*	*	2.7	95.2	93.3	39.4	23.4	13.4	12.5	14.2	*
2005 5/.....	40.0	*	*	2.5	90.1	92.0	40.0	21.9	11.8	13.0	14.3	*
2004 5/.....	37.6	*	*	2.4	83.7	87.5	36.2	20.6	14.9	14.8	16.3	*
2003 5/.....	39.1	*	*	2.6	90.5	91.1	36.1	21.6	13.1	12.0	13.6	*
2002.....	38.9	*	*	2.3	89.8	89.0	37.9	21.7	13.7	12.0	17.0	*
2001.....	37.3	*	*	1.8	93.1	80.4	34.1	19.3	13.6	13.8	17.7	*
2000.....	37.1	*	*	2.2	92.2	77.6	35.3	21.7	12.8	15.4	12.3	*
1999.....	37.1	*	*	2.4	95.0	73.8	33.1	20.7	16.8	14.3	18.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic black, female

2007 5/.....	4.0	*	*	0.9	7.3	7.3	6.1	3.6	1.7	*	*	*
2006 5/.....	4.2	*	*	0.9	7.9	8.7	5.8	3.3	1.7	*	*	*
2005 5/.....	3.8	*	*	0.9	6.9	7.3	5.2	3.3	1.8	2.1	*	*
2004 5/.....	3.8	*	*	0.7	7.1	7.4	4.5	3.4	2.6	*	*	*
2003 5/.....	4.0	*	*	0.9	7.6	7.6	5.2	3.2	2.0	2.1	*	*
2002.....	4.3	*	*	0.9	8.4	7.8	6.1	3.5	2.1	*	*	*
2001.....	4.0	*	*	*	6.3	7.7	6.7	3.3	1.5	*	*	*
2000.....	4.2	*	*	*	7.9	7.7	5.9	3.9	2.0	*	*	*
1999.....	4.5	*	*	0.9	9.3	7.6	6.2	3.8	1.6	*	*	*

\* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes races other than white and black.

5/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality."

6/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

7/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.