


Figure 11.1. Percent of persons of all ages who assessed their health as excellent or very good: United States, 1997-2004




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded persons with unknown health status (about 0.5% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the appendix for more details.

DATA SOURCE: Family Core component of the 1997-2004 National Health Interview Surveys. The estimate for 2004 was based on data collected from January through September.

- For the period January through September 2004, the percentage of persons who assessed their health as excellent or very good was 66.8% (95% confidence interval = 66.2%-67.4%), which is similar to the 2003 estimate of 67.5%.
- From 1997 through 2001, the annual percentage of persons who assessed their health as excellent or very good remained similar: 68.5% in 1997, 69.1% in 1998, 68.7% in 1999, 68.3% in 2000, and 68.4% in 2001. The estimates in 2002 (67.4%) and 2003 (67.5%) were significantly lower than that in 2001.



Figure 11.2. Percent distribution of respondent-assessed health status, by sex for all ages: United States, January-September 2004




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 297 persons (0.4%) with unknown health status.

DATA SOURCE: Based on data collected from January through September in the Family Core component of the 2004 National Health Interview Survey.

- For both sexes combined, most persons assessed their health as either excellent (35.5%) or very good (31.3%). A smaller percentage of persons assessed their health as good (23.9%), and even fewer assessed their health as fair (7.0%) or poor (2.4%).
- Compared with males, females were less likely to assess their health as excellent and more likely to assess their health as good, fair, or poor.



Figure 11.3. Percent of persons of all ages who assessed their health as excellent or very good, by age group and sex: United States, January-September 2004




NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 297 persons (0.4%) with unknown health status.

DATA SOURCE: Based on data collected from January through September in the Family Core component of the 2004 National Health Interview Survey.

- For both sexes combined, the percentage of persons who assessed their health as excellent or very good decreased with age: 82.0% for those under 18 years, 66.3% for those aged 18-64 years, and 37.2% for those aged 65 years and over.
- For persons aged 18-64 years, the percentage of persons who assessed their health as excellent or very good was higher for men than for women.



Figure 11.4. Age- sex-adjusted percent of persons of all ages who assessed their health as excellent or very good, by race/ethnicity: United States, January-September 2004



NOTES: Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses excluded 297 persons (0.4%) with unknown health status. Estimates are age- sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18-64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through September in the Family Core component of the 2004 National Health Interview Survey.

- After adjusting for age and sex, the percentage of persons who assessed their health as excellent or very good was 58.8% for Hispanic persons, 69.9% for non-Hispanic white persons, and 58.2% for non-Hispanic black persons.
- The health of Hispanic persons and non-Hispanic black persons was less likely to be assessed as excellent or very good than that of non-Hispanic white persons.



Data tables for figures 11.1-11.4:

Data table for figure 11.1. Percent of persons of all ages who assessed their

health as excellent or very good: United States, 1997-2004

| Year | Percent | 95% confidence interval |
|------------------------|---------|-------------------------|
| 1997 | 68.5 | 68.0-69.1 |
| 1998 | 69.1 | 68.6-69.6 |
| 1999 | 68.7 | 68.1-69.3 |
| 2000 | 68.3 | 67.7-68.8 |
| 2001 | 68.4 | 67.9-69.0 |
| 2002 | 67.4 | 66.8-68.0 |
| 2003 | 67.5 | 66.9-68.0 |
| January-September 2004 | 66.8 | 66.2-67.4 |

Data table for figure 11.2. Percent distribution of respondent-assessed health

status, by sex for all ages: United States, January-September 2004

| Health status and sex | Percent | 95% confidence interval |
|-----------------------|---------|-------------------------|
| Excellent | | |
| Total | 35.5 | 34.8-36.2 |
| Male | 36.7 | 35.9-37.5 |
| Female | 34.4 | 33.7-35.2 |
| Very good | | |
| Total | 31.3 | 30.7-31.9 |
| Male | 31.4 | 30.7-32.1 |
| Female | 31.1 | 30.5-31.8 |
| Good | | |
| Total | 23.9 | 23.4-24.4 |
| Male | 23.4 | 22.8-24.0 |
| Female | 24.3 | 23.7-24.9 |
| Fair | | |
| Total | 7.0 | 6.7-7.2 |
| Male | 6.4 | 6.1-6.7 |
| Female | 7.5 | 7.2-7.8 |
| Poor | | |
| Total | 2.4 | 2.2-2.5 |
| Male | 2.1 | 1.9-2.3 |
| Female | 2.6 | 2.4-2.8 |



Data table for figure 11.3. Percent of persons of all ages who assessed their health as excellent or very good, by age group and sex: United States, January-September 2004

| Age and sex | Percent | 95% confidence interval |
|-------------------------------------|---------|-------------------------|
| Under 18 years | | |
| Total | 82.0 | 81.0-82.9 |
| Male | 81.5 | 80.4-82.6 |
| Female | 82.4 | 81.3-83.6 |
| 18-64 years | | |
| Total | 66.3 | 65.6-67.0 |
| Male | 67.6 | 66.8-68.4 |
| Female | 65.1 | 64.2-65.9 |
| 65 years and over | | |
| Total | 37.2 | 35.8-38.5 |
| Male | 36.4 | 34.6-38.3 |
| Female | 37.7 | 36.2-39.3 |
| All ages: crude ¹ | | |
| Total | 66.8 | 66.2-67.4 |
| Male | 68.1 | 67.4-68.7 |
| Female | 65.6 | 64.8-66.3 |
| All ages: age-adjusted ² | | |
| Total | 66.7 | 66.1-67.3 |
| Male | 67.3 | 66.6-67.9 |
| Female | 66.1 | 65.4-66.8 |

¹Crude estimates are presented in the figure.

Data table for figure 11.4. Age- sex-adjusted percent of persons of all ages who assessed their health as excellent or very good, by race/ethnicity: United States, January-September 2004

| Race/ethnicity | Percent ¹ | 95% confidence interval |
|------------------------|----------------------|-------------------------|
| Hispanic or Latino | 58.8 | 57.1-60.6 |
| Not Hispanic or Latino | | |
| White, single race | 69.9 | 69.2-70.5 |
| Black, single race | 58.2 | 56.6-59.8 |

¹Estimates are age- sex-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18-64 years, and 65 years and over.

²Estimates are age-adjusted to the 2000 projected U.S. standard population using three age groups: under 18 years, 18-64 years, and 65 years and over.