


Figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997–March 2006




NOTES: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home. The analyses excluded persons with unknown information on personal care needs (about 0.01% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Release" and Table III in the Appendix for more details.

DATA SOURCE: Family Core component of the 1997–2006 NHIS. The estimate for 2006 was based on data collected from January through March. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ In early 2006, 6.7% (95% confidence interval = 5.48–7.83%) of adults aged 65 years and over needed help with personal care from other persons. This estimate was not significantly different from the 2005 estimate of 6.3%.

■ The annual percentage of adults who needed help with personal care from other persons was 6.6% in 1997, 6.3% in 1998, 6.4% in 1999, 6.4% in 2000, 6.5% in 2001, 6.2% in 2002, 6.5% in 2003, 6.3% in 2004, and 6.3% in 2005.

Figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January–March 2006




NOTE: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, adults aged 85 years and over (19.9%) were more than five times as likely as adults aged 65–74 years (3.6%) to need help with personal care from other persons.
- For all three age groups, there was no significant difference between women and men in the percentage of adults who needed help with personal care.

Figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January–March 2006



NOTES: Personal care needs, or activities of daily living (ADL), include eating, bathing, dressing, or getting around inside the person's home. Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: Based on data collected from January through March in the Family Core component of the 2006 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ The age-sex-adjusted percentage of persons who needed help with personal care from other persons was 9.9% for Hispanic persons, 6.0% for non-Hispanic white persons, and 12.3% for non-Hispanic black persons.

■ The percentage of those who needed help with personal care from other persons was higher for non-Hispanic black persons than for non-Hispanic white persons.

Data tables for figures 12.1–12.3:

Data table for figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997–March 2006

Year	Percent	95% confidence interval
1997	6.6	6.1-7.1
1998	6.3	5.9-6.8
1999	6.4	5.8-6.9
2000	6.4	5.9-6.9
2001	6.5	6.0-7.0
2002	6.2	5.7-6.7
2003	6.5	6.0-7.1
2004	6.3	5.8-6.8
2005	6.3	5.83-6.76
January–March 2006	6.7	5.48-7.83

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–March 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January–March 2006

Age and sex	Percent	95% confidence interval
65–74 years		
Total	3.6	2.44-4.68
Male	2.8	1.39-4.18
Female	4.2	2.51-5.90
75–84 years		
Total	6.6	4.71-8.51
Male	7.4	3.99-10.90
Female	6.0	3.67-8.36
85 years and over		
Total	19.9	14.59-25.22
Male	15.8	7.52-24.02
Female	22.1	15.67-28.45
65 years and over: crude¹		
Total	6.7	5.48-7.83
Male	5.7	4.04-7.30
Female	7.4	5.86-8.90
65 years and over: age-adjusted²		
Total	6.6	5.47-7.82
Male	6.0	4.32-7.74
Female	7.0	5.58-8.49

¹Crude estimates are presented in the figure.

²Estimates are age adjusted using the projected 2000 U.S. population as the standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January–March 2006

Race/ethnicity	Percent ¹	95% confidence interval
Hispanic or Latino	9.9	5.02-14.74
Not Hispanic or Latino:		
White, single race	6.0	4.68-7.30
Black, single race	12.3	7.86-16.72

¹Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population using three age groups: 65–74 years, 75–84 years, and 85 years and over.

DATA SOURCE: National Health Interview Survey, 2006. Data are based on household interviews of a sample of the civilian noninstitutionalized population.