


**Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–June 2013**




DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2013, Sample Adult Core component.

- For January–June 2013, the percentage of adults aged 65 and over who had ever received a pneumococcal vaccination was 60.7% (95% confidence interval = 58.55%–62.89%), which was higher than, but not significantly different from the 2012 estimate of 59.9%.
- The percentage of adults aged 65 and over who had ever received a pneumococcal vaccination increased rapidly from 42.4% in 1997 to 53.1% in 2000, then increased more slowly to 60.7% in January–June 2013.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • The analyses excluded those with unknown pneumococcal vaccination status (about 5% of respondents each year). • See [Technical Notes](#) for more details.

**Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2013**




DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2013, Sample Adult Core component.

- For both sexes combined, the percentage of adults who had ever received a pneumococcal vaccination was 56.7% for persons aged 65–74 and 66.2% for persons aged 75 and over.
- For both sexes combined, adults aged 75 and over were more likely than those adults aged 65–74 to have ever received a pneumococcal vaccination. This pattern was also seen in men.
- For adults aged 65–74, women were more likely than men to have ever received a pneumococcal vaccination.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • The analyses excluded the 3.9% of adults aged 65 and over with unknown pneumococcal vaccination status. • See [Technical Notes](#) for more details.

**Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2013**



DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2013, Sample Adult Core component.

- The percentage of adults aged 65 and over who had ever received a pneumococcal vaccination was 40.1% for Hispanic persons, 64.6% for non-Hispanic white persons, and 50.4% for non-Hispanic black persons.
- Non-Hispanic white adults were more likely than non-Hispanic black adults and Hispanic adults to have ever received a pneumococcal vaccination.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • The analyses excluded the 3.9% of adults aged 65 and over with unknown pneumococcal vaccination status. • See [Technical Notes](#) for more details.

## Data tables for Figures 5.1–5.3:

### Data table for Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–June 2013

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	42.4 (40.9-43.9)	42.6 (41.1-44.1)
1998	46.0 (44.5-47.5)	46.3 (44.8-47.8)
1999	49.7 (48.1-51.3)	49.9 (48.3-51.5)
2000	53.1 (51.5-54.7)	53.4 (51.8-55.0)
2001	54.0 (52.4-55.6)	54.2 (52.6-55.7)
2002	56.0 (54.5-57.5)	56.2 (54.8-57.7)
2003	55.6 (54.1-57.1)	55.7 (54.2-57.1)
2004	56.8 (55.3-58.3)	57.0 (55.5-58.5)
2005	56.2 (54.82-57.64)	56.3 (54.95-57.72)
2006	57.1 (55.25-58.88)	57.2 (55.43-59.04)
2007	57.7 (55.68-59.69)	57.8 (55.78-59.80)
2008	59.6 (57.68-61.48)	59.8 (57.98-61.64)
2009	60.6 (58.74-62.51)	61.0 (59.15-62.79)
2010	59.4 (57.61-61.17)	59.8 (58.02-61.51)
2011	62.3 (60.55-63.95)	62.7 (61.04-64.31)
2012	59.9 (58.29-61.49)	60.4 (58.86-62.01)
January–June 2013	60.7 (58.55-62.89)	61.3 (59.10-63.40)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with the 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2013, Sample Adult Core component.

**Data table for Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2013**

Age (years) and sex	Percent	95% confidence interval
65-74, total	56.7	53.88-59.57
65-74, male	53.2	48.96-57.50
65-74, female	59.8	56.42-63.08
75 and over, total	66.2	63.21-69.20
75 and over, male	68.5	63.88-73.18
75 and over, female	64.6	60.72-68.46
65 and over (crude <sup>1</sup> ), total	60.7	58.55-62.89
65 and over (crude <sup>1</sup> ), male	59.2	55.98-62.45
65 and over (crude <sup>1</sup> ), female	61.9	59.32-64.49
65 and over (age-adjusted <sup>2</sup> ), total	61.3	59.10-63.40
65 and over (age-adjusted <sup>2</sup> ), male	60.5	57.37-63.70
65 and over (age-adjusted <sup>2</sup> ), female	62.1	59.46-64.66

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2013, Sample Adult Core component.

**Data table for Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2013**

Race/ethnicity	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
Hispanic or Latino	40.1 (33.72-46.54)	40.8 (34.06-47.49)
Not Hispanic or Latino, single race, white	64.6 (62.16-66.94)	65.0 (62.63-67.37)
Not Hispanic or Latino, single race, black	50.4 (46.07-54.74)	49.4 (45.02-53.84)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2013, Sample Adult Core component.