


General health status

Figure 11.1. Percentage of persons of all ages who had excellent or very good health: United States, 1997– June 2017




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses exclude persons with unknown health status (about 0.1% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997– June 2017, Family Core component.

- For January–June 2017, the percentage of persons who had excellent or very good health was 66.2% (95% confidence interval = 65.47%–66.93%), which was not significantly different from the 2016 estimate of 66.4%.
- The percentage of persons who had excellent or very good health decreased, from 68.5% in 1997 to 65.9% in 2006. Since 2006, the percentage of persons who had excellent or very good health has remained stable.

Figure 11.2. Percent distribution of respondent–assessed health status for all ages, by sex: United States, January–June 2017




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses exclude the 0.1% of persons with unknown health status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.

- For both sexes combined, most persons’ health was either excellent (34.9%) or very good (31.3%). A smaller percentage of persons had good (23.7%), fair (8.0%), or poor (2.2%) health. This pattern held for males and females.
- Men were more likely to have excellent health compared with women. Women were more likely than men to have fair or poor health.

Figure 11.3. Percentage of persons of all ages who had excellent or very good health, by age group and sex: United States, January–June 2017




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses exclude the 0.1% of persons with unknown health status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.

- For both sexes combined, the percentage of persons with excellent or very good health decreased with age, 84.8% for those under age 18 years to 64.4% for those aged 18–64, and 45.3% for those aged 65 and over. This pattern was observed in both men and women.
- For persons of all ages and adults 18–64, males were more likely than females to have excellent or very good health.

Figure 11.4. Age-sex-adjusted percentage of persons of all ages who had excellent or very good health, by race and ethnicity: United States, January–June 2017



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Health status data were obtained by asking respondents to assess their own health and that of family members living in the same household as excellent, very good, good, fair, or poor. The analyses exclude the 0.1% of persons with unknown health status. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and three age groups: under 18 years, 18–64, and 65 and over. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.

- After adjustment for age and sex, the percentage of persons by race and ethnicity who had excellent or very good health was 61.3% for Hispanic persons, 70.8% for non-Hispanic white persons, and 59.3% for non-Hispanic black persons.
- Of the three race and ethnicity groups, non-Hispanic white persons were more likely to have excellent or very good health compared with non-Hispanic black persons and Hispanic persons.

Data tables for Figures 11.1–11.4:

Data table for Figure 11.1. Percentage of persons of all ages who had excellent or very good health: United States, 1997– June 2017

| Year | Percent | 95% confidence interval |
|-------------------|---------|-------------------------|
| 1997 | 68.5 | 68.0–69.1 |
| 1998 | 69.1 | 68.6–69.6 |
| 1999 | 68.7 | 68.1–69.3 |
| 2000 | 68.3 | 67.7–68.8 |
| 2001 | 68.4 | 67.9–69.0 |
| 2002 | 67.4 | 66.8–68.0 |
| 2003 | 67.5 | 66.9–68.0 |
| 2004 | 66.5 | 66.0–67.1 |
| 2005 | 66.5 | 65.98–67.11 |
| 2006 | 65.9 | 65.25–66.61 |
| 2007 | 66.0 | 65.18–66.88 |
| 2008 | 66.0 | 65.30–66.79 |
| 2009 | 66.3 | 65.46–67.10 |
| 2010 | 65.7 | 64.95–66.35 |
| 2011 | 65.6 | 64.98–66.24 |
| 2012 | 65.8 | 65.19–66.41 |
| 2013 | 66.3 | 65.70–66.93 |
| 2014 | 66.5 | 65.85–67.08 |
| 2015 | 66.1 | 65.37–66.73 |
| 2016 | 66.4 | 65.76–67.09 |
| January–June 2017 | 66.2 | 65.47–66.93 |

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997– June 2017, Family Core component.

Data table for Figure 11.2. Percent distribution of respondent–assessed health status, by sex for all ages: United States, January–June 2017

| Health status and sex | Percent | 95% confidence interval |
|-----------------------|---------|-------------------------|
| Excellent, total | 34.9 | 34.29–35.49 |
| Excellent, male | 36.3 | 35.46–37.11 |
| Excellent, female | 33.6 | 32.79–34.32 |
| Very good, total | 31.3 | 30.66–31.97 |
| Very good, male | 30.9 | 30.20–31.65 |
| Very good, female | 31.7 | 30.79–32.60 |
| Good, total | 23.7 | 23.01–24.34 |
| Good, male | 23.3 | 22.55–24.10 |
| Good, female | 24.0 | 23.25–24.77 |
| Fair, total | 8.0 | 7.62–8.32 |
| Fair, male | 7.5 | 7.06–8.03 |
| Fair, female | 8.4 | 7.96–8.83 |
| Poor, total | 2.2 | 2.00–2.33 |
| Poor, male | 1.9 | 1.75–2.15 |
| Poor, female | 2.4 | 2.16–2.59 |

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.

Data table for Figure 11.3. Percentage of persons of all ages who had excellent or very good health, by age group and sex: United States, January–June 2017

| Age (years) and sex | Percent | 95% confidence interval |
|---|---------|-------------------------|
| Under 18, total | 84.8 | 83.50–86.10 |
| Under 18, male | 85.0 | 83.46–86.48 |
| Under 18, female | 84.6 | 82.83–86.34 |
| 18–64, total | 64.4 | 63.68–65.18 |
| 18–64, male | 65.2 | 64.29–66.14 |
| 18–64, female | 63.7 | 62.78–64.55 |
| 65 and over, total | 45.3 | 43.92–46.78 |
| 65 and over, male | 45.4 | 43.15–47.74 |
| 65 and over, female | 45.3 | 43.66–46.90 |
| All ages (crude ¹), total | 66.2 | 65.47–66.93 |
| All ages (crude ¹), male | 67.2 | 66.39–68.01 |
| All ages (crude ¹), female | 65.2 | 64.35–66.13 |
| All ages (age-adjusted ²), total | 67.3 | 66.59–67.96 |
| All ages (age-adjusted ²), male | 67.8 | 67.01–68.63 |
| All ages (age-adjusted ²), female | 66.8 | 65.89–67.60 |

¹Crude estimates are presented.

²Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and three age groups: under 18 years, 18–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.

Data table for Figure 11.4. Age-sex-adjusted percentage of persons of all ages who had excellent or very good health, by race and ethnicity: United States, January–June 2017

| Race and ethnicity | Percent ¹ | 95% confidence interval |
|--|----------------------|-------------------------|
| Hispanic or Latino | 61.3 | 58.82–63.66 |
| Not Hispanic or Latino, single race, white | 70.8 | 69.96–71.72 |
| Not Hispanic or Latino, single race, black | 59.3 | 56.52–61.97 |

¹Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and three age groups: under 18 years, 18–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Family Core component.