

**Table 45. Tuberculosis Cases and Percentages, by Noninjecting Drug Use,<sup>1</sup> Ages ≥15 Years: Reporting Areas, 2016**

Reporting area	Total cases	Cases with information on noninjecting drug use		Cases reporting noninjecting drug use	
		No.	(%)	No.	(%)
<b>United States</b>	<b>8,878</b>	<b>8,759</b>	<b>(98.7)</b>	<b>599</b>	<b>(6.8)</b>
Alabama	108	108	(100.0)	15	(13.9)
Alaska	50	49	(98.0)	15	(30.6)
Arizona	185	182	(98.4)	30	(16.5)
Arkansas	86	86	(100.0)	5	(5.8)
California	2,011	1,966	(97.8)	119	(6.1)
Colorado	61	60	(98.4)	3	(5.0)
Connecticut	51	51	(100.0)	2	(3.9)
Delaware	15	15	(100.0)	1	(6.7)
District of Columbia	25	25	(100.0)	1	(4.0)
Florida	616	612	(99.4)	43	(7.0)
Georgia	283	279	(98.6)	31	(11.1)
Hawaii	115	113	(98.3)	1	(0.9)
Idaho	14	12	(85.7)	0	(0)
Illinois	326	323	(99.1)	21	(6.5)
Indiana	102	102	(100.0)	7	(6.9)
Iowa	48	45	(93.8)	1	(2.2)
Kansas	39	39	(100.0)	3	(7.7)
Kentucky	90	90	(100.0)	5	(5.6)
Louisiana	122	117	(95.9)	10	(8.5)
Maine	23	23	(100.0)	1	(4.3)
Maryland	214	210	(98.1)	2	(1.0)
Massachusetts	182	181	(99.5)	4	(2.2)
Michigan	125	124	(99.2)	7	(5.6)
Minnesota	149	149	(100.0)	1	(0.7)
Mississippi	57	56	(98.2)	6	(10.7)
Missouri	99	97	(98.0)	5	(5.2)
Montana	4	4	(100.0)	1	(25.0)
Nebraska	25	23	(92.0)	2	(8.7)
Nevada	53	53	(100.0)	5	(9.4)
New Hampshire	14	14	(100.0)	1	(7.1)
New Jersey	282	279	(98.9)	3	(1.1)
New Mexico	37	36	(97.3)	3	(8.3)
New York State <sup>2</sup>	193	190	(98.4)	8	(4.2)
New York City	554	546	(98.6)	33	(6.0)
North Carolina	206	206	(100.0)	21	(10.2)
North Dakota	21	21	(100.0)	2	(9.5)
Ohio	135	133	(98.5)	9	(6.8)
Oklahoma	72	71	(98.6)	9	(12.7)
Oregon	65	63	(96.9)	6	(9.5)
Pennsylvania	164	164	(100.0)	3	(1.8)
Rhode Island	11	11	(100.0)	0	(0)
South Carolina	87	83	(95.4)	7	(8.4)
South Dakota	12	12	(100.0)	1	(8.3)
Tennessee	98	98	(100.0)	12	(12.2)
Texas	1,181	1,181	(100.0)	108	(9.1)
Utah	18	18	(100.0)	0	(0)
Vermont	5	5	(100.0)	0	(0)
Virginia	196	196	(100.0)	8	(4.1)
Washington	199	190	(95.5)	9	(4.7)
West Virginia	14	13	(92.9)	1	(7.7)
Wisconsin	35	34	(97.1)	8	(23.5)
Wyoming	1	1	(100.0)	0	(0)
American Samoa <sup>3</sup>	1	1	(100.0)	0	(0)
Fed. States of Micronesia <sup>3</sup>	86	86	(100.0)	0	(0)
Guam <sup>3</sup>	63	59	(93.7)	0	(0)
Marshall Islands <sup>3</sup>	131	131	(100.0)	4	(3.1)
N. Mariana Islands <sup>3</sup>	18	18	(100.0)	0	(0)
Puerto Rico <sup>3</sup>	68	68	(100.0)	8	(11.8)
Republic of Palau <sup>3</sup>	20	20	(100.0)	0	(0)
U.S. Virgin Islands <sup>3</sup>	0	0	...	0	...

<sup>1</sup>Noninjecting drug use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia).

<sup>2</sup>Excludes New York City.

<sup>3</sup>Not included in U.S. totals.

**Note:** Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.